Data-driven analysis on the simulations of the spread of COVID-19 under different interventions of China

Ting Tian[1#], Jingwen Zhang[1#], Shiyun Lin[1#], Yukang Jiang[1#], Jianbin Tan[1#], Zhongfei Li[2], Xueqin Wang[3,1*]

[1]School of Mathematics, Sun Yat-sen University, Guangzhou, 510275, China

[2]School of Management, Sun Yat-sen University, Guangzhou, 510275, China

[3]School of Management, University of Science and Technology of China, Hefei, 230026, China

* Correspondence author: wangxq20@ustc.edu.cn

Dr. T. Tian, J. Zhang, S. Lin, Y. Jiang, and J. Tan contributed equally to this article.

Abstract

Objectives

Since February 2020, COVID-19 has spread rapidly to more than 200 countries in the world. During the pandemic, local governments in China implemented different interventions to efficiently control the spread of the epidemic. Characterizing transmission of COVID-19 under some typical interventions is to help countries develop appropriate interventions.

Methods

We established compartmental model that allowed the number of infected and infectious to be unknown and the effective reproduction number to change over time, thus the effects of policies could be reasonably reflected and estimated. By using the epidemic data of three representative cities of China (Wuhan, Wenzhou and Shenzhen), we migrated the estimated policy modes to other countries.

Results

The smallest expected cumulative confirmed cases under different interventions would be 5936 with 95% CI (5012,6966) for South Korea, 146012 with 95% CI (140504, 154264) for Italy, and 400642 with 95% CI (390331,409431) for the United States until May 31, 2020, respectively.

Conclusions

Based on the simulation of epidemic in South Korea, Italy and the United States, it is

reasonable that South Korea and Italy continue to maintain their current policies,

while the implementation of interventions of Wenzhou may significantly decrease the

magnitude of the outbreak of COVID-19 for the United States.

Keywords: Compartmental model; Time-varying reproduction number; Infected

individuals without isolation; Mode migration; Policy intervention

Introduction

Since December 2019, the first report of COVID-19 which was detected in Wuhan

was sent to the World Health Organization (WHO) by Chinese government (World

Health Organization, 2020a). The epidemic has rapidly spread to all provinces in

China and more than 200 countries around the world. WHO declared that COVID-19

became a pandemic on March 11, 2020 (World Health Organization, 2020b). China

issued a series of intervention policies to reduce the transmission of COVID-19, such

as suspending public transportations in areas of severe outbreak (e.g., Wuhan),

cancelling public gathering and delaying reopening of enterprises. These endeavors

have led to the significant suppression of the epidemic of COVID-19. There was no

domestically newly reported confirmed case for the first time in China on March 18,

2020 (Xinhuannet, 2020). The local and national responses of China have made

remarkable contributions to the prevention and control of COVID-19 at the expense

of the economic loss. Taking individual consumption industry as an example, the

amount of total retail sales of consumer goods decreased by about 20.5% with a sharp

drop of catering services by 43.1% from January to February 2020 compared to the

same duration of 2019 according to the National Bureau of Statistics in China (2020).

Consequently, policymakers need to anticipate the likely outcome of interventions in

terms of their own epidemic situations.

Predicting the scale of the epidemic is essential to inform policymakers the likely

outcomes of potential intervention options (Lipsitch et al., 2011), i.e. the expected

scale of cumulative confirmed cases for COVID-19 could provide insights into the

likely outcomes of different intervention policies. The estimation of the time-varying

reproduction numbers could indicate the effects of intervention policies (Cauchemez

et al., 2006, Wallinga and Teunis, 2004). Yang et al. (2020) showed that if the

implementation of intervention policies delayed 5 days, the expected scale of

cumulative confirmed cases would be over 3 times larger than actual ones in China. In

this study, researchers used epidemic data from Hubei province, Zhejiang province,

and Guangdong province. We selected the most severe cities in these three provinces

as our important examples, i.e. Wuhan, Wenzhou, and Shenzhen. In these cases, their

time-varying reproduction numbers were used to unfold the effects of interventions,

and thus to predict the expected scale of COVID-19 by using the real-time epidemic

data in various countries. In doing so, the possible courses of COVID-19 under these

interventions in South Korea, Italy, and the United States can be made.

Wuhan was the first city of the COVID-19 outbreak with the most severe

epidemic in China. It suspended public transportation, cancelled all outbound trains

and flights (The prevention and control headquarters of Wuhan, 2020) and started to

construct a specialist emergency hospital (i.e. Huoshenshan Hospital) for patients on

January 23. All residential areas were closed on February 11, 2020. Wenzhou was the

city with the largest number of cumulative confirmed cases in Zhejiang province. In

Wenzhou, the inter-provincial and municipal bus services were suspended on January

27(Wenzhou Municipal People's Government, 2020a), 14 expressway entrances were

temporarily closed from 10 p.m. onwards on January 29 (Wenzhou Municipal

People's Government, 2020b), and on February 1, it announced that every household

might send one person every two days outside for necessity purchases (Wenzhou

Municipal People's Government, 2020c). Shenzhen was the city of the first confirmed

case in Guangdong province, and its prevention and control strategy was adopted as a

classical case in the report of WHO (2020). On January 24, all the entertainment

places were closed. It announced that all the individuals returning from Hubei needed

to be isolated at home or other suitable facilities for 14 days on February 2 (Shenzhen

Municipal People's Government, 2020a), and the residential units with confirmed

cases were forced to implement "hard quarantine" for 14 days and checkpoints were

set up in all entrances of residential areas on February 7 (Shenzhen Municipal

People's Government, 2020b). All details about the interventions implemented in

these cities were given in Figure 1.

In this paper, we used the time-varying reproduction number to reflect the effects

of interventions implemented in Wuhan, Wenzhou, and Shenzhen, China. Our goal

was to predict the likely outcomes of these interventions in South Korea, Italy and the

United States so that the lessons can be learned from the key decisions made in the

representative cities.

Methods

Data sources

The epidemic data were extracted for Wuhan, Wenzhou, and Shenzhen from the

Chinese Center for Disease Control and Prevention (China CDC) (National Health

Commission of the People's Republic of China, 2020), and the data for South Korea,

Italy and the United States were downloaded through April 23, 2020 from COVID-19

Knowledge & Data Hub (Sciences, 2020).

Model assumptions and solutions

We introduced Susceptible-Infected and infectious without isolation-Hospitalized in

isolation-Removed (SIHR) model with four compartments: Susceptible (S), infected

and infectious without isolation (I), hospitalized in isolation (H), removed (R)(Hsieh

et al., 2004, Li, 2018). Based on the definitions, we made the following assumptions:

(a) The susceptible (S) individuals have no immunity to the disease;

(b) The infected and infectious without isolation (I) were assumed to be

infectious since they caught the virus, i.e., ignoring the latent period, and

would transform into the hospitalized in isolation eventually after symptoms

onset;

(c) The hospitalized (H) individuals were in isolation and treated carefully and

would not transmit COVID-19 to the susceptible;

(d) The removed (R) individuals would not be infected again.

It is crucial to evaluate the number of individuals who are infected and infectious

without isolation for the control of epidemic, which remain unknown and are needed

to be estimated in SIHR model. Based on the above assumptions, the transfer

relationships between S, I, H, R were defined (see Figure 1 in the supplements) and

the corresponding dynamic system of these four compartments along with parameters

α, β, γ were given in the supplements.

The basic reproduction number R_0, which is the expected number of secondary

cases produced by a single (typical) infection in a completely susceptible population

(Jones, 2007), is defined by SIHR model:

143
$$R_0 = \alpha/\beta,$$

where $1/\beta$ is the mean of the incubation period of COVID-19, and is assumed to be

5 days in terms of studies about COVID-19 (Backer et al., 2020, Guan et al., 2020, Li

et al., 2020) by ignoring the latent period. To obtain the time-varying reproduction

number R_t, we assumed that the parameter α was time variant and would decrease

by the influence of interventions. We used logistic function (Tan et al., 2020) to

simulate the decreasing trend of the α:

150
$$\alpha_{\alpha_0,d,m}(t) = \frac{\alpha_0}{1+\exp\left(\lambda_m\left(t-d-\frac{m}{2}\right)\right)},$$

where α_0 denotes the maximum contact numbers per person of COVID-19 during

the early outbreak, d is the time when the control measures start to be effective and

α starts to decline, m represents the duration of a process where the epidemic is to

nearly vanish, λ_m is chosen as $\frac{2\log\left((1-\varepsilon)/\varepsilon\right)}{m}$ and ε is fixed to be 0.01. The above

parameters α_0, d, m are estimated from real epidemic data.

The smaller the values of d and m, the earlier effectiveness and the stronger

intensity of interventions were implemented, respectively. The graph of time-varying

α is given in Figure 2 in supplements.

By the setting above, R_t was calculated as:

160
$$R_t = \frac{\alpha_{\alpha_0,d,m}(t)}{\beta}.$$

We introduce randomness to SIHR model and applied the Bayesian method

combined with Markov Chain Monte Carlo (MCMC) for the estimation of parameters

and missing data (Bishop, 2006, Soubeyrand, 2016). Once the prior distribution of the

parameters was given, we used Block Metropolis–Hastings algorithm (Andrieu et al.,

2000, Andrieu et al., 2003) to simulate the posterior distribution of the parameters and

the number of individuals in each class. The point estimates were presented as the

median of the posterior distribution while 95% credible intervals were constructed

with 2.5% and 97.5% quantiles.

Results

The estimation of policy pattern

To describe the effects of interventions in different cities, we used the data from the

starting dates of the COVID-19 outbreak to the dates of well-controlled in each

representative cities individually, indicating that the time range for each

representative city would be different (Green et al., 2002).

Among them, the starting date of the outbreak was defined as the day when the

number of hospitalized individuals (H) started to increase and continued to increase

for the next three consecutive days, while the date of well-controlled was at which the

number of hospitalized individuals (H) started to decrease and continued to decrease

for the next three consecutive days(Green et al., 2002). Based on these definitions, we

used the data from January 21 to February 9 in Wenzhou and from January 19 to

February 6 in Shenzhen. In Wuhan, the starting date was January 15, but the clinical

standard was changed on February 12, we used February 11 as the data cutoff for

Wuhan. The trends of R_t for representative cities were shown in Figure 2.

The initial value of R_t for Wuhan was greater than that for Shenzhen, followed

by Wenzhou. The effectiveness of intervention policies for Shenzhen started earlier

than that for Wuhan, followed by Wenzhou. The intensity of policies for Wenzhou

was stronger than that for Wuhan, followed by Shenzhen. Comparing with details

about the interventions implemented in these cities in Figure 1, the effects of these

policies could be reasonably reflected and estimated.

The mode migration

We applied the SIHR model whose R_t is time-invariant in South Korea, Italy and the

United States. We used the real-time epidemic data from February 26 to February 28

in South Korea, from March 28 to March 30 in Italy and from March 29 to March 31

in the United States and estimated the posterior distributions country by country. And

then we migrate the intensity of policy intervention of Wuhan, Wenzhou, and

Shenzhen to South Korea, Italy, and the United States, and predict the posterior

distributions of numbers of infected and infectious without isolation and cumulative

confirmed cases. The simulation of COVID-19 trend for constant R_t and

time-varying R_t of three policies of China in South Korea, Italy, and the United

States are given as follows (Figures 3-5).

From our prediction, if interventions were not implemented since 23 February,

the expected cumulative confirmed cases would be 7705971with corresponding 95%

credible intervals (CI) (2419211,17969284) on March 7 when the expected

cumulative confirmed cases were over 10% population for the first time in South

Korea (Figure 3A). Similarly, the expected cumulative confirmed cases would be

6680363 with 95% CI (442494,35236491) on March 29 when the expected

cumulative confirmed cases were over 10% population for the first time in Italy

(Figure 4A). And the expected cumulative confirmed cases will be 34179685 with 95%

CI (5274919,136469414) on April 17 when the expected cumulative confirmed cases

will over 10% population for the first time in the United States, if interventions were

not implemented since March 13 (Figure 5A). Overall, if interventions were not

implemented, it can be expected that how tremendously large populations would be

infected in these countries. Thus, the interventions are critical to control the fast

spread of the disease. In SIHR model, the estimated number of infected and infectious

without isolation (I) could be used to estimate the stage of the epidemic, while the

expected cumulative confirmed cases under different policy patterns reflected the

magnitude of the outbreak of infectious diseases.

Based upon our prediction, the actual cumulative confirmed cases of COVID-19

in South Korea from February 26 to March 9 were within the ranges of expected

cumulative confirmed cases for Shenzhen pattern, whose intensity was relatively mild

compared with both Wuhan and Wenzhou patterns. According to the report (Normile,

2020), South Korea has implemented similar interventions to Shenzhen pattern, where

it implemented policies closely following and isolating the close contact individuals

of confirmed cases to reduce the transmission of COVID-19. Also, as its estimated

number of I was 2639 with corresponding 95% CI (2306,3055) on February 26,

there is not much difference for the expected cumulative confirmed cases in the

simulation of three urban interventions in South Korea (Figure 3B). This indicated

that mild policy like Shenzhen pattern implemented at an early stage of epidemic

could effectively curb the outbreak of infectious diseases. It is reasonable that South

Korea would maintain its current intervention.

For Italy, there is a substantial difference in the simulation of three policies

implemented, the expected cumulative confirmed cases will be 179571 with 95% CI

(165868, 205749) under Wuhan pattern, 146012 with 95% CI (140504, 154264)

under Wenzhou pattern and 214881 with 95% CI (187566, 265198) under Shenzhen

pattern on May 31, respectively (Figure 4B). The likely outcomes of different

interventions were distinct since the estimated number of I was large (25768) on

March 28, which implies that it is cautious to carry out a policy at this stage of

epidemic. Note that the expected number of cumulative confirmed cases under the

interventions of Wenzhou for Italy would be the smallest. There were the most

intense efforts in the interventions of Wenzhou for both suspending inner and outer

public transportations (middle-distance prevention) and restricting residents out

(short-distance prevention). It was required that such a large-scale response mounted

to block the broad extent of transmission not only inside of city but also inside of

communities.

Comparing the actual situations of Italy, the cumulative confirmed cases were

187,327 on April 23, 2020, which were 28.7% more than the simulation of

COVID-19 under interventions of Wenzhou. If taking such highly stringent

interventions, the epidemic situation could be mitigated where the expected number

of cumulative confirmed cases would be 146,012 on May 31, decreasing by 19% and

32% compared to the implementation of policy of Wuhan and Shenzhen, respectively.

In the United States, there is a considerable difference from the policy of

Shenzhen compared with other two patterns. The expected cumulative confirmed

cases will be 694055 with 95% CI (665937,725910) under Wuhan pattern, 400642

with 95% CI (390331,409431) under Wenzhou pattern and 1096227 with 95% CI

(1037815,1172518) under Shenzhen pattern on May 31, respectively. One reason for

this distinction is that the number of I was 83232 (Figure 5B) on March 29, which

implied certain risks for long-term outbreak of epidemic. The cumulative confirmed

cases were 849094 on April 23, 2020, and were 113.3% more than the simulation of

COVID-19 under interventions of Wenzhou, which implied that the implementations

of interventions of Wenzhou may significantly decrease the magnitude of the

outbreak of COVID-19 for the United States. If taking the relatively mild

interventions of Shenzhen, the situation would get worse and the expected cumulative

confirmed cases would be over 1000000 in the United States on May 31.

Discussion

Our simulation results indicated that the mild interventions were difficult to control

the magnitude of outbreak of infectious diseases when the number of infected and

infectious without isolation (I) was quite large, but it was effective to implement at

the early stage of epidemic. Once the epidemic became severe, the highly stringent

interventions were needed to be implemented for the control of epidemic. Although it

still takes time to make a reasonable assessment of the patterns and effects of these

policies, the transmission patterns of COVID-19 under some typical intervention

measures could be learned from them, and the transmission trend of COVID-19 could

be predicted.

In general, there are no one-size-fits-all interventions applicable for all regions

(Wu et al., 2020). Taking any potential interventions depends on the specific setting

of the region. We did not quantitatively measure the details of interventions and did

not consider public health and economic costs of interventions in our analysis.

However, the likely outcomes of interventions were predicted to inform the

policymakers to consider the appropriate sets of proactive interventions.

Author contributions:

All authors attest that they meet the ICMJE criteria for authorship. TT and XW had

roles in the study design, data collection, data analysis, data interpretation, literature

search, and developed the idea and research. TT, JZ, SL, YJ and JT wrote the first

draft of the manuscript and all other authors discussed results and edited the

manuscript. JZ, SL, YJ and JT had roles in the data collection, data analysis, and ZL

had roles in the data interpretation. TT, JZ, SL, YJ, and JT contributed equally to this

article. All authors reviewed and approved the final version of the manuscript.

Conflict of interest

All authors declare no competing interests.

Acknowledgements

We would like to thank all individuals who are collecting epidemiological data of the

COVID-19 outbreak around the world.

References

1. Andrieu C, de Freitas N, Doucet A. Sequential Bayesian estimation and model

   selection for dynamic kernel machines. 2000.

2. Andrieu C, De Freitas N, Doucet A, Jordan MI. An introduction to MCMC for

   machine learning. Machine learning 2003;50(1-2):5-43.

3. Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel

   coronavirus (2019-nCoV) infections among travellers from Wuhan, China, 20-28

   January 2020. Euro Surveill 2020;25(5).

4. Bishop CM. Pattern recognition and machine learning: springer, 2006.

5. Cauchemez S, Boelle PY, Thomas G, Valleron AJ. Estimating in real time the

efficacy of measures to control emerging communicable diseases. Am J Epidemiol

2006;164(6):591-7.

6. Green MS, Swartz T, Mayshar E, Lev B, Leventhal A, Slater PE, et al. When is an

epidemic an epidemic? The Israel Medical Association journal: IMAJ

2002;4(1):3-6.

7. Guan W-j, Ni Z-y, Hu Y, Liang W-h, Ou C-q, He J-x, et al. Clinical Characteristics

of Coronavirus Disease 2019 in China. New England Journal of Medicine 2020.

8. Hsieh Y-H, Lee J-Y, Chang H-L. SARS epidemiology modeling. Emerging

infectious diseases 2004;10(6):1165.

9. Jones JH. Notes on R0. Califonia: Department of Anthropological Sciences 2007.

10. Li MY. An introduction to mathematical modeling of infectious diseases: Springer,

2018.

11. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission

Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia. New

England Journal of Medicine 2020.

12. Lipsitch M, Finelli L, Heffernan RT, Leung GM, Redd SC, Group HnS.

Improving the evidence base for decision making during a pandemic: the example

of 2009 influenza A/H1N1. Biosecur Bioterror 2011;9(2):89-115.

13. National Bureau of Statistics in China. National Economy Withstood the Impact

of COVID-19 in the First Two Months; 2020. Available from:

http://www.stats.gov.cn/english/PressRelease/202003/t20200316_1732244.html.

14. National Health Commission of the People's Republic of China. Distribution of

COVID-19 cases in China; 2020. Available from:

http://2019ncov.chinacdc.cn/2019-nCoV/index.html. [Accessed April 26, 2020].

15. Normile D. Coronavirus cases have dropped sharply in South Korea. What's the

secret to its success. Science 2020.

16. World Health Organization. Report of the WHO-China Joint Mission on

Coronavirus Disease 2019 (COVID-19). 2020.

17. Sciences CAo. COVID-19 Knowledge & Data Hub; 2020. Available from:

http://geodoi.ac.cn/covid-19/en/index.aspx.

18. Shenzhen Municipal People's Government. Circular of the prevention and control

headquarters of pneumonia epidemic caused by new coronavirus infection in

Shenzhen; 2020a. Available from:

http://www.sz.gov.cn/szzt2010/yqfk2020/szzxd/zczy/zcwj/fkzc/content/post_6725

195.html.

19. Shenzhen Municipal People's Government. Shenzhen issued three epidemic notice!

To the citizens living in shenzhen, to the personnel in shenzhen, the enterprises in

shenzhen; 2020b. Available from:

http://www.sz.gov.cn/cn/xxgk/zfxxgj/zwdt/content/post_6697629.html.

20. Soubeyrand S. Construction of semi-Markov genetic-space-time SEIR models and

inference. 2016.

21. Tan J, Jinag Y, Tian T, Wang X. P-SIHR probabilistic graphical model: An

applicable model of COVID-19 in estimating the number of infectious individuals

without isolation and time-varying reproduction number. Acta Mathematicae

Applicatae Sinica 2020;42(3).

22. The prevention and control headquarters of Wuhan. Circular of the prevention and

control headquarters of pneumonia epidemic caused by new cornoavirus infection

in Wuhan (No.1); 2020. Available from:

http://www.gov.cn/xinwen/2020-01/23/content_5471751.htm.

23. Wallinga J, Teunis P. Different Epidemic Curves for Severe Acute Respiratory

Syndrome Reveal Similar Impacts of Control Measures. American Journal of

Epidemiology 2004;160(6):509-16.

24. Wenzhou Municipal People's Government. Notice of the Leading Group for the

Prevention and Control of COVID-19 in Wenzhou (No. 3); 2020a. Available from:

http://www.wenzhou.gov.cn/art/2020/1/28/art_1219304_41858870.html.

[Accessed March 31, 2020].

25. Wenzhou Municipal People's Government. Notice of the Leading Group for the

Prevention and Control of COVID-19 in Wenzhou (No. 4); 2020b. Available from:

http://www.wenzhou.gov.cn/art/2020/1/30/art_1219304_41862456.html.

26. Wenzhou Municipal People's Government. Notice of the Leading Group for the

Prevention and Control of COVID-19 in Wenzhou (No. 6); 2020c. Available from:

http://www.wenzhou.gov.cn/art/2020/2/2/art_1219304_41867466.html. [Accessed

March 31, 2020].

27. World Health Organization. Rolling updates on coronavirus disease (COVID-19);

2020a. Available from:

https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-they

-happen. [Accessed April 20, 2020].

28. World Health Organization. WHO Director-General's opening remarks at the

media briefing on COVID-19-11 March 2020; 2020b. Available from:

https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-a

t-the-media-briefing-on-covid-19---11-march-2020.

29. Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic

and international spread of the 2019-nCoV outbreak originating in Wuhan, China:

a modelling study. The Lancet 2020.

30. Xinhuannet. No local newly confrimed cases for the first time in China; 2020.

Available from:

http://www.xinhuanet.com/politics/2020-03/18/c_1125730531.htm.

31. Yang Z, Zeng Z, Wang K, Wong S-S, Liang W, Zanin M, et al. Modified SEIR and

AI prediction of the epidemics trend of COVID-19 in China under public health

interventions. Journal of Thoracic Disease 2020.

411   **Figure 1. The timeline of interventions implemented in Wuhan, Wenzhou, and**

412   **Shenzhen.**

419   **Figure 2. The trends of R_t for Wuhan, Wenzhou and Shenzhen.**

A

B

Figure 3. The simulation of COVID-19 trend for constant R_t (A) and

time-varying R_t (B) of three policies in South Korea. The red curve with

corresponding dashed lines represents the expected cumulative confirmed cases with

95% CI and the blue curve with corresponding dashed lines represent the expected

number of infected and infectious without isolation (*I*) with 95% CI.

A

B

Figure 4. The simulation of COVID-19 trend for constant R_t (A) and

time-varying R_t (B) of three policies in Italy. The red curve with corresponding

dashed lines represents the expected cumulative confirmed cases with 95% CI and the

blue curve with corresponding dashed lines represent the expected number of infected

and infectious without isolation (I) with 95% CI.

A

B

Figure 5. The simulation of COVID-19 trend for constant R_t (A) and

time-varying R_t (B) of three policies in the United States. The red curve with

corresponding dashed lines represents the expected cumulative confirmed cases with

95% CI and the blue curve with corresponding dashed lines represent the expected

number of infected and infectious without isolation (*I*) with 95% CI.

452 **Supplementary Information:**

453 Stochastic SIHR model

454 COVID-19 prevalence

455 Model validity