Evaluate the risk of resumption of business for the states of New York, New Jersey and Connecticut via a pre-symptomatic and asymptomatic transmission model of COVID-19

Authors: Ting Tian[1]†, Jianbin Tan[1]†, Yukang Jiang[1]†, Xueqin Wang[2,1]*, Heping Zhang[3]*

Affiliations:

[1]School of Mathematics, Sun Yat-sen University, Guangzhou, Guangdong, China

[2]School of Management, University of Science and Technology of China, Hefei, Anhui, China

[3]School of Public Health, Yale University, New Haven, CT, USA

†Dr. T. Tian, J. Tan and Y. Jiang contributed equally to this article

*Dr. X. Wang and Dr. H. Zhang are corresponding authors

Abstract

Background

The United States has the highest numbers of confirmed cases of COVID-19, where they took up nearly half in the hot spot states of New York, New Jersey, Connecticut, and California. The workforce in these states was required to work from home except for essential services. It is necessary to evaluate an appropriate date for resumption of business since premature reopening of economy will lead to broader spread of COVID-19, while the opposite situation would cause greater loss of economy.

Methods

To consider pre-symptomatic and asymptomatic transmission of COIVD-19, it is crucial to evaluate the unobserved numbers of unidentified infectious individuals but not the observed number of confirmed cases, which reflect the real-time risks of different stage of infectious disease. We proposed an epidemic model in considering the pre-symptomatic transmission and asymptomatic transmission of COVID-19 to evaluate the real-time risk of epidemic for the states of New York, New Jersey and Connecticut, and compared with California state (where it effectively phased reopened on May 8) for assessments of the appropriate Monday for resumption of business.

Results

The predicted numbers of unidentified infectious individuals per 100,000 for states of New York, New Jersey and Connecticut which are close to those in California state are 12.23 with 95% CI (10.68, 13.57) on June 1, 25.65 with 95% CI (20.04, 30.43) on

June 15, 28.49 with 95% CI (19.10, 38.65) on June 22, respectively, which may cause

11.23%, 15.64% and 17.32% higher than the estimated number of cumulative

confirmed cases on July 11, if the second wave of the infection would happen after

people return to work.

Conclusions

It may be feasible for states of New York, New Jersey and Connecticut to reopen

business on June 1 (or even May 18), June 15 and June 22. For the period after

resumption of work, if the number of unidentified infectious individuals is still non-

zero, the risks for the second wave of the infection would never vanish.

Keywords

resumption of business, pre-symptomatic transmission, asymptomatic transmission,

Bayesian modeling, unidentified infectious individuals

Introduction

The outbreak of novel coronavirus disease (COVID-19) has spread over 200 countries since December 2019[1]. It is unprecedented to have over 7 million cumulative confirmed cases of COVID-19 worldwide[2]. The "battle" against COVID-19 in China has provided experience and likely outcomes of interventions to the ongoing hard-hit areas. As a novel and acute infectious disease, the transmissibility of COVID-19 was unclear at the early stage of epidemic, the Chinese government implemented relatively strict nonpharmaceutical interventions in the hot spot areas, where the public transportations were suspended within and outside of the city in Hubei province since January 23, 2020[3]. All nationwide residents were recommended to stay at home except for essential needs. The holiday season of the Chinese Spring Festival had been prolonged to late February when essential services were recommended operating gradually outside Hubei province[4]. In April, a comprehensive resumption of business is approaching in China[5].

Epidemic situations in the United States

Meanwhile, the confirmed case of COVID-19 was reported since late January 2020 in the United States[6]. There were over 1000 cumulative confirmed cases on March 13, 2020[7], when the White House declared a national emergency concerning COVID-19 outbreak[8] and issued a "call to action" coronavirus guidelines on March 16, 2020[9]. The United States has become the most severe country of COVID-19 with 366,346, 154,154, 40,468, and 94,743 cumulative confirmed cases in the states

of New York, New Jersey, Connecticut, and California until May 24, 2020[10],

respectively. Making things worse, New York and California are the two states that

contribute the most to the real gross domestic product (GDP) over years in the United

States (Figure 1)[11].

The state of New York had the first confirmed case of COVID-19 reported on

March 1, 2020, and it proclaimed an executive order on March 16, 2020, including

reducing half of the local government workforce, allowing the statewide nonessential

workforce to work from home starting on March 17, 2020, and closing all schools

starting on March 18, 2020[12]. However, as the rapidly increasing number of

additional cases of COVID-19 in the state, the governor announced an aggressive

policy of "New York State on PAUSE" on March 20, 2020[13], and required all

people in the state to wear masks or face covering in public since April 15, 2020[14].

On May 1, all statewide K-12 schools and college facilities continued to close for the

remaining academic year[15]. The guide of "NY Forward Reopening" Plan was

available on May 11, 2020 and there were three regions reopened businesses for phase

one on May 15, 2020[16]. Until May 24,2020, there had been seven regions reopened

for businesses[17].

The state of New Jersey had the first positive case of COVID-19 reported on

March 4, 2020[18]. The governor of New Jersey recommended the cancellation of

statewide public gatherings over 250 individuals since March 12, 2020, suspended

visiting state prisons and statewide halfway houses since March 14, 2020, and closed

restaurants, bars, movie theaters, gyms, casinos since March 16, 2020[19]. The

governor announced an order including the statewide stay at home and closure of

statewide non-essential retail industries on March 21, 2020[20]. States park and gold

courses were effectively reopened on May 2, 2020[21]. The car gathering provision

was permitted resumption on May 12, 2020 and construction and non-essential retail

provisions were resumed on May 18, 2020[22]. In-person sales were authorized to

reopen at car, motorcycle, and boat dealerships and bike shops by appointment only

and with social distancing measure on May 20,2020[23]. Three states of New York,

New Jersey and Connecticut signed a multi-state agreement on beaches reopened on

May 22, 2020[24].

The state of Connecticut had the first positive case as a Connecticut resident on

March 8, 2020[25]. Three states of New York, New Jersey, and Connecticut

announced measures to slow the spread of COVID-19 throughout the tri-state area on

March 16, 2020[26]. Governor signed an executive order for businesses and residents

"stay safe, stay home" on March 20, 2020[27]. Statewide K-12 schools which

remained closed for the rest of the academic year were announced on May 5,

2020[28]. The restaurants, offices, retail stores, outdoor museum and zoos as the first

phase were authorized to reopen on May 20, 2020[29] and hair salons and

barbershops were aligned to reopen in Early June[30].

The state of California had two positive cases of COVID-19 reported on January

26, 2020[31]. One month later, there was the first possible case of local transmission

of COIVD-19 in California[32]. The "stay home except for essential needs" order was

issued on March 19, 2020, where all individuals were required to stay at home except

for the workforce in 16 critical infrastructure sectors[33]. On May 7, 2020, the state

public health officer determined the entire state gradually moved to Stage 2 of

California's Pandemic Resilience Roadmap, i.e. reopening the lower-risk workplaces

and other spaces[34]. Therefore, California state had phased reopened on May 8.

The timelines of four states (New York, New Jersey, Connecticut and California)

of intervention details are shown in Figure 2. According to the capacity of medical

facilities in the website of COVID-19.direct[10], there could be 52200 beds and

219,000 hospitals, 25,300 beds and 5,300 hospitals, 8,200 beds and 46 hospitals, and

87,800 beds and 463,000 hospitals in the states of New York, New Jersey,

Connecticut, and California, respectively.

On April 13, 2020, there were 7 states joined together to form a multi-state

council to combat COVID-19 and to help reopen the economy including the states of

New York, New Jersey, and Connecticut[35]. As the epidemic of COVID-19 was

getting worse in the United States, the state governors issued a series of interventions

to slow the spread of virus, including requiring all individuals to stay at home and

reducing the workforce in the industries. The whole social system was slowed down

and the employment in total nonfarm was fell over the month change (Figure 3),

where the unemployment rates were 14.7% and 13.3% in April and May, 2020[36],

and thus, the time to restore the economy in the United States, especially for the states

of New York, New Jersey, Connecticut, has been the most significant and

consequential decision for the president of the United States and the governors of the

states. To recover the economy for people to return to work, the key concern is the

152   risk and severity of the second wave of the infection.

Methods

Data collection

To emphasize the effectiveness of our model for evaluating the real-time risk of

epidemic before the second wave of the infection, we focused on the data during the

period before the protest in United States[37]. We collected the epidemic data from

March 13, 2020 when the national emergency concerning COVID-19 was proclaimed

to May 24, 2020 in four states of New York, New Jersey, Connecticut, and California.

The data are available on the New York Times[38].

Model

Based on the report of WHO, the transmission of COVID-19 could be caused by the

individuals infected with the virus before significant symptoms developed[39], or

even the carriers who did not develop symptoms. Pre-symptomatic transmission and

asymptomatic transmission would interfere with our judgment on the current

magnitude of outbreak of infectious diseases because of the existence of the duration

between catching virus and being confirmed by testing. To consider this nature of

COIVD-19, it is crucial to evaluate the unobserved number of unidentified infectious

individuals but not the observed number of confirmed cases.

To understand the mechanism of COVID-19, we divided a concerned population

into four compartments: susceptible (S), unidentified infectious (I), self-healing

without being confirmed (H), and confirmed cases (C).

(a) The susceptible (S) individuals, who have no immunity to the disease, are the

majority of the population at an early stage of the epidemic.

(b) Unidentified infectious (I) individuals are infectious individuals without being

confirmed, which consist of two kinds of individuals: infectors who would have

symptoms eventually and asymptomatic carriers who would not develop

symptoms. We assumed that I would transfer to H or C eventually.

(c) Self-healing without being confirmed (H) individuals are assumed to be no longer

infectious and resistant to COVID-19.

(d) Confirmed cases (C) contain two kinds of individuals: patients in hospital and

asymptomatic carriers who are assumed to stay at home, and unable to transmit

the disease.

We introduced Susceptible (S) - Unidentified infectious (I) - Self-healing without

being confirmed (H) - Confirmed cases (C) (SIHC) model with four compartments:

Susceptible (S), unidentified infectious (I), self-healing without being confirmed (H),

confirmed cases (C). We divided I into two part: asymptomatic carriers without

being confirmed (I_1), infectors who would have symptoms eventually (I_2). Similarly,

we divided C into two part: confirmed cases who self-quarantine or self-isolate (C_1),

confirmed patients at hospitals (C_2). The transfer diagram of S, I_1, I_2, H, C_1, C_2 are

given in Figure 1 in the Supplementary.

The dynamic system of numbers of compartments S, I, H, and C at the time

t: $S(t)$, $I(t)$, $H(t)$, and $C(t)$ are assumed as:

$$\frac{dS(t)}{dt} = -\rho\theta(t)\frac{I(t)}{N}S(t),$$

$$\frac{dI(t)}{dt} = \rho\theta(t)\frac{I(t)}{N}S(t) - (\frac{1}{D_C} + \frac{1}{D_H})I(t),$$

$$\frac{dH(t)}{dt} = \frac{I(t)}{D_H},$$

$$\frac{dC(t)}{dt} = \frac{I(t)}{D_C}, \tag{1}$$

where ρ is transmissibility and $\theta(t)$ is the average contact numbers per person at

the time t[40], which is assumed to be time-varying; D_H is the average length of

duration between catching the virus and self-healing without being confirmed. D_c is

the average length of duration between catching virus and being confirmed by testing;

N is the total number of population.

Let $\alpha(t) = \rho\theta(t)$. Generally, $\theta(t)$ is controlled by policy interventions, which

is a constant at an early stage of the epidemic and decreases along with the

implementation of interventions until the control policies are completely carried out,

i.e., reaching the lowest level. Based on this point of view, we furthermore assumed

$\alpha(t)$ is monotonically decreasing curve[41] with four parameters α_0, d, m, and η:

$$\alpha(t) = \alpha_0(\frac{\eta}{1+\exp\left(\lambda_m\left(t-d-\frac{m}{2}\right)\right)} + 1 - \eta), \tag{2}$$

where α_0 denotes the maximum of $\alpha(t)$ at an early stage of outbreak of infectious

disease, d is the time when the control measures start to be effective and the $\alpha(t)$

starts to decline, m represents the duration of decreasing process of $\alpha(t)$, η is the

maximum decreasing proportion of $\alpha(t)$ comparing with α_0, i.e., the minimum of

$\alpha(t)$ is $\alpha_0(1 - \eta)$. λ_m is chosen as $\frac{2\log((1-\varepsilon)/\varepsilon)}{m}$ and ε is fixed to be 0.01. The

graph of $\alpha(t)$ is given in Figure 2 in the Supplementary.

By the setting above, we can derive the time-varying reproductive rate R_t[42,

43] by SIHC (susceptible, unidentified infectious, self-healing without being

confirmed, and confirmed cases) model:

220
$$R_t = \frac{\alpha_0 D_C D_H}{D_H + D_C} \cdot \left(\frac{\eta}{1 + \exp\left(\lambda_m \left(t - d - \frac{m}{2} \right) \right)} + 1 - \eta \right). \tag{3}$$

221   Let $\Omega = (\alpha_0, \eta, m, D_C)$ and $\Theta = (\Omega, d, D_H)$, we introduced randomness to the

222   epidemic model above: Let S_t, I_t, H_t, C_t be the observed values of S, I, H, C at the

223   time t with constraint:

224
$$N = S_t + I_t + H_t + C_t, \tag{4}$$

225   and $A_t = (I_t, H_t, C_t)$ is assumed to be a multi-dimensional Markov process, i.e., for

226    $g < t$ and given H_g:

227
$$p(A_t | A_{s \le g}) = p(A_t | A_g) = Poiss\left(A_t \middle| A_t(A_g, \Theta) \right), \tag{5}$$

where $A_t(A_g, \Theta)$ is the evolve values of I, H and C at the time t determined by

the deterministic dynamic system (1) with the initial value (S_g, I_g, H_g, C_g) and

parameters Θ. $Poiss(\cdot | v)$ is the density of multi-dimensional independent Poisson

distribution with mean vector v. Similarly, we define $I_t(A_g, \Theta)$, $C_t(A_g, \Theta)$,

$H_t(A_g, \Theta)$, which are the corresponding values at time t determined by the

deterministic dynamic system (1) with the initial value (S_g, I_g, H_g, C_g) and

parameters Θ.

Let $\alpha(t) = \alpha(k), k \le t < k + 1$, k is an integer. For simplicity, we could

assume that H_t $(g < t)$ is not random given the initial value (S_g, I_g, H_g, C_g) and

parameters Θ, i.e., $H_t | A_g =: H_t(A_g, \Theta)$ a.s., if we know D_C.

Notice that the observable data is $C_{1:T}, N$, where T is the length of time.

Assume that $H_1 = 0$ and treat $I_{1:T}$ and $H_{2:T}$ as the missing data, we applied

techniques of Markov Chain Monte Carlo (MCMC)[44-47] for approximations of

posterior distributions of parameters Ω and unknown numbers of I, H at each time

by given priors with values of d, and D_H. Once the prior distribution of Ω: $\pi(\Omega)$,

was given, we apply the Bayesian method combined with MCMC (Markov Chain

Monte Carlo) to simulate the posterior distribution:

245
$$\pi(\Omega, I_{1:T}, H_{2:T} | C_{1:T}, N, d, m, D_H, H_1) \propto$$

246
$$\pi(\Omega) \prod_{t=1}^{T-1} Poiss\big(I_{t+1} \big| I_{t+1}(A_t, \Theta)\big) Poiss\big(C_{t+1} \big| C_{t+1}(A_t, \Theta)\big) I(H_{t+1} =$$

247
$$H_{t+1}(A_t, \Theta)). \tag{6}$$

For the choice of prior distributions, notice that D_C is governed by the mean of

incubation period, in terms of studies about the incubation period of COVID-19[48],

we choose log-normal distribution with log-mean $\log(5.1)$ as the prior distribution

of D_C. Similarly, we choose log-normal distribution with log-mean $\log(30)$ as the

prior distribution of m, which implies a priori hypothesis that the control policies are

completely carried out one month later, while the priors of other parameters are

chosen non-informative or flat priors.

For the fixed parameters d, m, and D_H, d is taken the beginning of

implementation of important policy interventions, i.e., $d = 8$ for New York state

("New York State on PAUSE" in Figure 2 in the main text), $d = 9$ for New Jersey

state ("statewide stay at home" in Figure 2 in the main text), $d = 8$ for Connecticut

state ("stay safe, stay home" in Figure 2 in the main text), $d = 7$ for California state

("stay home except for essential needs" in Figure 2 in the main text). D_H is assumed

to be 9.5 according to the clinical study of asymptomatic cases[49]. The point

estimates of Ω, $I_{1:T}$ and $H_{2:T}$ were presented as the median of the posterior

distribution while 95% credible intervals were constructed with 2.5% and 97.5%

quantiles.

Moreover, since $\pi(\Omega, I_{1:T}, H_{2:T} \mid C_{1:T}, N, d, m, D_H. H_1)$ has no close form, we

apply Gibbs sampler imbedded Metropolis-Hastings steps for updating parameters.

Let $\Omega^{(k)}, I_{I:T}^{(k)}, H_{2:T}^{(k)}$ be the k-th sample, given initial $\Omega^{(0)}, I_{I:T}^{(0)}, H_{2:T}^{(0)}$, the details of

updating are given in Algorithm S1 and Algorithm S2 in Supplementary.

Results

To assess the risk and severity of the second wave of the infection after the

resumption of business, it is important to consider the unobserved numbers of

unidentified infectious individuals in the community. Our approach is based on

publicly available data to simulate the possible outcomes of an outbreak of COVID-

19 at different business reopening dates and to provide recommendations as to when it

is reasonable safe for people back to work.

The estimated results for four states are given in Figures 3 in the Supplementary.

The estimated cumulative numbers of confirmed cases on July 11 are 395638 with

95% CI (391461, 399830) for New York state, 179220 with 95% CI (173988, 185356)

for New Jersey state, 52720 with 95% CI (49306, 56948) for Connecticut state, and

213263 with 95% CI (192476, 232771) for California state, which estimated

proportions of cumulative infected individuals without being confirmed (including I

and H) are 20.11% with 95% CI (19.66%, 20.65%), 32.57% with 95% CI (31.77%,

34.48%), 34.72% with 95% CI (33.03%, 37.86), and 36.85% with 95% CI (35.33%,

38.62%), respectively. Normally, higher testing rate would lead to smaller proportion

of cumulative infected individuals without being confirmed. The estimated

proportions were consistent with the reported totally test per 100,000 of four states

(9312, 7434, 6321, and 4396 for states of New York, New Jersey, Connecticut, and

California, respectively)[50].

Focusing on states of New York, New Jersey, and Connecticut, the predicted

numbers of daily new infected individuals and new confirmed cases, are given in

Figure 4 and the details of infectious diseases maps for states of New York, New

Jersey, and Connecticut are given in Figures 4-6 in the Supplementary and the daily

trajectories of COVID-19 in combined three states were shown in an additional

graphics interchange format (gif) file 1. Note that there are gaps between the numbers

of daily new confirmed cases and new infected individuals per day in Figure 4, which

is caused by the pre-symptomatic transmission and asymptomatic transmission of

COVID-19. We called this phenomenon as lagging effect of daily new confirmed

cases.

For considering the potential risk of infectious disease, we present the estimated

numbers of unidentified infectious individuals per 100,000 for states of New York,

New Jersey, Connecticut, and California, which are given in Figure 5. Based on

Figure 5, the estimated timing of the peak of unidentified infectious individuals in the

states of New York, New Jersey, and Connecticut had passed. Compared with

California state, where it effectively reopened the economy on May 8, the numbers of

unidentified infectious individuals in the states of New York, New Jersey and

Connecticut was higher and showed different downward patterns.

We considered the possible timing for resuming business in the different

Mondays. The predicted numbers of unidentified infectious individuals per 100000

for states of New York, New Jersey, Connecticut, and California on May 11, May 18,

May 25, June 1, June 8, June 15, June 22 and June 29 are given in Table 1. The dates

that the predicted numbers of unidentified infectious individuals per 100000 for states

of New York, New Jersey and Connecticut close to those in California state are May

18, June 15 and June 22, respectively.

To realize the potential risk of resuming business on different Mondays, we

simulated the possible outcomes that the second wave of the infection would happen

after people return to work. There could be maximum contact degree once all

individuals back to work, and then the stringent interventions were implemented to

curb the wave of infectious disease a week later. In doing so, for the starting days of

June 1, June 8, June 15, June 22 and June 29, we modified the estimated $\hat{\alpha}(t)$ by

estimated $\hat{\alpha}_0$ to last 7 days and specifying to its minimum value afterward, i.e.,

$\hat{\alpha}(t) = \hat{\alpha}_0(1 - \hat{\eta})$ representing the most intensity of control policies for states of

New York, New Jersey, and Connecticut accordingly. The simulated results are given

in Figure 6.

For New York, the predicted numbers of cumulative confirmed cases per 100000

on July 11 for the simulated outbreak of epidemic when people would return to work

on June 1, June 8, June 15, June 22 and June 29 are 11.23%, 7.65%, 5.02%, 3.10%,

and 1.68% higher than the estimated number of cumulative confirmed cases per

100000 on July 11 if no resumption of business.

For New Jersey, the predicted numbers of cumulative confirmed cases per

100,000 on July 11 for the simulated outbreak of epidemic when people would return

to work on June 1, June 8, June 15, June 22 and June 29 are 31.05%, 22.51%,

15.64%, 10.13%, and 5.73% higher than the estimated number of cumulative

confirmed cases per 100000 on July 11 if no resumption of business.

For Connecticut, the predicted numbers of cumulative confirmed cases per

100000 on July 11 for the simulated outbreak of epidemic when people would return

to work on June 1, June 8, June 15, June 22 and June 29 are 46.44%, 35.27%,

25.63%, 17.32%, and 10.27% higher than the estimated number of cumulative

confirmed cases per 100000 on July 11 if no resumption of business.

Discussion

To understand the spread of COVID-19 for the states of New York, New Jersey,

Connecticut, we applied an epidemic model considering pre-symptomatic and

asymptomatic transmission of COVID-19 to estimate the numbers of daily new

infected individuals and new confirmed case for these states. Normally, the numbers

of new infected cases mirrored the actual degree of transmission, while the numbers

of new confirmed cases, often lagged the real situation by the existence of the

duration between catching virus and being confirmed by testing. According to Figure

4, such a lagging effect could not be neglected at an early stage of epidemic, would

tend to disappear over time.

For the control of epidemic, it was reasonable to alleviate the lagging effect in

the process of epidemic, which could be achieved by shortening the D_c in SIHC

model. Notice that D_c was also affected by the testing rate, except the length of

incubation period. The estimated $\widehat{D}_C$ of all four states were smaller than the reported

average length of incubation period: 5.1[48] (2.37 with 95% CI (2.32, 2.44) for New

York, 4.53 with 95% CI (4.31, 4.89) for New Jersey, 4.94 with 95% CI (4.55, 5.46)

for Connecticut, and 4.99 with 95% CI (4.61, 5.37) for California), it implied the

policy enhancing the testing rate may be implemented in all four states, which was

effective to reduce the number of unidentified infectious individuals.

The decision for the resumption of business is not only a public health issue, but

also economic issue. What we are focusing on is the epidemiological feasibility of

returning to work at an early date, there are still many other factors needed to be paid

attention[51]. The higher numbers of unidentified infectious individuals were

obtained, the higher risks for the resumption of business would have been. According

to Figure 6, we concluded that there could be risks for the resumption of businesses

on June 1 for the states of New York, New Jersey and Connecticut. If governors of

states delayed one week, or more, for the resumption of businesses, that would

significantly scale down the simulated magnitude of the second wave of the infection.

But the benefit of delaying decision would be smaller and smaller when the current

control policies continue to be effective.

Conclusion

Due to the fact that California state had been in process of reopening economy

on May 8, compared with it, it may be feasible for states of New York, New Jersey

and Connecticut to reopen business on June 1 (or even May 18), June 15 and June 22,

which may cause 11.23%, 15.64% and 17.32% higher than the estimated number of

cumulative confirmed cases per 100,000 on July 11, if the second wave of the

infection would happen after people return to work. Meanwhile, for the period after

resumption of work, if the number of unidentified infectious individuals is still non-

zero, the risks for the second wave of the infection would never vanish, which

underscore the necessity of maintaining epidemic prevention measures, such as,

wearing a mask and keeping a social distance.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The data that support the findings of this study are available from the New York

Times

Competing interests

The authors declare that they have no competing interests

Funding

Not applicable

Authors' contributions

TT, XW and HZ developed the idea and research. TT, JT and YJ wrote the first draft

of the manuscript and all other authors discussed results and edited the manuscript.

TT, JT and YJ collected and validated epidemiological data and census data. All

authors read and approved the final manuscript

Acknowledgements

We would like to thank all individuals who are collecting epidemiological data of the

COVID-19 outbreak around the world.

References

1.   **Distribution of COVID-19 cases in the world**

     [http://2019ncov.chinacdc.cn/2019-nCoV/global.html]

2.   **Rolling updates on coronavirus disease (COVID-19)**

[https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-

they-happen]

3.   Chinese Center for disease control and prevention: **Epidemic update and risk**

assessment of 2019 Novel Coronavirus. In.: Chinese Center for disease

control and prevention; 2020.

4.   **State Council stresses orderly resumption of production and operation**

[http://english.www.gov.cn/policies/latestreleases/202002/22/content_WS5e50

f888c6d0595e03c213c0.html]

5.   **China's progress on business resumption on April 16**

[http://english.www.gov.cn/news/topnews/202004/16/content_WS5e9869f5c6

d0b3f0e949598e.html]

6.   Holshue ML, DeBolt C, Lindquist S, Lofy KH, Wiesman J, Bruce H, Spitters

C, Ericson K, Wilkerson S, Tural A *et al*: **First Case of 2019 Novel**

Coronavirus in the United States. *New England Journal of Medicine* 2020.

7.   World Health Organization: **Coronavirus disease 2019 (COVID-19)**

Situation Report-53. In.; 2020.

8.   **Proclamation on Declaring a National Emergency Concerning the Novel**

Coronavirus Disease (COVID-19) Outbreak

[https://www.whitehouse.gov/presidential-actions/proclamation-declaring-

national-emergency-concerning-novel-coronavirus-disease-covid-19-

outbreak/]

9.   The White House: **30 Days to slow the spread**. In.; 2020.

10.   Johns Hopkins University Center for Systems Science and Engineering (JHU

CSSE): **2019 Novel Coronavirus COVID-19 (2019-nCoV) Data**. In.; 2020.

11.   The United States Census Bureau: **Table of United States counties**. In.; 2020.

12.   **Video, Audio, Photos & Rush Transcript: During Coronavirus Briefing,**

Governor Cuomo Issues Executive Order

13.   **Governor Cuomo Signs the 'New York State on PAUSE' Executive Order**

[https://www.governor.ny.gov/news/governor-cuomo-signs-new-york-state-

pause-executive-order]

14.   **Video, Audio, Photos & Rush Transcript: Amid Ongoing COVID-19**

Pandemic, Governor Cuomo Issues Executive Order Requiring All People

in New York to Wear Masks or Face Covering in Public

[https://www.governor.ny.gov/news/video-audio-photos-rush-transcript-amid-

ongoing-covid-19-pandemic-governor-cuomo-issues-1]

15.   **Amid Ongoing COVID-19 Pandemic, Governor Cuomo Announces**

Schools and College Facilities Statewide Will Remain Closed for the Rest

of the Academic Year [https://www.governor.ny.gov/news/amid-ongoing-

covid-19-pandemic-governor-cuomo-announces-schools-and-college-

facilities]

16.   **Amid Ongoing COVID-19 Pandemic, Governor Cuomo Announces Three**

Regions of New York State Ready to Begin Reopening May 15th

[https://www.governor.ny.gov/news/amid-ongoing-covid-19-pandemic-

governor-cuomo-announces-three-regions-new-york-state-ready]

17. **Video, Audio, Photos & Rush Transcript: Amid Ongoing COVID-19 Pandemic, Governor Cuomo Announces Seventh Region Hits Benchmark to Begin Reopening Tomorrow** [https://www.governor.ny.gov/news/video-audio-photos-rush-transcript-amid-ongoing-covid-19-pandemic-governor-cuomo-announces-29]

18. **Governor Murphy, Acting Governor Oliver, and Commissioner Persichilli Announce First Presumptive Positive Case of Novel Coronavirus in New Jersey**

    [https://www.state.nj.us/governor/news/news/562020/approved/20200304e.shtml]

19. **Governor Murphy Signs Executive Order**

    [https://www.state.nj.us/governor/news/news/562020/approved/news_archive.shtml]

20. **Governor Murphy Announces Statewide Stay at Home Order, Closure of All Non-Essential Retail Businesses**

    [https://www.state.nj.us/governor/news/news/562020/approved/20200320j.shtml]

21. **Governor Murphy Signs Executive Order Reopening State Parks and Golf Courses**

    [https://nj.gov/governor/news/news/562020/approved/20200429c.shtml]

22. **Governor Murphy Signs Executive Order Permitting Resumption of Non-Essential Construction, Curbside Pickup at Non-Essential Retail**

Businesses, and Gatherings in Cars

[https://nj.gov/governor/news/news/562020/approved/20200513a.shtml]

23. **Governor Murphy and Superintendent Callahan Announce Administrative Order Authorizing In-Person Sales at Car Dealerships, Motorcycle Dealerships, Boat Dealerships, and Bike Shops**

[https://nj.gov/governor/news/news/562020/approved/20200519d.shtml]

24. **Governor Murphy, Governor Cuomo, Governor Lamont, Governor Carney Announce Multi-State Agreement on Beaches Ahead of Memorial Day Weekend**

[https://nj.gov/governor/news/news/562020/approved/20200515a.shtml]

25. **Governor Lamont Announces First Positive Case of Novel Coronavirus Involving a Connecticut Resident** [https://portal.ct.gov/Office-of-the-Governor/News/Press-Releases/2020/03-2020/Governor-Lamont-Announces-First-Positive-Case-of-Novel-Coronavirus-Involving-a-Connecticut-Resident]

26. **Northeast Governors From Connecticut, New York, New Jersey Announce Collective Measures to Combat Spread of COVID-19**

[https://portal.ct.gov/Office-of-the-Governor/News/Press-Releases/2020/03-2020/Northeast-Governors-From-Connecticut-New-York-New-Jersey-Announce-Collective-Measures]

27. **Governor Lamont Signs Executive Order Asking Connecticut Businesses and Residents: 'Stay Safe, Stay Home'** [https://portal.ct.gov/Office-of-the-Governor/News/Press-Releases/2020/03-2020/Governor-Lamont-Signs-

Executive-Order-Asking-Connecticut-Businesses-and-Residents-Stay-Safe]

28.   **Governor Lamont Announces Classes at K-12 Public Schools Will Remain**

Canceled for the Rest of the Academic Year [https://portal.ct.gov/Office-of-

the-Governor/News/Press-Releases/2020/05-2020/Governor-Lamont-

Announces-Classes-at-Schools-Will-Remain-Canceled-for-the-Rest-of-the-

Academic-Year]

29.   **Governor Lamont Releases Rules for Businesses Under First Phase of**

Connecticut's Reopening Plans Amid COVID-19

[https://portal.ct.gov/Office-of-the-Governor/News/Press-Releases/2020/05-

2020/Governor-Lamont-Releases-Rules-for-Businesses-Under-First-Phase-of-

Reopening-Plans]

30.   **Governor Lamont and Governor Raimondo to Align Hair Salon and**

Barber Shop Reopenings in Early June [https://portal.ct.gov/Office-of-the-

Governor/News/Press-Releases/2020/05-2020/Governor-Lamont-and-

Governor-Raimondo-to-Align-Hair-Salon-and-Barber-Shop-Reopenings-in-

Early-June]

31.   **Two Confirmed Cases of Novel Coronavirus in California**

[https://www.cdph.ca.gov/Programs/OPA/Pages/NR20-001.aspx]

32.   **CDC Confirms Possible First Instance of COVID-19 Community**

Transmission in California

[https://www.cdph.ca.gov/Programs/OPA/Pages/NR20-006.aspx]

33.   **Order   of the state public health officer**

[https://www.cdph.ca.gov/Programs/CID/DCDC/CDPH%20Document%20Li

brary/COVID-19/Health%20Order%203.19.2020.pdf]

34. **Order of the state public health officer**

[https://www.cdph.ca.gov/Programs/CID/DCDC/CDPH%20Document%20Li

brary/COVID-19/SHO%20Order%205-7-2020.pdf]

35. **Massachusetts Joins New York, New Jersey, Connecticut, Pennsylvania, Delaware and Rhode Island's Multi-State Council to Get People Back to Work and Restore the Economy**

[https://www.governor.ny.gov/news/massachusetts-joins-new-york-new-jersey-connecticut-pennsylvania-delaware-and-rhode-islands]

36. U.S. Department of Labor: **News Release Bureau of Labor Statistics**. In.; 2020.

37. **The last days of May: A visual timeline of the George Floyd Protests**

[https://www.mndaily.com/article/2020/06/pf-the-george-floyd-protests-a-visual-timeline]

38. **Coronavirus in the U.S.: Latest Map and Case Count**

[https://www.nytimes.com/interactive/2020/us/coronavirus-us-cases.html]

39. World Health Organization: **Coronavirus disease 2019 （COVID-19） Situation Report-73**. In.; 2020.

40. Blackwood JC, Childs LM: **An introduction to compartmental modeling for the budding infectious disease modeler**. *Letters in Biomathematics* 2018, **5**(1):195-221.

541   41.   Tan J, Jinag Y, Tian T, Wang X: **P-SIHR probabilistic graphical model: An**

542   **applicable model of COVID-19 in estimating the number of infectious**

543   **individuals without isolation and time-varying reproduction number**. *Acta*

544   *Mathematicae Applicatae Sinica* 2020, **42**(3).

545   42.   Anderson RM, Anderson B, May RM: **Infectious diseases of humans:**

546   **dynamics and control**: Oxford university press; 1992.

547   43.   Jones JH: **Notes on R0**. *Califonia: Department of Anthropological Sciences*

548   2007.

549   44.   Ghosh JK, Delampady M, Samanta TJAItBAT, Methods: **Bayesian Inference**

550   **and Decision Theory**. 2006:29-63.

551   45.   Andrieu C, De Freitas N, Doucet A, Jordan MIJMl: **An introduction to**

552   **MCMC for machine learning**. 2003, **50**(1-2):5-43.

553   46.   Chib S, Greenberg EJTas: **Understanding the metropolis-hastings**

554   **algorithm**. 1995, **49**(4):327-335.

555   47.   Soubeyrand S: **Construction of semi-Markov genetic-space-time SEIR**

556   **models and inference**. 2016.

557   48.   Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, Meredith HR, Azman AS,

558   Reich NG, Lessler J: **The Incubation Period of Coronavirus Disease 2019**

559   **(COVID-19) From Publicly Reported Confirmed Cases: Estimation and**

560   **Application**. *Annals of Internal Medicine* 2020.

561   49.   Hu Z, Song C, Xu C, Jin G, Chen Y, Xu X, Ma H, Chen W, Lin Y, Zheng Y *et*

562   *al*: **Clinical characteristics of 24 asymptomatic infections with COVID-19**

screened among close contacts in Nanjing, China. *Sci China Life Sci* 2020.

50.   **Live tracker: How many coronavirus cases have been reported in each**

U.S. state? [https://www.politico.com/interactives/2020/coronavirus-testing-

by-state-chart-of-new-cases/]

51.   **Return to Work for Healthcare Personnel with Confirmed or Suspected**

COVID-19 [https://www.cdc.gov/coronavirus/2019-ncov/hcp/return-to-

work.html]

Table 1. The estimated numbers of unidentified infectious individuals per 100,000 in the four states in the different Mondays.

State \\ Date	New York	New Jersey	Connecticut	California
May 11	30.68 (28.48, 32.61)	82.15(74.17, 88.56)	75.41 (64.16, 87.44)	22.5 (20.94, 24.29)
May 18	22.64 (20.87, 24.31)	65.4 (57.62, 71.66)	64.94 (53.86, 77.03)	23.66 (22.18, 25.47)
May 25	16.71 (14.8, 18.32)	51.74 (43.37, 58.56)	55.2 (42.8, 67.95)	24.68 (22.6, 26.86)
June 1	12.23 (10.68, 13.57)	40.98 (33.59, 47.06)	46.92 (35.31, 59.07)	25.64 (23.1, 28.16)
June 8	8.96 (7.66, 10.06)	32.45 (25.98, 37.9)	39.8 (28.88, 51.07)	26.63 (23.47, 29.64)
June 15	6.55 (5.49, 7.46)	25.65 (20.04, 30.43)	33.72 (23.64, 44.36)	27.62 (23.76, 31.22)
June 22	4.77 (3.92, 5.53)	20.23 (15.4, 24.47)	28.49 (19.1, 38.65)	28.58 (24.06, 32.91)
June 29	3.48 (2.79, 4.11)	15.97 (11.85, 19.66)	24.09 (15.43, 33.45)	29.53 (24.26, 34.67)

Real Gross Domestic Product in 2018

Figure 1. The contributions of Gross Domestic Product for four states of New

York, New Jersey, Connecticut, and California.

Figure 2. The timeline of interventions against COVID-19 in the states of New

York, New Jersey, Connecticut and California.

Figure 3. The month over change of employment change in the total nonfarm

sectors in the United States

Figure 4. **The magnitude of outbreak of COVID-19 in the three states of New York (Figure 2-a), New Jersey (Figure 2-b), and Connecticut (Figure 2-c) from March 13 to July 11, 2020.** The daily confirmed new cases form March 13 to May 24, 2020, were observed data, combing with the projected cases until July 11, 2020.

Figure 5. The trend of the numbers of unidentified infectious individuals per 100000 for states of New York, New Jersey, Connecticut, and California from March 13 to July 11, 2020. The corresponding 95% credible intervals were represented in each state.

Figure 6. The possible outbreak of epidemic when people returned to work in the forthcoming Mondays for states of New York, New Jersey, and Connecticut (per 100,000). The points represent the observed cumulative confirmed cases in the three states of New York, New Jersey and Connecticut.

Additional files

Additional graphics interchange format (gif) file 1: **A COVID-19 distribution map of three states of New York, New Jersey, and Connecticut.** The daily cumulative confirmed cases from March 8 to June 1, 2020 were presented.