COVID 19: Real-time Forecasts of confirmed cases, active cases, and health infrastructure requirements for India and its states using the ARIMA model.

Rishabh Tyagi[1*] Mahadev Bramhankar[1*] Mohit Pandey[1] Kishore M[1]

1: IIPS, Mumbai; *: Co-first authors

Abstract

Background: COVID-19 is an emerging infectious disease which has been declared a Pandemic by the World Health Organization (WHO) on 11th March 2020. The Indian public health care system is already overstretched, and this pandemic is making things even worse. That is why forecasting cases for India is necessary to meet the future demands of the health infrastructure caused due to COVID-19.

Objective: Our study forecasts the confirmed and active cases for COVID-19 until July mid, using time series Autoregressive Integrated Moving Average (ARIMA) model. Additionally, we estimated the number of isolation beds, Intensive Care Unit (ICU) beds and ventilators required for the growing number of COVID-19 patients.

Methods: We used ARIMA model for forecasting confirmed and active cases till the 15th July. We used time-series data of COVID-19 cases in India from 14th March to 22nd May. We estimated the requirements for ICU beds as 10%, ventilators as 5% and isolation beds as 85% of the active cases forecasted using the ARIMA model.

Results: Our forecasts indicate that India will have an estimated 7,47,772 confirmed cases (95% CI: 493943, 1001601) and 296,472 active cases (95% CI:196820, 396125) by 15th July. While Maharashtra will be the most affected state, having the highest number of active and confirmed cases, Punjab is expected to have an estimated 115 active cases by 15th July. India needs to prepare 2,52,001 isolation beds (95% CI: 167297, 336706), 29,647 ICU beds (95% CI: 19682, 39612), and 14,824 ventilator beds (95% CI: 9841, 19806).

Conclusion: Our forecasts show an alarming situation for India, and Maharashtra in particular. The actual numbers can go higher than our estimated numbers as India has a limited testing facility and coverage.

1. Introduction

COVID-19 is an emerging infectious disease caused by severe acute respiratory syndrome Corona Virus 2 (SARS-CoV-2). The first human cases of COVID-19 were first reported by officials in Wuhan City, China, in December 2019. Retrospective investigations by Chinese authorities have identified human cases with onset of symptoms in early December 2019. While some of the earliest known cases had a link to a wholesale food market in Wuhan, some did not (WHO, 2020). The World Health Organization (WHO) has declared the outbreak of the novel Coronavirus (COVID-19) as a pandemic on 11th March 2020.

This pandemic is spreading very quickly throughout the world, and the number of confirmed cases reached 6.44 million on 2nd June. Globally, the recovery rate is around 46.7 percent and case fatality rate of 5.9 percent (Worldometer, 2nd June 2020). On 2nd June, India's confirmed cases of COVID-19 crossed 2 Lakhs. The recovery rate was 48.31 percent amongst COVID-19 patients. The case fatality rate of COVID-19 in India declined to 2.8 percent on 2nd June in India. They also told that 73 percent of COVID-19 deaths in India are people with co-morbidities (Press Information Bureau 2020 c).

In India, the first positive case of COVID-19 was detected on 30th January 2020, in Kerala (India Today, 2020). With the experience of the overwhelming burden of COVID-19 in Europe and China, the Government of India implemented a complete nationwide lockdown on 25th March for 21 days (Press Information Bureau 2020 a). The purpose of the nationwide lockdown was to contain the spread of the Coronavirus so that the Government could take a multi-prong strategy: add more beds in its network of hospitals, scale up the production of the testing kits for COVID-19 and personal protection equipment (PPE) for the health workers. On 3rd June, with respect to the health infrastructure in the country for the management of COVID-19, 952 dedicated COVID hospitals with 1,66,332 Isolation beds, 21,393 ICU beds, and 72,762 Oxygen supported beds are available. 2,391 dedicated COVID Health Centres with 1,34,945 Isolation beds; 11,027 ICU beds and 46,875 Oxygen supported beds have been operationalized. (Press Information Bureau 2020 c).

A few studies have attempted to predict numbers of confirmed and active cases at the national and state level for India to assess the burden of COVID-19 in future. Tiwari et al. (2020) made their prediction for India based on the pattern of China using a machine learning approa0ch.

They predicted that the peak of the cases for India would be attained between the third and fourth weeks of April 2020 in India. The infection was likely to be controlled by the end of May 2020. Chakraborty (2020) used a hybrid approach, based on an ARIMA and Wavelet-based forecasting model, to make short-term forecasts of the number of daily confirmed cases in Canada, France, India, South Korea, and the United Kingdom. The ARIMA model has been used in many studies for forecasting case count of epidemic diseases based on the time series modelling (Gupta and Pal 2020; Tandon et al., 2020; Kumar et al. 2020; Perone, 2020).

In India, the healthcare infrastructure is not up to the mark. The number of hospital beds per 1000 population is less than one - it is just one indicator to cite the vulnerable situation of India's health care systems (World Bank Database). Several studies from countries like China, Italy and Spain show the requirements of different healthcare facilities required to fight this pandemic. Guan et al. (2020) in his study on data regarding 1099 patients with laboratory-confirmed COVID-19 from 552 hospitals in 30 provinces, in mainland China through 29th January 2020, found that 5.0% were admitted to the ICU, 2.3% who underwent invasive mechanical ventilation, and 1.4% who died. A study by Lazzerini et al. (2020) for Italy and Spain suggested that 40–55% of COVID-19 positive cases have been hospitalized, with 7–12% requiring admission to intensive care units. The COVID-19 Cases in Italy shows that 10–25% of patients will require ventilation, and some patients will need ventilation for several weeks. Remuzzi et al., (2020) find that the percentage of patients in intensive care reported daily in Italy between 1st March and 11th March 2020, has consistently been between 9% and 11% of patients who are actively infected.

In the Indian scenario, a press release by PIB on 8th May stated that of the total 35,902 active cases, 4.8 per cent patients are in ICU, 1.1 per cent on ventilators and 3.3 per cent are on oxygen support (Press Information Bureau 2020 b). With the rising number of cases, it is crucial to forecast the requirements of healthcare infrastructure like isolation beds, ICU beds and ventilators at national and state level to make the respective authorities aware of the situation they might be facing shortly. In this study, we use the well-known ARIMA time-series model to forecast the confirmed and active cases for India at national and state level. Another objective of this study is to estimate the requirements of healthcare infrastructure in the future based on the forecasts of active cases.

2. Data and Methods

2.1 Data

Data on COVID-19 was obtained from the data-sharing portal covid19india.org. Information is collected on daily confirmed and active cases at the national and state level from 14[th] March to 22[nd] May 2020. This dataset provides excel of the patient database, which is used to build a required time-series. From this dataset, we have used cumulative confirmed and active cases for India and selected states. Selection of states is based on the criterion that states should have at least 100 confirmed cases on 3rd May 2020. By using this selection criterion, India and 17 other states selected which are Andhra Pradesh, Bihar, Delhi, Gujarat, Haryana, Jammu & Kashmir, Odisha, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Punjab, Rajasthan, Tamil Nadu, Telangana, Uttar Pradesh, West Bengal.

2.2 Method

In this study, we applied the ARIMA model to our considered time series data of COVID-19 cases by using R- studio, for forecasting cumulative confirmed and active cases. This model has been preferred for the time series forecasting in various fields as the model predictions are based on different parameters. The required parameters for the ARIMA model are *(p, d, q)* which evaluate autoregressive term, integrated moving average and past lag term for stationary time series respectively. The degree of parameters p, d and q are determined based on the partial Auto-correlation function (PACF), Augmented Dickey-Fuller Test to test the stationary time series and Complete Auto-Correlation Function (ACF) respectively. These parameters help to capture overall fluctuations in earlier time-series which helps to predict the future. ARIMA has some advantages over the other models as it not only captures the overall picture in earlier trend, but it also provides a 95% confidence interval (CI) for our point estimates.

The model for forecasting future confirmed and active cases of COVID-19 cases is represented as,

$$ARIMA\ (p, d, q): \qquad X_t = \alpha 1\ X_{t\ -1} + \alpha 2\ X_{t\ -2} + \beta 1 Z_{t-1} + \beta 2\ Z_{t\ -2} + Z_t \qquad (i)$$

$$\text{Where,} \qquad Z_t = X_t - X_{t-1} \qquad (ii)$$

Here, X_t is the predicted number of cumulative confirmed and active COVID-19 cases at t^{th} day; $\alpha 1, \alpha 2, \beta 1$ and $\beta 2$ are parameters whereas Z_t is the residual term for t^{th} day.

Finally, we estimate the requirement of health infrastructure, i.e., the requirement of Intensive care unit (ICU), Ventilator support, and isolation beds using the following equations based on forecasted active cases.

Required number of beds = Forecasted active cases at ith day*(85/100)

Required number of ICU = Forecasted active cases at ith day*(10/100)

Required number of Ventilators = Forecasted active cases at ith day*(5/100)

Based on these equations, infrastructure estimates till the mid of July.

3. Results

3.1 ARIMA model fit for confirmed and active cases of COVID-19 in India

Fig 1(a) & 1(b), shows the ARIMA model fitted correlogram for the active and confirmed cases. In these figures, we see four subfigures which reveal the trend for the earlier and forecasted values for both confirmed and active cases. Forecasting based on PACF and ACF graphs helps to determine parameters p and q. Moreover, the best ARIMA model fit is considered having the lowest Akaike Information Criterion (AIC) value.

Fig 1(a) also shows the fitted model for total confirmed cases with ARIMA (2, 3, 3) having low AIC. With the help of this model, we have predicted future confirmed cases with 95% CI till 15th July 2020. Similarly, for total active cases, we have a suitable model with ARIMA (3, 3, 1) which helps to predict active cases for the same duration.

Fig. 1(a) Correlogram and ARIMA forecast for the Confirmed COVID-19 Cases in India

Fig. 1(b) Correlogram and ARIMA Forecast for the Active COVID-19 Cases in India

Source: Based on the author's calculations using R-studio

3.2 Forecasted Confirmed and active cases of COVID-19 for India:

The analysis result for India in **Table 1 and Fig. 2** shows the total confirmed and active cases for India. It has observed that with time, confirmed cases will increase at a faster rate. India will be having 5,19,258 confirmed cases (95% CI: 389542, 648794) at June 30 and 7,47,772 confirmed cases (95% CI: 493943, 1001601) till July 15.

Similarly, based on our forecasts for active cases for India, we found that active cases will be 2,19,793 (95% CI: 162014, 277571) at 30th June and 2,96,472 active cases (95% CI: 196820, 396125) at 15th July. The combined number of recovered and death cases are also increasing rapidly and are expected to be around 60 percent of total confirmed cases by 15th July based on our forecasts.

Fig.2 Trend of Forecasted Confirm and Active COVID-19 Cases over the Period in India

Source: Based on the author's calculations using R-studio

3.3 Forecasted Confirmed cases of COVID-19 for the selected states of India:

Table 2, shows the forecast of confirmed for selected states of India on 15th June, 30th June and 15th July. Our forecasts show that that confirmed cases for states vary a lot from one state to another. Our forecasts for 15th June for confirmed cases show that Haryana, Punjab and Kerala will have around 2000 confirmed cases. However, states like Maharashtra 108712 confirmed cases (95% CI: 89631, 127793), Tamil Nadu 36759 confirmed cases (95% CI: 26703, 46816),

Delhi 26257 confirmed cases (CI: 20794, 31720) & Gujarat 22860 confirmed cases (95% CI: 18187, 27533) would have a high number of confirmed cases at the 15th June. On 30th June, only Haryana and Punjab will be having around 2000 confirmed cases based on our forecasts. Though, states like Maharashtra 149072 confirmed cases (95% CI: 109237, 188908), Tamil Nadu 54263 confirmed cases (95% CI: 31834, 76692), Delhi 34968 confirmed cases (95% CI: 24090, 45847) & Gujarat 28852 confirmed cases (95% CI: 19770, 37934) will have to occur confirmed cases on 30th June. On 15th July, Punjab will be having the least number of confirmed cases around 4000 confirmed cases. However, other states like Maharashtra 189433 confirmed cases (95% CI: 124301, 254565), Tamil Nadu 74637 confirmed cases (95% CI: 35282, 113991), Delhi (43680 confirmed cases (95% CI: 26271, 61089) & Bihar 38027 confirmed cases (95% CI: 14862, 61192) will have a high number of confirmed cases at the 15th July.

3.4 Forecasted active cases of COVID-19 for the selected states of India:

Table 3 shows the state-wise active cases for India. On 15th June, Punjab, Kerala and Andhra Pradesh will be having less than 1000 active cases based on our forecasts. However, states like Maharashtra (67984 active cases (95% CI: 56127, 79840)), Delhi (10907 active cases (95% CI: 5590, 16224)) & Tamil Nadu (8534 active cases (95% CI: 0, 21421)) will have a high number of active cases on the 15th June.

On 30th June, Haryana, Punjab, Andhra Pradesh and Kerala will be having fewer than 2000 active cases based on our forecasts. However, states like Maharashtra (91428 active cases (95% CI: 67487, 115639), Delhi (13840 active cases (95% CI: 3386, 24294) & Bihar (11680 active cases (95% CI: 3864, 19496) will have a high number of active cases on the 30th June. On 15th July, Punjab, Andhra Pradesh and Kerala will be having under 2000 active cases based on our forecasts. However, states like Maharashtra (114873 active cases (95% CI: 76308, 153438), Bihar (17479 active cases (95% CI: 2997, 31961) & Delhi (16773 active cases (95% CI: 142, 33404) will have a high number of active cases on 15th July.

3.5 Estimates of required healthcare infrastructure for India

Table 4, 5, and 6 show the required number of isolation beds, ICU beds and ventilators for active cases of COVID-19 on 15th June, 30th June and 15th July, respectively for India. The

total number of isolation beds required is 130500 (95% CI: 107894, 153106). It also shows that 15353 ICU beds (95% CI: 12693, 18012) and 7676 ventilators (95% CI: 6347, 9006) will be required in India by 15th June. On 30th June. The total number of isolation beds required for India is 186824 (95% CI: 137712, 235935). It also shows that 21979 ICU beds (95% CI: 16201, 27757) and 10990 ventilators (95% CI: 8101, 13879) will be required in India by 30th June. India will require 252001 isolation beds (95% CI: 167297, 336706), 29647 ICU beds (95% CI: 19682, 39612) and 14824 ventilators (95% CI: 9841, 19806) on July 15.

3.6 Estimates of required healthcare infrastructure at state-level

Table 4, 5, and 6 show the required number of isolation beds, ICU beds and ventilators for patients that will be suffering from COVID-19 on 15[th], 30th June and 15th July respectively for Indian states. The overall picture for estimates shows a huge variation across the states. By 15th June, Kerala, Haryana, Punjab and Andhra Pradesh will recover most of the active cases and requires less than 1000 isolation beds, around 100 ICU beds and 50 ventilator beds according to our point estimates. While, on the other hand, Maharashtra, with the largest number of active cases shows an alarming situation and will require 57786 isolation beds (95% CI: 47708, 67864), 6798 ICU beds (95% CI: 5613, 7984) and 3399 ventilator beds (95% CI: 2806, 3992).

At the end of June month, Kerala, Haryana, Punjab, Odisha, Telangana and Andhra Pradesh will require less than 2000 isolation beds, under 200 ICU beds and less than 100 ventilator beds. While, in Maharashtra, will require 77714 isolation beds (95% CI: 57364, 98064), 9143 ICU beds (95% CI: 6749, 11537) and 4571 ventilator beds (95% CI: 3374, 5768).

At the mid of July, Kerala, Haryana, Punjab, Odisha, Telangana and Andhra Pradesh will require less than 2000 isolation beds, around 200 ICU beds and 100 ventilator beds. While, on the other hand, Maharashtra, with the largest number of active cases and will require 97642 isolation beds (95% CI: 64862, 130422), 11487 ICU beds (95% CI: 7631, 15344) and 5744 ventilators (95% CI: 3815, 7672) will be required by 15th July.

4. Discussion and Conclusion

Our current forecasts for confirmed and active cases are in line with the actual number of cases. On 2nd June, India has 207,186 confirmed cases, while our forecasts suggested 206,764 confirmed cases (95% CI: 194194, 219334). India had 101,070 active cases on 2nd June, while

our forecasts suggested that India will have 104,526 active cases (95% CI: 96145, 112907). Our results also show that daily confirmed cases will increase at a faster pace as around mid-July India will be getting around 16,000-17,000 daily confirmed cases. By 15th July, we forecast that the total confirmed cases will be around 7.5 lakhs, whereas total active cases will be close to 2.96 lakhs based on our point estimates for India. Maharashtra will be the most affected state even at the 15th July with 1.9 Lakh confirmed cases and 1.15 Lakh active cases while Punjab will be least affected having 2648 confirmed cases and only 115 active cases based on our point estimates. The total number of isolation beds required for India is 2,52,001 (95% CI: 167297, 336706). Our estimates also show that 29,647 ICU beds (95% CI: 19682, 39612) and 14,824 ventilators (95% CI: 9841, 19806) will be required in India by the 15th July. When it comes to states, Maharashtra will be the most affected state and will require 97,642 isolation beds (95% CI: 64862, 130422).

On 3rd June, with respect to the health infrastructure in the country for the management of COVID-19, 952 dedicated COVID hospitals with 1,66,332 Isolation beds, 21,393 ICU beds and 72,762 Oxygen supported beds are available. 2,391 dedicated COVID Health Centres with 1,34,945 Isolation beds; 11,027 ICU beds and 46,875 Oxygen supported beds have also been operationalized. Based on the current preparedness of the Indian healthcare infrastructure, it seems likely that India will not be facing a shortage at the national level. However, the health infrastructure stress will likely be acute in high-burden states, led by Maharashtra, Tamil Nadu and Delhi as they have more than 10,000 active cases on 2nd June. In particular, Maharashtra is the most affected state with around 40,000 active cases as on 2nd June. On 29th May, BrihanMumbai Municipal Corporation (BMC) reported that almost all intensive care unit (ICU) beds for COVID-19 patients were occupied in Mumbai. They also tell that the city is already using 72% of the total 373 ventilators available for COVID-19 treatment (Mint, 2020). This shows an alarming situation for Maharashtra and in particular, Mumbai. The most affected state Maharashtra needs to prepare at least 1 lakh isolation beds, 12,000 ICU beds and 6000 ventilators by 15th July to fight this pandemic efficiently.

Our forecasts are based on the data which grossly under-reported the cases as found by many studies for India. Rao *et al.,* (2020), projected that India might be detecting 1 out of 4 cases of COVID-19 based on his mathematical model. Another study by (Goli and James, 2020) also find

that India is detecting just 3.6% of the total number of infections of COVID-19 with a huge variation across its states. They also suggest that India must increase its testing capacity and go for widespread testing to know the real picture of the pandemic. Lack of coverage of testing is not revealing the true prevalence of COVID-19. (Subramanian and James, 2020) suggested that for detection of the true prevalence of COVID-19 infections in the country, India can adopt the well-established National Family Health Survey (NFHS) framework as a solution to ascertain the true prevalence of COVID-19. Understanding the urgency of the situation, on 8th May, The Indian Council of Medical Research (ICMR) announced that it will conduct a study in 75 affected districts across the country to identify people who were exposed to the novel coronavirus infection and yet showed mild or no symptoms. The study can help ascertain whether there has been community transmission of the respiratory disease in those areas or not. This much-needed study by ICMR will reveal the true prevalence of asymptomatic COVID-19 infected people to check for community spread (The Economic Times, 2020). The data issues of COVID-19 such as under-reporting, lack of coverage etc. results in an underestimation of the cases, which in turn will lead to underestimation of the forecasts in our study.

Our forecasts show an alarming situation for India in the future. In coming months, the actual numbers can go higher than our forecasts of confirmed cases, active cases and healthcare infrastructure as migrant workers are returning home in large numbers currently in large numbers (Tyagi *et al.,* 2020). Currently, India has lifted lockdown with restrictions applying only on the containment zones. This will lead to a surge in the number of daily confirmed and active cases. The requirement of isolation beds, ICUs and ventilators will also be increased in that scenario. So, India and its majorly affected states like Maharashtra, Gujarat, Tamil Nadu, Bihar and Delhi need to be well prepared for the pandemic challenge in future and focus on increasing their healthcare infrastructure, and other states should also remain alert till the pandemic completely recedes.

Table 1: Forecast of Confirmed and Active Cases of COVID-19 for India from 23rd May to 15th July, 2020,

Date	Confirmed Cases Point Estimates	95% Confidence Interval Lower	95% Confidence Interval Upper	Active Cases Point Estimates	95% Confidence Interval Lower	95% Confidence Interval Upper	Death & Recovered (%)
23 May 2020	131258	130705	131810	73186	71515	72858	44
25 May 2020	144843	143116	146570	78244	76558	79930	46
27 May 2020	159100	155473	162727	84551	81574	87528	47
29 May 2020	174321	168258	180384	91021	86476	95566	48
31 May 2020	190164	181148	199179	97682	91332	104032	49
02 June 2020	206764	194194	219334	104526	96145	112907	49
04 June 2020	224206	207521	240891	111556	100930	122182	50
06 June 2020	242329	220948	263711	118771	105691	131851	51
08 June 2020	261231	234534	287928	126171	110433	141909	52
10 June 2020	280918	248292	313545	133756	115161	152351	52
12 June 2020	301326	262133	340520	141527	119877	163176	53
14 June 2020	322510	276089	368930	149482	124584	174380	54
16 June 2020	344458	290144	398772	157623	129282	185963	54
18 June 2020	367148	304251	430044	165949	133974	197924	55
20 June 2020	390607	318423	462791	174460	138659	210260	55
22 June 2020	414824	332635	497013	183156	143340	222972	56
24 June 2020	439792	346862	532722	192037	148015	236060	56
26 June 2020	465524	361103	569946	201104	152686	249522	57
28 June 2020	492013	375335	608692	210356	157352	263359	57
30 June 2020	519258	389542	648974	219793	162014	277571	58
02 July 2020	547264	403715	690814	229415	166672	292158	58
04 July 2020	576027	417835	734219	239222	171325	307119	58
06 July 2020	605547	431888	779206	249215	175973	322456	59
08 July 2020	635827	445863	825790	259392	180616	338169	59
10 July 2020	666863	459742	873984	269755	185253	354257	60
12 July 2020	698658	473515	923801	280303	189884	370722	60
14 July 2020	731212	487167	975256	291036	194509	387563	60
15 July 2020	747772	493943	1001601	296472	196820	396125	60

Table 2: Forecast of Confirmed Cases of COVID-19 for India and its States on 15th June, 30th June and 15th July, 2020, India

Dates	June mid			June end			July mid		
	PE	95% CI		PE	95% CI		PE	95% CI	
STATES		LB	UB		L.B.	UB		L.B.	UB
India	333384	283103	383665	519258	389542	648974	747772	493943	1001601
Andhra Pradesh	4104	3019	5190	5002	2814	7191	5900	2377	9424
Bihar	12036	8245	15826	23135	12143	34126	38027	14862	61192
Delhi	26257	20794	31720	34968	24090	45847	43680	26271	61089
Gujarat	22860	18187	27533	28852	19770	37934	34843	20475	49212
Haryana	1901	1107	2694	2422	839	4005	2943	407	5479
Jammu & Kashmir	2802	2014	3591	3621	2064	5178	4441	1959	6923
Karnataka	4973	3552	6394	6996	4107	9886	9019	4350	13688
Kerala	2276	1236	3316	4014	1072	6955	6345	240	12449
Madhya Pradesh	11192	8547	13837	14342	9240	19443	17491	9452	25531
Maharashtra	108712	89631	127793	149072	109237	188908	189433	124301	254565
Odisha	2910	1978	3843	3986	2174	5798	5062	2196	7927
Punjab	2262	0	6123	2441	0	10786	2648	0	16885
Rajasthan	12358	9907	14808	16025	11133	20917	19693	11854	27531
Tamil Nadu	36759	26703	46816	54263	31834	76692	74637	35282	113991
Telangana	2924	1798	4050	3650	1404	5897	4377	778	7976
Uttar Pradesh	11888	9273	14503	15733	10514	20952	19579	11218	27940
West Bengal	6289	4774	7804	8137	5191	11084	9986	5322	14649

Source: Based on the author's calculations using R-studio

Table 3: Forecast of Active Cases of COVID-19 for India and its States on 15th June, 30th June and 15th July, 2020, India

Dates	June mid			June end			July mid		
		95% CI			95% CI			95% CI	
STATES	PE	LB	UB	PE	L.B.	UB	PE	L.B.	UB
India	153529	126934	180125	219793	162014	277571	296472	196820	396125
Andhra Pradesh	1000	0	2180	1065	0	3327	1131	0	4685
Bihar	6922	3684	10160	11680	3864	19496	17479	2997	31961
Delhi	10907	5590	16224	13840	3386	24294	16773	142	33404
Gujarat	9133	5175	13091	10716	3213	18219	12299	572	24027
Haryana	1057	0	2470	1589	0	4618	2200	0	7351
Jammu & Kashmir	1758	1079	2437	2372	1040	3705	2986	867	5105
Karnataka	4203	2579	5826	6121	2820	9422	8039	2707	13372
Kerala	918	102	1734	1362	0	3035	1807	0	4520
Madhya Pradesh	5109	1629	8589	6487	0	13379	7864	0	18866
Maharashtra	67984	56127	79840	91428	67487	115369	114873	76308	153438
Odisha	1285	330	2241	1627	0	3573	1969	0	5116
Punjab	115	0	2305	115	0	2967	115	0	3502
Rajasthan	6586	3884	9288	8962	3571	14353	11338	2703	19972
Tamil Nadu	8534	0	21421	9163	0	35097	9793	0	51497
Telangana	1399	100	2697	1854	0	4393	2310	0	6339
Uttar Pradesh	5006	2093	7919	6716	936	12496	8426	0	17660
West Bengal	3274	2298	4250	4157	2361	5954	5041	2275	7807

Table 4: Forecast of Active Cases of COVID-19 and Required Health Infrastructure for India and its States on 15th June, 2020, India

STATES	Active Cases	95% CI		ICU	95% CI		Ventilators	95% CI		Beds	95% CI	
	PE	LB	UB	PE	LB	UB	PE	LB	UB	PE	LB	UB
India	153529	126934	180125	15353	12693	18012	7676	6347	9006	130500	107894	153106
Andhra Pradesh	1000	0	2180	100	0	218	50	0	109	850	0	1853
Bihar	6922	3684	10160	692	368	1016	346	184	508	5884	3131	8636
Delhi	10907	5590	16224	1091	559	475	545	279	238	9271	4751	4039
Gujarat	9133	5175	13091	913	518	1309	457	259	655	7763	4399	11127
Haryana	1057	0	2470	106	0	247	53	0	124	899	0	2100
Jammu & Kashmir	1758	1079	2437	176	108	244	88	54	122	1494	918	2071
Karnataka	4203	2579	5826	420	258	583	210	129	291	3572	2192	4952
Kerala	918	102	1734	92	10	173	46	5	87	780	87	1474
Madhya Pradesh	5109	1629	8589	511	163	859	255	81	429	4343	1385	7301
Maharashtra	67984	56127	79840	6798	5613	7984	3399	2806	3992	57786	47708	67864
Odisha	1285	330	2241	129	33	224	64	16	112	1093	280	1905
Punjab	115	0	2305	12	0	231	6	0	115	98	0	1959
Rajasthan	6586	3884	9288	659	388	929	329	194	464	5598	3302	7895
Tamil Nadu	8534	0	21421	853	0	214	427	0	1071	7254	0	18208
Telangana	1399	100	2697	140	10	270	70	5	135	1189	85	2293
Uttar Pradesh	5006	2093	7919	501	209	792	250	105	396	4255	1779	6731
West Bengal	3274	2298	4250	327	230	425	164	115	213	2783	1953	3613

Source: Based on the author's calculations using R-studio

Table 5: Forecast of Active Cases of COVID-19 and Required Health Infrastructure for India and its States on 30th June, 2020, India

STATES	Active Cases	95% CI		ICU	95% CI		Ventilators	95% CI		Beds	95% CI	
	PE	LB	UB	PE	LB	UB	PE	LB	UB	PE	LB	UB
India	219793	162014	277571	21979	16201	27757	10990	8101	13879	186824	137712	235935
Andhra Pradesh	1065	0	3327	107	0	333	53	0	166	906	0	2828
Bihar	11680	3864	19496	1168	386	1950	584	193	975	9928	3284	16572
Delhi	13840	3386	24294	1384	339	288	692	169	144	11764	2878	2446
Gujarat	10716	3213	18219	1072	321	1822	536	161	911	9109	2731	15486
Haryana	1589	0	4618	159	0	462	79	0	231	1350	0	3925
Jammu & Kashmir	2372	1040	3705	237	104	370	119	52	185	2016	884	3149
Karnataka	6121	2820	9422	612	282	942	306	141	471	5203	2397	8008
Kerala	1362	0	3035	136	0	304	68	0	152	1158	0	2580
Madhya Pradesh	6487	0	13379	649	0	1338	324	0	669	5514	0	11372
Maharashtra	91428	67487	115369	9143	6749	11537	4571	3374	5768	77714	57364	98064
Odisha	1627	0	3573	163	0	357	81	0	179	1383	0	3037
Punjab	115	0	2967	12	0	297	6	0	148	98	0	2522
Rajasthan	8962	3571	14353	896	357	1435	448	179	718	7618	3036	12200
Tamil Nadu	9163	0	35097	916	0	351	458	0	1755	7789	0	29833
Telangana	1854	0	4393	185	0	439	93	0	220	1576	0	3734
Uttar Pradesh	6716	936	12496	672	94	1250	336	47	625	5709	796	10621
West Bengal	4157	2361	5954	416	236	595	208	118	298	3534	2007	5061

Source: Based on the author's calculations using R-studio

Table 6: Forecast of Active Cases of COVID-19 and Required Health Infrastructure for India and its States on 15th July, 2020, India

STATES	Active Cases	95% CI		ICU	95% CI		Ventilators	95% CI		Beds	95% CI	
	PE	LB	UB	PE	LB	UB	PE	LB	UB	PE	LB	UB
India	296472	196820	396125	29647	19682	39612	14824	9841	19806	252001	167297	336706
Andhra Pradesh	1131	0	4685	113	0	468	57	0	234	961	0	3982
Bihar	17479	2997	31961	1748	300	3196	874	150	1598	14857	2547	27167
Delhi	16773	142	33404	1677	14	12	839	7	6	14257	121	102
Gujarat	12299	572	24027	1230	57	2403	615	29	1201	10455	486	20423
Haryana	2200	0	7351	220	0	735	110	0	368	1870	0	6248
Jammu & Kashmir	2986	867	5105	299	87	510	149	43	255	2538	737	4339
Karnataka	8039	2707	13372	804	271	1337	402	135	669	6833	2301	11366
Kerala	1807	0	4520	181	0	452	90	0	226	1536	0	3842
Madhya Pradesh	7864	0	18866	786	0	1887	393	0	943	6685	0	16036
Maharashtra	114873	76308	153438	11487	7631	15344	5744	3815	7672	97642	64862	130422
Odisha	1969	0	5116	197	0	512	98	0	256	1674	0	4349
Punjab	115	0	3502	12	0	350	6	0	175	98	0	2977
Rajasthan	11338	2703	19972	1134	270	1997	567	135	999	9637	2297	16977
Tamil Nadu	9793	0	51497	979	0	515	490	0	2575	8324	0	43772
Telangana	2310	0	6339	231	0	634	115	0	317	1963	0	5388
Uttar Pradesh	8426	0	17660	843	0	1766	421	0	883	7162	0	15011
West Bengal	5041	2275	7807	504	228	781	252	114	390	4285	1934	6636

References

1. Chakraborty, T., & Ghosh, I. (2020). Real-time forecasts and risk assessment of novel coronavirus (COVID-19) cases: A data-driven analysis. *Chaos, solitons, and fractals*, *135*, 109850. Advance online publication. https://doi.org/10.1016/j.chaos.2020.109850

2. Goli, S and James, K.S. (2020) How much India detecting SARS-CoV-2 Infections? A model-based estimation. medRxiv preprint doi: https://doi.org/10.1101/2020.04.09.20059014.

3. Guan, W. J., Ni, Z. Y., Hu, Y., Liang, W. H., Ou, C. Q., He, J. X., ... & Du, B. (2020). Clinical characteristics of coronavirus disease 2019 in China. *New England journal of medicine*, *382*(18), 1708-1720.

4. Gupta R., Pal S. K. (2020), Trend Analysis and Forecasting of COVID-19 outbreak in India. MedRxiv. https://doi.org/10.1101/2020.03.26.200445110

5. Kumar P., Kalita H., Patairiya S., Sharma Y. D., Nanda C., Rani M., Rahmai J., Bhagavathula A. S. (2020), Forecasting the dynamics of COVID-19 Pandemic in Top 15 countries in April 2020 through ARIMA Model with Machine Learning Approach. MedRxiv. https://doi.org/10.1101/2020.03.30.20046227

6. Perone, G. (2020). An ARIMA model to forecast the spread and the final size of COVID-2019 epidemic in Italy. *medRxiv*.

7. Rao, A. S., Krantz, S. G., Kurien, T., Bhat, R., & Kurapati, S. (2020). Model-based retrospective estimates for covid-19 or Coronavirus in India: continued efforts required to contain the virus Spread. *Current Science*, *118*(7), 1023-1025.

8. Remuzzi, A., & Remuzzi, G. (2020). COVID-19 and Italy: what next? *The Lancet*.

9. Singh, R. K., Rani, M., Bhagavathula, A. S., Sah, R., Rodriguez-Morales, A. J., Kalita, H., Nanda, C., Sharma, S., Sharma, Y. D., Rabaan, A. A., Rahmani, J., & Kumar, P. (2020). Prediction of the COVID-19 Pandemic for the Top 15 Affected Countries: Advanced Autoregressive Integrated Moving Average (ARIMA) Model. *JMIR public health and surveillance*, *6*(2), e19115. https://doi.org/10.2196/19115

10. Subramanian, S. V., & James, K. S. (2020). Use of the Demographic and Health Survey framework as a population surveillance strategy for COVID-19. *The Lancet Global Health*.

11. Tandon, H., Ranjan, P., Chakraborty, T., & Suhag, V. (2020). Coronavirus (COVID-19): ARIMA based time-series analysis to forecast near future. arXiv preprint arXiv:2004.07859.

12. Tiwari, S., Kumar, S., & Guleria, K. (2020). Outbreak Trends of Coronavirus Disease-2019 in India: A Prediction. *Disaster medicine and public health preparedness*, 1–6. Advance online publication. https://doi.org/10.1017/dmp.2020.115

13. Tyagi, R., Dwivedi, L.K., & Sanzgiri, A. (2020). Estimation of Effective Reproduction Numbers for COVID-19 using Real-Time Bayesian Method for India and its States. Paper: 13 IIPS Analytical Series on COVID-19

14. World Health Organization. (2020). Coronavirus disease 2019 (COVID-19): situation report, 94.

Web-based Sources

15. Forecasting COVID-19 cases in India https://towardsdatascience.com/forecasting-COVID-19-cases-in-india-c1c410cfc730

16. India Today (2020). Kerala reports first confirmed coronavirus case in India. https://www.indiatoday.in/india/story/kerala-reports-first-confirmed-novel-coronavirus-case-in-india-1641593-2020-01-30

17. Livemint (2020). Mumbai coronavirus update: 99% ICU beds in hospitals are occupied, says BMC. https://www.livemint.com/news/india/coronavirus-in-mumbai-99-icu-beds-in-hospitals-is-occupied-says-bmc-11590742717577.html

18. Press Information Bureau (2020). 'Government of India issues Orders prescribing lockdown for containment of COVID-19 Epidemic in the country', 24th March.

19. Press Information Bureau (2020). 'PIB'S DAILY BULLETIN ON COVID-19', 8th May.

20. Press Information Bureau (2020). 'PIB'S DAILY BULLETIN ON COVID-19', 3rd June.

21. The Economic Times (2020). ICMR to study prevalence of asymptomatic COVID-19 infected people to check for community spread. Retrieved from https://economictimes.indiatimes.com/news/politics-and-nation/icmr-to-study-prevalence-of-asymptomatic-covid-19-infected-people-to-check-for-community-spread/articleshow/75631365.cms?from=mdr

Fig.4 Forecast trend of Confirmed cases of COVID-19 for Indian States

Fig.4 Forecast trend of Active cases of COVID-19 for Indian States

Trend for the future requirements of Beds facility in India

Trend for the future requirements of ICU facility India

Trend for the future requirements of Ventilators in India