Plasma concentration and safety of lopinavir/ritonavir in patients with Covid-19: a

retrospective cohort study

<u>**Running title**</u>: Lopinavir/r concentration and safety in Covid-19

Laurent Chouchana PhD[a] [ǁ], Sana Boujaafar PharmD[b] [ǁ], Ines Gana PhD[b], Laure-Hélène Preta

PharmD[a], Lucile Regard PhD[c], Paul Legendre PhD[d], Celia Azoulay MD[d], Etienne Canouï MD[e],

Jeremie Zerbit PharmD[f], Nicolas Carlier MD[c], Benjamin Terrier PhD[d], Solen Kerneis PhD[e], Rui

Batista PharmD[f], Jean-Marc Treluyer PhD[a,b], Yi Zheng PhD[b], Sihem Benaboud PhD[b]

[a] Regional Pharmacovigilance Center, Pharmacology Department, Cochin Hospital, AP-

HP.Centre – Université de Paris. Paris, France.

[b] Pharmacology Department, Cochin Hospital, AP-HP.Centre – Université de Paris. Paris,

France.

[c] Pneumology Department, Cochin Hospital, AP-HP.Centre – Université de Paris. Paris, France.

[d] Internal Medicine Department, Cochin Hospital, AP-HP.Centre – Université de Paris. Paris,

France.

[e] Antimicrobial Stewardship, Cochin Hospital, AP-HP.Centre – Université de Paris. Paris,

France.

[f] Pharmacy Department, Cochin Hospital, AP-HP.Centre – Université de Paris. Paris, France.

[ǁ] These authors equally contribute to this work; order is based on internal agreement.

Corresponding author: Dr Laurent Chouchana. Pharmacovigilance Center, Cochin Hospital, 27

rue du Faubourg Saint Jacques, 75014 Paris. Tel +33158414210; Mail

laurent.chouchana@aphp.fr

Number of words: 2784

Number of tables/figures: 2/1

ABSTRACT

Background

There is an urgent need of active treatment for coronavirus disease 2019 (Covid-19). Although

efficacy have not been proven, lopinavir/ritonavir 400 mg/100 mg twice daily has been proposed

as a treatment of moderate to severe Covid-19. Previously published cohorts showed Covid-19 is

associated with major inflammation. To date, no data are available regarding lopinavir/ritonavir

plasma concentration and its safety in Covid-19 patients.

Methods

Real-world Covid-19 experience based on a retrospective cohort study.

Results

On the cohort of 31 patients treated by lopinavir/ritonavir for Covid-19, we observed very high

lopinavir plasma concentrations, increased of 4.6-fold (IQR 2.9-6.4), with regards to average

plasma concentrations in HIV treatment. All except two patients were above the upper limit of

the concentration ranges of HIV treatment. In this cohort, about one over four to five patients

prematurely stopped lopinavir/ritonavir therapy due to a moderate adverse drug reaction, mainly

hepatic and gastrointestinal disorders.

Conclusion

Patients with Covid-19 pneumonitis treated with lopinavir/ritonavir have plasma concentrations

dramatically higher than expected. Owing to that high plasma concentration may be required for

antiviral activity against SARS-CoV-2, it appears that lopinavir dosage should not be reduced in

the absence of adverse effect. About 80% of the patients well tolerated lopinavir/ritonavir therapy

under these plasma concentrations. However, cautious is necessary as drug repurposing can be

associated with a new drug safety profile.

Funding

None

Keywords : Covid-19; SARS-CoV-2; plasma concentration; lopinavir; ritonavir; therapeutic

drug monitoring; pharmacovigilance; drug safety

INTRODUCTION

Since early December 2019, a pandemic infectious disease due to a novel coronavirus, called severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), is spreading all over the world. To date, no specific therapeutic agent has proven its clinical efficacy against this outbreak. However, due to the urgent need for a treatment, several antiviral drugs are being repurposed and used off-label to treat Covid-19, including lopinavir/ritonavir (LPV/r) (1). The latter is a human immunodeficiency virus (HIV) protease inhibitor approved since years. LPV is prescribed in patients with HIV in association with another protease inhibitor, ritonavir (RTV), used as a potent P450 3A4 cytochrome (CYP) inhibitor in order to dramatically increase LPV plasma exposure. LPV/r has been proposed in previous coronavirus outbreaks in 2003 and 2012 due to SARS-CoV-1 and to Middle East respiratory syndrome coronavirus (MERS-CoV), respectively (2, 3). In vitro activity and data from rodent models showed LPV has an antiviral activity against MERS-CoV(4, 5). It has also proven its potency to inhibit SARS-CoV-2 replication in vitro (6). Furthermore, in an open-label study comparing the addition of LPV/r to ribavirin in patients with SARS-CoV, the authors showed a reduced risk of acute respiratory distress syndrome (ARDS) or death by comparison with a historical control group treated by ribavirin alone (2). Viral load was also reduced in the LPV/r treated patients.

Based on these results, this drug has been considered to be potentially useful in patients with SARS-CoV-2. To date, efficacy of LPV in SARS-CoV-2 patients has been assessed in only one clinical trial in China, the LOTUS trial (7). This open-label randomized controlled clinical trial was performed in order to evaluate the efficacy and safety of LPV/r in severe Covid-19 patients. No benefit was observed with LPV/r treatment beyond standard care. However, in these patients with severe Covid-19, LPV/r treatment was started lately after a median time of 13 days after the

first symptoms. Furthermore, no therapeutic drug monitoring was performed for LPV/r to assess

an ideal drug exposure. Nevertheless, due to the absence of specific treatment for Covid-19,

LPV/r is still a therapeutic option used for patients with moderate to severe Covid-19 pneumonia.

To date, more than 40 clinical trials assessing the efficacy of LPV/r against Covid-19 are ongoing

around the world (8).

In order to assess LPV/r plasma concentration and its safety in Covid-19 patients, we conducted a

retrospective cohort study as part of routine care in adult patients hospitalized for a Covid-19

infection and requiring oxygen.

91
92

RESULTS

Patients

A total of 31 patients have been included during a 14 day period, all receiving at least one dose of LPV/r for a Covid-19 pneumonia. The median age of patients was 63 (interquartile range [IQR], 51-78) years and 71% were men (Table 1). All patients had a typical Covid-19 pneumonia according to radiological pattern and required oxygen with a median oxygen saturation of 92.5% (IQR, 90-96%) in room air. Pulmonary injury at CT scan was mostly moderate to extensive. Four (12%) patients had a doubtful or negative nasopharyngeal swab SARS-CoV-2 PCR, contrasting with a typical chest CT scan for Covid-19. Median C-reactive protein (CRP) and IL-6 levels at admission were 94.1 (45.4-176.0) mg/L and 60.4 (29.7-164.5) ng/mL. Twenty-one (68%) patients did not receive any antibiotic at admission or before being hospitalized. LPV/r was started within the first 48 hours after hospitalization and median time to starting LPV/r after symptom onset was 8 days (IQR, 7-10 days). No patient had drug-drug interaction involving RTV and 13 (42%) patients did not received concomitant therapy. LPV/r treatment duration was 7 (IDR, 3-8) days. At the end of the LPV/r course, five patients had recovered, eight patients were transferred to intensive care unit or died, 17 were still hospitalized with oxygen dependency. Clinical outcome is unknown for one patient that has been transferred to another hospital during treatment.

LPV/r plasma concentrations

Of the 31 patients who received LPV/r for Covid-19, plasma assays were not available for three patients due to technical issues and a total of 28 patients were analyzed. LPV plasma concentrations ranges from 4,732 to 35,012 ng/mL. Median level for trough (Cmin) and peak (Cmax) were 20,153 (IQR, 16,633-26,505) ng/mL and 22,550 (IQR 12,720-33,612) ng/mL,

respectively. In all patients, considering either trough or peak for LPV and RTV, plasma

concentrations were above the average concentrations observed during HIV treatment (4,000

ng/mL and 9,000 ng/mL for trough and peak concentrations, respectively) (Figure 1A). All

except two patients had plasma concentration over the upper limit of the concentration ranges

observed in HIV. With regards with these plasma concentration ranges, LPV concentrations in

Covid-19 patients were increased of about 4.6-fold (IQR 2.9-6.4) (Figure 1B). Extent of LPV

plasma concentration increase was not associated with CRP or IL-6 levels (data not shown).

Safety

LPV/r was discontinued before the end of scheduled course for 14 (45%) patients (Table 2).

Reasons for early ending LPV/r therapy were the occurrence of adverse drug reaction, therapeutic

limitation or patient deceased and poor efficacy in seven (22%), four (13%) and three (10%)

patients, respectively. Suspected adverse drug reaction were assessed as possibly related to LPV/r

therapy. They consisted in four cases of liver injuries (three cases moderate cytolytic hepatitis

between three and six times above the upper limit of normal range and one case of isolated

hyperbilirubinemia), two cases of gastrointestinal disorders (nausea/vomiting and diarrhea) and

one case of psychiatric disorders (agitation/anxiety). These adverse effects were mild and all

patients recovered after drug withdrawn. In patients with cytolytic hepatitis, LPV plasma

concentrations were 6.1-fold above average plasma concentrations in HIV patients.

DISCUSSION

To our knowledge, this is the first study showing that Covid-19 patients had very high LPV plasma exposure with regards to the therapeutic levels in HIV. Although LPV/r regimen was 400/100 mg bid as in HIV treatment, LPV plasma concentrations were in median about 4.6-fold higher, and up to 8-fold higher in some patients. All patients of this retrospective cohort had a LPV plasma concentration over the average concentrations observed in HIV patients. As LPV is extensively metabolized by the hepatic cytochrome P450 system, almost exclusively by the CYP3A isozyme, LPV/r therapy was not started in the event of drug-drug interaction. None was detected during therapy and no patient had liver cirrhosis, which could have impaired LPV/r clearance. Only two patients had a medical history of hepatitis, including one patient with an active hepatitis B without hepatic insufficiency.

Some clinical evidence show that patients with moderate to severe Covid-19 have a major inflammation. Elevated levels of blood IL-6 are commonly seen in severe Covid-19 illness and could led to a cytokine storm syndrome (9). Previous studies in macaques inoculated with SARS-CoV found that defects in T-cell and B-cell function and the excess production of type 2 cytokines could lead to a deficiency in control of viral replication and prolonged proinflammatory responses (10). In Covid-19 patients, a persistent blood virus load and an exacerbated inflammatory response has been shown to be partially driven by the transcriptional factor NFκB (11). These results are in line with our cohort where patients had very high CRP and IL-6 levels at starting therapy. Besides, it is long-standing known that inflammatory responses and infections decrease drug metabolism capacity in human and experimental animals (12). In cultured human hepatocytes, direct treatments with tumor necrosis factor-α (TNF-α), interleukin-1β (IL-1β), interleukin-6 (IL-6), interferon-γ (IFN-γ) and transforming growth factor-β (TGF-β) can reduce the expression of CYP1A2, CYP2C8 and CYP3A4 (13, 14). In animals and humans, several

studies or reports have shown the role of inflammation in reducing drug metabolism and CYP450

inhibition (15, 16). CYP3A4 expression is transcriptionally regulated by pregnane X receptor

(PXR), which is a ligand-dependent transcription factor (17). NF-κB activation by tumor necrosis

factor-α, in a setting of inflammatory response, has been shown to suppress PXR activation and

PXR-mediated gene expression (17, 18). This results in a reduced CYP450 expression and a

decreased drug metabolism, especially CYP450-mediated metabolism. Other genes regulated by

PXR, including ABCB1, coding for P-glycoprotein transporter, could also have their expression

decreased, contributing to decrease non-renal drug clearance, such as LPV/r. Owing to its low

oral bioavailability and extensive metabolism by CYP3A4 isoenzyme, LPV needs to be co-

administered with low doses of RTV, a potent CYP3A inhibitor, to achieve drug concentrations

high enough to inhibit viral replication (19). An added reduced expression of CYP3A4 could

dramatically increase LPV plasma concentrations in Covid-19 patients.

Interestingly, nevertheless very high plasma concentrations in all assayed patients, LPV/r was

overall well tolerated. The main adverse effects were moderate hepatobiliary disorders that have

been attributed to LPV/r therapy. In HIV, moderate-to-severe elevations in serum

aminotransferase levels (>5 times the upper limit of normal) are found in 3% to 10% of patients,

although rates may be higher in patients with HIV-HCV coinfection (20). These elevations are

usually asymptomatic and self-limited and can resolve even with continuation of the medication.

Drug causality could be hardly assessable as about 20% of the patients with Covid-19 have been

reported to have increased transaminases (9). However, in our cohort, transaminases quickly

decreased after stopping lopinavir, making drug causality probable. Overall, adverse effects

reported in this Covid-19 cohort are in line with LPV/r safety profile (21). Adverse effects led to

stop LPV/r treatment in seven (22%) patients, which could appear elevated, even though, due to

the relative limited number of patients treated, conclusions are difficult to draw. In the LOTUS

trial, assessing LPV/r efficacy in Covid-19, treatment had been stopped for adverse events in

13.8% of the patients in the LPV/r group.(7) As a comparison, in previous clinical trials with

LPV/r in HIV patients, about 11% to 14% of the patients experienced grade 2-4 adverse effects,

mainly gastro-intestinal or hepatic disorders (22, 23). However, physicians should be aware that

drug repurposing is a high risk condition, which can be associated with a different drug safety

profile (24). The drug is used in a setting not having been correctly assessed, that could led to an

increase of adverse effects and an unfavorable risk-to-benefits ratio (25). Finally, in other words,

78% of the patients in our cohort did not experienced adverse drug reaction, which could be

surprisingly in this context of very high plasma concentrations. One could hypothesize that the

relationship between the upper threshold of 8,000 ng/mL observed in HIV patients and lopinavir

safety is uncertain, at least during a short treatment as in Covid-19.

Keeping high LPV plasma concentration appears essential considering a possible clinical efficacy

of LPV in Covid-19. Antiviral activity assays on cultured cells showed that the 50% effective

concentration (EC_{50}) of LPV varied between 8.0 and 11.6 μM (i.e. between 5,030 and 7,290

ng/mL) for MERS-CoV.(5, 26, 27) Considering SARS-CoV, in vitro EC_{50} was 17.1 μM (i.e.

10,800 ng/mL), and a concentration as high as 25,000 ng/mL has been reported to inhibit viral

replication depending on cell type used (5, 28). For SARS-CoV-2, an intro study showed that

LPV but not RTV inhibits viral replication with EC_{50} at 26.6 μM (i.e. 16,800 ng/mL).(6) In our

cohort, median trough plasma level was 20,153 (IQR 16,633-26,505) ng/mL barely at these

concentrations or just above. The question of whether these plasma concentrations are effective

to inhibit *in vivo* SARS-CoV-2 replication, especially in lungs, is unknown. Furthermore, Covid-

19 associated vasculopathy or thrombosis could limit pulmonary diffusion. Altogether, this

suggest that high plasma concentrations are required to have an antiviral activity. Finally, as most

HIV protease inhibitors, LPV has very high protein binding of about 98.5% (27). Therefore, in

considering whether a dose is appropriate for SARS-CoV-2, it is critical to consider free drug

concentrations. For standard HIV-1, Cmin after 75 ng/mL that is substantially below EC_{50}

reported for coronaviruses (27). Therefore, targeting SARS-CoV-2 antiviral activity in patients is

challenging. It appears that LPV/r dosage should not be reduced on the basis of therapeutic drug

monitoring in the absence of adverse drug reaction.

Our study has several limitations. First, due to the retrospective design using data from routine

care, plasma assays were drawn at different times that prevent from an accurate pharmacokinetic

estimation. Second, according to different practices in wards, Covid-19 severity at baseline was

heterogeneous between patients. However, most importantly, in all except three due to drawing

issue, LPV plasma concentrations regardless assays timing were unexpectedly high comparing

HIV experience. Finally, unbound concentrations, which are the active ones, were not estimated.

In conclusion, in our single center Covid-19 experience, we found that LPV/r treated patients had

unexpected very high plasma concentrations, well above those observed in HIV patients.

However, LPV dosage should not be reduced in the absence of adverse effect, owing that to high

plasma concentration may be required for an *in vivo* antiviral activity as suggested by previous

studies. We observed that about one over four to five patients stopped LPV/r therapy in relation

with moderate adverse drug reactions attributed to LPV therapy. Cautious is needed in this

context of drug repurposing, which can be associated with a new drug safety profile. These early

data need to be confirmed with prospective studies.

MATERIALS AND METHODS

Study population

We used data collected from routine care of patients hospitalized for Covid-19 in a single center setting (Cochin Hospital, Paris). Patients hospitalized in medicine wards and starting a treatment by LPV/r for Covid-19 pneumonitis between March 18[th] and April 1[st] have been included in the cohort. Covid-19 pneumonia was considered in the presence of CT scan injuries evocating a SARS-CoV-2 infection and in the absence of other diagnosis.(29) Covid-19 patients were considered as confirmed or suspected if they had a positive or negative PCR, respectively. Patients with ARDS (defined by the need for an invasive or non-invasive mechanical ventilation) at admission were not included.(30) Data regarding patient demographics, type of Covid-19 infection, patient outcome and LPV/r clinical and biological safety were collected. Patients have been identified according to LPV/r dispensing by the hospital pharmacy. The study has been performed in accordance with the declaration of Helsinki and received approval by the Cochin Hospital Institutional Review Board (number 2020-08019).

Treatment strategy

According to our local protocol at the time of the study, patients were eligible to receive a LPV/r treatment as specific anti-Covid-19 therapy if they (i) had a pneumonitis evocating a SARS-CoV-2 infection at CT scan and (ii) were requiring oxygen (minimal flow rate 2-3 liters per minute) . LPV/r was started within 24-48 hours after admission. LPV/r therapy consisted in a dose of 400/100 mg twice a day in tablets during seven to ten days, according to ward's practices. Drug-drug interactions were carefully monitored before initiation and during LPV/r treatment. Patients

were follow-up until the end of LPV/r treatment. For each patient, a therapeutic drug monitoring

was performed in a routine care setting within the first three days of therapy.

LPV and ritonavir assays

Plasma assays were drawn in a routine care setting, according to ward's practices. Plasma

concentrations of LPV and ritonavir were quantified using high-performance liquid

chromatography tandem mass spectrometry (Xevo TQD, Waters®) by the department of clinical

pharmacology (Cochin Hospital, Paris, France). A volume of 100 μL of plasma sample was

mixed with 300 μL of methanol. Then, the mixture was centrifuged at 21,000 g during 10

minutes. A volume of 50 μL of supernatant was diluted by 50 μL of waters (0.05% formic acid,

v/v), and 10 μL was injected into the chromatographic system. The separation was carried out on

a BEH C18 analytical column (1.7μm, 1.7μm, 50*2.1 mm, Waters, Saint-Quentin, France). The

phase mobile composed of 60% of water (0.05 % formic acid, v/v) and 40% of methanol (0.05%

formic acid, v/v). The method was validated according to the EMA guideline for bioanalytical

method   validation   (https://www.ema.europa.eu/en/ich-m10-bioanalytical-method-validation).

The calibration range was from 160 (limit of quantification, LOQ) to 40,000 ng/mL for LPV and

10 (LOQ) to 2,500 ng/mL for ritonavir. The inter- and intra-assay bias and CV were inferior to

15% for LPV and ritonavir. No significant matrix effect on quantification of LPV and ritonavir

was found in this method. LPV plasma concentrations were considered as peak level (Cmax) and

as through level (Cmin) for assays drawn between one hour to five hours, and from eight hours

after tablet intake, respectively. In HIV treatment, observed Cmin and Cmax ranges are 1,000-

8,000 ng/mL (target > 4,000 ng/mL for non-naïve patients) and 7,000-11,000 ng/mL, respectively

(31, 32).

LPV/r treatment safety

278 LPV/r treatment safety was assessed in a routine care setting. Cases of suspected adverse drug

279 reaction were spontaneously notified to the Pharmacovigilance regional center. After case-by-

280 case review assessment and review by a senior pharmacologist, cases were reported to the French

281 pharmacovigilance system.

Statistics

284 Quantitative and qualitative data are presented using non-parametric settings. Descriptive

285 analyses have been performed using Prism GraphPad 7.0 and R statistical package version 3.6.1

286 or later (The R Foundation for Statistical Computing, https://www.R-project.org/).

AUTHORS' CONTRIBUTIONS

Study design: LC, SBo, IG, JMT, SBe

Data collection: LC, SBo, IG, LHP, LR, PL, CA, EC, JZ, NC, BT, SK, RB, JMT, YZ, SBe

Data analysis: LC, SBo; LHP, SBe

Draft the manuscript: LC, SBo, SBe

Critically review the manuscript: LC, SBo, EC, NC, BT, SK

Approved the manuscript: LC, SBo, IG, LHP, LR, PL, CA, EC, JZ, NC, BT, SK, RB, JMT, YZ,

SBe

CONFLICT OF INTEREST STATEMENTS

None

ROLE OF FUNDING SOURCE

None

REFERENCES

1. Lu C-C, Chen M-Y, Chang Y-L. 2020. Potential therapeutic agents against COVID-19: What we know so far. J Chin Med Assoc JCMA.

2. Chu CM, Cheng VCC, Hung IFN, Wong MML, Chan KH, Chan KS, Kao RYT, Poon LLM, Wong CLP, Guan Y, Peiris JSM, Yuen KY, HKU/UCH SARS Study Group. 2004. Role of lopinavir/ritonavir in the treatment of SARS: initial virological and clinical findings. Thorax 59:252–256.

3. Kim UJ, Won E-J, Kee S-J, Jung S-I, Jang H-C. 2016. Combination therapy with lopinavir/ritonavir, ribavirin and interferon-α for Middle East respiratory syndrome. Antivir Ther 21:455–459.

4. Chan JF-W, Yao Y, Yeung M-L, Deng W, Bao L, Jia L, Li F, Xiao C, Gao H, Yu P, Cai J-P, Chu H, Zhou J, Chen H, Qin C, Yuen K-Y. 2015. Treatment With Lopinavir/Ritonavir or Interferon-β1b Improves Outcome of MERS-CoV Infection in a Nonhuman Primate Model of Common Marmoset. J Infect Dis 212:1904–1913.

5. de Wilde AH, Jochmans D, Posthuma CC, Zevenhoven-Dobbe JC, van Nieuwkoop S, Bestebroer TM, van den Hoogen BG, Neyts J, Snijder EJ. 2014. Screening of an FDA-approved compound library identifies four small-molecule inhibitors of Middle East respiratory syndrome coronavirus replication in cell culture. Antimicrob Agents Chemother 58:4875–4884.

6. Choy K-T, Yin-Lam Wong A, Kaewpreedee P, Sia S-F, Chen D, Yan Hui KP, Wing Chu DK, Wai Chan MC, Pak-Hang Cheung P, Huang X, Peiris M, Yen H-L. 2020. Remdesivir,

lopinavir, emetine, and homoharringtonine inhibit SARS-CoV-2 replication in vitro.

Antiviral Res 104786.

7. Cao B, Wang Y, Wen D, Liu W, Wang J, Fan G, Ruan L, Song B, Cai Y, Wei M, Li X, Xia

J, Chen N, Xiang J, Yu T, Bai T, Xie X, Zhang L, Li C, Yuan Y, Chen H, Li H, Huang H,

Tu S, Gong F, Liu Y, Wei Y, Dong C, Zhou F, Gu X, Xu J, Liu Z, Zhang Y, Li H, Shang L,

Wang K, Li K, Zhou X, Dong X, Qu Z, Lu S, Hu X, Ruan S, Luo S, Wu J, Peng L, Cheng

F, Pan L, Zou J, Jia C, Wang J, Liu X, Wang S, Wu X, Ge Q, He J, Zhan H, Qiu F, Guo L,

Huang C, Jaki T, Hayden FG, Horby PW, Zhang D, Wang C. 2020. A Trial of Lopinavir-

Ritonavir in Adults Hospitalized with Severe Covid-19. N Engl J Med.

8.   Search of: COVID | lopinavir - List Results - ClinicalTrials.gov.

9. Guan W-J, Ni Z-Y, Hu Y, Liang W-H, Ou C-Q, He J-X, Liu L, Shan H, Lei C-L, Hui DSC,

Du B, Li L-J, Zeng G, Yuen K-Y, Chen R-C, Tang C-L, Wang T, Chen P-Y, Xiang J, Li S-

Y, Wang J-L, Liang Z-J, Peng Y-X, Wei L, Liu Y, Hu Y-H, Peng P, Wang J-M, Liu J-Y,

Chen Z, Li G, Zheng Z-J, Qiu S-Q, Luo J, Ye C-J, Zhu S-Y, Zhong N-S, China Medical

Treatment Expert Group for Covid-19. 2020. Clinical Characteristics of Coronavirus

Disease 2019 in China. N Engl J Med.

10. Smits SL, de Lang A, van den Brand JMA, Leijten LM, van IJcken WF, Eijkemans MJC,

van Amerongen G, Kuiken T, Andeweg AC, Osterhaus ADME, Haagmans BL. 2010.

Exacerbated innate host response to SARS-CoV in aged non-human primates. PLoS Pathog

6:e1000756.

11. Hadjadj J, Yatim N, Barnabei L, Corneau A, Boussier J, Pere H, Charbit B, Bondet V,

Chenevier-Gobeaux C, Breillat P, Carlier N, Gauzit R, Morbieu C, Pene F, Marin N, Roche

N, Szwebel T-A, Smith N, Merkling S, Treluyer J-M, Veyer D, Mouthon L, Blanc C,

Tharaux P-L, Rozenberg F, Fischer A, Duffy D, Rieux-Laucat F, Kerneis S, Terrier B.

2020. Impaired type I interferon activity and exacerbated inflammatory responses in severe

Covid-19 patients. medRxiv 2020.04.19.20068015.

12. Morgan ET. 2009. Impact of infectious and inflammatory disease on cytochrome P450-

mediated drug metabolism and pharmacokinetics. Clin Pharmacol Ther 85:434–438.

13. Aitken AE, Morgan ET. 2007. Gene-specific effects of inflammatory cytokines on

cytochrome P450 2C, 2B6 and 3A4 mRNA levels in human hepatocytes. Drug Metab

Dispos Biol Fate Chem 35:1687–1693.

14. Dickmann LJ, Patel SK, Rock DA, Wienkers LC, Slatter JG. 2011. Effects of interleukin-6

(IL-6) and an anti-IL-6 monoclonal antibody on drug-metabolizing enzymes in human

hepatocyte culture. Drug Metab Dispos Biol Fate Chem 39:1415–1422.

15. Lee JH, Cho YK, Jung YS, Kim YC, Lee MG. 2008. Effects of Escherichia coli

lipopolysaccharide on telithromycin pharmacokinetics in rats: inhibition of metabolism via

CYP3A. Antimicrob Agents Chemother 52:1046–1051.

16. Maezono S, Sugimoto K, Sakamoto K, Ohmori M, Hishikawa S, Mizuta K, Kawarasaki H,

Watanabe Y, Fujimura A. 2005. Elevated blood concentrations of calcineurin inhibitors

during diarrheal episode in pediatric liver transplant recipients: involvement of the

suppression of intestinal cytochrome P450 3A and P-glycoprotein. Pediatr Transplant

9:315–323.

17. Gu X, Ke S, Liu D, Sheng T, Thomas PE, Rabson AB, Gallo MA, Xie W, Tian Y. 2006.

Role of NF-kappaB in regulation of PXR-mediated gene expression: a mechanism for the

suppression of cytochrome P-450 3A4 by proinflammatory agents. J Biol Chem 281:17882–

17889.

18. Okamura M, Shizu R, Hosaka T, Sasaki T, Yoshinari K. 2019. Possible involvement of the

competition for the transcriptional coactivator glucocorticoid receptor-interacting protein 1

in the inflammatory signal-dependent suppression of PXR-mediated CYP3A induction

in vitro. Drug Metab Pharmacokinet 34:272–279.

19. Sham HL, Kempf DJ, Molla A, Marsh KC, Kumar GN, Chen CM, Kati W, Stewart K, Lal

R, Hsu A, Betebenner D, Korneyeva M, Vasavanonda S, McDonald E, Saldivar A,

Wideburg N, Chen X, Niu P, Park C, Jayanti V, Grabowski B, Granneman GR, Sun E,

Japour AJ, Leonard JM, Plattner JJ, Norbeck DW. 1998. ABT-378, a highly potent inhibitor

of the human immunodeficiency virus protease. Antimicrob Agents Chemother 42:3218–

3224.

20. 2012. LopinavirLiverTox: Clinical and Research Information on Drug-Induced Liver

Injury. National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda (MD).

21. Kanters S, Socias ME, Paton NI, Vitoria M, Doherty M, Ayers D, Popoff E, Chan K,

Cooper DA, Wiens MO, Calmy A, Ford N, Nsanzimana S, Mills EJ. 2017. Comparative

efficacy and safety of second-line antiretroviral therapy for treatment of HIV/AIDS: a

systematic review and network meta-analysis. Lancet HIV 4:e433–e441.

22. Aboud M, Kaplan R, Lombaard J, Zhang F, Hidalgo JA, Mamedova E, Losso MH,

Chetchotisakd P, Brites C, Sievers J, Brown D, Hopking J, Underwood M, Nascimento MC,

Punekar Y, Gartland M, Smith K. 2019. Dolutegravir versus ritonavir-boosted lopinavir

both with dual nucleoside reverse transcriptase inhibitor therapy in adults with HIV-1

infection in whom first-line therapy has failed (DAWNING): an open-label, non-inferiority,

phase 3b trial. Lancet Infect Dis 19:253–264.

23. Meraviglia P, Schiavini M, Castagna A, Viganò P, Bini T, Landonio S, Danise A, Moioli

MC, Angeli E, Bongiovanni M, Hasson H, Duca P, Cargnel A. 2004. Lopinavir/ritonavir

treatment in HIV antiretroviral-experienced patients: evaluation of risk factors for liver

enzyme elevation. HIV Med 5:334–343.

24. Pushpakom S, Iorio F, Eyers PA, Escott KJ, Hopper S, Wells A, Doig A, Guilliams T,

Latimer J, McNamee C, Norris A, Sanseau P, Cavalla D, Pirmohamed M. 2019. Drug

repurposing: progress, challenges and recommendations. Nat Rev Drug Discov 18:41–58.

25. Ferner RE, Aronson JK. 2020. Chloroquine and hydroxychloroquine in covid-19. BMJ

369:m1432.

26. Sheahan TP, Sims AC, Leist SR, Schäfer A, Won J, Brown AJ, Montgomery SA, Hogg A,

Babusis D, Clarke MO, Spahn JE, Bauer L, Sellers S, Porter D, Feng JY, Cihlar T, Jordan

R, Denison MR, Baric RS. 2020. Comparative therapeutic efficacy of remdesivir and

combination lopinavir, ritonavir, and interferon beta against MERS-CoV. Nat Commun

11:222.

27. Smith PF, Dodds M, Bentley D, Yeo K, Rayner C. 2020. Dosing will be a key success

factor in repurposing antivirals for COVID-19. Br J Clin Pharmacol.

28. Wu C-Y, Jan J-T, Ma S-H, Kuo C-J, Juan H-F, Cheng Y-SE, Hsu H-H, Huang H-C, Wu D,

Brik A, Liang F-S, Liu R-S, Fang J-M, Chen S-T, Liang P-H, Wong C-H. 2004. Small

molecules targeting severe acute respiratory syndrome human coronavirus. Proc Natl Acad

Sci U S A 101:10012–10017.

29. Shi H, Han X, Jiang N, Cao Y, Alwalid O, Gu J, Fan Y, Zheng C. 2020. Radiological

findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study.

Lancet Infect Dis 20:425–434.

30. ARDS Definition Task Force, Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND,

Caldwell E, Fan E, Camporota L, Slutsky AS. 2012. Acute respiratory distress syndrome:

the Berlin Definition. JAMA 307:2526–2533.

31. Morlat P. 2019. Prise en charge du VIH - Recommandations du groupe d'experts. France.

32. Breilh D, Pellegrin I, Rouzés A, Berthoin K, Xuereb F, Budzinski H, Munck M, Fleury

HJA, Saux M-C, Pellegrin J-L. 2004. Virological, intracellular and plasma pharmacological

parameters predicting response to lopinavir/ritonavir (KALEPHAR study). AIDS Lond Engl

18:1305–1310.

TABLES

Table 1. Demographic and clinical characteristics of the patients at baseline

Patient characteristics	
Age – year	63 (51-78)
Sex – male:female sex	22:9
Comorbidities	
Hypertension	11 (35%)
Diabetes	8 (26%)
Cardiovascular diseases (others)	7 (23%)
Malignancy or immunosuppression	6 (19%)
Chronic respiratory disease (including asthma)	3 (10%)
Hepatitis or liver cirrhosis (Child-Pugh B or more)	2 (6%)
Rheumatic disease	2 (6%)
Chronic kidney failure	1 (3%)
None	6 (19%)
SARS-CoV-2 PCR	
- Positive	25 (80%)
- Doubtful*	2 (6%)
- Negative*	2 (6%)
- N/A	2 (6%)
Extent of pneumonia at CT scan at admission	
- Minor (less than 10%)	1 (3%)
- Moderate (between 10-25%)	9 (29%)
- Extensive (between 25-50%)	12 (39%)
- Severe (more than 50%)	7 (23%)
- N/A	2 (6%)
Oxygen saturation at admission (without oxygen) – %	92.5 (90-96)
C-reactive protein (CRP) at admission – mg/L	94.1 (45.4-176.0)
Interleukin-6 (IL-6) at admission – ng/mL	60.4 (29.7-164.5)
Type of antibiotics therapy at patient admission	
- None	21 (68%)
- Cefotaxime and rovamycin	5 (16%)
- Macrolide alone**	3 (10%)
- Penicillin or cephalosporin alone***	2 (6%)

435 Data are presented as median (IQR) or as no (%).

436 * PCR resulting in doubtful or negative results have been repeated twice for each patient

437 ** azithromycin or rovamycin *** piperacillin/tazobactam or cefotaxime

Table 2. Lopinavir/ritonavir therapy characteristics

Lopinavir/ritonavir therapy characteristics	
Time from symptom onset to starting lopinavir/ritonavir – days	8 (7-10)
Type of specific anti-COVID-19 drug associated with lopinavir/ritonavir	
- None	13 (42%)
- Cephalosporin or penicillin*	6 (19%)
- Cephalosporin* and macrolide*	3 (10%)
- Macrolide*	3 (10%)
- Corticosteroids	2 (6%)
- Corticosteroids and antibiotics	2 (6%)
- Sarilumab (anti-IL-6)	2 (6%)
Duration of lopinavir/ritonavir therapy – days	7 (3-8)
Lopinavir plasma concentration– ng/mL	
- at trough	20,153 (IQR, 16,633-26,505)
- at peak	22,550 (IQR 12,720-33,612)
Lopinavir plasma concentration – fold increase**	4.6 (IQR 2.9-6.4)
Reasons for therapy termination	
- Scheduled end of treatment	17 (52%)
- Adverse drug reaction	7 (22%)
- Therapeutic limitation or deceased	4 (13%)
- Poor efficacy	3 (10%)
Types of adverse drug reaction	
- Cytolytic hepatitis	3
- Isolated hyperbilirubinemia	1
- Nausea and vomiting	1
- Diarrhea	1
- Agitation/anxiety	1
Clinical outcome at the end of lopinavir/ritonavir therapy	
- Hospitalized with oxygen dependency	17
- Recovered	5
- Transferred to intensive care unit	4
- Deceased	4
- Unknown***	1

* penicillin: piperacillin/tazobactam; cephalosporin: cefotaxime; macrolide: azithromycin or

rovamycin

** fold increase with regards of average plasma concentration observed in HIV patients.

*** one patient being transferred in another hospital during lopinavir/ritonavir time course.

 FIGURE LEGENDS

Figure 1. Lopinavir/ritonavir plasma concentration and magnitude lopinavir plasma

concentration increase in COVID-19 patients compared to HIV patients.

A. Lopinavir (left) and ritonavir (right) plasma concentrations in COVID-19 patients.

Horizontal red line represents the peak and residual concentrations observed in HIV patients after

400 mg/100 mg lopinavir/ritonavir twice daily.

Lopinavir plasma concentration observed in HIV patients were 1,000-8,000 ng/mL and 7,000-

11,000 ng/mL, respectively. Ritonavir plasma concentrations observed in HIV patients were 100-

250 ng/mL and 300-500 ng/mL, respectively.

B. Magnitude of lopinavir plasma concentration increase compared to average plasma

concentration observed in HIV patients.

Average plasma concentrations were 4,000 and 9,000 ng/mL for trough and peak, respectively.

A

B