TITLE: Neural Network Predicts Need for Red Blood Cell Transfusion for Patients with Acute Gastrointestinal Bleeding Admitted to the Intensive Care Unit

AUTHORS: Dennis Shung[1]*, Egbert Castro[2]*, Jessie Huang[3], J. Kenneth Tay[5], Michael Simonov[1], Loren Laine[1,4], Smita Krishnaswamy[3,6]

*Both are first authors

AFFILIATIONS:
[1] Yale School of Medicine, New Haven, Connecticut

[2] Yale University, Computational Biology and Bioinformatics, New Haven, Connecticut

[3] Yale University, Department of Computer Science, New Haven, Connecticut

[4] VA Connecticut Healthcare System, West Haven, Connecticut

[5] Stanford University, Palo Alto, California

[6] Yale University, Department of Genetics, New Haven, Connecticut

Funding: DS is supported by NIH grant T32 DK007017.

(i) Guarantor of the article: Dennis Shung, M.D.

(ii) Specific author contributions: DS was responsible for designing the study, performing data

extraction and analysis, and writing the manuscript. EC co-designed the study, performed data

extraction, data analysis, and revised the manuscript. JH performed data analysis and provided

revisions to the manuscript. JKT assisted with statistical analysis and comparison and revised

the manuscript. MS assisted with data extraction and revised the manuscript. LL revised the

manuscript and provided guidance on study design. SK provided guidance on study design,

advised on strategies for data preparation and analysis, and revised the manuscript.

All authors have approved the final version of the manuscript.

(iii) All authors have approved the final version of the manuscript.

Data Access: All clinical data from MIMIC-III was approved under the oversight of the

Institutional Review Boards of Beth Israel Deaconess Medical Center (Boston, MA) and the

Massachusetts Institute of Technology (Cambridge, MA). Requirement for individual patient

consent was waived because the project did not impact clinical care and all protected health information was deidentified. The data was available on PhysioNet were derived from protected health information that has been de-identified and not subject to HIPAA Privacy Rule restrictions. All use of the data was performed with credentialed access under the oversight of the data use agreement through PhysioNet and the Massachusetts Institute of Technology.

Abbreviations: Recurrent Neural Network (RNN), Long-Short Term Memory (LSTM), Electronic Health Record (EHR), Medical Information Mart for Intensive Care III (MIMIC-III), Area Under the Receiver Operating Curve (AUROC), Glasgow-Blatchford Score (GBS)

Structured Summary

Background:

Acute gastrointestinal bleeding is the most common gastrointestinal cause for hospitalization. For high risk patients requiring intensive care unit stay, predicting transfusion needs during the first 24 hours using dynamic risk assessment may improve resuscitation.

Aims:

Provide dynamic risk prediction for red blood cell transfusion in admitted patients with severe acute gastrointestinal bleeding.

Methods:

A patient cohort admitted for acute gastrointestinal bleeding (N = 2,524) was identified from the Medical Information Mart for Intensive Care III (MIMIC-III) critical care database, separated into training (N = 2,032) and validation (N = 492) sets. 74 demographic, clinical, and laboratory test features were consolidated into 4-hour time intervals over the first 24 hours from admission. The outcome measure was the transfusion of red blood cells during each 4-hour time interval. A long short-term memory (LSTM) model, a type of Recurrent Neural Network (RNN), was compared to the Glasgow-Blatchford Score (GBS).

Results:

The LSTM model performed better than GBS overall (AUROC 0.81 vs 0.63;P<0.001$)$ and at each 4-hour interval (P<0.01). At high sensitivity and high specificity cutoffs, the LSTM model outperformed GBS (P<0.001). The LSTM model performed better in patients directly admitted from the ED to ICU (0.82 vs 0.63;P<0.001), upper GIB (0.84 vs 0.68;P<0.001), lower GIB (0.77 vs 0.58;P<0.001), and unspecified GIB (0.85 vs 0.64;P<0.001).

Conclusions:

A LSTM model can be used to predict the need for transfusion of packed red blood cells over the first 24 hours from admission to help personalize the care of high-risk patients with acute gastrointestinal bleeding.

Keywords: Outcomes Research, Non-Variceal Hemorrhage, Varices, Endoscopy

Introduction:

Acute gastrointestinal bleeding accounts for over 2.2 million hospital days and 19.2 billion dollars of medical charges annually in the United States and frequently requires red-blood cell transfusion.[1] The management of severe acute gastrointestinal bleeding begins with resuscitation using intravenous fluids and transfusion of packed red blood cells, which are given to 43% of patients hospitalized with upper gastrointestinal bleeding in the United Kingdom and 21% of patients hospitalized with lower gastrointestinal bleeding in the United States.[2,3]

Transfusion needs may change during the hospital stay, but a tool to dynamically predict transfusion needs over time does not yet exist in clinical care. Patients with severe acute gastrointestinal bleeding who require care in the intensive care setting generally have higher transfusion needs and may benefit most from a predictive tool to guide resuscitation efforts. Current guidelines are based on a restrictive transfusion strategy using a hemoglobin threshold for transfusion of 7g per deciliter, which has been associated with a significant 45% relative risk reduction in 45-day mortality as compared to a threshold of 9g per deciliter in patients with upper gastrointestinal bleeding.[4]

Dynamic risk prediction, where predictions are generated in real time every hour based on clinical and laboratory values, may help guide resuscitation strategies and help in timing endoscopic intervention, particularly in severely ill patients who require intensive care. Existing clinical risk scores used to screen for risk of needing transfusion of packed red blood cells, such as the Glasgow-Blatchford Score, are static models that only use clinical information at the time of admission (e.g. initial systolic blood pressure).[5] Machine learning approaches to model risk for gastrointestinal bleeding have shown promise in outperforming existing clinical risk scores, but are also static models.[6,7] Electronic health records (EHRs) can capture clinical data in real

time, and have been used to create automated tools to model adverse events, such as sepsis, post-operative complications, and acute kidney injury.[8-11] Neural networks, computing systems inspired by the human brain, consisting of collections of layered neurons that process and transmit information to the next layer of neurons, are a transformative machine learning model that have the potential to dynamically predict the need for blood transfusion for patients over time.[12] Recurrent neural networks have been demonstrated to be better than state-of-the-art risk models for continuous prediction of acute kidney injury up to 48 hours, the onset of septic shock 28 hours before onset, and all-cause inpatient mortality.[13-15] We propose the use of a Long-Short-Term Memory (LSTM) Network, an advanced recurrent neural network that learns what to store in its memory, to process data from electronic health records with an internal memory that stores relevant information over time and can generate a probability of transfusion within the 4 hour intervals for patients with severe acute gastrointestinal bleeding. Figure 1 shows the use of our model in an example patient from the MIMIC test set with generated risk predictions throughout the first 24 hours from admission. (Figure 1)

Methods:

Data Source:

A patient cohort presenting with acute gastrointestinal bleeding was identified from the Medical Information Mart for Intensive Care III (MIMIC-III) critical care database.[16,17] The database contains data for over 40,000 patients in the Beth Israel Deaconess Medical Center from 2001 to 2012 requiring an ICU stay.

Patients were included if they had an admission diagnosis containing the terms "gastro", "bleed", "melena", "hematochezia". The diagnoses were collated and then manually reviewed. Patients were excluded if vital signs were only available greater than 24 hours from time of admission. The data included information that was updated over time during the course of hospitalization, including laboratory results and vital signs. For laboratory values, any negative

entry or non-quantizable (e.g. >=, <) was converted to missing. Medications, current procedural terminology codes, and ICD9/10 codes from the visit were excluded from the analysis. The dataset had a total of 74 features: 5 clinical and demographic variables and 69 laboratory variables.

Study Design

The final cohort included 2,524 hospital admissions and was randomly split into a training set with 2,032 hospital admissions and a validation set with 492 hospital admissions. (Table 1) We chose to compare the model to the Glasgow-Blatchford Score, which was designed to predict need for transfusion of packed red blood cells or intervention to control bleeding, rebleeding, or dying.[5] A Glasgow-Blatchford Score was calculated through manual review of Admission Notes extracted from MIMIC-III for all patients in the validation set. The performance of the neural network model and the GBS were compared by applying the same GBS for each 4-hour time interval for the first 24 hours and comparing the neural network output at each 4-hour interval.

Comparison of High Sensitivity and High Specificity Thresholds

Pre-defined cutoff of >90% was defined for sensitivity and specificity of the LSTM neural network model and the Glasgow Blatchford Score. A high sensitivity threshold means that the algorithm would not miss time periods when patients need packed red blood cell transfusions. A high specificity threshold means that the algorithm would only indicate need for packed red blood cell transfusion during a time period where the patient actually needs one, minimizing false alerts. The specific high sensitivity and high specificity threshold for the Glasgow Blatchford Score was matched to the corresponding cutoff for the LSTM Neural Network.

Sensitivity Analysis

Patients who are admitted directly from the emergency department to the intensive care unit may be different from patients who are transferred to the ICU from an inpatient ward. We applied our model to a validation set with only patients who were admitted to the ICU directly from the emergency department, as defined by arrival to ICU less than 1 hour from leaving the

emergency department. We also perform this sensitivity analysis on subsets of patients presenting with upper GIB, lower GIB, and GIB of unspecified source.

Input Variables:

A total of 74 input variables were used and included age, gender, vital signs (systolic blood pressure, diastolic blood pressure, heart rate), and 69 unique laboratory values. (Table 2) The vital signs and laboratory values were extracted and then consolidated into 4-hour time intervals over the first 24 hours from admission.

Outcome Variable

The predicted outcome measure was the transfusion of packed red blood cells, calculated as binary 0 (no transfusion) or 1 (transfusion given). At the beginning of each 4-hour time interval, the model makes a prediction on whether a transfusion will be needed by the end of the 4-hour interval.

Data Pre-Processing

Each patient was represented by a sequence of events with each 4-hour period containing information recorded in the vitals and laboratory values. Information for each patient was encoded into 4-hour time intervals up to the first 24 hours. After excluding lab values with greater than 90% missingness, remaining lab values with greater than 50% missingness in the dataset were converted to missing indicator variables, with 1 as present and 0 as missing. To harmonize the input variables across patients, the first timepoint for each patient was fixed at the first recording of heart rate, systolic blood pressure, and diastolic blood pressure. Consolidation of vital signs and laboratory values in each 4-hour interval was performed by taking the mean of each value. All continuous values were normalized and centered. Age was maintained as a continuous variable, with patients greater than 89 years old coded as 89 years old.

Missing Values

To examine the role of the data imputation method used, we compared 4 different imputation strategies. The first was imputation of the mean value of for any missing value. The second was a carryforward approach, or using the previously recorded value if a value was present at a previous time point but no subsequent value was measured. This assumes that the laboratory value is constant until the next time point in clinical decision-making.[18] The third was mean imputation with a new variable that served as a missingness indicator for every variable. The fourth was carryforward with a missingness indicator for every variable.

Model background

Recurrent neural networks allow for processing of sequential information by storing information as internal states over multiple time points. Long short-term memory (LSTM) networks are a type of RNN that can be useful for clinical measurements because they carefully tune the information passed between subsequent time-iterations of the model. Given a series of EHR data, $x^{(0)}, x^{(1)}, ..., x^{(T-1)}$, where $x^{(t)}$ represents the input variables for the $(t+1)$th 4-hour interval, at the beginning of each 4-hour interval our goal is to predict whether transfusion is needed in the next 4 hours. The output is a sequence of probability predictions $\widehat{y^{(1)}}, \widehat{y^{(2)}}, ..., \widehat{y^T}$, where $\widehat{y^{(t)}} \in [0,1]$ is the prediction for whether transfusion is needed in the tth 4-hour interval. The LSTM model consists of 2 layers of 128 LSTM cells each, followed with a linear layer that maps from hidden state space to the prediction space. We obtain the log-probabilities by adding a LogSoftmax later in the last layer of the network. Thus the output of the neural network is a sequence $\widehat{p^{(1)}}, \widehat{p^{(2)}}, ..., \widehat{p^{(T)}}$, where $\widehat{p^{(t)}}$ is the log-probability of $\hat{y}$ being either of the target classes, and our decision rule is to administer transfusion if $\widehat{p^{(t)}} > threshold$, where the threshold is determined by desired sensitivity or specificity. We use the negative log likelihood for the output at each time of interest as the loss function. The model is trained for up to 100 epochs with hyperparameters corresponding to the lowest validation loss recorded and used to obtain testing accuracy.

Statistical Analysis

Two-tailed t tests and chi-squared test were used to compare baseline characteristics between the training and validation sets. We assessed model performance using the area under the curve (AUROC) and compared it to the performance of GBS using the nonparametric DeLong test.[19] McNemar's test was used to compare specificity and sensitivity at specified cutoffs between the two risk assessment models.

Results

Demographics were similar between training and validation sets with the median age 69 for both, proportion of men (41% in training, 39% on validation), and predominantly white (70% in training, 77% in validation). Both sets had similar percentage of patients with upper gastrointestinal bleeding (training 33% vs validation 41%). The training set had more patients with gastrointestinal bleeding and unspecified source (46% vs 26% P<0.01), while the validation set had more patients with lower gastrointestinal bleeding (33% vs 21% P=0.02). Vital signs and laboratory values were similar in the training and validation sets. (Table 1)

The performance of the LSTM model on the four different imputation strategies were similar and all significantly better than GBS. (Table 5) The results we subsequently present are for the strategy with the highest AUROC (mean imputation and missing indicators). For the main analysis of all patients with acute gastrointestinal bleeding who were transferred to the ICU, the LSTM performed significantly better than GBS in predicting packed red blood cell transfusion across the entire 24 hour period (AUROC 0.81 CI 0.80-0.83 vs 0.63 CI 0.61-0.65;P<0.001). The model also consistently outperformed GBS in each 4-hour interval. (Table 3, Figure 2) At the cutoff for high specificity (>90%), the LSTM performed better than GBS (LSTM sensitivity 42% versus GBS sensitivity 12%;P<0.001). At the cutoff for high sensitivity (>90%), the LSTM performed better than GBS (LSTM sensitivity 37% versus GBS sensitivity 15%;P<0.001). (Table 6)

For the sensitivity analysis of only patients admitted directly from the ED to the ICU, the LSTM network was also significantly better than GBS (AUROC 0.82 CI 0.80-0.83 vs. 0.63 CI 0.61-0.66, $P<0.001$). The LSTM network was also significantly better than GBS for upper GIB (0.84 CI 0.81-0.86 vs 0.68 CI 0.64-0.71;$P<0.001$), lower GIB (0.77 CI 0.74-0.80 vs 0.58 CI 0.54-0.61;$P<0.001$), and GIB of unspecified origin (0.85 CI 0.82-0.88 vs 0.64 CI 0.59-0.68;$P<0.001$). (Table 4) There was no clear difference in performance as measured by AUROC in patients who were directly admitted to the ICU from the ED.

Discussion

Predicting the need for transfusion of packed red blood cells has direct relevance to guiding the management of patients with acute gastrointestinal bleeding. This is the first study to show that a LSTM network model is able to predict the need for packed red blood cell transfusion for patients with severe acute gastrointestinal bleeding. By anticipating needs for transfusion, this is a first step towards personalizing treatment and tailoring appropriate resuscitation to reduce clinical decompensation and death for patients with severe acute gastrointestinal bleeding. While endoscopic evaluation is important, adequate resuscitation is the cornerstone of management prior to endoscopy.[20-23]

Previous risk scores capture information from specific points in time at admission, and do not incorporate new clinical data over the course of hospitalization. Electronic health records contain longitudinal information on patients admitted to the hospital and reflect real-world practice, which can be used to develop risk prediction models.[24] For patients who have severe disease requiring intensive care unit stay, mortality may be more due to end organ damage due to inadequate perfusion.[3,25,26] Despite the significant computing requirements necessary to run neural networks, existing electronic health records are now deploying cloud computing infrastructure able to perform computationally intensive tasks. The emerging capabilities of

cloud infrastructure in electronic health records, such as the Cognitive Computing platform for Epic Systems, make the deployment of neural networks for clinical care feasible.

We envision the future of care for all patients to be enhanced by customized machine learning decision support tools that will provide both initial risk stratification and ongoing risk assessment to provide treatment at the right time for the right patient. Using a dynamic risk assessment, resuscitation needs could be estimated early and optimized in preparation for endoscopic evaluation and intervention. This individualized decision-making potentially will minimize organ damage from inadequate resuscitation, which drives the risk for mortality in these patients.[25] The LSTM model can be tuned for provider preference with either a high sensitivity threshold or a high specificity threshold. Alert fatigue is particularly relevant in the ICU, since clinically irrelevant alerts can have an impact on patient safety.[27] In order to minimize alert fatigue, a high specificity threshold could be set for the algorithm. However, if providers do not want to miss any time periods when patients need packed red blood cell transfusions, a high sensitivity threshold can be set to minimize false negatives. Although the LSTM network model is much better than a currently used tool (GBS), it still falls short of optimal performance. More work will be needed to develop and validate neural network models.

Strengths of this study include modeling patients with severe illness requiring intensive care unit stay, which may benefit disproportionately from timely transfusion and resuscitation and the use of vital signs and laboratory tests that are standardized and can be easily mapped across electronic health record systems.

Limitations include the absence of prospective and independent validation in other datasets and proportion of missing data that required imputation methods, which may introduce bias to the data. Furthermore, our comparison to Glasgow-Blatchford Score may be suboptimal, since we

studied patients with any overt gastrointestinal bleeding regardless of source, but the GBS was primarily developed for patients with upper gastrointestinal bleeding.

In summary, we present the first application of recurrent neural networks to dynamically predict need for packed red blood cell transfusion over time using electronic health record data. We report superior performance compared to the Glasgow Blatchford Score. Our approach may lead to delivery of earlier resuscitation with packed red blood cells to minimize ischemic end organ damage in patients with severe acute gastrointestinal bleeding. Future directions include external validation of the model on other cohorts of high-risk patients with gastrointestinal bleeding, along with prospective implementation and deployment in the electronic health record system for high risk patients with gastrointestinal bleeding.

Table 1: Demographics and Baseline Data for the Training and Validation Set

	Training Set N = 2,032		Validation Set N = 492		
	N	Proportion	N	Proportion	p-value
Demographic Information					
Male	836	41%	190	39%	0.31
Age					
>89	144	7%	42	9%	0.29
75-89	629	31%	168	34%	0.24
50-75	935	46%	200	41%	0.14
25-50	316	16%	71	14%	0.43
<25	8	0%	4	1%	0.27
Ethnicity					
White	1429	70%	380	77%	0.08
African American	244	12%	52	11%	0.35
Hispanic	75	4%	22	4%	0.37
Asian American	74	4%	15	3%	0.42
Other	210	10%	23	5%	0.05
Clinical Features					
Upper Gastrointestinal Bleeding	679	33%	203	41%	0.07
Lower Gastrointestinal Bleeding	428	21%	162	33%	0.02
Unspecified Location	925	46%	127	26%	<0.01
Outcomes					
Packed Red Blood Cells	1542	76%	381	77%	0.39
In-Hospital Mortality	156	8%	32	6.5%	0.35
	Mean	Std Dev	Mean	Std Dev	p-value
Vital Signs					
Heart Rate (beats per minute)	88.9	18	88.1	16.6	0.35
Systolic Blood Pressure	126.9	22.9	127.1	22.2	0.86
Diastolic Blood Pressure	64.2	16.9	65.7	16.6	0.07
Laboratory Tests					
Alanine Aminotransferase (ALT)	41	86.9	39.8	152	0.87
Albumin	3.1	0.63	3.1	0.65	1.00
Alkaline Phosphatase	117.2	152.8	118.4	130.5	0.86
Anion Gap	15	4.3	14.8	4	0.33
Aspartate Aminotransferase (AST)	75.8	159	75	446	0.97
Bicarbonate	23.8	4.5	24	4.4	0.37
Bilirubin, Total	1.52	3	1.47	2.5	0.70

Calcium, Total	8.1	0.88	8.2	0.8	0.01
Chloride	104.4	6.1	104.4	6.5	1.00
Creatinine	1.43	1.2	1.5	1.4	0.31
Glucose	147	77.5	143.8	60.6	0.32
Magnesium	1.9	0.71	1.9	0.4	1.00
Phosphate	3.5	1.2	3.5	1.2	1.00
Potassium	4.3	0.76	4.4	0.8	0.01
Sodium	138.8	4.6	138.7	4.8	0.68
Urea Nitrogen	37.5	28	39.5	30.4	0.18
Basophils	0.39	0.98	0.34	0.34	0.06
Eosinophils	1.4	1.9	1.4	1.7	1.00
Hematocrit	28	9.5	27.7	6.5	0.41
Hemoglobin	9.5	2.5	9.3	2.4	0.10
International Normalized Ratio (INR)	1.8	2.5	1.8	1.7	1.00
Lymphocytes (%)	17.1	10.2	16.6	10.1	0.33
Mean Corpuscular Hemoglobin (MCH)	30.2	3	29.9	2.9	0.04
Mean Corpuscular Hemoglobin Concentration (MCHC)	33.5	1.7	33.3	1.8	0.03
Mean Corpuscular Volume (MCV)	90	7.6	89.7	7.3	0.42
Monocytes (%)	4.6	2.6	4.6	2.3	1.00
Neutrophils (%)	75.1	12.3	76.5	11.7	0.02
Platelet Count (x1000)	231.8	139.3	238.9	127.4	0.28
Prothrombin Time (PT)	17.1	11.4	19.1	15.7	0.01
Partial Thromboplastin Time (PTT)	31.3	13.7	31.3	14.2	1.00
Red Blood Cell Distribution Width (RDW)	16.1	2.5	15.9	2.2	0.08

Table 2: Input Variables (N = 74)

Category	Input Variables
Demographic (2)	Gender Age
Vital Signs (3)	Heart Rate Systolic Blood Pressure Diastolic Blood Pressure
Laboratory Variables (69)	Blood Gas (Chloride, Free Calcium, Glucose, Hematocrit, Hemoglobin, Lactate, Oxygen, Oxygen Saturation, pH, Arterial Pressure of Oxygen, Potassium, Sodium) White Blood Cells, Neutrophils, Basophils, Eosinophils, Lymphocytes, Monocytes, Hemoglobin, Hematocrit, Mean Corpuscular Hemoglobin, Mean Corpuscular Hemoglobin Concentration, Mean Corpuscular Volume, Red Blood Cell Distribution Width, Platelet Count, International Normalized Ratio, Prothrombin Time, Partial Thromboplastin Time Sodium, Potassium, Chloride, Bicarbonate, Anion Gap, Magnesium, Phosphate, Calcium, Creatinine, Urea Nitrogen, Glucose Alanine Aminotransferase, Aspartate Aminotransferase, Alkaline Phosphatase, Albumin, Amylase, Lipase, Direct Bilirubin, Total Bilirubin Creatine Kinase, Creatine Kinase-MB, Ferritin, Total Iron, Iron Binding Capacity, Lactate Dehydrogenase, Thyroid Stimulating Hormone, Transferrin, Troponin T, Vancomycin, Fibrinogen Urine Studies (Creatinine, Sodium, Red Blood Cells, White Blood Cells, pH, Protein, Specific Gravity, Urobilinogen)

Table 3: Performance of the Long-Short Term Memory (LSTM) Model and the Glasgow Blatchford Score in Predicting Transfusion of Packed Red Blood Cells by Comparison of Area Under the Receiver Operating Curve (AUROC) for all patients (N = 492) overall and by each 4 hour interval.

	Long-Short Term Memory Network Model Validation AUROC 95% Confidence Interval	Glasgow-Blatchford Score Validation AUROC 95% Confidence Interval	p-value
Overall	**0.81 (0.80-0.83)**	**0.63 (0.61-0.65)**	**<0.001**
Hour 4	0.79 (0.75-0.83)	0.70 (0.65-0.74)	<0.001
Hour 8	0.76 (0.72-0.80)	0.64 (0.59-0.69)	<0.001
Hour 12	0.72 (0.67-0.77)	0.62 (0.56-0.67)	<0.001
Hour 16	0.73 (0.68-0.79)	0.65 (0.59-0.71)	0.005
Hour 20	0.78 (0.72-0.84)	0.62 (0.55-0.69)	<0.001
Hour 24	0.78 (0.72-0.84)	0.60 (0.52-0.69)	<0.001

Table 4: Sensitivity Analyses of Overall Performance of the Long-Short Term Memory (LSTM) Model and the Glasgow Blatchford Score in Predicting Transfusion of Packed Red Blood Cells by Comparison of Area Under the Receiver Operating Curve (AUROC)

	Long-Short Term Memory Network Model Validation AUROC 95% Confidence Interval	Glasgow-Blatchford Score Validation AUROC 95% Confidence Interval	p-value
Patients from ED to ICU (N = 412)	0.82 (0.80-0.83)	0.63 (0.61-0.66)	<0.001
Upper GIB (N = 203)	0.84 (0.81-0.86)	0.68 (0.64-0.71)	<0.001
Lower GIB (N = 162)	0.77 (0.74-0.80)	0.58 (0.54-0.61)	<0.001
GIB with Unspecified Source (N = 127)	0.85 (0.82-0.88)	0.64 (0.59-0.68)	<0.001

Table 5: Comparison of the overall performance of Long-Short Term Memory network model with different imputation methods to address missingness in the first 24 hours after admission for all patients admitted to the Intensive Care Unit with Acute Gastrointestinal Bleeding. The Glasgow-Blatchford Score was able to be calculated with no missing values at presentation.

Imputation Method	Validation Set AUROC 95% Confidence Interval	p-value (vs. Glasgow-Blatchford Score)
Glasgow-Blatchford Score	0.63 (0.58-0.67)	
Mean Imputation	0.80 (0.78-0.82)	<0.001
Carryforward Imputation	0.80 (0.79-0.82)	<0.001
Mean Imputation and Missing Indicators	0.81 (0.80-0.83)	<0.001
Carryforward Imputation and Missing Indicators	0.81 (0.79-0.83)	<0.001

Table 6: Performance of LSTM Neural Network model Compared to the Glasgow Blatchford Score at matched cutoffs of 91% sensitivity and 93% for the internal validation dataset

Risk Stratification Tool	Sensitivity	Specificity
High Specificity Cutoff		
Glasgow Blatchford Score = 14	12%	93%
LSTM Neural Network	42%	93%
High Sensitivity Cutoff		
Glasgow Blatchford Score = 5	91%	22%
LSTM Neural Network	91%	46%

Figure 1: Example of neural network prediction for the first 24 hours of a 62 year old man with Hepatitis C cirrhosis presenting with 2 days of intermittent coffee ground emesis and lethargy. Glasgow Blatchford Score = 14 a) Continuous risk prediction of the neural network through the first 24 hours with the threshold set above 0.5 for detecting need for transfusion. The arrows indicate need for transfusion during that time period. b) Measurements of Heart Rate, Systolic Blood Pressure, and Hemoglobin occurring during the first 24 hours.

Figure 2: Comparison of the overall Area Under the Receiver Operating Curve (AUROC) as a measure of performance of the Long-Short Term Memory (LSTM) Neural Network model and Glasgow-Blatchford Score (GBS).

REFERENCES:
1. Peery AF, Crockett SD, Murphy CC, et al. Burden and Cost of Gastrointestinal, Liver, and Pancreatic Diseases in the United States: Update 2018. *Gastroenterology.* 2019;156(1):254-272.e211.
2. Hearnshaw SA, Logan RF, Lowe D, Travis SP, Murphy MF, Palmer KR. Acute upper gastrointestinal bleeding in the UK: patient characteristics, diagnoses and outcomes in the 2007 UK audit. *Gut.* 2011;60(10):1327-1335.
3. Strate LL, Ayanian JZ, Kotler G, Syngal S. Risk factors for mortality in lower intestinal bleeding. *Clin Gastroenterol Hepatol.* 2008;6(9):1004-1010; quiz 1955-.
4. Villanueva C, Colomo A, Bosch A, et al. Transfusion Strategies for Acute Upper Gastrointestinal Bleeding. *New England Journal of Medicine.* 2013;368(1):11-21.
5. Blatchford O, Murray WR, Blatchford M. A risk score to predict need for treatment for uppergastrointestinal haemorrhage. *The Lancet.* 2000;356(9238):1318-1321.
6. Shung DL, Au B, Taylor RA, et al. Validation of a Machine Learning Model That Outperforms Clinical Risk Scoring Systems for Upper Gastrointestinal Bleeding. *Gastroenterology.* 2019.
7. Shung D, Simonov M, Gentry M, Au B, Laine L. Machine Learning to Predict Outcomes in Patients with Acute Gastrointestinal Bleeding: A Systematic Review. *Dig Dis Sci.* 2019;64(8):2078-2087.
8. Herasevich V, Pieper MS, Pulido J, Gajic O. Enrollment into a time sensitive clinical study in the critical care setting: results from computerized septic shock sniffer implementation. *Journal of the American Medical Informatics Association.* 2011;18(5):639-644.
9. Umscheid CA, Betesh J, VanZandbergen C, et al. Development, implementation, and impact of an automated early warning and response system for sepsis. *J Hosp Med.* 2015;10(1):26-31.
10. Mohamadlou H, Lynn-Palevsky A, Barton C, et al. Prediction of Acute Kidney Injury With a Machine Learning Algorithm Using Electronic Health Record Data. *Canadian Journal of Kidney Health and Disease.* 2018;5:2054358118776326.
11. Bihorac A, Ozrazgat-Baslanti T, Ebadi A, et al. MySurgeryRisk: Development and Validation of a Machine-learning Risk Algorithm for Major Complications and Death After Surgery. *Annals of Surgery.* 2019;269(4).
12. LeCun Y, Bengio Y, Hinton G. Deep learning. *Nature.* 2015;521(7553):436-444.
13. Tomašev N, Glorot X, Rae JW, et al. A clinically applicable approach to continuous prediction of future acute kidney injury. *Nature.* 2019;572(7767):116-119.
14. Henry KE, Hager DN, Pronovost PJ, Saria S. A targeted real-time early warning score (TREWScore) for septic shock. *Science Translational Medicine.* 2015;7(299):299ra122.
15. Rajkomar A, Oren E, Chen K, et al. Scalable and accurate deep learning with electronic health records. *npj Digital Medicine.* 2018;1(1):18.
16. Goldberger AL, Amaral LA, Glass L, et al. PhysioBank, PhysioToolkit, and PhysioNet: components of a new research resource for complex physiologic signals. *Circulation.* 2000;101(23):E215-220.
17. Johnson AEW, Pollard TJ, Shen L, et al. MIMIC-III, a freely accessible critical care database. *Scientific Data.* 2016;3(1):160035.
18. Lin Y-W, Zhou Y, Faghri F, Shaw MJ, Campbell RH. Analysis and prediction of unplanned intensive care unit readmission using recurrent neural networks with long short-term memory. *PloS one.* 2019;14(7):e0218942-e0218942.
19. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics.* 1988;44(3):837-845.

20.   Oakland K, Chadwick G, East JE, et al. Diagnosis and management of acute lower gastrointestinal bleeding: guidelines from the British Society of Gastroenterology. *Gut.* 2019;68(5):776.

21.   Laine L, Jensen DM. Management of Patients With Ulcer Bleeding. *American Journal of Gastroenterology.* 2012;107(3).

22.   Sung JJY, Chiu PWY, Chan FKL, et al. Asia-Pacific working group consensus on non-variceal upper gastrointestinal bleeding: an update 2018. *Gut.* 2018;67(10):1757.

23.   Gralnek IM, Dumonceau JM, Kuipers EJ, et al. Diagnosis and management of nonvariceal upper gastrointestinal hemorrhage: European Society of Gastrointestinal Endoscopy (ESGE) Guideline. *Endoscopy.* 2015;47(10):a1-46.

24.   Goldstein BA, Navar AM, Pencina MJ, Ioannidis JPA. Opportunities and challenges in developing risk prediction models with electronic health records data: a systematic review. *Journal of the American Medical Informatics Association.* 2016;24(1):198-208.

25.   Baradarian R, Ramdhaney S, Chapalamadugu R, et al. Early intensive resuscitation of patients with upper gastrointestinal bleeding decreases mortality. *Am J Gastroenterol.* 2004;99(4):619-622.

26.   Oakland K, Guy R, Uberoi R, et al. Acute lower GI bleeding in the UK: patient characteristics, interventions and outcomes in the first nationwide audit. *Gut.* 2018;67(4):654.

27.   Kizzier-Carnahan V, Artis KA, Mohan V, Gold JA. Frequency of Passive EHR Alerts in the ICU: Another Form of Alert Fatigue? *J Patient Saf.* 2019;15(3):246-250.

AUROC for GBS versus LSTM Neural Network