Air Pollution and Lipometabolic Disturbance: A Systematic Review and Meta-Analysis

Haohao Chen[a], Liuhua Shi[b], Andrew Rosenberg[b], Ye Zhu[e], Lixin Tao[c,d], Zhiwei Sun[e,f], Ji Wang[e,f, *]

a Eighth Clinical School, Fuxing Hospital, Capital Medical University, Beijing 100038, China

b Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory University, Atlanta Georgia 30322, United States.

c Department of Epidemiology and Health Statistics, School of Public Health, Capital Medical University, Beijing 100069, China

d Beijing Municipal Key Laboratory of Clinical Epidemiology, Capital Medical University, Beijing 100069, P.R. China

e Department of Toxicology and Sanitary Chemistry, School of Public Health, Capital Medical University, Beijing 100069, P.R. China

f Beijing Key Laboratory of Environmental Toxicology, Capital Medical University, Beijing 100069, P.R. China

* To whom correspondence should be addressed:

Ji Wang, Department of Toxicology and Sanitary Chemistry, School of Public Health, Capital Medical University, Beijing 100069, P.R. China, Tel.: (86) 10-83911776, E-mail: wangji@ccmu.edu.cn

Abstract

Background: Growing evidence suggests that long-term exposure to air pollutants is associated with cardiovascular morbidity, including lipometabolic disturbance.

Objectives: To explore the chronic effects of air pollutants on lipametabolic disturbance via detectable lipoprotein parameters.

Methods: Seven online databases were searched to conduct a meta-analysis of epidemiological studies examining the relationship between air pollution and lipid parameter levels. Subgroup analysis was additionally carried out for each air pollutant studied.

Results: A total of 2,274 records were retrieved, resulting in 10 studies included in the final quantitative meta-analysis, comprising seven studies in Europe and the United States and three studies in mainland China. Using a random-effect model, the results showed that for each 10 $\mu g/m^3$ increase in $PM_{2.5}$, TC, LDL-C, and HDL-C levels and metabolic syndrome (MetS) incidence increased by 3.31% (95% CI: 2.29%, 8.91%), 2.34% (95% CI: 1.30%, 3.39%) , -1.57% (95% CI: -1.85%, -1.28%), and 4.33% (95% CI: 2.69%, 5.98%), respectively; for each 10 $\mu g/m^3$ increase in PM_{10}, TG, HDL-C, and LDL-C levels increased by 5.27% (95% CI: 2.03%, 8.50%), -0.24% (95% CI: -0.95%, -0.47%), and 0.45% (95% CI: -0.57%, 1.47%), respectively; for each 10 $\mu g/m^3$ increase in NO_2, TG and HDL-C levels increased by 4.18% (95% CI: 1.12%, 7.23%) and -0.51% (95% CI: -2.61%, 1.58%), respectively. No significant associations were detected for combinations of air pollutants on lipometabolic disturbance.

Conclusion: Increased air pollutant exposure was significantly associated with changes in blood lipid parameter levels, which can be an indicator of the body's lipometabolic disturbance.

Keywords: air pollutant, particulate matter, lipoprotein, meta-analysis

1. Introduction

Air pollution is among the greatest environmental risks to global health(Cohen et al., 2017; Lelieveld, Evans, Fnais, Giannadaki, & Pozzer, 2015; Shi et al., 2016). The World Health Organization (WHO) estimates that ambient air pollution is responsible for nearly seven million premature deaths worldwide every year(Landrigan et al., 2018). Major components of atmospheric pollution include particulate matter of different diameter sizes, such as $PM_{2.5}$, PM_{10}, UFP (ultrafine particulate matter), and gas compounds such as CO, NO_x, SO_2, O_3, among others. Primarily deriving from fuel combustion, industrial emissions, transportation and ground dust, these pollutants can cause serious harm to the human body(Runzhi, 2020 (In China)). A growing body of evidence demonstrates the links between air pollution and cardiovascular diseases(Akintoye et al., 2016; Combes & Franchineau, 2019; Lin et al., 2019; Zheng, Liu, Zhang, & Chen, 2018), respiratory diseases(Dagenais et al., 2020; H. C. Huang et al., 2020; Schraufnagel et al., 2019; Zu et al., 2017), and neurological disorders(Myung, Lee, & Kim, 2019; Seelen et al., 2017; Thomson, 2019), among which cardiovascular disease is the leading cause of death in low-income and middle-income countries(Yusuf et al., 2020). Conventionally, risk factors for cardiovascular disease (CVD) have been associated with abnormal levels in lipoprotein-lipid parameters, including low density lipoprotein cholesterol (LDL-C), high density lipoprotein cholesterol (HDL-C), total cholesterol (TC), and triglycerides (TG), which may indicate lipometabolic disturbance (Yuanning et al., 2019 (In China)). The co-occurrence of these established risk factors, known collectively as metabolic syndrome (MetS), may therefore amplify susceptibility to the CVD risks associated with air pollution exposure(Dabass et al., 2018; Eze et al., 2015; Tsai, Lin, Chen, & Chu, 2018; Yang, Qian, et al., 2018).

The biological mechanism of exposure to air pollution and cardiovascular diseases remains

unclear. Five possible biologically plausible mechanisms(Weidong et al., 2018 (In China))

underscore the pathways through which exposure may lead to cardiovascular morbidity: first,

oxidative stress and inflammatory reactions; second, endothelial injury and dysfunction; third, an

imbalance in the function of the autonomic nervous system, fourth, changes in the function of

blood clotting; and fifth, lipid internal balance disorder. Among them, oxidative stress and

inflammatory reaction, as well as lipid internal balance disorder, most likely account for changes

in multiple lipid parameters associated with air pollution.

In recent years, studies have investigated the associations between air pollution and lipid

profile parameters, adding to evidence that suggests air pollution may contribute to changes in

blood lipid parameters(Cai et al., 2017; Hou et al., 2020; Lee et al., 2019; Vânia, Rita, Matias, &

Baltazar, 2019; Yang, Bloom, et al., 2018). However, most previous studies have been limited to

single air pollutants. To date, no systematic review has been found which offers a comprehensive

synthesis of the links between various atmospheric pollutants and lipid parameters. The aim of this

meta-analysis is to provide a comprehensive and quantitative overview of the literature regarding

the association between air pollutants ($PM_{2.5,}PM_{10,}NO_2$), lipid parameters (HDL-C, LDL-C, TC,

TG), and MetS morbidity.

2. Materials and Methods

2.1 Search Strategy

This study was conducted using standard methods which follow the Preferred Reporting

Items for Systematic Review and Meta-Analysis (PISMA)(J et al., 2020; "Preferred reporting

items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and

explanation," 2016). Seven electronic databases, including China National Knowledge

Infrastructure (CNKI), Wanfang, Vip, SinoMed, Pubmed, EMBASE, and the Cochrane Library,

were searched for peer-reviewed articles published from 2000 to January 28, 2020. Keywords

searched included: ("Air Pollution" OR "Pollution, Air" OR "Air Quality" OR "Ultrafine Fibers"

OR "Airborne Particulate Matter" OR "Particulate Matter, Airborne" OR "Air Pollutants,

Particulate" OR "Particulate Air Pollutants" OR "Ambient Particulate Matter" OR "Particulate

Matter, Ambient" OR "Ultrafine Particulate Matter" OR "Particulate Matter, Ultrafine" OR

"Ultrafine Particles" OR "Particles, Ultrafine") AND ("Cardiovascular Disease" OR "Disease,

Cardiovascular" OR "Diseases, Cardiovascular" OR "HDL" OR "LDL" OR "cholesterol" OR

" cholesterin" OR "cholesteric" OR "TC" OR "TG" OR "dyslipidaemia" OR "HDL-C" OR

"LDL-C") AND ("Cohort" OR "Cross-sectional" OR "Case control" OR "Case-control" OR

"Epidemiology OR "Epidemiological"). Additionally, references of included literature and one

previously published systematic review were manually retrieved.

2.2 Eligibility Criteria

2.2.1 Inclusion Criteria

Studies were included if: (1) The study includes all population with no age, gender, or race

restrictions. (2) The study describes the relationship between particulate air pollutants and

metabolic disease/cardiovascular disease. (3) Results quantify changes in lipid parameters,

including HDL-C, LDL-C, TC, TG, etc. (4) Cross-sectional studies, cohort studies, or case control

studies. (5) The Literature is published in either Chinese or English.

2.2.2 Exclusion Criteria

Studies were excluded if: (1) The research was based on animal experiments or cytological

experiments. (2) The study was a review or meta-analysis. (3) The study was a duplicate. (4)

Effect estimates were not reported as a percent change in lipid metabolism per 10 µg/m^3 change in

particulate air pollutants. (5) The data reported resulted from an independent study. If inconsistent

results have been reported for a given population or period at the time publications were retrieved,

older studies were superseded by the most recent publication.

2.3 Literature Screening and Quality Evaluation

Two researchers independently retrieved and screened the literature following the above

order and criteria. Upon review, a third researcher was consulted when discrepancies occurred.

The methodological quality of each cohort study was evaluated according to the

Newcastle-Ottawa Scale (NOS) literature quality evaluation tool(Luchini, Stubbs, Solmi, &

Veronese, 2017), which includes three aspects of population selection, comparability, and results.

A total of 8 items belonging to NOS could be graded by an "*" for each item, and the top score

can include up to 9 "*". Cross-sectional studies were assessed by the American Health Care

Quality and Research Institute (AHRQ) scale for quality evaluation(B, 2017). An 11-point scale is

used to calculate responses of either "Yes", "No", or "Not clear", for a total score of 11.

2.4 Data extraction and statistical analysis

Information was independently extracted by two researchers based on a pre-set data

extraction form which included: author name(s), year of publication, study period, location,

population, median age, sample size, experimental design type, exposure time, and contaminants.

Divergent interpretations were reconciled by a third researcher.

When the literature data were extracted, the percent changes in lipid metabolism per 10

$\mu g/m^3$ change in particulate air pollutants were directly utilized. As such, the results presented in

this meta-analysis are expressed as the percentage change in lipid parameters and MetS incidence

for each 10 $\mu g/m^3$ increment in air pollutant. The between-study heterogeneity was evaluated

using both the I^2 statistic and Q-test(Huedo-Medina, Sanchez-Meca, Marin-Martinez, & Botella,

2006). If $I^2 \geq 50\%$ or the p-value for a Q-test was < 0.01, then heterogeneity was considered "high"

and the Der Simonian and Laird random effects model was used to pool effect estimates.

Otherwise, heterogeneity was considered "moderate" or "low" and the Mantel-Haenszel fixed

effects model was used to pool effect estimates. The Meta-analysis was performed using Stata

software version 14.0. A p-value < 0.05 was considered statistically significant.

3. Results

3.1 Literature Retrieval and Characteristics

A total of 2,274 records were obtained from multiple databases. After duplicates were

removed, the titles and abstracts of 1,714 records were screened. 1,154 full-text publications were

subsequently assessed using the eligibility criteria. As a result, ten records met all criteria and

were included in the final quantitative meta-analysis(Cai et al., 2017; Hou et al., 2020; Lee et al.,

2019; Shanley et al., 2016; Sorensen et al., 2015; Wallwork et al., 2017; Wang et al., 2018; Wu et

al., 2019; Yang, Bloom, et al., 2018; Yitshak Sade, Kloog, Liberty, Schwartz, & Novack, 2016). A

flow diagram of the study selection process is shown in the Fig. 1.

Overall, this study includes a total of ten records comprising eight cohort studies(Cai et al.,

2017; Lee et al., 2019; Shanley et al., 2016; Wallwork et al., 2017; Wang et al., 2018; Wu et al.,

2019; Yang, Bloom, et al., 2018; Yitshak Sade et al., 2016) and two cross-sectional studies(Hou et

al., 2020; Sorensen et al., 2015). An exposure time of less than or equal to thirty days was

recorded as short-term exposure, whereas an exposure time greater than thirty days was recorded

as long-term exposure. Study characteristics are summarized in Table 1.

3.2 Quality Evaluation

Using the NOS scale to evaluate eight cohort studies, one study was less representative, three

studies were less comparable, six mentioned follow-ups, eight mentioned the specific

measurement methods from which the results were derived, and six were subject to the integrity of

follow-up. For this quality evaluation, a study was awarded one "*", corresponding to one point,

for each NOS item it contained. Using the AHRQ scale to evaluate two cross-sectional studies,

both mentioned the source of data, patient identification time period, follow-up situation, and

continuity of the subject. For evaluation, "yes" was recorded as one point, whereas "no" or

"uncertain" was recorded as zero for each component of the AHRQ scale. Final scores for each

respective study is summarized in Table 1.

3.3 Data Extraction and Conversion

The results of the data extraction of the ten records are shown in Table 2. The growth rate of

particulate air pollutants varied from study to study. As such, in order to pool data for further

analysis, original data from each study was uniformly converted to the rate of change of each lipid

parameter per 10 μg/m^3 increment change in air pollutant. A complete list of converted results is

reported in Table 3.

3.4 Quantitative Meta-analysis

Based on a quantitative comparison of the ten studies included in the analysis, a total of

twelve groups were identified to characterize the relationship between particulate air pollutants

and lipid parameters, including PM$_{2.5}$/TC; PM$_{2.5}$/HDL-C; PM$_{2.5}$/LDL-C; PM$_{2.5}$/MetS; PM$_{10}$/TG;

PM$_{10}$/LDL-C; PM$_{10}$/HDL-C; NO$_2$/TC; NO$_2$/TG; NO$_2$/HDL-C; PM$_{10}$/HDL-C (diabetic population);

PM$_{10}$/LDL-C(diabetic population).After data conversion, these effect estimates were analyzed

using twelve different analyses for each pollutant/outcome combination.

3.4.1 PM$_{2.5}$ with TC, HDL-C, LDL-C, and MetS(Lee et al., 2019; Sorensen et al., 2015; Wu et al.,

2019; Yang, Bloom, et al., 2018)

PM$_{2.5}$ exposure was significantly associated with TC, HDL-C, LDL-C, and MetS.

Specifically, for each 10 μg/m^3 increase in PM$_{2.5}$, TC levels increased by 3.31% (95% CI:

-0.0229~0.0891, P=0.046), LDL-C levels increased by 2.34% (95% CI: 0.0130~0.0339, P=0.001),

HDL-C levels decreased 1.57% (95% CI: -0.0185~-0.0128, P=0.001), and the MetS increased by

4.33% (95% CI: 0.0269~0.0598, P<0.001) (See Fig. 2).

3.4.2 PM$_{10}$ with TG, HDL-C, and LDL-C(Cai et al., 2017; Shanley et al., 2016; Yang, Bloom, et

al., 2018; Yitshak Sade et al., 2016)

PM$_{10}$ exposure was significantly associated with TG, HDL-C, LDL-C. Specifically, for each

10 μg/m^3 increase in PM$_{10}$, TG levels increased by 5.27% (95% CI: 0.0203~0.0850, P<0.001),

HDL-C levels decreased by 0.24% (95% CI: -0.0095~0.0047, P=0.002), and LDL-C levels

increased by 0.45% (95% CI: -0.057~0.0147, P<0.001) (See Fig. 3).

3.4.3 NO_2 with TG, HDL-C, and TC(Cai et al., 2017; Sorensen et al., 2015; Yang, Bloom, et al.,

2018)

NO_2 exposure was associated with TG and HDL-C; however, was not significantly associated

with TC. Specifically, for each 10 μg/m^3 increase in NO_2, TG levels increased by 4.18% (95% CI:

0.0112~0.0723, P=0.020), and HDL-C levels decreased by 0.51% (95% CI: -0.0261~0.0158,

P<0.001). TC levels increased by 1.01% (95% CI: 0.0035~0.0167, P=0.858) (See Fig. 4).

3.4.4 PM_{10} with LDL-C and HDL-C in diabetic population(Wang et al., 2018; Yitshak Sade et al.,

2016)

Three studies included subjects with diabetes mellitus, suggesting differential modification of

LDL-C and HDL-C levels associated with PM_{10} exposure. PM_{10} exposure was significantly

associated with HDL-C levels, but not with LDL-C levels. For each 10 μg/m^3 increase in PM_{10},

HDL-C levels decreased 0.45% (95% CI: -0.0072~-0.0017, P=0.009), while LDL-C increased by

1.14% (95% CI: 0.0090~0.00138, P=0.265) (See Fig. 5).

4 Discussion

This meta-analysis found significant associations between $PM_{2.5}$, PM_{10}, and NO_2, and

lipoprotein parameters, with evidence of substantial between-study heterogeneity. These findings

add to evidence that MetS may increase susceptibility to cardiovascular morbidity associated with

air pollution exposure.

Gaio et al.(Vânia et al., 2019) reported that LDL-C and TG levels increased by 0.12% (95%

CI: 0.0178~0.0206) and 3.14% (95% CI: 0.0136~0.0495) respectively, per 10

µg/m^3 increase in PM$_{10}$, which is consistent with our study, although effect estimates were smaller

than the current study. They also observed a decrease of 0.45% (95% CI: -0.0265~-0.0181) in

HDL-C levels per 10 µg/m^3 increase in NO$_2$, but no statistically significant changes in TC. When

considering the differences between the two studies, such factors as the use of differential

inclusion and exclusion criteria for screening may have resulted in more or less robust records for

final selection. To the extent that studies focused on associations between particulate air pollutants

and lipometabolic disturbance were much less based on the pre-search, the inclusion criteria in our

study were broader, while the exclusion criteria were similar to the meta-analysis by Gaio et al. As

a result, our study included an additional study by Maayan et al.(Yitshak Sade et al., 2016) in the

subgroup category for studies investigating the relationship between PM$_{10}$ exposures and LDL-C

and TG levels. In this study, PM$_{2.5}$ was more significantly associated with multiple lipid

parameters compared to PM$_{10}$, which is consistent with findings by Hazrije et al.(Mustafić et al.,

2012); however, Hazrije et al.(Mustafić et al., 2012) reported a stronger association between

myocardial infarction and PM$_{2.5}$ compared to PM$_{10}$. This may result from the differential

characteristics of PM$_{2.5}$ such as smaller diameter size, larger surface area, slower sedimentation

rate, and longer retention time in the air compared to PM$_{10}$.

This paper reported that elevated PM$_{10}$ exposure on the diabetic population was associated

with higher HDL-C levels, but not significantly associated with LDL-C levels. This was only

partially consistent with the study by Wang et al.(Minzhen & Shan, 2018 (In China)), which

reported significant associations between PM$_{10}$ exposure and both HDL-C levels and LDL-C

levels. Such discrepancy may be due to differences in the characteristics of the study population.

Further, two studies(Wang et al., 2018; Yitshak Sade et al., 2016) included in this meta-analysis

contained subjects with diabetes using data obtained from medical institutions, whereas the Wang

et al.(Minzhen & Shan, 2018 (In China)) had the advantage of follow-up throughout the study

period, which may have provided a more precise measure of changes in lipid parameters.

The present study suggests that air pollution may adversely affect lipometabolic balance by

promoting LDL oxidation, disrupting the scavenger receptor and LDL receptor body function,

accelerating the accumulation of lipids in plaque(Fuertes, Plaat, & Minelli, 2020; Miller, 2020).

HDL, as a protective factor for lipid metabolic disorders, decreased in levels when exposed to air

pollutants(Bell et al., 2017).

Despite a considerably inclusive search strategy, with few restrictions on study design and

population characteristics, our final selection yielded 10 qualified studies for this meta-analysis.

We detected high between-study heterogeneity in most of the subgroups studied, which, taken

together, may attenuate credibility with cumulative evidence. Yet, heterogeneity may be captured

by unmeasured variables that were not reported such as population susceptibility, temperature,

distribution of air pollutants or other potential sources(Liu et al., 2019; Ward-Caviness, 2019).

There are some limitations that should be addressed. First, most of the literature failed to

consider whether subjects had taken lipid-lowering drugs or related drugs using history before or

during the study period, which may bias results and lead to less positive change in blood lipid

parameter levels(Ho & Hee, 2015). Furthermore, evidence suggests that noise intensity can

directly impact lipid parameters levels(Cai et al., 2017; Y.-K. Huang, Hanneke, & Jones, 2019;

Jørgensen et al., 2019; Klompmaker et al., 2019; Shapovalova, Ryzhova, & Ryzhov, 2010;

Sorensen et al., 2015). Considering noise has obvious geographical distribution characteristics, i.e.,

urban residential areas typically have higher levels of noise intensity than rural or remote areas,

and that distance between residential areas and main roadways could fluctuate multiple lipid

parameters, future analyses should further examine noise intensity as a possible confounding risk

factor. Similarly, green spaces near residential areas or workplaces have been shown to influence

lipometabolic balance and should additionally be included as a confounding variable(Ji, Zhu, Lv,

& Shi, 2019; Klompmaker et al., 2019; Yang et al., 2019). Moreover, residential greenness has

been shown elsewhere to have a beneficial effect on MetS and diabetes; accordingly, this

association could be attenuated after adjustment for air pollution(Thiering et al., 2016; Yang et al.,

2020). To quantify air pollution exposure, many studies(Cai et al., 2017; Hou et al., 2020; Lee et

al., 2019; Shanley et al., 2016; Sorensen et al., 2015; Wang et al., 2018) use ground-based

measurements derived from fixed air quality monitoring sites. While other studies(Wallwork et al.,

2017; Wu et al., 2019; Yang, Bloom, et al., 2018; Yitshak Sade et al., 2016) incorporate predictive

models estimated from spatiotemporal hybrid modeling. More specifically, models may utilize

low-cost GPS on mobile devices to monitor an average individual exposure to air pollutants in real

time, over the study period. Such methods greatly increase the accuracy of measurements and may

reflect a better estimate of true exposure to air pollutants. Despite the limitations, previous studies

have focused on the relationship between blood lipid parameters levels and a single air pollutant.

Examining the effects of air pollution exposure one pollutant at a time underestimates the

complexity of atmospheric chemical mixing and the multiple pollutant sources implicated in the

link between air pollution exposure and adverse health, and more specifically, the nuanced effects

of various pollutants on blood lipid parameters.

5 Conclusion

This study found that PM_{10}, $PM_{2.5}$, and NO_2 were significantly associated with HDL-C,

LDL-C, TC, and TG levels, suggesting a link between air pollution and cardiovascular morbidity.

Therefore, improving air quality may yield substantial health benefits.

Declaration of Competing Interest

The authors declare no competing interests.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (81872649),

National Key R&D Program of China (2017YFC0211600), Scientific Research Project of Beijing

Educational Committee (KM201910025023), and HERCULES Exposome Research Centre

(P30ES019776).

References

Akintoye, E., Shi, L., Obaitan, I., Olusunmade, M., Wang, Y., Newman, J. D., & Dodson, J. A.

(2016). Association between fine particulate matter exposure and subclinical

atherosclerosis: A meta-analysis. *European journal of preventive cardiology, 23*(6),

602-612. doi:10.1177/2047487315588758

B, B. A. (2017). The Agency for Healthcare Research and Quality and the Development of a

Learning Health Care System. *JAMA internal medicine, 177*(7).

doi:10.1001/jamainternmed.2017.2589

Bell, G., Mora, S., Greenland, P., Tsai, M., Gill, E., & Kaufman, J. D. (2017). Association of Air

Pollution Exposures With High-Density Lipoprotein Cholesterol and Particle Number:

The Multi-Ethnic Study of Atherosclerosis. *Arteriosclerosis, thrombosis, and vascular*

biology, 37(5), 976-982. doi:10.1161/atvbaha.116.308193

Cai, Y., Hansell, A. L., Blangiardo, M., Burton, P. R., de Hoogh, K., Doiron, D., . . . Hodgson, S.

(2017). Long-term exposure to road traffic noise, ambient air pollution, and

cardiovascular risk factors in the HUNT and lifelines cohorts. *European heart journal,*

38(29), 2290-2296. doi:10.1093/eurheartj/ehx263

Cohen, A. J., Brauer, M., Burnett, R., Anderson, H. R., Frostad, J., Estep, K., . . . Forouzanfar,

M. H. (2017). Estimates and 25-year trends of the global burden of disease

attributable to ambient air pollution: an analysis of data from the Global Burden of

Diseases Study 2015. *Lancet (London, England), 389*(10082), 1907-1918.

doi:10.1016/s0140-6736(17)30505-6

Combes, A., & Franchineau, G. (2019). Fine particle environmental pollution and cardiovascular diseases. *Metabolism: Clinical and Experimental, 100.* doi:10.1016/j.metabol.2019.07.008

Dabass, A., Talbott, E. O., Rager, J. R., Marsh, G. M., Venkat, A., Holguin, F., & Sharma, R. K. (2018). Systemic inflammatory markers associated with cardiovascular disease and acute and chronic exposure to fine particulate matter air pollution (PM2.5) among US NHANES adults with metabolic syndrome. *Environmental research, 161*, 485-491. doi:10.1016/j.envres.2017.11.042

Dagenais, G. R., Leong, D. P., Rangarajan, S., Lanas, F., Lopez-Jaramillo, P., Gupta, R., . . . Yusuf, S. (2020). Variations in common diseases, hospital admissions, and deaths in middle-aged adults in 21 countries from five continents (PURE): a prospective cohort study. *The Lancet, 395*(10226), 785-794. doi: 10.1016/S0140-6736(19)32007-0

Eze, I. C., Schaffner, E., Foraster, M., Imboden, M., von Eckardstein, A., Gerbase, M. W., . . . Probst-Hensch, N. (2015). Long-Term Exposure to Ambient Air Pollution and Metabolic Syndrome in Adults. *PloS one, 10*(6), e0130337. doi:10.1371/journal.pone.0130337

Fuertes, E., Plaat, D. A. v. d., & Minelli, C. (2020). Antioxidant genes and susceptibility to air pollution for respiratory and cardiovascular health. *Free Radical Biology and Medicine.* doi:10.1016/j.freeradbiomed.2020.01.181

Ho, A. C., & Hee, C. S. (2015). New drugs for treating dyslipidemia: beyond statins. *Diabetes & metabolism journal, 39*(2). doi:10.4093/dmj.2015.39.2.87

Hou, J., Liu, X., Tu, R., Dong, X., Zhai, Z., Mao, Z., . . . Wang, C. (2020). Long-term exposure

to ambient air pollution attenuated the association of physical activity with metabolic

syndrome in rural Chinese adults: A cross-sectional study. *Environment international,*

136, 105459. doi:10.1016/j.envint.2020.105459

Huang, H. C., Tantoh, D. M., Hsu, S. Y., Nfor, O. N., Frank, C. F. L., Lung, C. C., . . . Liaw, Y. P.

(2020). Association between coarse particulate matter (PM 10-2.5) and

nasopharyngeal carcinoma among Taiwanese men. *Journal of Investigative Medicine,*

68(2), 419-424. doi:10.1136/jim-2019-001119

Huang, Y.-K., Hanneke, R., & Jones, R. M. (2019). Bibliometric analysis of cardiometabolic

disorders studies involving NO(2), PM(2.5) and noise exposure. *BMC public health,*

19(1), 877-877. doi:10.1186/s12889-019-7195-1

Huedo-Medina, T. B., Sanchez-Meca, J., Marin-Martinez, F., & Botella, J. (2006). Assessing

heterogeneity in meta-analysis: Q statistic or I2 index? *Psychol Methods, 11*(2),

193-206. doi:10.1037/1082-989x.11.2.193

J, P. M., E, M. J., M, B. P., Isabelle, B., Tammy, H., D, M. C., . . . David, M. (2020). Mapping of

reporting guidance for systematic reviews and meta-analyses generated a

comprehensive item bank for future reporting guidelines. *Journal of clinical*

epidemiology, 118. doi:10.1016/j.jclinepi.2019.11.010

Ji, J. S., Zhu, A., Lv, Y., & Shi, X. (2019). Residential greenness and air pollution mortality

using the Chinese Longitudinal Healthy Longevity Survey: a longitudinal analysis. *The*

Lancet, 394, S16. doi:10.1016/S2542-5196(20)30027-9

Jørgensen, J. T., Bräuner, E. V., Backalarz, C., Laursen, J. E., Pedersen, T. H., Jensen, S.

S., . . . Andersen, Z. J. (2019). Long-term exposure to road traffic noise and incidence

of diabetes in the danish nurse cohort. *Environmental health perspectives, 127*(5). doi:10.1289/EHP4389

Klompmaker, J. O., Janssen, N. A. H., Bloemsma, L. D., Gehring, U., Wijga, A. H., Brink, C. V., . . . Hoek, G. (2019). Associations of combined exposures to surrounding green, air pollution, and road traffic noise with cardiometabolic diseases. *Environmental health perspectives, 127*(8). doi:10.1289/EHP3857

Landrigan, P. J., Fuller, R., Acosta, N. J. R., Adeyi, O., Arnold, R., Basu, N., . . . Zhong, M. (2018). The Lancet Commission on pollution and health. *The Lancet, 391*(10119), 462-512. doi: 10.1016/S0140-6736(17)32345-0

Lee, S., Park, H., Kim, S., Lee, E.-K., Lee, J., Hong, Y. S., & Ha, E. (2019). Fine particulate matter and incidence of metabolic syndrome in non-CVD patients: A nationwide population-based cohort study. *International journal of hygiene and environmental health, 222*(3), 533-540. doi:10.1016/j.ijheh.2019.01.010

Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., & Pozzer, A. (2015). The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature, 525*(7569), 367-371. doi:10.1038/nature15371

Lin, Z., Chen, R., Jiang, Y., Xia, Y., Niu, Y., Wang, C., . . . Kan, H. (2019). Cardiovascular Benefits of Fish-Oil Supplementation Against Fine Particulate Air Pollution in China. *Journal of the American College of Cardiology, 73*(16), 2076-2085. doi:10.1016/j.jacc.2018.12.093

Liu, M., Xue, X., Zhou, B., Zhang, Y., Sun, B., Chen, J., & Li, X. (2019). Population susceptibility differences and effects of air pollution on cardiovascular mortality:

epidemiological evidence from a time-series study. *Environmental science and pollution research international, 26*(16), 15943-15952. doi:10.1007/s11356-019-04960-2

Luchini, C., Stubbs, B., Solmi, M., & Veronese, N. (2017). Assessing the quality of studies in meta-analyses: Advantages and limitations of the Newcastle Ottawa Scale. *World Journal of Meta-Analysis, 5*(04), 80-84. doi: 10.1093/humrep/dew244

Miller, M. R. (2020). Oxidative stress and the cardiovascular effects of air pollution. *Free Radical Biology and Medicine*. doi:10.1016/j.freeradbiomed.2020.01.004

Minzhen, W., & Shan, Z. (2018 (In China)). Research progress on the effects of atmospheric PM_(2.5) exposure on type 2 diabetes in people. *Journal of Environmental and Occupational Medicine, 35*(02), 137-142. doi:10.13213/j.cnki.jeom.2018.17454

Mustafić, H., Jabre, P., Caussin, C., Murad, M. H., Escolano, S., Tafflet, M., . . . Jouven, X. (2012). Main Air Pollutants and Myocardial Infarction: A Systematic Review and Meta-analysis. *JAMA: The Journal of the American Medical Association, 307*(7). doi:10.1001/jama.2012.126

Myung, W., Lee, H., & Kim, H. (2019). Short-term air pollution exposure and emergency department visits for amyotrophic lateral sclerosis: A time-stratified case-crossover analysis. *Environment international, 123*, 467-475. doi:10.1016/j.envint.2018.12.042

Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. (2016). *BMJ (Clinical research ed.), 354*, i4086. doi:10.1136/bmj.i4086

Runzhi, C. (2020 (In China)). The effects of major atmospheric pollutants on population health

and its response progress. *Chinese Journal of Public Health Management, 36*(01), 51-53. doi:10.19568/j.cnki.23-1318.2020.02.037

Schraufnagel, D. E., Balmes, J. R., Cowl, C. T., De Matteis, S., Jung, S. H., Mortimer, K., . . . Wuebbles, D. J. (2019). Air Pollution and Noncommunicable Diseases: A Review by the Forum of International Respiratory Societies' Environmental Committee, Part 2: Air Pollution and Organ Systems. *Chest, 155*(2), 417-426. doi:10.1016/j.chest.2018.10.041

Seelen, M., Toro Campos, R. A., Veldink, J. H., Visser, A. E., Hoek, G., Brunekreef, B., . . . Vermeulen, R. C. H. (2017). Long-Term Air Pollution Exposure and Amyotrophic Lateral Sclerosis in Netherlands: A Population-based Case-control Study. *Environmental health perspectives, 125*(9), 097023-097023. doi:10.1289/EHP1115

Shanley, R. P., Hayes, R. B., Cromar, K. R., Ito, K., Gordon, T., & Ahn, J. (2016). Particulate Air Pollution and Clinical Cardiovascular Disease Risk Factors. *Epidemiology (Cambridge, Mass.), 27*(2), 291-298. doi:10.1097/ede.0000000000000426

Shapovalova, V. P., Ryzhova, T. V., & Ryzhov, V. M. (2010). Lipid metabolism state under exposure to noise and aluminum dust. *Meditsina truda i promyshlennaia ekologiia*(7), 18-20. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/20853625

Shi, L., Zanobetti, A., Kloog, I., Coull, B. A., Koutrakis, P., Melly, S. J., & Schwartz, J. D. (2016). Low-Concentration PM2.5 and Mortality: Estimating Acute and Chronic Effects in a Population-Based Study. *Environmental health perspectives, 124*(1), 46-52. doi:10.1289/ehp.1409111

Sorensen, M., Hjortebjerg, D., Eriksen, K. T., Ketzel, M., Tjonneland, A., Overvad, K., &

Raaschou-Nielsen, O. (2015). Exposure to long-term air pollution and road traffic noise in relation to cholesterol: A cross-sectional study. *Environment international, 85*, 238-243. doi:10.1016/j.envint.2015.09.021

Thiering, E., Markevych, I., Brüske, I., Fuertes, E., Kratzsch, J., Sugiri, D., . . . Heinrich, J. (2016). Associations of Residential Long-Term Air Pollution Exposures and Satellite-Derived Greenness with Insulin Resistance in German Adolescents. *Environmental health perspectives, 124*(8), 1291-1298. doi:10.1289/ehp.1509967

Thomson, E. M. (2019). Air Pollution, Stress, and Allostatic Load: Linking Systemic and Central Nervous System Impacts. *Journal of Alzheimer's Disease, 69*(3), 597-614. doi:10.3233/JAD-190015

Tsai, S.-S., Lin, Y.-S., Chen, S.-T., & Chu, P.-H. (2018). Metabolic syndrome positively correlates with the risks of atherosclerosis and diabetes in a Chinese population. *European Journal of Internal Medicine*. doi:10.1016/j.ejim.2018.04.009

Vânia, G., Rita, R., Matias, D. C., & Baltazar, N. (2019). Ambient air pollution and lipid profile: Systematic review and meta-analysis. *Environmental pollution (Barking, Essex : 1987), 254*(Pt B). doi:10.1016/j.envpol.2019.113036

Wallwork, R. S., Colicino, E., Zhong, J., Kloog, I., Coull, B. A., Vokonas, P., . . . Baccarelli, A. A. (2017). Ambient fine particulate matter, outdoor temperature, and risk of met. *American journal of epidemiology, 185*(1), 30-39. doi:10.1093/aje/kww157

Wang, M., Zheng, S., Nie, Y., Weng, J., Cheng, N., Hu, X., . . . Bai, Y. (2018). Association between short-term exposure to air pollution and dyslipidemias among type 2 diabetic patients in Northwest China: A population-based study. *International journal of*

environmental research and public health, 15(4). doi:10.3390/ijerph15040631

Ward-Caviness, C. K. (2019). A review of gene-by-air pollution interactions for cardiovascular disease, risk factors, and biomarkers. *Hum Genet, 138*(6), 547-561. doi:10.1007/s00439-019-02004-w

Weidong, W., Zhen, A., Shuangjun, G., Jie, X., Wei, F., & Jie, S. (2018 (In China)). Research progress on the development of atherosclerosis induced by PM_(2.5) exposure. *Journal of Shandong University (Health Science), 56*(11), 18-26. Retrieved from http://www.cnki.com.cn/Article/CJFDTotal-SDYB201811004.htm

Wu, X. M., Broadwin, R., Basu, R., Malig, B., Ebisu, K., Gold, E. B., . . . Green, S. (2019). Associations between fine particulate matter and changes in lipids/lipoproteins among midlife women. *Science of the Total Environment, 654*, 1179-1186. doi:10.1016/j.scitotenv.2018.11.149

Yang, B. Y., Bloom, M. S., Markevych, I., Qian, Z. M., Vaughn, M. G., Cummings-Vaughn, L. A., . . . Dong, G. H. (2018). Exposure to ambient air pollution and blood lipids in adults: The 33 Communities Chinese Health Study. *Environment international, 119*, 485-492. doi:10.1016/j.envint.2018.07.016

Yang, B. Y., Liu, K. K., Markevych, I., Knibbs, L. D., Bloom, M. S., Dharmage, S. C., . . . Dong, G. H. (2020). Association between residential greenness and metabolic syndrome in Chinese adults. *Environment international, 135*. doi:10.1016/j.envint.2019.105388

Yang, B. Y., Markevych, I., Heinrich, J., Bloom, M. S., Qian, Z., Geiger, S. D., . . . Dong, G. H. (2019). Residential greenness and blood lipids in urban-dwelling adults: The 33 Communities Chinese Health Study. *Environmental Pollution, 250*, 14-22.

doi:10.1016/j.envpol.2019.03.128

Yang, B. Y., Qian, Z. M., Li, S., Fan, S., Chen, G., Syberg, K. M., . . . Dong, G. H. (2018). Long-term exposure to ambient air pollution (including PM1) and metabolic syndrome: The 33 Communities Chinese Health Study (33CCHS). *Environmental research, 164*, 204-211. doi:10.1016/j.envres.2018.02.029

Yitshak Sade, M., Kloog, I., Liberty, I. F., Schwartz, J., & Novack, V. (2016). The Association Between Air Pollution Exposure and Glucose and Lipids Levels. *The Journal of clinical endocrinology and metabolism, 101*(6), 2460-2467. doi:10.1210/jc.2016-1378

Yuanning, L., Aiqun, H., Hui, Z., Wenqin, W., Yanfen, L., & Mei, X. (2019 (In China)). Application of combined detection of TC, TG, HDL-C, LDL -C, ApoA1 and ApoB in cardiovascular and cerebrovascular diseases. *Medical Journal of Liaoning, 33*(01), 50-52. Retrieved from http://www.cnki.com.cn/Article/CJFDTOTAL-LNYX201901019.htm

Yusuf, S., Joseph, P., Rangarajan, S., Islam, S., Mente, A., Hystad, P., . . . Dagenais, G. (2020). Modifiable risk factors, cardiovascular disease, and mortality in 155□722 individuals from 21 high-income, middle-income, and low-income countries (PURE): a prospective cohort study. *The Lancet, 395*(10226), 795-808. doi:10.1016/S0140-6736(19)32008-2

Zheng, Q., Liu, H., Zhang, J., & Chen, D. (2018). The effect of ambient particle matters on hospital admissions for cardiac arrhythmia: A multi-city case-crossover study in China. *Environmental Health: A Global Access Science Source, 17*(1). doi:10.1186/s12940-018-0404-z

Zu, K., Liu, X., Shi, L., Tao, G., Loftus, C. T., Lange, S., & Goodman, J. E. (2017). Concentration-response of short-term ozone exposure and hospital admissions for asthma in Texas. *Environment international, 104*, 139-145. doi:10.1016/j.envint.2017.04.006

Figure Legend

Fig.1. Flow diagram of study selection process.

Fig.2. Forest plot of $PM_{2.5}$ exposure (per 10 μg/m^3 increase) with changes of TC, HDL-C, LDL levels and MetS morbidity. (* % change in the outcome per 10 mg/m^3 pollutant increase. Abbreviations: I^2 =measure of between-study heterogeneity; P=Cochran's Q test P-value.)

Fig.3. Forest plot of PM_{10} exposure (per 10 μg/m^3 increase) with changes of TG, LDL-C, and HDL-C levels. (* % change in the outcome per 10 mg/m^3 pollutant increase. Abbreviations: I^2 =measure of between-study heterogeneity; P=Cochran's Q test P-value.)

Fig.4. Forest plot of NO_2 exposure (per 10 μg/m^3 increase) with changes of TC, TG and HDL-C levels. (* % change in the outcome per 10 mg/m^3 pollutant increase. Abbreviations: I^2 =measure of between-study heterogeneity; P=Cochran's Q test P-value.)

Fig.5. Forest plot of PM_{10} exposure (per 10 μg/m^3 increase) with changes of HDL-C and LDL-C levels in diabetic population. (* % change in the outcome per 10 mg/m^3 pollutant increase. Abbreviations: I^2 =measure of between-study heterogeneity; P=Cochran's Q test P-value.)

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5