Association of Renin-Angiotensin-Aldosterone System Inhibition with Risk of

COVID-19, Inflammation Level, Severity and Death in Patients With COVID-19: A

Rapid Systematic Review and Meta-Analysis

Running title: ACEI/ARB on COVID-19

Xiao Liu1*, MD, Chuyan Long1*, MD, Qinmei Xiong1, MD, Chen Chen1, MD, Jianyong

Ma1, MD, Yuhao Su1, M.D., Kui Hong1,3, MD, PhD

1. Cardiology Department, the Second Affiliated Hospital of Nanchang University, Jiangxi,

330006, China

2. Jiangxi Key Laboratory of Molecular Medicine, Jiangxi, 330006, China

*Dr. Liu, and Dr. Long are co-first authors.

Address for Correspondence

Dr. Kui Hong, MD, PhD;

Cardiovascular Department, the Second Affiliated Hospital of Nanchang University;

Minde Road No. 1, Nanchang, Jiangxi, 330006, China;

Email: hongkui88@163.com;

Tel: +86-0791-86312917

Introduction

The coronavirus disease 2019 (COVID-19) pandemic is becoming one of the most far-reaching

public health crises in recorded history. At right time of writing this review, on April 29[th],

2020, the number of infected persons worldwide has exceeded 3.01 million, with more than

200,000 reported deaths (http://2019ncov.chinacdc. cn/2019-nCoV/global.html). At present,

there is no specifically targeted, effective treatment for patients with COVID-19. There is an

urgent need, therefore, to determine how to alleviate the severe clinical symptoms and reduce

the morbidity and mortality due to this disease. Severe acute respiratory syndrome coronavirus

2 (SARS-CoV-2) shares the same cell entry receptor as SARS-CoV, in which the viruses' spike

proteins bind to the host cell surface. Angiotensin converting enzyme 2 (ACE2) receptors,

which are an essential regulator of renin–angiotensin–aldosterone system (RAAS) activity[1, 2].

The RAAS is a vital regulator of cardiovascular and renal function, including blood pressure,

and which plays a vital role in regulating acute lung injury[3, 4]. The ACE2 level is significantly

reduced in patients following SARS-CoV infection, resulting in RAAS system imbalance, and

eventually caused severe acute lung injury. Angiotensin I converting enzyme inhibitors

(ACEIs) and Angiotensin II receptor blockers (ARBs) are ACE2 receptor antagonists that can

reduce the activity of the RAAS system and which have been widely used in the past several

decades to treat cardiovascular diseases such as hypertension and hypertrophic

cardiomyopathy[3]. However, some animal experiments and clinical trials have shown that

ACEIs potentially result in an increase in ACE2 receptors[5]. Some concern has therefore been

expressed that the use of ACEIs/ARBs might increase the risk of COVID-19 infection after

exposure to SARS-CoV-2 and result in a poor prognosis[6]. On April 12th, 2020, the European

Society of Hypertension COVID-19 Task Force issued the following statement: the current

evidence does not support that RAAS inhibitors aggravate the condition of COVID-19

patients[5].

Accumulating evidence revealed that inflammatory cytokine storm and dysfunction of

immune system largely contribute to the severe COVID-19, even cause death[7]. The latest

evidence shows that peripheral blood interlekin-6 (IL-6) levels, a critical mediator of

respiratory failure, shock, and multiorgan dysfunction, were significantly increased in

COVID-19 patients who had used a RAAS inhibitor[8, 9]. Moreover, CD3 and CD8 cell counts

were significantly reduced, suggesting that the level of inflammation in patients treated with

ACEIs is relatively low[8]. Several retrospective studies have shown that the use of

ACEIs/ARBs prior to COVID-19 infection could reduce the severity of COVID-19 and was

associated with a better prognosis, especially in patients with both hypertension and

COVID-19 [10-12]. However, other studies have shown that there is no significant correlation

between the use of ACEIs/ARBs and the severity and mortality of patients with COVID-19[10,

13-17]. Because the affection of RAAS inhibitors on the clinical prognosis of COVID-19 patients

is unclear, in this article, we will review the current evidence and assess the clinical prognosis

of COVID-19 patients with or without hypertension treated with ACEs/ARBs.

Methods

This study was performed according to PRISMA guidelines (http://www.prisma-

statement.org; **Table S1** Online Data Supplement)[18].

Literature Search

Two authors (L X. and Y.L) independently searched the PubMed and Embase databases for published articles and the preprint platforms medRxiv (https://www.medrxiv.org/) and bioRxiv (https://www.biorxiv.org/) (since many studies are available on these websites prior to publication, which allows for collection of the latest data) without language restrictions. Furthermore, the research references were traced and cross-checked. In the case of any discrepancy, it was resolved by consensus with the third author (H.K). The databases were searched for articles from February 1, 2019 to May 1, 2020 using the following search terms: ACE inhibitor, angiotensin II receptor blocker, angiotensin-converting enzyme inhibitor, 2019-novel coronavirus, SARS-CoV-2, COVID-19, and 2019-nCoV.

Study Selection

Studies were considered eligible for inclusion if they 1) were designed as a randomized controlled trial, case-control study, or cohort study and 2) assessed the relationship between ACEI/ARB use and the level of inflammation, disease severity, and mortality in patients with COVID-19. If multiple studies used the same population, we selected the most recent publication. Certain publication types (e.g., reviews, editorials, letters, conference abstracts, and animal studies) or studies with insufficient data were excluded from this analysis.

Data Extraction and Quality Assessment

Two authors (L X. and Y.L) independently extracted the study information and the basic characteristics of the articles using a standardized form, including the first author, year of

publication, country, sample size, sex ratio, age, sample size, RAAS type, length of follow-up,

adjustments for confounders and adjusted odds ratio (OR) with 95% confidence intervals

(CIs). Two authors independently used modified Jada scores and Newcastle-Ottawa Scale

(NOS)[19] scores to evaluate the quality of the RCTs (randomized controlled trials) and

observational studies, respectively. A Jada score >5 and an NOS score >7 were considered high

quality scores[20].

Statistical Analyses

RevMan5.3 (Review Manager [RevMan], version 5.3, Cochrane Collaboration) software was

used for statistical data processing, and the OR and 95% CIs were used to estimate the effect.

Heterogeneity among the studies was analyzed using the I^2 test with the following

interpretation: low heterogeneity, defined as $I^2 < 50\%$; moderate heterogeneity, defined as

$I^2 = 50\%$ to 75%; and high heterogeneity, defined as $I^2 > 75\%$[21]. The meta-analysis was

performed using the random effects model when the heterogeneity was more than 25%,

otherwise, the fixed effects model was applied. If there were ACEI and ARB subgroups in the

study, the combined OR and the 95% CIs were pooled using RevMan5.3 software. If the study

did not report the OR directly, count data for outcomes were used to generate unadjusted ORs

and 95% CIs. We also excluded reports with unadjusted ORs in the sensitivity analysis to

assess the robustness of our results. A funnel plot was used to test for the presence of

publication bias, and P value < 0.05 was considered statistically significant.

Results

Study Selection

The systematic search of the electronic databases identified 343 articles (PubMed=54,

EMBASE=112, Medrxiv=132, ArXiv=45). After excluding duplicates and title/abstracts

screened, 22 articles underwent a more detailed full-text assessment, after which a total of 11

articles with 33,483 patients were included[8, 10-17, 22, 23] (**Figure 1**).

Study Characteristics and Quality

Table 1 shows the basic characteristics of the included studies. Overall, the sample sizes

included in the articles ranged from 42 to 8,910, and ages ranged from 49 to 67 years old.

Among the 11 studies, 7 were published [8, 10, 12, 15-17, 23] and 4 were found on a preprint server[11,

[13, 14, 22]; 3 studies were based on the general population of COVID-19 patients [11, 12, 15] and 8

were based on patients with COVID-19 and hypertension[8, 10, 13, 14, 16, 17, 22, 23]. Four articles

reported the effect of ACEIs/ARBs on the level of inflammation[8, 10, 13, 17], two studies assessed

the risk of COVID-19 inflection[15, 16], and all included studies evaluated the severity of disease

or/and mortality[8, 10-17, 22, 23]. All included studies were observational studies. The NOS score of

all of the observational studies was > 6, indicating that all of the studies were of high quality

(**Table S2** Online Data Supplement).

Inflammation Level

There were 4 articles that evaluated the relationship between ACEI/ARB and level of

inflammation in patients with COVID-19[8, 13, 17, 22], but since the data were not reported in the

same unit across studies, they were not pooled; therefore, we conducted a systematic review

instead (**Table 2**). Meng et al[8]. first assessed the impact of RAAS inhibitors on the level of

inflammation in patients with COVID-19, and although they found no significant change in

CRP levels from the peripheral blood in patients receiving ACEI/ARB treatment, IL-6 levels

were significantly reduced, and the level of immune cells (CD3, CD8) counts were

significantly increased.

In contrast, another study showed that there was no difference in IL-6 levels, while

high-sensitivity C-reactive protein(CRP) test were significantly decreased[13]. A study that

included 342 COVID-19 patients with hypertension who had taken ACEI/ARB drugs found

that IL-6 levels were significantly reduced, while CRP level show no significantly different[17].

More recently, a multi-center study including 1,128 patients with hypertension and COVID-19

reported that the number of patients with high CRP values who had received ACEI/ARB

therapy showed no significantly different than in the patients who had not received ACEI/ARB

therapy[10].

Risk of Covid-19 Infection

Two[15, 16] studies including a total of 10,629 patients reviewed the risk of COVID-19 infection,

including one general population-based study and one study including patients with

hypertension. There was no significant increase in the risk of COVID-19 infection in patients

receiving ACEI/ARB therapy (OR=0.95, 95% CI: 0.89-1.02; P=0.14, I^2=0%), with no

significant heterogeneity (**Figure 2A**). Subgroup analysis based on population showed that the

results did not change in the general population (OR=0.96, 95% CI: 0.89-1.05; P=0.45, I^2=0%)

or in the hypertensive population (OR=1.00, 95% CI: 0.90-1.12; P=1.00) (**Figure 3A**).

Risk of Severe COVID-19

Eight studies assessed the relationship between the use of ACEI/ARB therapy and severe COVID-19[8, 11, 13-17, 22], with two reports in the general population[11, 15], and 6 reports in patients with hypertension[8, 13, 14, 16, 17, 22]. Compared with the non-ACEI/ARB group, the risk of severe COVID-19 infection decreased by 35% (OR=0.75, 95% CI: 0.59-0.96; P=0.02) in patients treated with an ACEI/ARB, with moderate heterogeneity (I^2=54%) (**Figure 2B**). However, in the sensitivity analysis, the results were not statistically significant when the unadjusted studies were excluded (OR=0.85, 95% CI: 0.66-1.08; P=0.19).

Subgroup analysis showed there was no statistical significant association between the risk of severe COVID-19 infection and ACEI/ARB use either in the general population (OR=0.85, 95% CI: 0.66-1.08; P=0.19, I^2=61%) or in hypertensive patients with (OR=0.73, 95% CI: 0.51-1.03; P=0.07, I^2=55%) (**Figure 3B**).

All-Cause Mortality

Seven studies reported all-cause mortality in 11,509 patients with COVID-19[8, 10, 12, 13, 17, 22, 23], and ACEI/ARB therapy was associated with a decreased risk of all-cause mortality (OR: 0.52, 95% CI: 0.35-0.79; P=0.002) with modest of heterogeneity (I^2=46%) (**Figure 2C**).There is no evidence of heterogeneity (I^2=1%) when one[23] study was excluded, with no change of the conclusion (OR: 0.46, 95% CI: 0.33-0.64; P<0.001). In the sensitivity analysis, the results were not statistically significant when the unadjusted studies were excluded (OR=0.34, 95% CI: 0.18-0.63; P=0.007, I^2=0%).

Only one[12] study reported all-cause mortality in COVID-19 patients in the general

population, and which found that ACEI/ARB therapy significantly decreased the mortality rate

of COVID-19 patients (OR: 0.31, 95% CI: 0.13-0.75; *P*=0.009). In hypertensive patients, the

use of an ACEI/ARB was associated with a decreased risk of all-cause mortality (OR: 0.57,

95% CI: 0.37-0.87; *P*=0.009) (**Figure 3C**).

Publication Bias

Publication bias was not detected in this study, as tests of publication bias are unsuitable when

the number of included studies for each outcome is less than 10[24].

Discussion

This study was the first systemic assessment of ACEI/ARB therapy and the clinical prognosis

of patients infected with COVID-19. We did not find an association between ACEI/ARB

therapy and the increased risk of COVID-19 infection either in general population or in

hypertensive patients. Moreover, we found that ACEI/ARB therapy could reduce the risk of

severe COVID-19; in a subgroup analysis, RAAS inhibitors reduced the risk of severe

COVID-19 by 15% and 27% in the general population and the hypertensive population,

respectively, although this benefit did not achieve statistical significance. Finally, ACEI/ARB

therapy reduced the rate of all-cause death in patients with COVID-19 by 48%, and in the

subgroup analysis, use of an ACEI/ARB reduced the all-cause mortality by 69% and 43% for

the general population and the hypertensive population, respectively.

Studies have shown that, similar to SARS-CoV, SARS-CoV-2 directly binds to the host

cell ACE2 receptor which is highly expressed in human lung tissue, gastrointestinal tract,

vascular endothelial cells, and arterial smooth muscle cells *in vivo*[1, 25]. Animal tests have

shown that the expression of ACE2 receptors is significantly increased during ACEI/ARB

therapy[5], leading some scholars to worry that the use of RAAS system inhibitors may

contribute to the spread of COVID-19 in the population[6]. On April 13th, 2020, the European

Society of Hypertension COVID-19 Task Force updated their recommendations on the use of

RAAS inhibitors in patients with COVID-19 and stated that the available evidence does not

support a deleterious effect of RAAS blockers in COVID-19 infections[5]. Our research also

determined that there is no significantly increased risk of infection with SARS-CoV-2 in

patients receiving ACE/ARB therapy. There are several possible explanations for these

findings. First, although some clinical studies have reported that RAAS inhibitors might

increase the expression of ACE2 receptors, these results are inconsistent[5] and most are small

studies with small sample sizes[26-28]. Furthermore, these studies detected circulating ACE2 or

soluble ACE2 receptor levels in the urine, and the level of soluble ACE2 receptors may not

accurately reflect the true level of ACE2 receptors in the organs and tissues[26-28]. One study

assessed the expression of ACE2 receptors in the duodenal and ileal tissues of 21 patients with

RAAS inhibition; the researchers found that the expression of ACE2 receptor mRNA and

protein increased significantly in the brush border of the small intestinal epithelial cells[29].

However, we know that SARS-CoV-2 is mainly transmitted through the respiratory tract

ACE2 receptors[1, 30], and to date, there have been no reports published on the expression of

ACE2 receptors in lung tissue after ACEI/ARB treatment, which is an important direction for

future study.

Our study found that RAAS inhibitors reduced the risk of severe COVID-19 infection by

47%, which might be related to the previously confirmed anti-inflammatory effects of RAAS

inhibitors[4]. ACEIs and AT1R inhibitors are commonly used RAAS system inhibitors that have

been widely used in the treatment of hypertension, diabetic nephropathy, and congestive heart

failure[3]. Lot of studies have shown that RAAS system is an important target for the treatment

of acute lung injury[31]. For example, a retrospective analysis showed that ACEIs and AT1R

inhibitors can also reduce the incidence of radiation pneumonitis[32]. In patients with severe

COVID-19, the levels of many pro-inflammatory factors (e.g., IL-6, IL-2, and TNF-α) were

significantly elevated, and levels of regulatory T cells decreased significantly[7]. A

non-randomized controlled clinical trial showed that the IL-6 inhibitor tocilizumab could

significantly reduce oxygen consumption, imaging abnormalities, and clinical prognosis in

patients with COVID-19[33]. Moreover, in patients receiving ACEI/ARB therapy, IL-6 and CRP

levels were significantly decreased, and the level of CD3 andCD8 increased significantly[8].

Overall, this evidence suggests that ACEI/ARB therapy might reduce lung injury and infection

severity by downregulating inflammation levels in patients with COVID-19.

Although we found use of ACEI/ARB was associated with the decreased risk of severe

COVID-19 in the overall analysis, this benefit did not reach statistical significance in the

sensitive analysis (with excluding unadjusted studies).Furthermore, subgroup analysis showed

the benefit was also not statistically significant either the general population or in patients with

hypertension. The possible reason for this result may be related to the limited sample size and

existence of many confounding factors in the baseline characteristics such as cardiovascular

disease, and the use of drugs may affect these results. Age has been proven to be a strong risk

factor for both infection with COVID-19 and a higher severity of COVID-19 infection[12].

Additionally, population-based studies have estimated that the prevalence of hypertension

range 9.5%-31.2% in COVID-19 patients, and many studies have shown that hypertension can

also significantly increase the severity of COVID-19[34]. Therefore, more studies with large

sample sizes with adjusted for these potential confounders are needed to clarify the impact of

ACEI/ARB use on the severity of COVID-19 in both the general population and in

hypertensive patients. Recently, one multi-center study that included 1,128 adult patients with

hypertension also showed that ACEI/ARB use reduced the risk of septic shock by 62% without

reducing the risk of other serious complications such as acute respiratory distress syndrome

and disseminated intravascular coagulation after matching baseline factors, suggesting that

ACEI/ARB therapy led to a different type of severity of COVID-19[10]. Therefore, the effect of

RAAS inhibitors on the severity of COVID-19 remains controversial and requires further

study.

In addition, we found that ACEI/ARB therapy reduced the risk of mortality by 47%, and

in the subgroup analysis, this mortality benefit remained significant for both in the general

population and in hypertensive patients. These results remained consistent in the sensitivity

analysis, which verified the robustness of our study. The number of studies included in the

subgroup of general population, however, was relatively limited (N=1)[12], and the case-control

study design increases bias (e.g., recall bias). Finally, we observed that 26.3% of the population

carriedhypertension[12], and studies have shown that COVID-19 patients with hypertension

receiving ACEI/ARB therapy have lower mortality compared with those non-hypertensive

individuals with COVID-19. It is therefore unclear whether the benefit was attributable to the

high proportion of patients with hypertension. Moreover, the study only adjusted for age and

sex and did not correct for enough of the potential confounding factors, such as drug use and

hypertension. Therefore, the benefits of RAAS inhibition in the general population with a

lower prevalence of hypertension still needs to be confirmed in future research.

Study Limitations

Our research has several limitations. First, all of the included articles were observational and

therefore cannot confirm the cause-and-effect relationship between ACEI/ARB therapy and

the clinical prognosis of patients with COVID-19; a large-scale RCT is needed to confirm our

results. Second, coexisting conditions such as hypertension have been shown a key prognostic

determinant (e.g. severity and mortality) in patients with COVID-19. The guidelines

recommend that patients with hypertension and COVID-19 continue their ACEI/ARB therapy,

and our study reinforced this recommendation and further showed that the use of ACEI/ARB

therapy might be associated with better clinical outcomes in the general population with

COVID-19 and in hypertensive patients with COVID-19. However, the benefit of RAAS

inhibitors in non-hypertensive patients might differ from those with hypertensive patients. Due

to data limitations, we cannot analyze the severity and clinical prognosis of ACEI/ARB

therapy in patients with COVID-19 without hypertension. Third, studies have shown that

ACEIs and ARBs may play different roles in patients with COVID-19, due to limited data, we

were unable to perform subgroup analyses of ACEIs and ARBs. Fourth, some characteristic

clinical values (e.g., drug variables) were missing. For example, the specific details of RAAS

inhibitors were lacking in all studies, which might have impact on our results. Fifth,

considering that all of the included studies were retrospective, the introduction of recall bias

was inevitable and may affect the reliability of the conclusions. Sixth, although we did not

register the protocol of this meta-analysis in the PROSPERO database, no relevant protocols of

this topic were found in this database at the writing of this review.

Conclusions

On the basis of the available evidence, we found that the use of ACEI/ARB therapy was not

associated with an increased risk of COVID-19 and that there was a decreased trend between

ACEI/ARB use and severe COVID-19 infection in both the general population and in patients

with hypertension, although this association was not statistically significant. The risk of

all-cause death was decreased with ACEI/ARB therapy in both the general population and in

patients with hypertension. Overall, RAAS inhibitors should be continued use in patients who

are at risk for, are being evaluated for, or have COVID-19. Our results need to be interpreted

with caution considering the potential for residual confounders, and more well-designed

studies that control clinical confounders are necessary to confirm our foundlings.

Acknowledgements

K.H. was responsible for the entire project and revised the draft. X.L. and Y.L. performed the

systematic literature review and drafted the first version of the manuscript. M.X., H.S, C.C

and Y.M reviewed, interpreted, and checked data. All authors took part in the interpretation

of the results and prepared the final version of the manuscript.

Sources of Funding

This work is supported by the National Natural Science Foundation of China [81530013, K.H.;

81600243, M.X.], Scientific and Technical Innovation Group of Jiangxi Province

[20181BCB24012, K.H.], Postgraduate Innovation Foundation of Jiangxi Province

[CX-2017198, X.L.].

Conflict of Interest

None

References

1. Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, Schiergens TS, Herrler G, Wu N-H, Nitsche A. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. Cell 2020.

2. Li W, Moore MJ, Vasilieva N, Sui J, Wong SK, Berne MA, Somasundaran M, Sullivan JL, Luzuriaga K, Greenough TC. Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. Nature 2003;**426**(6965):450-454.

3. Romero CA, Orias M, Weir MR. Novel RAAS agonists and antagonists: clinical applications and controversies. Nature Reviews Endocrinology 2015;**11**(4):242-252.

4. Paul M, Poyan Mehr A, Kreutz R. Physiology of local renin-angiotensin systems. Physiological reviews 2006;**86**(3):747-803.

5. Kreutz R, Algharably EAE, Azizi M, Dobrowolski P, Guzik T, Januszewicz A, Persu A, Prejbisz A, Riemer TG, Wang JG, Burnier M. Hypertension, the renin-angiotensin system, and the risk of lower respiratory tract infections and lung injury: implications for COVID-19. Cardiovasc Res 2020.

6. Garami AR. Preventing a covid-19 pandemic–Is there a magic bullet to save COVID-19 patients? We can give it a try. BMJ 2020;**368**:m810.

7. Qin C, Zhou L, Hu Z, Zhang S, Yang S, Tao Y, Xie C, Ma K, Shang K, Wang W. Dysregulation of immune response in patients with COVID-19 in Wuhan, China. Clinical Infectious Diseases 2020.

8. Meng J, Xiao G, Zhang J, He X, Ou M, Bi J, Yang R, Di W, Wang Z, Li Z, Gao H, Liu L, Zhang G. Renin-angiotensin system inhibitors improve the clinical outcomes of COVID-19 patients with hypertension. Emerg Microbes Infect 2020;**9**(1):757-760.

9. Coomes EA, Haghbayan H. Interleukin-6 in COVID-19: a systematic review and meta-analysis. MedRxiv 2020.

10. Zhang P, Zhu L, Cai J, Lei F, Qin JJ, Xie J, Liu YM, Zhao YC, Huang X, Lin L, Xia M, Chen MM, Cheng X, Zhang X, Guo D, Peng Y, Ji YX, Chen J, She ZG, Wang Y, Xu Q, Tan R, Wang H, Lin J, Luo P, Fu S, Cai H, Ye P, Xiao B, Mao W, Liu L, Yan Y, Liu M, Chen M, Zhang XJ, Wang X, Touyz RM, Xia J, Zhang BH, Huang X, Yuan Y, Rohit L, Liu PP, Li H. Association of Inpatient Use of Angiotensin Converting Enzyme Inhibitors and Angiotensin II Receptor Blockers with Mortality Among Patients With Hypertension Hospitalized With COVID-19. Circ Res 2020.

11. Bean D, Kraljevic Z, Searle T, Bendayan R, Pickles A, Folarin A, Roguski L, Noor K, Shek A, o'gallagher K, Zakeri R, Shah A, Teo J, Dobson RJB. Treatment with ACE-inhibitors is associated with less severe disease with SARS-Covid-19 infection in a multi-site UK acute Hospital Trust. 2020.

12. Mehra MR, Desai SS, Kuy S, Henry TD, Patel AN. Cardiovascular Disease, Drug Therapy, and Mortality in Covid-19. N Engl J Med 2020.

13. Yang G, Tan Z, Zhou L, Yang M, Peng L, Liu J, Cai J, Yang R, Han J, Huang Y, He S. Angiotensin II Receptor Blockers and Angiotensin-Converting Enzyme Inhibitors Usage is

Associated with Improved Inflammatory Status and Clinical Outcomes in COVID-19 Patients With Hypertension. 2020.

14. Liu Y, Huang F, Xu J, Yang P, Qin Y, Cao M, Wang Z, Li X, Zhang S, Ye L, Lv J, Wei J, Xie T, Gao H, Xu K-F, Wang F, Liu L, Jiang C. Anti-hypertensive Angiotensin II receptor blockers associated to mitigation of disease severity in elderly COVID-19 patients. 2020.

15. Mancia G, Rea F, Ludergnani M, Apolone G, Corrao G. Renin-Angiotensin-Aldosterone System Blockers and the Risk of Covid-19. N Engl J Med 2020.

16. Reynolds HR, Adhikari S, Pulgarin C, Troxel AB, Iturrate E, Johnson SB, Hausvater A, Newman JD, Berger JS, Bangalore S, Katz SD, Fishman GI, Kunichoff D, Chen Y, Ogedegbe G, Hochman JS. Renin-Angiotensin-Aldosterone System Inhibitors and Risk of Covid-19. N Engl J Med 2020.

17. Li J, Wang X, Chen J, Zhang H, Deng A. Association of Renin-Angiotensin System Inhibitors With Severity or Risk of Death in Patients With Hypertension Hospitalized for Coronavirus Disease 2019 (COVID-19) Infection in Wuhan, China. JAMA Cardiol 2020.

18. Moher D, Liberati A, Tetzlaff J, Altman DG. The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA Statement. Open Medicine 2009;**3**(3):e123-30.

19. Mcpheeters ML. Newcastle-Ottawa Quality Assessment Scale. 2012.

20. Liu X, Ma J, Huang L, Zhu W, Yuan P, Wan R, Hong K. Fluoroquinolones increase the risk of serious arrhythmias: A systematic review and meta-analysis. Medicine (Baltimore) 2017;**96**(44):e8273.

21. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. Bmj 2003;**327**(7414):557-560.

22. Zeng Z, Sha T, Zhang Y, Wu F, Hu H, Li H, Han J, Song W, Huang Q, Chen Z. Hypertension in patients hospitalized with COVID-19 in Wuhan, China: A single-center retrospective observational study. 2020.

23. Huang Z, Cao J, Yao Y, Jin X, Luo Z, Xue Y, Zhu C, Song Y, Wang Y, Zou Y, Qian J, Yu K, Gong H, Ge J. The effect of RAS blockers on the clinical characteristics of COVID-19 patients with hypertension. Ann Transl Med 2020;**8**(7):430.

24. Higgins JP, Green S. *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]The Cochrane Collaboration, 2011. Available from http://handbook.cochrane.org.chanpter: 10.4.3.1 Recommendations on testing for funnel plot asymmetry*; 2011.

25. Lan J, Ge J, Yu J, Shan S, Zhou H, Fan S, Zhang Q, Shi X, Wang Q, Zhang L. Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. Nature 2020:1-6.

26. Abe M, Oikawa O, Okada K, Soma M. Urinary angiotensin-converting enzyme 2 increases in diabetic nephropathy by angiotensin II type 1 receptor blocker olmesartan. Journal of the Renin-Angiotensin-Aldosterone System 2015;**16**(1):159-164.

27. Furuhashi M, Moniwa N, Mita T, Fuseya T, Ishimura S, Ohno K, Shibata S, Tanaka M, Watanabe Y, Akasaka H. Urinary angiotensin-converting enzyme 2 in hypertensive patients

may be increased by olmesartan, an angiotensin II receptor blocker. American journal of hypertension 2015;**28**(1):15-21.

28. Mariana CP, Ramona PA, Ioana BC, Diana M, Claudia RC, Stefan VD, Maria KI. Urinary angiotensin converting enzyme 2 is strongly related to urinary nephrin in type 2 diabetes patients. International urology and nephrology 2016;**48**(9):1491-1497.

29. Vuille-dit-Bille RN, Camargo SM, Emmenegger L, Sasse T, Kummer E, Jando J, Hamie QM, Meier CF, Hunziker S, Forras-Kaufmann Z. Human intestine luminal ACE2 and amino acid transporter expression increased by ACE-inhibitors. Amino acids 2015;**47**(4):693-705.

30. Guo Y-R, Cao Q-D, Hong Z-S, Tan Y-Y, Chen S-D, Jin H-J, Tan K-S, Wang D-Y, Yan Y. The origin, transmission and clinical therapies on coronavirus disease 2019 (COVID-19) outbreak–an update on the status. Military Medical Research 2020;**7**(1):1-10.

31. Imai Y, Kuba K, Rao S, Huan Y, Guo F, Guan B, Yang P, Sarao R, Wada T, Leong-Poi H. Angiotensin-converting enzyme 2 protects from severe acute lung failure. Nature 2005;**436**(7047):112-116.

32. Pinter M, Kwanten WJ, Jain RK. Renin–Angiotensin System Inhibitors to Mitigate Cancer Treatment–Related Adverse Events. Clinical Cancer Research 2018;**24**(16):3803-3812.

33. Xu X, Han M, Li T, Sun W, Wang D, Fu B, Zhou Y, Zheng X, Yang Y, Li X. Effective treatment of severe COVID-19 patients with tocilizumab. Proceedings of the National Academy of Sciences 2020.

34. Zheng Y-Y, Ma Y-T, Zhang J-Y, Xie X. COVID-19 and the cardiovascular system. Nature Reviews Cardiology 2020;**17**(5):259-260.

Figure Legends

Figure 1. PRISMA Flow Diagram

Figure 2. Summary of the associations between use of ACEI/ARB and clinical outcomes among patients with COVID-19. A. Risk of COVID-19 infection; B. Risk of severe COVID-19 infection; C. All-cause death.

*severe COVID-19 or death

Figure 3. Subgroup analysis of the associations between use of ACEI/ARB and clinical outcomes among patients with COVID-19 stratified by general population and hypertensive population: A. Risk of COVID-19 infection; B. Risk of severe COVID-19 infection; C. All-cause death.

1   *severe COVID-19 or death

3   **Nonstandard Abbreviations and Acronyms:** COVID-19: Coronavirus disease 2019; OR:

4   Odd risk; CI: Confidence interval; ACE2: angiotensin-converting enzyme 2; ACEI: ACE

5   inhibitors; ARB: ACE angiotensin-receptor blockers; NOS: Newcastle-Ottawa Scale score;

6   RAAS: renin–angiotensin–aldosterone system; SARS: Severe acute respiratory syndrome.

A: COVID-19 inflection

B:Severe COVID-19

C: All-cause death

A: COVID-19 inflection

B: Severe COVID-19

C: All-cause death