

---

# AVOIDING COVID-19: AEROSOL GUIDELINES

---

**Matthew J. Evans**

Department of Physics  
Massachusetts Institute of Technology  
Cambridge, MA 02139  
m3v4n5@mit.edu

May 21, 2020

## ABSTRACT

The COVID-19 pandemic has brought into sharp focus the need to understand respiratory virus transmission mechanisms. In preparation for an anticipated influenza pandemic, a substantial body of literature has developed over the last few decades showing that the short-range aerosol route is an important, though often neglected transmission path. We develop a simple mathematical model for COVID-19 transmission via aerosols, apply it to known outbreaks, and present quantitative guidelines for ventilation and occupancy in the workplace.

**Keywords** SARS-CoV-2 · COVID-19 · coronavirus

## 1 Introduction

The world is learning to navigate the COVID-19 pandemic and a great deal of information about the disease is already available [1, 2, 3]. In order to adjust to this new reality it is important to understand what can be done to avoid infection, and to avoid infecting others.

SARS-CoV-2, the coronavirus which causes COVID-19, is thought to be transmitted via droplets, surface contamination and aerosols [4, 5, 6, 7, 8, 9]. Contact tracing of COVID-19 outbreaks points to aerosols as a possibly dominant mechanism of transmission since most outbreaks occur in closed environments, while they are rare in environments which disfavor aerosol transmission (e.g., outdoors) [10]. Furthermore, infection via inhalation of aerosols and small droplets dominates large droplet exposure in most cases for physical reasons linked to complex, but well understood fluid dynamics [11, 12, 13, 14]. Despite this fact, there is a common misconception that aerosol transmission implies efficient long-range transmission (as in measles), and thus that the absence of long-range transmission implies an absence of aerosol transmission. The truth is more nuanced, and includes the possibility of a dominant short-range aerosol path limited by pathogen dilution, deposition, and decay [15].

The production of aerosolized viruses by a contagious individual occurs naturally as a result of discrete events (e.g., coughing or sneezing), or through a continuous process like breathing or talking [16]. Droplets which are formed with diameters less than  $\sim 50 \mu\text{m}$  quickly lose most of their water to evaporation, shrinking by a factor of 2-3

in diameter and becoming “droplet nuclei”. These fine particles settle very slowly and mix with the air in the environment [17].

Aerosol transmission happens when a susceptible individual inhales these now sub- $20 \mu\text{m}$  droplet nuclei that are suspended in the air around them [18, 19]. This is thought to be the dominant transmission mechanism for influenza and rhinovirus [18, 20] and possibly also for SARS and MERS [21]. For influenza, it has been shown that aerosol particles as small as  $1.5 \mu\text{m}$  are sufficient for transmission [22]. Furthermore, it is known from other viral respiratory diseases that aerosol exposure can result in infection and illness at much lower doses than other means (e.g., nasal inoculation) [23].

In line with current recommendations [24], we will assume that hand-hygiene protocols are being followed sufficiently well to ensure that surface contamination is sub-dominant. We will also assume that contagious individuals are wearing some kind of face covering which is sufficient to disrupt the momentum of any outgoing airflow [25] and catch large droplets [26]. These actions leave aerosols, investigated here, as the dominant transmission mechanism.

In the following sections we present a simple model for aerosol transmission, apply this model to known outbreaks, and develop guidelines for reducing the probability of transmission in the workplace. This analysis is done under the assumption that all workers are wearing some form of face covering (“mask”), however, since effective filtration is difficult to achieve, we assign no protective value to the wearer for mask use.

## 2 Infection Model

The probability of developing an infection  $P_{\text{inf}}$  given exposure to a volume  $V$  of saliva with a concentration  $\rho$  of virions is

$$P_{\text{inf}} = 1 - e^{-\rho V/N_{\text{inf}}} \quad (1)$$

where the infectivity  $N_{\text{inf}}$  is the number of virions needed to make infection likely in the lungs for aerosol inhalation [23, 27]. The infectivity value  $N_{\text{inf}}$  includes probable deposition location (i.e., upper vs. lower respiratory tract) and local deposition efficiency (e.g., not all inhaled droplets are deposited in the lungs [23, 28]). It should be noted, however, that small doses are less likely to cause *illness* than what is indicated by Eqn. 1 [23, 29].

Rather than work explicitly with the probability of infection  $P_{\text{inf}}$ , we will use the product of the viral dose and the infectivity

$$D_{\text{inf}} = \rho V/N_{\text{inf}} \quad (2)$$

as a proxy, and refer to it as the “infective dose”. Note that the infectivity and the viral concentration always appear together here, so only their ratio  $\rho/N_{\text{inf}}$  is important in our model. While  $\rho$  has been measured for COVID-19 [30, 31],  $N_{\text{inf}}$  has not been directly measured. We estimate  $N_{\text{inf}}$  from influenza and other coronaviruses [27], and check this value by applying our model to known outbreaks (see appendix A).

## 3 Aerosol Infective Dose Model

In this section we compute the infective dose which results from the presence of a contagious individual in various scenarios. This informs prescriptions for the maximum time a susceptible individual may be exposed to potentially contaminated air, or the minimum time between occupants in the same space. Each of these calculations uses the values in Table 1 and concludes with an exposure time limit for an infectious dose less than  $D_{\text{inf}}^{\text{max}} = 0.1$  such that the accumulation of 4 such does results in a 10% probability of infection (see section 6).

### 3.1 Steady State in a Room

A contagious individual will shed virions into the room they occupy through breathing, talking, coughing, etc. The aerosolized virion concentration in a room can be expressed in the form of a differential equation as

$$\frac{d\rho_A(t)}{dt} = \rho_0 \frac{r_{\text{src}}}{V_{\text{room}}} - \rho_A(t) \left( \frac{1}{\tau_{\text{room}}} + \frac{1}{\tau_a} \right) \quad (3)$$

where  $\rho_0$  is the nominal viral concentration in saliva, which is emitted in aerosol form at a rate of  $r_{\text{src}}$ , and  $\tau_a$  is the viral decay time in aerosol form. The air cycle-time in the room is

$$\tau_{\text{room}} = V_{\text{room}}/r_{\text{room}} \quad (4)$$

where  $r_{\text{room}}$  is rate at which air is removed from the room (or filtered locally). The steady-state concentration is

$$\rho_A^{\text{SS}} = \rho_0 \frac{r_{\text{src}}}{V_{\text{room}}} \frac{\tau_{\text{room}}\tau_a}{\tau_{\text{room}} + \tau_a}. \quad (5)$$

and has units of viral copies per liter of air.

Breathing air in a room with an aerosol virus concentration  $\rho_A$  will cause an accumulation of exposure  $N_A$  (number virions) proportional to the time passed in that room  $t$

$$N_A = \rho_A r_b t \quad (6)$$

such that

$$D_{\text{inf}}^{\text{SS}} = \frac{\rho_0}{N_{\text{inf}}} \frac{r_{\text{src}}}{V_{\text{room}}} \frac{\tau_{\text{room}}\tau_a}{\tau_{\text{room}} + \tau_a} r_b t. \quad (7)$$

If the room is well ventilated (i.e.,  $\tau_{\text{room}} \ll \tau_a$ ) the infective dose is approximately

$$D_{\text{inf}}^{\text{SS}} \simeq 0.01 \frac{r_{\text{src}}}{1 \text{ nL/min}} \frac{1 \text{ m}^3/\text{min}}{r_{\text{room}}} \frac{t}{1 \text{ min}} \quad (8)$$

for an office, lab or bathroom occupied by a contagious individual.

The infective dose crosses our example event exposure threshold of  $D_{\text{inf}}^{\text{max}} = 0.1$  in an office-like space ( $r_{\text{room}} \sim 10 \text{ m}^3/\text{min} \sim 350 \text{ cfm}$ ) after 100 minutes,

$$t_{\text{max}}^{\text{Room}} \simeq 100 \text{ min} \frac{1 \text{ nL/min}}{r_{\text{src}}} \frac{r_{\text{room}}}{10 \text{ m}^3/\text{min}}, \quad (9)$$

while in small spaces with less ventilation (e.g., a car or bathroom with  $r_{\text{room}} \sim 1 \text{ m}^3/\text{min}$ ) the dose would cross the threshold after only 10 minutes. We have assumed that the contagious individual is quietly working such that  $r_{\text{src}} = V_b$ , but if they are talking (e.g., in video conference), or coughing occasionally, the source term may be higher by an order of magnitude or more (e.g.,  $r_{\text{src}} \gtrsim V_t$ ).

### 3.2 Steady State in a Building

When a susceptible individual occupies a room that is connected via the HVAC system to a room occupied by a contagious individual, there is the possibility of aerosols passing through the HVAC system [7, 38].

However, the volume of a droplet is proportional to its diameter *cubed*, which leads to the majority of the viral dose being delivered in the  $\sim 10 \mu\text{m}$  droplet nuclei [17, 22], which are filtered by HVAC systems. Most HVAC filters will remove the majority of the viral load associated with COVID-19 transmission, and a high quality filter (i.e., MERV 12) is sufficient to remove  $> 90\%$  of the larger droplet nuclei [39]. The remaining aerosols are further diluted by fresh make-up air, and then spread among all of the air spaces associated with the HVAC system. The associated infective dose in rooms connected to a room occupied by a contagious individual will be roughly

$$D_{\text{inf}}^{\text{Sys}} = (1 - f_{\text{sys}}) \frac{\rho_0}{N_{\text{inf}}} \frac{r_{\text{src}}}{V_{\text{sys}}} \frac{\tau_{\text{sys}}\tau_a}{\tau_{\text{sys}} + f_{\text{sys}}\tau_a} r_b t \quad (10)$$

where  $f_{\text{sys}}$  is the fraction of the viral dose which is removed by the filters or displaced by make-up air, and  $\tau_{\text{sys}} = r_{\text{sys}}/V_{\text{sys}}$  is the time required to cycle the entire air volume through the HVAC system.

Description	Symbol	Value	Reference
Viral Load in Saliva	$\rho_0$	1000 /nL	[30, 31, 32]
Sneeze Aerosol Volume	$V_s$	1 $\mu$ L	[33, 34]
Cough Aerosol Volume	$V_c$	100 nL	[33, 34]
Talking Aerosol Volume	$V_t$	10 nL/minute	[33, 34]
Breathing Aerosol Volume	$V_b$	1 nL/minute	[33, 34]
Aerosol Decay Time	$\tau_a$	2 h	[35, 36]
Breathing Rate	$r_b$	10 L/minute	[37]
Respiratory Infectivity	$N_{\text{inf}}$	1000	[27, 23]
Maximum Exposure and Minimum Wait Times			
Room Steady State	$t_{\text{max}}^{\text{Room}} \simeq 100 \text{ min}$	$\frac{1 \text{ nL/min}}{r_{\text{src}}} \frac{r_{\text{room}}}{10 \text{ m}^3/\text{min}}$	Eqn. 9
System Steady State	$t_{\text{max}}^{\text{Sys}} \simeq 100 \text{ min}$	$\frac{0.1}{1-f_{\text{sys}}} \frac{10 \text{ nL/min}}{r_{\text{src}}} \frac{f_{\text{sys}} r_{\text{sys}}}{100 \text{ m}^3/\text{min}}$	Eqn. 12
Occupancy Wait Time	$t_{\text{min}}^{\text{TE}} \simeq \tau_{\text{room}} \ln \left( 100 \frac{V_{\text{src}}}{1 \mu\text{L}} \frac{1 \text{ m}^3/\text{min}}{r_{\text{room}}} \right)$		Eqn. 15
Passage Wait Time	$t_{\text{min}}^{\text{PS}} \simeq \tau_{\text{room}} \ln \left( 100 \frac{V_{\text{src}}}{1 \mu\text{L}} \frac{10 \text{ m}^3}{V_{\text{room}}} \frac{\Delta t}{1 \text{ min}} \right)$		Eqn. 17
Flow Rate Conversion		$1 \text{ m}^3/\text{min} \simeq 35 \text{ cfm}$	

Table 1: Parameters used in the aerosol transmission model. See section 5 for more information.

In the well ventilated limit, as above, the infective dose is

$$D_{\text{inf}}^{\text{Sys}} \simeq 10^{-4} \frac{1-f_{\text{sys}}}{0.1} \frac{100 \text{ m}^3/\text{min}}{f_{\text{sys}} r_{\text{sys}}} \frac{r_{\text{src}} t}{10 \text{ nL}} \quad (11)$$

and the maximum occupancy time for  $D_{\text{inf}}^{\text{Sys}} < 0.01$  is

$$t_{\text{max}}^{\text{Sys}} \simeq 100 \text{ min} \frac{0.1}{1-f_{\text{sys}}} \frac{10 \text{ nL}/\text{min}}{r_{\text{src}}} \frac{f_{\text{sys}} r_{\text{sys}}}{100 \text{ m}^3/\text{min}}. \quad (12)$$

We have set the  $r_{\text{src}}$  scale at 10 nL/min to allow for talking and an occasional coughs from the contagious individual. The maximum dose used in this example is 0.01 instead of 0.1 due to the likely larger cohort exposed (see section 6).

To understand why transmission through HVAC systems appears to be rare we note that even medium quality air filters (MERV 9) provide better than 99% filtering after some loading [39]. The value we use here assumes little or no fresh make-up air and is representative of an unloaded (i.e., clean) air filter, which is the *worst case*. Also, in buildings where the HVAC system does not recirculate air this type of transmission cannot occur (i.e.,  $f_{\text{sys}} = 1$ ).

### 3.3 Transient Occupancy and Events

Some spaces are occupied briefly by many people (e.g., bathrooms) and many not have time to come to steady state. Coughing and sneezing events can also cause a transient increase in the viral concentration in a room. The viral dose associated with these types of transients can be quantified as

$$D_{\text{inf}}^{\text{TE}} = \frac{\rho_0}{N_{\text{inf}}} \frac{V_{\text{src}}}{V_{\text{room}}} r_b \int_{t_1}^{t_2} e^{-t \frac{\tau_{\text{room}} + \tau_a}{\tau_{\text{room}} \tau_a}} dt \quad (13)$$

where the transient event occurs at time  $t = 0$  and exposure is from time  $t_1$  to  $t_2$ .

We can estimate when a room is “safe” for a new occupant by computing the dose in the limit of  $t_2 \rightarrow \infty$ ,

$$D_{\text{inf}}^{\text{TE}} \simeq \frac{\rho_0}{N_{\text{inf}}} \frac{V_{\text{src}}}{V_{\text{room}}} \tau_{\text{room}} r_b e^{-t_1/\tau_{\text{room}}} \quad (14)$$

where we have again assumed that  $\tau_{\text{room}} \ll \tau_a$ . This leads to a minimum wait time of

$$t_{\text{min}}^{\text{TE}} \simeq \tau_{\text{room}} \ln \left( 100 \frac{V_{\text{src}}}{1 \mu\text{L}} \frac{1 \text{ m}^3/\text{min}}{r_{\text{room}}} \right) \quad (15)$$

where we have set the  $V_{\text{src}}$  scale at 1  $\mu$ L to allow for some coughing or a sneeze from the previous occupant.

As a concrete example, for a 70 square-foot bathroom with a 70 cfm fan in operation (i.e.,  $\tau_{\text{room}} \simeq 10$  minutes and  $r_{\text{room}} = 2 \text{ m}^3/\text{min}$ ), Eqn. 15 gives a 40 minute wait time between occupants.

Going back to Eqn. 13, we can also compute the exposure due to brief passage through a common space in which  $\Delta t = t_2 - t_1 \ll \tau_{\text{room}}$  (e.g., a halls, stairwells and elevators),

$$D_{\text{inf}}^{\text{PS}} \simeq \frac{\rho_0}{N_{\text{inf}}} \frac{V_{\text{src}}}{V_{\text{room}}} r_b \Delta t e^{-t_1/\tau_{\text{room}}}. \quad (16)$$

The minimum time between occupants for  $D_{\text{inf}}^{\text{PS}} < 0.01$  is

$$t_{\text{min}}^{\text{PS}} \simeq \tau_{\text{room}} \ln \left( 100 \frac{V_{\text{src}}}{1 \mu\text{L}} \frac{10 \text{ m}^3}{V_{\text{room}}} \frac{\Delta t}{1 \text{ min}} \right), \quad (17)$$

which will give negative values for large spaces or short passage times, indicating that no wait is necessary. However, this assumes well mixed air, so some mixing time is necessary to avoid a “close encounter” as described in the next section. For small spaces with poor circulation (e.g., elevators), on the other hand, long wait times are required.

### 3.4 Close Encounters

Mixing of aerosols into an airspace may require a few air cycle-times before a fairly uniform concentration can be assumed [40, 41]. In order to understand the potential infection risk associated with close proximity, we can make a very rough estimate of the exposure in the vicinity of a mask-wearing cougher. The cough momentum will be disrupted by the mask, but the aerosols will exit the mask on essentially all sides [25, 16] resulting in a cloud around the cougher that will then either move with local air currents or rise with the cougher's body plume [42]. We assume that the large droplets are trapped in the mask and the small droplets settle out around the cougher, while aerosols are dispersed into the air around the cougher. The infective dose at a distance  $d$  for spherical dispersion is

$$D_{\text{inf}}^{\text{CE}} = \frac{\rho_0 V_c}{N_{\text{inf}}} \frac{3}{4\pi d^3} r_b t. \quad (18)$$

For a typical cough this leads to

$$D_{\text{inf}}^{\text{CE}} \simeq 0.04 \left( \frac{1 \text{ m}}{d} \right)^3 \frac{t}{10 \text{ sec}} \quad (19)$$

which suggests that an infection risk is still present at short distances, even if the cougher is wearing a mask [43]. This analysis is clearly oversimplified, as details of the cough, the mask, and local air currents will prevent isotropic dispersion.

## 4 Mitigation Measures

The calculations in the previous section suggest a variety of possible mitigation measures. This section briefly describes some means of reducing the probability of aerosol infection.

### 4.1 Masks and Respirators

Masks and respirators can be used to reduce the probability of infection, both for the wearer and for the people they interact with. Standard surgical masks provide some level of protection [44, 45], but they are far less effective filters than N95 respirators (a factor of 2–10 for surgical masks, and 8–80 for N95s) [46]. N95 masks have been demonstrated to be effective in preventing COVID-19 in a hospital setting [47], but benefiting from them requires proper fit and user compliance [48, 49, 50, 51], which suggests that widespread use is likely to be ineffective [52, 53].

Arguably more important than the mask one wears are the masks worn by the people one interacts with, both directly (e.g., in conversation) and indirectly (e.g., by sharing a common space) [54]. Droplets produced while breathing, talking and coughing can be significantly reduced by mask usage [55, 56, 16], but much of the air expelled while coughing and sneezing goes around the mask [25].

### 4.2 HVAC and Portable Air Filters

Increasing air exchange rates in an HVAC system, and avoiding recirculation can both be used to reduce aerosol concentrations indoors. Consistent use of local ventilation (e.g., bathroom fans) can also help to avoid infection. In buildings and spaces where these measures are not available or not sufficient, local air filters (a.k.a., air purifiers) can be used to increase the production of clean air into the space. Small stand-alone units which filter  $10 \text{ m}^3/\text{min}$  or more are readily available and could help to increase safety in elevators, bathrooms and offices.

As noted in sections 3.2 and 5, the particle size of interest is greater than  $1 \mu\text{m}$ , so special filtering technology is *not* required [39]. Care should be taken, however, when changing filters both in stand-alone units and HVAC systems as they may contain significant viral load. Stand-alone units should be disabled for at least 3 days before changing the filter to allow time for viral infectivity to decay [35]. Building HVAC filters should be changed with proper personal protective equipment, as recommended by the CDC [39].

### 4.3 Clean Rooms and HEPA Filters

Many laboratory spaces are outfitted with HEPA filters to provide clean air for laboratory operations. Clean rooms offer a clear advantage over other spaces as they are designed to provide air which is free of small particles.

If the HEPA filters are part of a recirculating air cycle, Eqn. 12 can be used with  $f_{\text{sys}} \gtrsim 0.999$ , allowing for essentially indefinite exposure times. It should be noted, however, that HEPA filters require lower air-speeds than those offered by typical HVAC systems and as such are not a “drop in replacement” option.

Laboratories that use portable clean rooms in large spaces can be treated in a similar manner, using Eqn. 11 for the air inside the clean room and Eqn. 8 to represent the clean air supplied to the laboratory space outside of the portable clean room.

Any clean room environment is likely to provide sufficient air flow to make aerosol infection very unlikely in the well-mixed approximation used in the equations suggested above. This will leave close encounters, as described in section 3.4, as the dominant infection path. If there are multiple occupants in a clean room they should avoid standing close to each other, or being “down wind” of each other [57].

### 4.4 UV-C Lighting

Illumination of the air-space in patient rooms with ultraviolet light has been shown to dramatically reduce viral longevity in aerosol form, and thereby prevent infection [15, 58]. This could be used to decrease  $\tau_a$  in spaces where increasing air circulation (i.e., reducing  $\tau_{\text{room}}$ ) is impractical. Care is required to avoid exposing occupants to UV-C radiation, which is required for viral deactivation but can




Figure 1: **Left:** Probability that the actual infective dose in any given encounter exceeds the nominal dose by some factor (i.e.,  $D_{inf} > CD_{inf}^0$  as a function of the dose ratio  $C$ ). The three curves allow some or all transmission parameters to vary randomly (see section 5 and appendix B). For instance, allowing all parameters to vary (blue trace) the probability of having  $D_{inf} > D_{inf}^0$  is about 10%,  $D_{inf} > 10 D_{inf}^0$  will happen in about 4% of cases, while 1% of cases will have  $D_{inf} \gtrsim 100 D_{inf}^0$ . Choosing the nominal viral load  $\rho_0$  equal to the median value of 10 /nL (rather than the 90<sup>th</sup> percentile as we have) would shift the blue curve to the right by a factor of 100. **Right:** Probability of infection as a function of nominal infective dose  $D_{inf}^0$ . The three solid curves relate  $P_{inf}$  to  $D_{inf}^0$  allowing some or all transmission parameters to vary randomly (as in the left figure), while the dashed curve (grey) shows  $P_{inf}$  with fixed parameters as in Eqn. 1. Allowing all parameters to vary, the probability of infection for a nominal dose of 1 is 14%. A factor of 10 lower  $D_{inf}^0$  reduces the probability of infection to 6%, and  $D_{inf}^0 \lesssim 3 \times 10^{-3}$  is required to reduce the probability of infection below 1%. Choosing the nominal viral load  $\rho_0$  equal of 10 /nL rather than 1000 /nL would reduce  $D_{inf}^0$  by 100 relative to our calculations in section 3 and shift the blue curve to the left by 100, such that  $P_{inf}$  would remain unchanged. The “random  $V_{src} N_{inf}$ ” curve is used in appendix A to evaluate the probability of infection in cases where  $\rho$  is assumed to be high.

be harmful to the eyes and skin. This could be done geometrically (e.g., only illuminate spaces above 2.5 m), or actively with motion sensors.

## 5 Model Parameters

The parameters used in this model are presented in Table 1. All of these parameters vary between individuals and events, and as such the values given here are intended to serve as a means of making rough estimates which can guide decision making. This section describes the provenance of these values and their variability.

The volume of saliva produced in a variety of activities is used to understand the emission of virions into the environment (known as “viral shedding”). The values given here are for “typical” individuals and behaviors, and actual values for any given person or event may vary by an order of magnitude [16, 33, 34, 59, 60]. Droplets produced while speaking, for instance, depend on speech loudness; speaking loudly, yelling or singing can produce an order of magnitude more saliva than speaking normally [61]. We are careful to avoid quantifying saliva production in terms

of the number of droplets produced, since the large droplet-nuclei are relatively small in number but carry most of the virions (i.e., slope of the number distribution is too shallow to compensate the fact that volume goes with diameter cubed). For a viral load of 1000 /nL, a 1  $\mu$ m droplet nucleus, for example, has only a 1% probability of containing a single virion [61].

The volume of air exchanged with the environment while breathing (“Breathing rate”  $r_b$ ) scales roughly with the weight of an individual and may vary by a factor of a few. As such, this is a relatively well determined parameter and we use only the nominal value in our calculations. The assumption is that an individual’s tidal volume is about 0.5 L, and the breathing cycle has a 3 s period [37].

The infectivity decay time of SARS-CoV-2 in aerosol form has been measured and found to be at least a few hours. Note that we are using the  $1/e$  decay time, not the half-life, for ease of computation. A separate decay time-scale is set by the settling time of the larger aerosol particles which contain the majority of the viral load. Using the “continuous fallout” model presented in [11] and the data presented in [61] we estimate this as

$\tau_{\text{fall}} \simeq u_{\text{settle}} V_{\text{room}} / A_{\text{room}} \sim 1$  hour for a room with a 3 m ceiling. In most cases, both the infectivity decay time and the settling decay time are long enough that they become irrelevant as the ventilation system is responsible for removing most of the aerosols.

The viral load in saliva,  $\rho$ , has been seen to vary by more than 8 orders of magnitude in individuals that test positive for COVID-19 [30, 31]. Roughly 90% of individuals tested have a viral load less than  $\rho_0 = 1000$  /nL, while 1% may have a viral load above  $3 \times 10^4$  /nL [31]. Viral load is seen to decline after symptom onset, so the pre-symptomatic viral load may be on the high-end of the distribution [30, 32, 62]. We use  $\rho_0 = 1000$  /nL for our nominal dose calculations because it results in a rough estimate of the probability of infection relative to the full distribution for probabilities of a few percent (i.e., the “random  $\rho V_{\text{src}} N_{\text{inf}}$ ” curve is close to the other curves around  $D_{\text{inf}}^0 \sim 0.03$  in Fig. 1, right). This choice does not impact the final probability of infection shown in Fig. 1, since a different choice of  $\rho_0$  would shift the “random  $\rho V_{\text{src}} N_{\text{inf}}$ ” curve to compensate.

The respiratory infectivity,  $N_{\text{inf}}$ , is not well known for SARS-CoV-2, but it has been measured for SARS [27], other coronaviruses, and influenza [23]. Variation by an order of magnitude among individuals is observed both for coronaviruses and influenza. The number we use,  $N_{\text{inf}} = 1000$ , is intended to be “typical” for coronaviruses and represents roughly 100 “plaque forming units” (PFU) each of which is roughly 10 virions [27].

## 6 Risk Assessment

Many alternatives can be expressed in terms of ratio of the resulting infective dose (e.g., doubling the airflow in a room, or comparing the dose in an office with that of a stairwell), such that the exact parameter values used are not important. For computations that require an absolute evaluation of risk, we compute the probability of infection given the distribution of  $D_{\text{inf}} \propto \rho_0 V_{\text{src}} / N_{\text{inf}}$ , with  $V_{\text{src}} = 1$  nL, which is common to all dose calculations (see Fig. 1).

The “random  $\rho V_{\text{src}} N_{\text{inf}}$ ” curve in Fig. 1 can be used to estimate the probability of a contagious individual causing infection in the people they interact with, given a “nominal infective dose”  $D_{\text{inf}}^0$  computed with the parameters in Table 1. For instance, if  $D_{\text{inf}}^0$  is the sum of 10 encounters like those described in section 3, each of which has  $D_{\text{inf}} \lesssim 2 \times 10^{-3}$  such that  $D_{\text{inf}}^0 \sim 0.02$ , then the probability of infection is 3%. The infection probability is fairly large despite the tiny *nominal* infective dose because of the non-negligible probability of the contagious individual having a high value of  $\rho$  and thereby delivering  $D_{\text{inf}} \gg D_{\text{inf}}^0$  (see Fig. 1, left). If the contagious individual delivers  $D_{\text{inf}}^0 \sim 0.02$  to each of 10 susceptible individuals (i.e. an 11 person cohort), the probability of transmission

to at least one person is nearly 30%:

$$P_{\text{trn}} = 1 - (1 - P_{\text{inf}})^{N_{\text{coh}} - 1} \simeq (N_{\text{coh}} - 1) P_{\text{inf}} \quad (20)$$

where  $N_{\text{coh}}$  is the number of people in the cohort and  $P_{\text{inf}}$  comes from Fig. 1, right.

The above example shows that limiting the number of potential transmission events between members of a cohort, and limiting the size of a cohort, are both important to minimizing  $P_{\text{trn}}$ . If only 4 encounters are allowed per person, and the cohort is kept to 3 people,  $D_{\text{inf}}^0 \lesssim 0.1$  per encounter gives a 20% probability of transmission.

The distribution of virions via a building’s HVAC system, discussed in section 3.2, should also be considered as it represents an interaction with a possibly unintended cohort (occupants of rooms connected via the HVAC system). In terms of the previous examples, this behaves as a single encounter with a large cohort.

An acceptable workplace infection risk could be chosen such that an infected employee has only a small chance of infecting another employee during their pre-symptomatic contagious period. The objective is to ensure that the contribution to the epidemic’s  $R_0$  value from the workplace is small, meaning that an employee is more likely to be infected in other activities (i.e., at home). The second example described above (with  $P_{\text{trn}} = 0.2$ ) would contribute 0.2 to  $R_0$ , meaning that for a decaying epidemic with  $0.5 < R_0 < 1$ , an employee would be 2 to 5 times more likely to be infected away from work than at work.

## 7 Conclusions

The new world of COVID-19 is here and we will all have to learn to live in it. Understanding the dangers of this new world and how to navigate them will be necessary as we come out of our houses and return to our workplaces.

The calculations presented here suggest that keeping the risk of infection low in the workplace will require both mitigation techniques which lower the viral dose encountered by susceptible individuals, and small interaction cohorts to avoid large outbreaks should a highly-contagious individual come to work (i.e., a pre-symptomatic “super-spreader”).

Some broad guidelines which we draw from the above analysis are:

1. Aerosol build-up in closed spaces should be treated with care. Avoiding infection requires good ventilation and/or short exposure times. Generally, office spaces should not be occupied by more than one person (see Eqn. 7).
2. Small or poorly ventilated common spaces where many people spend time (i.e., bathrooms and elevators) are of particular concern. At least 4 ventilation cycle times should be allowed to pass between occupants (see Eqns. 13 and 16). Con-

sider increasing airflow and adding local air filters to increase the rate of introduction of clean air.

3. Recirculation in HVAC systems should be avoided if possible. High quality filters (e.g., MERV 12) should be used in recirculating HVAC systems and office use should be kept to a minimum, to avoid transmission through the air circulation (see Eqn. 10).
4. Mask use is critical when a minimum of 2 m interpersonal distance cannot be maintained, but is not sufficient to prevent infection at distances less than 1 m. Leakage of aerosols around the mask may contaminate the air volume around a contagious individual (see Eqn. 19).
5. If possible, workers should be divided into groups (“cohorts”) consisting of less than 5 members.

Potential transmission events (e.g., sharing a common space, such as a bathroom or elevator) should be eliminated between members of different cohorts (see Eqn. 20).

## 8 Acknowledgements

The author would like to thank Dr. Lisa Barsotti and Prof. Paola Cappellaro for their help with construction and editing of this article. The analysis presented here would not have been possible without the inspiration provided by Dr. Dawn Mautner, the information supplied by Prof. Peter Fisher, and the many hours of quiet time for which I have Lisetta Turini to thank. An early draft of [57] and feedback from the authors of that work were both very helpful and much appreciated. The author is also very grateful for the computing support provided by The MathWorks, Inc.

## A Outbreaks

Documented outbreaks offer a means for checking the plausibility of our aerosol transmission model. In this section we select a few outbreaks where transmission via aerosols appears likely (e.g., distances were too large for droplets, or the pattern of infection followed the airflow).

Information about the outbreaks is, however, limited, and the viral load of the contagious individual is not known, so these comparisons only provide weak constraints on the model parameters. In particular, reported outbreaks are likely to involve unusually contagious individuals (a.k.a. “super-spreaders”), with  $\rho_0 \gtrsim 1000/\text{nL}$  (roughly 10% of cases), so we use that value in these computations. Since this fixes the value of  $\rho$ , we use the “random  $V_{\text{src}}N_{\text{inf}}$ ” curve in Fig. 1 to compute the probability of infection.

### A.1 Guangzhou Restaurant

[63] documents a COVID-19 outbreak associated with air flow in a restaurant (Guangzhou, China). The space defined by the wall mounted AC unit is  $16\text{ m}^2$ . Assuming a modest air cycle-time of 30 minutes ( $r_{\text{room}} \sim 2\text{ m}^3/\text{min}$ ), a somewhat talkative contagious individual ( $r_{\text{src}} \sim 4\text{ nL}/\text{min}$ ), and a 30 minute overlap between dinners at different tables, the other dinners were exposed to  $D_{\text{inf}} \sim 0.6$ . In fact, roughly half of the other dinners contracted COVID-19.

### A.2 Hunan Coach

[64] (SCMP article) documents an outbreak on a long distance coach (Hunan, China). This 45 person coach should have  $r_{\text{room}} \simeq 8\text{ m}^3/\text{min}$  [10], and we will estimate the ride duration as 2 hours. The contagious individual did not interact with others, so we assign a source rate of  $r_{\text{src}} = r_b = 1\text{ nL}/\text{min}$ . The resulting infective dose of  $D_{\text{inf}} \sim 0.15$  is in reasonable agreement with the fact that 8 of the 45 passengers were infected. The infections were somewhat localized, possibly due to incomplete mixing of the air in the bus.

This outbreak also resulted in the infection of a passenger who boarded 30 minutes after the contagious passenger had disembarked. This could have been due to surface contamination, but then one must wonder why no passengers on later voyages were infected (since fomites last for days) [35]. More likely it indicates that a few air cycle-times passed while the bus was stopped resulting in an order of magnitude reduction in infective dose, and continued during the voyage thereby clearing the air. (Outbreaks on vehicles were common in China, possibly due to poor ventilation [10].)

### A.3 Seattle Choir

[65, 66] Of 60 singers 87% infected after singing together for 2.5 hours in a church the size of a volleyball court ( $\sim 400\text{ m}^3$ ). An air cycle time of 40 minutes gives  $r_{\text{room}} \sim 10\text{ m}^3/\text{min}$ , and singing can be approximated by

$r_{\text{src}} \sim 30$  nL/min. The 150 minute exposure yields an infective dose of  $D_{\text{inf}} \sim 4.5$ . Sadly, this makes the probability of infection quite high. Even more sadly, this is not the only choir outbreak [67, 68].

#### A.4 Diamond Princess

[69] concludes from the outbreak on the Diamond Princess cruise ship that long-range airborne transmission is unlikely since SARS-CoV-2 did not pass through the ship’s air conditioning system. Two Okinawa taxi-drivers were, however, infected by passengers during a shore visit. This outbreak is different from the others in that the contagious individual which initiated the outbreak was not involved in the taxi-driver infections. At the time of the shore visit in Okinawa there were roughly 25 pre-symptomatic infected passengers on the Diamond Princess at least a few of whom disembarked.

If we assume that the taxi ride lasted 10 minutes (the port in Naha is close to the main attractions), that the contagious passenger spent the ride talking to another passenger (or coughed once), and that the taxi had the fan on low ( $r_{\text{room}} \sim 1$  m<sup>3</sup>/min), the driver was exposed to an infective dose of  $D_{\text{inf}} \sim 1$  according to Eqn. 7. Using the “random  $\rho V_{\text{src}} N_{\text{inf}}$ ” curve in Fig. 1 since these were secondary infections, we find that the probability of transmission was about 15% for each contagious passenger who disembarked. This probability is relatively insensitive to our assumptions and would only change by a factor of 2 if the dose changed by an order of magnitude (in either direction). If both infections in Naha were caused by the same contagious individual (one on the ride in, one on the ride out) it would be appropriate to use the “random  $V_{\text{src}} N_{\text{inf}}$ ” probability of 60%.

Notably, there were no other COVID-19 infections linked to the Diamond Princess’ stop in Okinawa, indicating that the closed environment and long exposure time of the taxi was likely a key ingredient for transmission. That is, changing the air-flow rate in the above calculation to  $r_{\text{room}} \gtrsim 100$  m<sup>3</sup>/min to represent shops and more open spaces would reduce the probability of infection to less than 2%. And, while surface transmission could explain transmission to taxi drivers, it does not explain the absence of transmission to shop keepers or other people with whom the passengers of the Diamond Princess interacted.

#### A.5 Hospital Air Sampling

There was no outbreak in the Nebraska Medical Center, but air sampling in and around COVID-19 patient rooms [70] offers a further cross-check of the calculations presented here. Approximately 3 viral RNA copies per liter of air sampled were found in a patient’s room (and in the hallway outside the patient’s room after the door was opened). These rooms have  $\tau_{\text{room}} \simeq 8$  min and  $V_{\text{room}} \simeq 30$  m<sup>3</sup>, indicating an airflow rate of  $r_{\text{room}} \sim 4$  m<sup>3</sup>/min [40]. Equation 5 indicates a steady-state concentration of  $\rho_{\text{A}}^{\text{SS}} \simeq 0.25$  /L. This suggests that either this patient had a very high viral load ( $\rho \sim 10^{10}$  /mL, which is in the top 3%), or that they were occasionally talking or coughing and had  $\rho \sim \rho_0$ .

Air sampling data is also available from hospitals and public areas in China. [71] reports quantitative viral deposition rate of roughly 2 /m<sup>2</sup> per minute in a patient’s room. The area sampled was 3 m from the patient’s bed, so too far for most droplets [11]. If we assume that this is due to slow settling of the larger droplet-nuclei with characteristic a speed of  $u_{\text{settle}} \sim 0.1$  m/min (see section 5 and [11, 61]), this implies a aerosol concentration of roughly 20 /m<sup>3</sup>. (Oddly, air samplers in patient rooms did not detect concentrations above their detection threshold, but the patient’s bathroom and other areas in the hospital had concentrations near 20 /m<sup>3</sup>).

A concentration of 20 /m<sup>3</sup> is more than a factor of 100 lower than that reported in Nebraska [70], but still implies  $\rho$  above the median of the distribution function shown in Fig. 2. If we assume an air flow rate in the room of  $r_{\text{room}} \sim 4$  m<sup>3</sup>/min and viral shedding dominated by breathing (i.e.,  $V_{\text{src}} \sim V_b$ ), for instance,  $\rho$  is roughly  $10^8$  /mL =  $\rho_0/10$ .

## B Parameter Probability Distributions

As described in section 5 several parameters used in this work vary significantly between individuals and events. This appendix describe the details of the probability density functions use for  $\rho$ ,  $V_{\text{src}}$  and  $N_{\text{inf}}$ .

Figure 2 shows the distribution for  $\rho$ . The estimated distribution used in this work is derived from [31] with the low end of the distribution pushed up somewhat to account for the downward trend in viral load after symptom onset. The log-normal distribution centered at  $10^6$  /mL with  $\sigma = 10^2$  is estimated from the linear fit in figure 2 in [32]. These distributions both have a median viral load of  $10^6$  /mL and roughly 10% of cases above  $10^9$  /mL. We have computed  $P_{\text{inf}}$  as in Fig. 1 for both distributions and they give quantitatively similar, and qualitatively identical, results.

To account for individual variability, we use a log-normal distributions for  $N_{\text{inf}}$  which is centered around 1000 and has a width of a factor of 2. This makes the 95% confidence interval 250–4000 which is in reasonable agreement with [27] and [23].



Similarly, an order of magnitude variation among individuals and events is also expected in in droplet and aerosol production [59, 61]. For this we also use a log-normal distribution centered around the  $V_{src}$  values given in Table 1 with a width of a factor of 3.




Figure 2: Probability density functions for  $\rho$ . The estimated distribution used in this work is shown along with a log-normal distribution centered at  $10^6$  /mL. The vertical line at  $10^9$  /mL indicates our “nominal  $\rho_0$ ” of 1000 /nL. 10% of the estimated  $\rho$  distribution lies above this line and 7% of the log-normal distribution lies above it.

## References

- [1] Michael Kochis. *Medical Student COVID-19 Curriculum*. <https://curriculum.covidstudentresponse.org/>.
- [2] Center for Disease Control and Prevention. Coronavirus (COVID-19). <https://www.cdc.gov/coronavirus/2019-ncov/index.html>.
- [3] Carlos del Rio and Preeti N Malani. COVID-19—New Insights on a Rapidly Changing Epidemic. *JAMA*, 323(14):1339–1340, April 2020. <https://doi.org/10.1001/jama.2020.3072>.
- [4] Sima Asadi, Nicole Bouvier, Anthony S Wexler, and William D Ristenpart. The coronavirus pandemic and aerosols: Does COVID-19 transmit via expiratory particles? *Aerosol Science and Technology*, 54(6):635–638, April 2020. <https://doi.org/10.1080/02786826.2020.1749229>.
- [5] Juan Wang and Guoqiang Du. COVID-19 may transmit through aerosol. *Irish Journal of Medical Science (1971 -)*, 2020. <https://doi.org/10.1007/s11845-020-02218-2>.
- [6] Elizabeth L Anderson, Paul Turnham, John R Griffin, and Chester C Clarke. Consideration of the Aerosol Transmission for COVID-19 and Public Health. *Risk Analysis*, 40(5):902–907, May 2020. <https://doi.org/10.1111/risa.13500>.
- [7] Leslie Dietz, Patrick F Horve, David A Coil, Mark Fretz, Jonathan A Eisen, and Kevin Van Den Wymelenberg. 2019 Novel Coronavirus (COVID-19) Pandemic: Built Environment Considerations To Reduce Transmission. *mSystems*, 5(2):e00245–20, April 2020. <https://doi.org/10.1111/risa.13500>.
- [8] Matthew Meselson. Droplets and aerosols in the transmission of sars-cov-2. *New England Journal of Medicine*, 0(0):null, 0. <https://doi.org/10.1056/NEJMc2009324>.
- [9] Center for Disease Control and Prevention. How COVID-19 Spreads. <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/how-covid-spreads.html>.
- [10] Hua Qian, Te Miao, Li LIU, Xiaohong Zheng, Danting Luo, and Yuguo Li. Indoor transmission of SARS-CoV-2. *medRxiv*, 2020. <https://www.medrxiv.org/content/early/2020/04/07/2020.04.04.20053058>.

- [11] Lydia Bourouiba, Eline Dehandschoewercker, and John W. M. Bush. Violent expiratory events: on coughing and sneezing. *Journal of Fluid Mechanics*, 745:537–563, 2014. <https://doi.org/10.1017/jfm.2014.88>.
- [12] Wenzhao Chen, Nan Zhang, Jianjian Wei, Hui-Ling Yen, and Yuguo Li. Short-range airborne route dominates exposure of respiratory infection during close contact. *Building and Environment*, 176:106859, 2020. <https://doi.org/10.1016/j.buildenv.2020.106859>.
- [13] L. Liu, Y. Li, P. V. Nielsen, J. Wei, and R. L. Jensen. Short-range airborne transmission of expiratory droplets between two people. *Indoor Air*, 27(2):452–462, 2017. <https://doi.org/10.1111/ina.12314>.
- [14] Prateek Bahl, Con Doolan, Charitha de Silva, Abrar Ahmad Chughtai, Lydia Bourouiba, and C Raina MacIntyre. Airborne or Droplet Precautions for Health Workers Treating Coronavirus Disease 2019? *The Journal of Infectious Diseases*, April 2020. <https://doi.org/10.1093/infdis/jiaa189>.
- [15] Raymond Tellier. Review of aerosol transmission of influenza A virus. *Emerging Infectious Disease journal*, 12(11):1657–1662, November 2006. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3372341/>.
- [16] Rajat Mittal, Rui Ni, and Jung-Hee Seo. The flow physics of COVID-19. *Journal of Fluid Mechanics*, 894:F2, 2020. <https://doi.org/10.1017/jfm.2020.330>.
- [17] X Xie. How far droplets can move in indoor environments – revisiting the Wells evaporation–falling curve. *Indoor Air*, 17(3):211–225, 2007. <https://doi.org/10.1111/j.1600-0668.2007.00469.x>.
- [18] Raymond Tellier. Aerosol transmission of influenza a virus: a review of new studies. *Journal of The Royal Society Interface*, 6(suppl\_6):S783–S790, 2009. <https://doi.org/10.1098/rsif.2009.0302.focus>.
- [19] Jasmin S Kutter, Monique I Spronken, Pieter L Fraaij, Ron AM Fouchier, and Sander Herfst. Transmission routes of respiratory viruses among humans. *Current Opinion in Virology*, 28:142 – 151, 2018. <https://doi.org/10.1016/j.coviro.2018.01.001>.
- [20] Elliot C Dick, Lance C Jennings, Kathy A Mink, Catherine D Wartgow, and Stanley L Inborn. Aerosol Transmission of Rhinovirus Colds. *The Journal of Infectious Diseases*, 156(3):442–448, September 1987. <https://doi.org/10.1093/infdis/156.3.442>.
- [21] Raymond Tellier, Yuguo Li, Benjamin J Cowling, and Julian W Tang. Recognition of aerosol transmission of infectious agents: a commentary. *BMC Infectious Diseases*, 19(1):101, 2019. <https://doi.org/10.1186/s12879-019-3707-y>.
- [22] Jie Zhou, Jianjian Wei, Ka-Tim Choy, Sin Fun Sia, Dewi K. Rowlands, Dan Yu, Chung-Yi Wu, William G. Lindsley, Benjamin J. Cowling, James McDevitt, Malik Peiris, Yuguo Li, and Hui-Ling Yen. Defining the sizes of airborne particles that mediate influenza transmission in ferrets. *Proceedings of the National Academy of Sciences*, 115(10):E2386–E2392, 2018. <https://doi.org/10.1073/pnas.1716771115>.
- [23] Peter F.M. Teunis, Nicole Brienens, and Mirjam E.E. Kretzschmar. High infectivity and pathogenicity of influenza a virus via aerosol and droplet transmission. *Epidemics*, 2(4):215 – 222, 2010. <https://doi.org/10.1016/j.epidem.2010.10.001>.
- [24] Center for Disease Control and Prevention. How to Protect Yourself Others. <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/prevention.html>.
- [25] Julian W. Tang, Thomas J. Liebner, Brent A. Craven, and Gary S. Settles. A schlieren optical study of the human cough with and without wearing masks for aerosol infection control. *Journal of The Royal Society Interface*, 6(suppl\_6):S727–S736, 2009. <https://doi.org/10.1098/rsif.2009.0295.focus>.
- [26] Alexander Rodriguez-Palacios, Fabio Cominelli, Abigail Basson, Theresa Pizarro, and Sanja Ilic. Textile masks and surface covers - a 'universal droplet reduction model' against respiratory pandemics. *medRxiv*, 2020. <https://www.medrxiv.org/content/early/2020/04/10/2020.04.07.20045617>.
- [27] Toru Watanabe, Timothy A Bartrand, Mark H Weir, Tatsuo Omura, and Charles N Haas. Development of a Dose-Response Model for SARS Coronavirus. *Risk Analysis*, 30(7):1129–1138, April 2010. <https://doi.org/10.1111/j.1539-6924.2010.01427.x>.
- [28] Theodore F. Hatch. Distribution and deposition of inhaled particles in respiratory tract. *Microbiology and Molecular Biology Reviews*, 25(3):237–240, 1961.
- [29] Mark Nicas and Gang Sun. An integrated model of infection risk in a health-care environment. *Risk Analysis*, 26(4):1085–1096, 2006. <https://doi.org/10.1111/j.1539-6924.2006.00802.x>.
- [30] Roman Wölfel, Victor M Corman, Wolfgang Guggemos, Michael Seilmaier, Sabine Zange, Marcel A Müller, Daniela Niemeyer, Terry C Jones, Patrick Vollmar, Camilla Rothe, Michael Hoelscher, Tobias Bleicker, Sebastian Brünink, Julia Schneider, Rosina Ehmann, Katrin Zwirgmaier, Christian Drosten, and Clemens Wendtner.

- Virological assessment of hospitalized patients with COVID-2019. *Nature*, 2020. <https://doi.org/10.1038/s41586-020-2196-x>.
- [31] Terry C Jones. An analysis of SARS-CoV-2 viral load by patient age. [https://virologie-ccm.charite.de/fileadmin/user\\_upload/microsites/m\\_cc05/virologie-ccm/dateien\\_upload/Weitere\\_Dateien/analysis-of-SARS-CoV-2-viral-load-by-patient-age-v2.pdf](https://virologie-ccm.charite.de/fileadmin/user_upload/microsites/m_cc05/virologie-ccm/dateien_upload/Weitere_Dateien/analysis-of-SARS-CoV-2-viral-load-by-patient-age-v2.pdf).
- [32] Kelvin Kai-Wang To, Owen Tak-Yin Tsang, Wai-Shing Leung, Anthony Raymond Tam, Tak-Chiu Wu, David Christopher Lung, Cyril Chik-Yan Yip, Jian-Piao Cai, Jacky Man-Chun Chan, Thomas Shiu-Hong Chik, Daphne Pui-Ling Lau, Chris Yau-Chung Choi, Lin-Lei Chen, Wan-Mui Chan, Kwok-Hung Chan, Jonathan Daniel Ip, Anthony Chin-Ki Ng, Rosana Wing-Shan Poon, Cui-Ting Luo, Vincent Chi-Chung Cheng, Jasper Fuk-Woo Chan, Ivan Fan-Ngai Hung, Zhiwei Chen, Honglin Chen, and Kwok-Yung Yuen. Temporal profiles of viral load in posterior oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-2: an observational cohort study. *The Lancet Infectious Diseases*, 20(5):565–574, 2020.
- [33] J. P. Duguid. The size and the duration of air-carriage of respiratory droplets and droplet-nuclei. *Epidemiology and Infection*, 44(6):471–479, 1946. <https://doi.org/10.1017/S0022172400019288>.
- [34] J Atkinson, Y Chartier, and CL Pessoa-Silva. Natural Ventilation for Infection Control in Health-Care Settings, 2009. <https://www.ncbi.nlm.nih.gov/books/NBK143281/>.
- [35] Neeltje van Doremalen, Trenton Bushmaker, Dylan H. Morris, Myndi G. Holbrook, Amandine Gamble, Brandi N. Williamson, Azaibi Tamin, Jennifer L. Harcourt, Natalie J. Thornburg, Susan I. Gerber, James O. Lloyd-Smith, Emmie de Wit, and Vincent J. Munster. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *New England Journal of Medicine*, 382(16):1564–1567, 2020. <https://doi.org/10.1056/NEJMc2004973>.
- [36] Alyssa C Fears, William B Klimstra, Paul Duprex, Amy Hartman, Scott C. Weaver, Ken S. Plante, Divya Mirchandani, Jessica Plante, Patricia V. Aguilar, Diana Fernandez, Aysegul Nalca, Allison Totura, David Dyer, Brian Kearney, Matthew Lackemeyer, J. Kyle Bohannon, Reed Johnson, Robert F Garry, Doug S Reed, and Chad J Roy. Comparative dynamic aerosol efficiencies of three emergent coronaviruses and the unusual persistence of sars-cov-2 in aerosol suspensions. *medRxiv*, 2020. <https://doi.org/10.1101/2020.04.13.20063784>.
- [37] *Physiology, Tidal Volume*. StatPearls. <https://www.ncbi.nlm.nih.gov/books/NBK482502/>.
- [38] Y Li, G M Leung, J W Tang, X Yang, C Y H Chao, J Z Lin, J W Lu, P V Nielsen, J Niu, H Qian, A C Sleigh, H J J Su, J Sundell, T W Wong, and P L Yuen. Role of ventilation in airborne transmission of infectious agents in the built environment ? a multidisciplinary systematic review. *Indoor Air*, 17(1):2–18, February 2007. <https://doi.org/10.1111/j.1600-0668.2006.00445.x>.
- [39] Pauline Elliott. Filtration and Air-Cleaning Systems to Protect Building Environments. pages 1–78, April 2003. <https://www.cdc.gov/niosh/docs/2003-136/pdfs/2003-136.pdf?id=10.26616/NIOSHPUB2003136>.
- [40] Angela L Hewlett, Scott E Whitney, Shawn G Gibbs, Philip W Smith, and Hendrik J Viljoen. Mathematical Modeling of Pathogen Trajectory in a Patient Care Environment. *Infection Control and Hospital Epidemiology*, 34(11):1181–1188, 2013. <https://doi.org/10.1086/673451>.
- [41] Luis A. Anchordoqui and Eugene M. Chudnovsky. A physicist view of the airborne infection. 2020. <https://arxiv.org/abs/2003.13689>.
- [42] Brent A Craven and Gary S Settles. A Computational and Experimental Investigation of the Human Thermal Plume. *Journal of Fluids Engineering*, 128(6):1251–1258, March 2006. <https://doi.org/10.1115/1.2353274>.
- [43] Seongman Bae, Min-Chul Kim, Ji Yeun Kim, Hye-Hee Cha, Joon Seo Lim, Jiwon Jung, Min-Jae Kim, Dong Kyu Oh, Mi-Kyung Lee, Seong-Ho Choi, Minki Sung, Sang-Bum Hong, Jin-Won Chung, and Sung-Han Kim. Effectiveness of Surgical and Cotton Masks in Blocking SARS-CoV-2: A Controlled Comparison in 4 Patients. *Annals of Internal Medicine*, April 2020. <https://annals.org/aim/fullarticle/2764367/effectiveness-surgical-cotton-masks-blocking-sars-cov-2-controlled-comparison>.
- [44] MacIntyre, Cauchemez, Dwyer, Seale, Cheung, and Browne. Face Mask Use and Control of Respiratory Virus Transmission in Households. *Emerg Infect Dis*, 15:233–241, 2009. <https://dx.doi.org/10.3201/eid1502.081167>.
- [45] C Raina MacIntyre and Abrar Ahmad Chughtai. Facemasks for the prevention of infection in healthcare and community settings. *BMJ*, 350, 2015. <https://www.bmj.com/content/350/bmj.h694>.
- [46] Shu-An Lee, Sergey A Grinshpun, and Tiina Reponen. Respiratory Performance Offered by N95 Respirators and Surgical Masks: Human Subject Evaluation with NaCl Aerosol Representing Bacterial and Viral Particle

- Size Range. *The Annals of Occupational Hygiene*, 52(3):177–185, March 2008. <https://doi.org/10.1093/annhyg/men005>.
- [47] X. Wang, Z. Pan, and Z. Cheng. Association between 2019-nCoV transmission and N95 respirator use. *Journal of Hospital Infection*, 2020. <https://doi.org/10.1016/j.jhin.2020.02.021>.
- [48] Jieliang Chen. Pathogenicity and transmissibility of 2019-nCoV, quick overview and comparison with other emerging viruses. *Microbes and Infection*, 22(2):69 – 71, 2020. Special issue on the new coronavirus causing the COVID-19 outbreak. <https://doi.org/10.1016/j.micinf.2020.01.004>.
- [49] Youlin Long, Tengyue Hu, Liqin Liu, Rui Chen, Qiong Guo, Liu Yang, Yifan Cheng, Jin Huang, and Liang Du. Effectiveness of N95 respirators versus surgical masks against influenza: A systematic review and meta-analysis. *Journal of Evidence-Based Medicine*, n/a(n/a). <https://doi.org/10.1111/jebm.12381>.
- [50] Shu-An Lee, Atin Adhikari, Sergey A. Grinshpun, Roy McKay, Rakesh Shukla, Haoyue Li Zeigler, and Tiina Reponen. Respiratory Protection Provided by N95 Filtering Facepiece Respirators Against Airborne Dust and Microorganisms in Agricultural Farms. *Journal of Occupational and Environmental Hygiene*, 2(11):577–585, 2005. PMID: 16234218. <https://doi.org/10.1080/15459620500330583>.
- [51] Christopher C. Coffey, Robert B. Lawrence, Donald L. Campbell, Ziqing Zhuang, Catherine A. Calvert, and Paul A. Jensen. Fitting Characteristics of Eighteen N95 Filtering-Facepiece Respirators. *Journal of Occupational and Environmental Hygiene*, 1(4):262–271, 2004. PMID: 15204866 <https://doi.org/10.1080/15459620490433799>.
- [52] C.R. MacIntyre, D. Dwyer, H. Seale, M. Fasher, R. Booy, P. Cheung, N. Ovdin, and G. Browne. The First Randomized, Controlled Clinical Trial of Mask Use in Households to Prevent Respiratory Virus Transmission. *International Journal of Infectious Diseases*, 12:e328, 2008. 13th International Congress on Infectious Diseases, Abstracts. <https://doi.org/10.1016/j.ijid.2008.05.877>.
- [53] Tom Jefferson, Chris B Del Mar, Liz Dooley, Eliana Ferroni, Lubna A Al-Ansary, Ghada A Bawazeer, Mieke L van Driel, Sreekumaran Nair, Mark A Jones, Sarah Thorning, and John M Conly. Physical interventions to interrupt or reduce the spread of respiratory viruses. *The Cochrane database of systematic reviews*, 2011(7):CD006207–CD006207, July 2011. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6993921/>.
- [54] Babak Javid, Michael P Weekes, and Nicholas J Matheson. Covid-19: should the public wear face masks? *BMJ*, 369, 2020. <https://www.bmj.com/content/369/bmj.m1442>.
- [55] Nancy H L Leung, Daniel K W Chu, Eunice Y C Shiu, Kwok-Hung Chan, James J McDevitt, Benien J P Hau, Hui-Ling YEN, Yuguo Li, Dennis K M Ip, J S Malik Peiris, Wing-Hong Seto, Gabriel M Leung, Donald K Milton, and Benjamin J Cowling. Respiratory virus shedding in exhaled breath and efficacy of face masks. *Nature Medicine*, 2020. <https://doi.org/10.1038/s41591-020-0843-2>.
- [56] Donald K Milton, M Patricia Fabian, Benjamin J Cowling, Michael L Grantham, and James J McDevitt. Influenza virus aerosols in human exhaled breath: particle size, culturability, and effect of surgical masks. *PLoS pathogens*, 9(3):e1003205–e1003205, March 2013. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3591312/>.
- [57] Augenbraun, Lasner, Mitra, Prabhu, Raval, Sawaoka, and J Doyle I.
- [58] Wladyslaw Kowalski. *Ultraviolet germicidal irradiation handbook: UVGI for air and surface disinfection*. Springer science & business media, 2010.
- [59] Z. Y. Han, W. G. Weng, and Q. Y. Huang. Characterizations of particle size distribution of the droplets exhaled by sneeze. *Journal of The Royal Society Interface*, 10(88):20130560, 2013. <https://doi.org/10.1098/rsif.2013.0560>.
- [60] W F Wells. On Air-borne Infection: Study II. Droplets and Droplet Nuclei. *American Journal of Epidemiology*, 20(3):611–618, November 1934. <https://doi.org/10.1093/oxfordjournals.aje.a118097>.
- [61] Valentyn Stadnytskyi, Christina E Bax, Adriaan Bax, and Philip Anfinrud. The airborne lifetime of small speech droplets and their potential importance in SARS-CoV-2 transmission. *Proc Natl Acad Sci USA*, page 202006874, May 2020. [www.pnas.org/cgi/doi/10.1073/pnas.2006874117](http://www.pnas.org/cgi/doi/10.1073/pnas.2006874117).
- [62] Xi He, Eric H Y Lau, Peng Wu, Xilong Deng, Jian Wang, Xinxin Hao, Yiu Chung Lau, Jessica Y Wong, Yujuan Guan, Xinghua Tan, Xiaoneng Mo, Yanqing Chen, Baolin Liao, Weilie Chen, Fengyu Hu, Qing Zhang, Mingqiu Zhong, Yanrong Wu, Lingzhai Zhao, Fuchun Zhang, Benjamin J Cowling, Fang Li, and Gabriel M Leung. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nature Medicine*, 26(5):672–675, 2020. <https://doi.org/10.1038/s41591-020-0869-5>.
- [63] Jianyun Lu, Jieni Gu, Kuibiao Li, Conghui Xu, Wenzhe Su, Zhisheng Lai, Deqian Zhou, Chao Yu, Bin Xu, and Zhicong Yang. COVID-19 Outbreak Associated with Air Conditioning in Restaurant, Guangzhou, China, 2020. *Emerging Infectious Disease journal*, 26(7), 2020. <https://doi.org/10.3201/eid2607.200764>.

- [64] Stefen Chen. Coronavirus can travel twice as far as official ‘safe distance’ and stay in air for 30 minutes. <https://www.scmp.com/news/china/science/article/3074351/coronavirus-can-travel-twice-far-official-safe-distance-and-stay>.
- [65] L Hamner, P Dubbel, and I Capron. High SARS-CoV-2 Attack Rate Following Exposure at a Choir Practice — Skagit County, Washington, March 2020. *Morb Mortal Wkly Rep*, 69(14):606–610, 2020. <http://dx.doi.org/10.15585/mmwr.mm6919e6>.
- [66] Richard Read. A choir decided to go ahead with rehearsal. <https://www.latimes.com/world-nation/story/2020-03-29/coronavirus-choir-outbreak>.
- [67] Ellen Kamphorst. Choir Heerde lost six members to corona: 40 per cent fell ill. <https://www.omroep gelderland.nl/nieuws/2448308/Koor-Heerde-verloor-zes-leden-aan-corona-40-procent-werd-ziek>.
- [68] Peter van der Lint. That one Passion that did go on, with disastrous consequences. <https://www.trouw.nl/verdieping/die-ene-passion-die-wel-doorging-met-rampzalige-gevolgen~b4ced33e/>.
- [69] Pengcheng Xu, Hua Qian, Te Miao, Hui-ling Yen, Hongwei Tan, Benjamin J. Cowling, and Yuguo J Li. Transmission routes of covid-19 virus in the diamond princess cruise ship. *medRxiv*, 2020. <http://www.medrxiv.org/content/early/2020/04/14/2020.04.09.20059113>.
- [70] Joshua L Santarpia, Danielle N Rivera, Vicki Herrera, M. Jane Morwitzer, Hannah Creager, George W. Santarpia, Kevin K Crown, David Brett-Major, Elizabeth Schnaubelt, M. Jana Broadhurst, James V. Lawler, St. Patrick Reid, and John J. Lowe. Transmission Potential of SARS-CoV-2 in Viral Shedding Observed at the University of Nebraska Medical Center. *medRxiv*, 2020. <https://www.medrxiv.org/content/early/2020/03/26/2020.03.23.20039446.1>.
- [71] Yuan Liu, Zhi Ning, Yu Chen, Ming Guo, Yingle Liu, Nirmal Kumar Gali, Li Sun, Yusen Duan, Jing Cai, Dane Westerdahl, Xinjin Liu, Kin-fai Ho, Haidong Kan, Qingyan Fu, and Ke Lan. Aerodynamic Characteristics and RNA Concentration of SARS-CoV-2 Aerosol in Wuhan Hospitals during COVID-19 Outbreak. *bioRxiv*, 2020. <https://www.biorxiv.org/content/early/2020/03/10/2020.03.08.982637>.