

## **Association of country-wide coronavirus mortality with demographics, testing, lockdowns, and public wearing of masks (Update June 2, 2020).**

Christopher T. Leffler, MD, MPH.<sup>1,2 \*</sup>

Edsel Ing MD, MPH, CPH, MIAD.<sup>3</sup>

Joseph D. Lykins V, MD.<sup>4,5</sup>

Matthew C. Hogan, MS, MPH.<sup>6</sup>

Craig A. McKeown, MD.<sup>7</sup>

Andrzej Grzybowski, MD, PhD, MBA.<sup>8,9</sup>

1. Department of Ophthalmology. Virginia Commonwealth University. Richmond, VA 23298.  
chrislefflermd@gmail.com

2. Department of Ophthalmology. Hunter Holmes McGuire VA Medical Center, Richmond, VA.

3. Department of Ophthalmology & Vision Sciences, University of Toronto.

4. Department of Internal Medicine, Virginia Commonwealth University. Richmond, VA 23298.

5. Department of Emergency Medicine. Virginia Commonwealth University. Richmond, VA 23298.

6. School of Medicine, Virginia Commonwealth University, Richmond, VA 23298.

7. Bascom Palmer Eye Institute, University of Miami, Miller School of Medicine.

8. Department of Ophthalmology, University of Warmia and Mazury, Olsztyn, Poland.

9. Institute for Research in Ophthalmology, Poznan, Poland.

\*Corresponding author: Christopher T. Leffler, MD, MPH.

Department of Ophthalmology. Virginia Commonwealth University. 401 N. 11th St., Box 980209,  
Richmond, VA 23298. chrislefflermd@gmail.com.

June 2, 2020.

None of the authors has any conflicts of interest to disclose.

## **Abstract.**

**Background.** Wide variation between countries has been noted in per-capita mortality from the disease (COVID-19) caused by the SARS-CoV-2 virus. Determinants of this variation are not fully understood.

**Methods.** Potential predictors of country-wide per-capita coronavirus-related mortality were examined, including age, sex ratio, obesity prevalence, temperature, urbanization, smoking, duration of infection, lockdowns, viral testing, contact tracing policies, and public mask-wearing norms and policies. Multivariable linear regression analysis was performed.

**Results.** In univariate analyses, the prevalence of smoking, per-capita gross domestic product, urbanization, and colder average country temperature were positively associated with coronavirus-related mortality. In a multivariable analysis of 194 countries, the duration of infection in the country, and the proportion of the population 60 years of age or older were positively associated with per-capita mortality, while duration of mask-wearing by the public was negatively associated with mortality (all  $p < 0.001$ ). The prevalence of obesity was independently associated with mortality in models which controlled for testing levels or policy. International travel restrictions were independently associated with lower per-capita mortality, but other containment measures and viral testing and tracing policies were not. In countries with cultural norms or government policies supporting public mask-wearing, per-capita coronavirus mortality increased on average by just 9.0% each week, as compared with 47% each week in remaining countries. On multivariable analysis, lockdowns tended to be associated with less mortality ( $p = 0.44$ ), and increased per-capita testing with higher reported mortality ( $p = 0.72$ ), though neither association was statistically significant.

**Conclusions.** Societal norms and government policies supporting the wearing of masks by the public, as well as international travel controls, are independently associated with lower per-capita mortality from COVID-19.

## **Introduction.**

The COVID-19 global pandemic caused by infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has presented a major public health challenge. For reasons that are not completely understood, the per-capita mortality from COVID-19 varies by several orders of magnitude between countries.<sup>1</sup> Numerous sources of heterogeneity have been hypothesized. Higher mortality has been observed in older populations and in men.<sup>2,3</sup> Patient-level behaviors, such as smoking, might also have an impact.<sup>3</sup> Other potentially relevant factors include economic activity, and environmental variation, such as temperature.<sup>4</sup> More urban settings and increased population density would be expected to enhance viral transmission.<sup>5</sup>

In addition, public health responses to the COVID-19 pandemic may influence per-capita mortality. Various strategies have been implemented, ranging from robust testing programs to lockdown or stay-at-home orders, to mandates regarding social distancing and face mask usage. Practices with theoretical benefit, such as social distancing, stay-at-home orders, and implementation of mandates regarding use of masks in public spaces, must be assessed quickly, as implementation has the potential to reduce morbidity and mortality.

Mask usage by the public is postulated to decrease infection by blocking the spread of respiratory droplets,<sup>1</sup> and was successfully implemented during other coronavirus outbreaks (i.e. SARS and MERS).<sup>6</sup> In the context of the ongoing pandemic, we assessed the impact of masks on per-capita COVID-19-related mortality, controlling for the aforementioned factors. We hypothesized that in countries where mask use was either an accepted cultural norm or favored by government policies on a national level, the per-capita mortality might be reduced, as compared with remaining countries.

## **Methods.**

### **Data acquisition.**

Country-wide coronavirus mortality data was retrieved from the publicly available Worldometers Database on May 9, 2020.<sup>7</sup> Countries were included if either: 1) coronavirus testing data were available by May 9, 2020,<sup>7</sup> or if: 2) testing and lockdown policies had been graded by the University of Oxford Coronavirus Government Response Tracker.<sup>54,55</sup>

Oxford University defined and scored several composite government response indices. The stringency index was defined in terms of containment policy and public information.<sup>55</sup> The government response index incorporated containment, economic measures, public information, and testing and tracing policies.<sup>55</sup> The containment and health index was defined in terms of containment measures, public information, and testing and tracing policies.<sup>55</sup>

Archived viral testing data for April 2020 were also downloaded.<sup>8</sup> The date of the country's first reported infection and first death were obtained from the European Centre for Disease Prevention and Control.<sup>9</sup>

Mean temperature in each country during the pandemic was estimated using the average monthly temperature in the country's largest city from public sources.<sup>10,11</sup>

Online news reports, including those cited in a public database,<sup>56</sup> were searched to identify countries in which the public wore masks early in the outbreak based on tradition, as well as countries in which the national government mandated or recommended mask-wearing by the public before April 16, 2020.

For each country, the population,<sup>12</sup> fraction of the population age 60 years and over, and age 14 and under, male: female ratio per country,<sup>13</sup> surface area,<sup>12,13</sup> gross domestic product per capita,<sup>14</sup> percent urbanization,<sup>12,15</sup> adult smoking prevalence<sup>16-19</sup> and prevalence of adult obesity<sup>57-76</sup> were tabulated. Whether a nation was an isolated political entity on an island was also recorded.

### **Statistical analysis.**

The prevalence of an infectious process undergoing exponential growth (or decay) appears linear over time when graphed on a logarithmic scale.<sup>1</sup> Therefore, we postulated that the logarithm of the country-wide infection prevalence would be linearly related with the duration of the infection in each country. In addition, our analysis postulated that deaths from coronavirus would follow infections with some delay.

On average, the time from infection with the coronavirus to onset of symptoms is 5.1 days,<sup>20</sup> and the time from symptom onset to death is on average 17.8 days.<sup>21</sup> Therefore, the time from infection to death is expected to be 23 days.<sup>1,22</sup> These incubation and mortality times were prespecified.<sup>1,22</sup> Therefore, the date of each country's initial infection was estimated as the earlier of: 5 days before the first reported infection, or 23 days before the first death.<sup>8,9,23</sup> Deaths by May 9, 2020 would typically reflect infections beginning 23 days previously (by April 16). Therefore, we recorded the time from the first infection in a country until April 16. We also recorded the period of the outbreak: 1) from the mandating of activity restrictions until April 16, and 2) from when public mask-wearing was recommended until April 16. In addition, we calculated the mean time-weighted score for each lockdown and testing policy as graded by the University of Oxford for the duration of the country's outbreak, from beginning through April 16.<sup>54</sup> For instance, if the school closure score was 1 for half the outbreak and 2 for the other half, then the mean score was 1.5.

In univariate analysis, characteristics of countries with above-median per-capita mortality were compared with the remaining (lower mortality) countries by the two-group t-test. The odds ratio for being in the high-mortality group was calculated by logistic regression.

Significant predictors of per-capita coronavirus mortality in the univariate analysis were analyzed by stepwise backwards multivariable linear regression analysis. The dependent variable was the logarithm (base 10) of per-capita coronavirus-related mortality. Because of the importance relative to public health, the weeks the country spent in lockdown and using masks, and per-capita testing levels, were retained in the model. In addition, because of their biological plausibility and presumed importance, urbanization, prevalence of obesity, and average ambient temperature were retained in most of the multivariable models presented below. Statistical analysis was performed with xlstat 2020.1 (Addinsoft, New York). An alpha (p value) of 0.05 was deemed to be statistically significant. The study was approved by the Virginia Commonwealth University Office of Research Subjects Protection.

## Results.

We studied coronavirus mortality in 198 countries, of which 183 had testing data,<sup>7</sup> 161 had government policies scored by Oxford University,<sup>54</sup> and 146 fell into both categories.

The 99 lower-mortality countries had 1.1 deaths per million population, in contrast with an average of 94.2 deaths per million population in the 99 higher-mortality countries ( $p < 0.001$ , Table 1). The median value was 3.7 deaths per million population.

We assumed that island nations might find it less challenging to isolate and protect their populations. However, 20 of 99 low-mortality countries were isolated on islands, compared with 27 of 99 high-mortality countries ( $p = 0.32$ ). Country surface area and population were not associated with coronavirus mortality (Table 1).

### Population characteristics.

Countries with older populations suffered higher coronavirus mortality. Countries with low mortality had on average 8.9% of their population over age 60, as compared with 18.9% in the high-mortality countries (Table 1). Sex ratio was not associated with country-wide mortality ( $p = 0.53$ , Table 1). Smoking prevalence was on average 13.8% in low mortality countries and 18.5% in high-mortality countries ( $p < 0.001$ , Table 1). The prevalence of obesity was on average 14.7% in low-mortality countries and 24.0% in high-mortality countries ( $p < 0.001$ , Table 1).

### Temperature.

Colder countries were associated with higher coronavirus mortality in univariate analysis. The mean temperature was 22.3 C (SD 7.6 C) in the low-mortality countries, and 14.0 C (SD 9.1 C) in the high-mortality countries ( $p < 0.001$ , Table 1).

### Economics.

Urbanization was associated with coronavirus mortality in univariate analysis. In low-mortality countries, on average 53% of the population was urban, as compared with 71% of the population in the high-mortality countries ( $p < 0.001$ , Table 1). Richer countries suffered a higher coronavirus related mortality. The mean GDP per capita was \$9,350 in the low-mortality countries, and was \$27,380 in the high-mortality countries (Table 1,  $p < 0.001$ ).

**Table 1.** Characteristics of countries with low and high per-capita coronavirus mortality by May 9, 2020 in 198 countries.

	Odds ratio (95% CI)	Mean (SD)		p value
		Low Mortality	High Mortality	
Deaths (per million)	--	1.12 (1.18)	94.2 (183.4)	<0.001
Deaths (per capita, log)	--	-6.46 (0.76)	-4.54 (0.64)	<0.001
Duration infection (weeks)	1.22 (1.08 to 1.37)	6.51 (2.87)	7.82 (2.37)	<0.001
Duration infection without masks (weeks)	1.34 (1.18 to 1.53)	5.10 (2.41)	6.83 (2.53)	<0.001
Duration infection without lockdown (weeks)	1.08 (0.97 to 1.20)	2.83 (2.93)	3.34 (2.41)	0.18
Temperature, mean (C)	0.90 (0.86 to 0.93)	22.3 (7.6)	14.0 (9.1)	<0.001
Urban population (%)	1.04 (1.02 to 1.05)	53.2 (22.4)	70.6 (19.9)	<0.001
GDP per capita (\$1000)	1.05 (1.03 to 1.07)	9.350 (17.140)	27.380 (27.530)	<0.001
Age 14 & under (% of pop.)	0.85 (0.81 to 0.89)	32.2 (9.7)	20.1 (6.6)	<0.001
Age 60 & over (% of pop.)	1.20 (1.14 to 1.27)	8.9 (5.3)	18.4 (7.9)	<0.001
Surface area (million km <sup>2</sup> )	1.08 (0.92 to 1.26)	0.557 (1.12)	0.804 (2.455)	0.37
Population (million)	0.998 (0.995 to 1.001)	53.1 (20.1)	24.9 (48.2)	0.18
Sex ratio (males / 100 females)	1.004 (0.990 to 1.018)	101.0 (11.0)	102.9 (27.8)	0.53
Smoking prevalence, adult (%)	1.09 (1.04 to 1.13)	13.8 (7.8)	18.5 (7.7)	<0.001
Obesity prevalence, adult (%)	1.14 (1.10 to 1.19)	14.7 (9.0)	24.0 (7.3)	<0.001
Tests per cap. (log) by Apr 4	2.99 (1.92 to 4.68)	-3.73 (1.20)	-2.65 (0.76)	<0.001
Tests per cap. (log) by Apr 16	3.60 (2.31 to 5.80)	-3.09 (0.87)	-2.31 (0.67)	<0.001
Tests per cap. (log) by May 9	4.69 (2.85 to 7.73)	-2.76 (0.85)	-1.91 (0.61)	<0.001

Durations run from the estimated date of first infection in the country until 23 days before May 9, 2020 (i.e. April 16), or the stated event (mask recommendation or lockdown). Obesity data available for 194 countries. Testing data available for 135 countries by April 4, 162 countries by April 16, and 183 countries by May 9.

## Masks.

In some Asian countries, masks were used extensively by the public, essentially from the beginning of the outbreak,<sup>1,24</sup> including: Thailand,<sup>25</sup> Japan,<sup>24,26</sup> South Korea,<sup>24,27</sup> Taiwan,<sup>28</sup> Macau,<sup>77</sup> Hong Kong,<sup>24,29,30</sup> Vietnam,<sup>31</sup> Malaysia,<sup>32</sup> Cambodia,<sup>33,34</sup> and the Philippines<sup>35</sup> (Table 2).<sup>78-81</sup> In fact, in Mongolia<sup>36</sup> and Laos,<sup>37</sup> the public began wearing masks before any cases were confirmed in their countries. Despite the fact that the outbreak tended to appear in these 12 countries quite early, the countries had experienced a low per-capita coronavirus mortality by May 9 (mean 1.8 per million, SD 2.4, Table 2).

The World Health Organization initially advised against widespread mask wearing by the public, as did the United States CDC.<sup>1,24</sup> Nonetheless, some Western

**Table 2.** Countries in which masks were widely used by the public or recommended by the government early in their outbreak.

	First Case Date	Per-capita mortality (per mil. pop.) by May 9.	Comment.
China	Dec. 1 <sup>78</sup>	3.2	Mask use is traditional. By the time human-to-human transmission was confirmed on Jan. 20, many in Beijing were already wearing masks. <sup>79</sup> The government required masks in public in Wuhan on Jan. 22. <sup>80</sup> Masks were ordered throughout China when around others in public on Jan. 31. <sup>81</sup>
Thailand	Jan. 13	0.8	Mask use was traditional, and was recommended by the Ministry of Public Health on Jan. 31. <sup>25</sup>
Japan	Jan. 16	4.8	Public use of masks is traditional, <sup>24</sup> and was manifest by Jan. 31. <sup>26</sup> The government initially recommended them when in “confined, badly ventilated spaces”. <sup>24</sup>
South Korea	Jan. 20	5.0	Use of masks is traditional. <sup>24</sup> South Korea initially had trouble obtaining enough masks, but at the end of February the government began to control the distribution of masks to the public. <sup>27</sup>
Taiwan	Jan. 21	0.3	Use of masks is traditional. By January 27, the government had to limit mask exports and limit sales from pharmacies to those needed for personal use. <sup>28</sup>
Macau	Jan. 22	0.0	Mask use is traditional. By Jan. 23, the government had implemented a mask distribution program for the public. <sup>77</sup>
Hong Kong	Jan. 23 <sup>29</sup>	0.5	Surgical masks were traditionally used, and also were recommended on public transport and in crowded places, on January 24, 2020. <sup>24,30</sup>
Vietnam	Jan. 23	0.0	Masks were widely used by the public immediately, <sup>31</sup> and were mandated by the government on March 16.
Malaysia	Jan. 25	3.3	Masks were used by the public almost immediately. <sup>32</sup>
Cambodia	Jan. 27 <sup>33</sup>	0.0	Masks were used by the public almost immediately after the outbreak in Wuhan. <sup>34</sup>
Philippines	Jan. 30	6.4	Masks were used from the start of the outbreak, <sup>35</sup> and were mandated on April 2.
Mongolia	Mar. 10	0.0	Mongolians began wearing masks in January. <sup>36</sup>
Laos	Mar. 24	0.0	The public in Laos began wearing masks even before any cases were reported in the country. <sup>37</sup>

governments mandated or recommended mask-wearing in public in March 2020. Masks were required in public in Venezuela beginning March 13.<sup>38,39</sup> In Slovakia, masks were mandated in shops and transit on March 15,<sup>40</sup> and more broadly in public on March 25.<sup>41</sup> Masks were dispensed by pharmacists for free in Sudan by March 16.<sup>82-83</sup> Masks were required in public in Czechia on March 19,<sup>42</sup> in Uzbekistan on March 25, and in Bosnia on March 29.<sup>43,84</sup> Masks were mandated in indoor public spaces in Slovenia on March 29.<sup>44</sup> In Austria, a mandate to wear masks in shops was announced on March 30, with the expectation that masks would be available by April 1.<sup>45</sup> In addition, the requirement to wear masks on public transit was announced there on April 6.<sup>46</sup> Masks were recommended for the public in Bulgaria on March 30.<sup>47</sup>

Government mandates or recommendations for mask wearing by the public were issued by April 16 in multiple countries, including: Israel and Sierra Leone on April 1; in Cuba,<sup>85</sup> the Philippines, and Mauritius on April 2; the United States, Turkey, Singapore,<sup>86-88</sup> and Cyprus on April 3; India,<sup>89</sup> Kenya,<sup>90</sup> Malawi,<sup>91</sup> Colombia, and the UAE on April 4; Indonesia,<sup>92</sup> Lebanon, Antigua and Barbuda,<sup>93</sup> and Trinidad and Tobago on April 5; Ukraine, Canada, Tunisia,<sup>94</sup> and Morocco on April 6; Honduras, Peru, and Panama on April 7; Ecuador, El Salvador, Lithuania, Chile, Benin,<sup>95</sup> and Mozambique<sup>96</sup> on April 8; Guatemala and Bahrain<sup>97</sup> on April 9; Gabon<sup>98</sup> and South Africa on April 10; Spain, Ethiopia,<sup>99</sup> and Sri Lanka on April 11; Ethiopia on April 12; Cameroon on April 13;<sup>100</sup> Equatorial Guinea, Sint Maarten, Mongolia, and Finland on April 14; Grenada on April 15; the Dominican Republic, Poland, Libya, Zambia, and Estonia on April 16.

### **Initial multivariable analyses.**

An initial multivariable analysis was conducted including all 198 countries. By multivariable linear regression, significant predictors of the logarithm of each country's per-capita coronavirus mortality included: duration of infection in the country, duration of wearing masks ( $p < 0.001$ ), percentage of the population over age 60, and urbanization (all  $p \leq 0.007$ , Appendix Table A1).

We also prepared a multivariable model to predict the logarithm of per-capita coronavirus mortality in the 194 countries with obesity data. In this model, lockdown, obesity, temperature, and urbanization were retained due to their plausibility as important factors (Table 3).

By multivariable linear regression, significant predictors of the logarithm of each country's per-capita coronavirus mortality included: duration of infection in the country, duration of wearing masks, and percentage of the population over age 60 (all  $p < 0.001$ , Table 3). With the duration of infection in the country controlled for, there was a nonsignificant trend for time in lockdown to be negatively associated with mortality (Table 3).

**Table 3.** Predictors of (log) Country-wide Per-capita Coronavirus Mortality by May 9 by Multivariable Linear Regression in 194 Countries.

	$10^{\text{coefficient}}$	Coefficient (SE)	95% CI	P
Duration in country (wks)	1.4739	0.168 (0.032)	0.105 to 0.232	<0.001
Time wearing masks (wks)	0.7398	-0.131 (0.028)	-0.185 to -0.076	<0.001
Time in lockdown (wks)	0.9409	-0.026 (0.035)	-0.095 to 0.042	0.44
Population, age $\geq$ 60 (%)	1.1220	0.050 (0.010)	0.029 to 0.071	<0.001
Urbanization (%)	1.0144	0.00623 (0.00368)	-0.00103 to 0.01348	0.09
Obesity prevalence (%)	1.0378	0.016 (0.009)	-0.002 to 0.034	0.08
Temperature, ambient (C)	0.9857	-0.0063 (0.009)	-0.024 to 0.011	0.48
Constant	--	-7.716 (0.381)	-8.47 to -6.96	<0.001

Duration of infection in country from estimated date of first infection until 23 days before May 9, 2020 (i.e. April 16). Mask and lockdown durations run from the stated event (mask recommendation or lockdown) or estimated date of first infection in the country (whichever was later) until 23 days before May 9, 2020 (i.e. April 16).

In a countries not recommending masks, the per-capita mortality tended to increase each week by a factor of 1.474, or 47.4%. In contrast, in countries recommending masks, the per-capita mortality tended to increase each week by a factor of  $1.4739 * 0.7398 = 1.090$ , or just 9.0%. Under lockdown (without masks), the per-capita mortality increased each week by  $(1.4739)(0.9409) = 1.387$ , or 38.7%, i.e. slightly less than the baseline condition (Table 3).

A country with 10% more of its population living in an urban environment than another country tended to suffer a mortality 15% higher ( $10^{0.0623} = 1.15$ , Table 3). A country in which the percentage of the population age 60 or over is 10% higher than in another country tended to suffer mortality 216% higher ( $10^{0.50} = 3.16$ , Table 3). A country with a prevalence of obesity 10% higher tended to suffer mortality 45% higher ( $10^{0.16} = 1.45$ , Table 3).

### Numbers of Viral Tests.

Among the 183 countries with viral (PCR) testing data by May 9, per-capita testing performed at all 3 time points was positively associated with per-capita mortality in univariate analysis (all  $p < 0.001$ , Table 1). By May 9, 2020, low-mortality countries had performed 1 test for every 575 members of the population, while high-mortality countries had performed 1 test for every 81 members of the population ( $p < 0.001$ , Table 1).

To the multivariable model (Table 3), we added testing by May 9, using data from 179 countries with both testing and obesity data. Duration of infection in the country, the duration that masks were recommended, and age at least 60 years continued to be significant predictors of per-capita mortality (all  $p \leq 0.001$ , Appendix Table A2). Each week the infection persisted in a country was associated with a 49.1% increase in per-capita mortality without masks (Table A2). In contrast, in countries where masks were recommended, the per-capita mortality tended to increase by 13.1% (because  $(1.4907)(0.7589) = 1.131$ , Table A2). In this model, the prevalence of obesity was significantly associated with country-wide per-capita mortality ( $p = 0.03$ ). If the

prevalence of obesity increased by 10% (e.g. from 10% to 20% of a population), the per-capita mortality tended to increase by 63% (Appendix Table A2)

In this model, a 10-fold increase (i.e. one logarithm) in per-capita testing tended to be associated with a 10.2% increase in reported per-capita mortality, though the trend was not close to reaching statistical significance ( $p=0.72$ , Appendix Table A2). Duration of the infection in the country, time during the outbreak in which masks were recommended, and the fraction of the population over age 60 were all still significant predictors of mortality (all  $p \leq 0.001$ , Appendix Table A2).

If early testing lowers mortality, one might expect negative regression coefficients. Testing on both April 16 and May 9 were added to the multivariable model of Table 3, using data from the 158 countries with both obesity and testing data by these dates. Per-capita testing (log) by April 16 was not negatively associated with per-capita mortality (log) by May 9 (coefficient 0.366, 95% CI -0.235 to 0.967,  $p=0.23$ ).

Likewise, testing on both April 4 (the earliest archived data) and May 9 were added to the multivariable model of Table 3, using data from the 131 countries with both obesity and testing data by these dates. Per-capita testing (log) by April 4 was not significantly associated with per-capita mortality (log) by May 9 (coefficient -0.006, 95% CI -0.332 to 0.320,  $p=0.97$ ). Given the coefficient, a 10-fold (one log) increase in early testing would be associated with a (non-significant) decrease in per-capita mortality of 1.4%.

Only 5 countries had performed over 1 test for every 10 people in the country by May 9, 2020 (in order of most testing to least): the Faeroe Islands, Iceland, the Falkland Islands, the UAE, and Bahrain. The Faeroe and Falkland Islands reported no coronavirus-related deaths. The remaining 3 countries had per-capita mortality above the median value. The highest per-capita mortality among this group was 29.0 per million population (or 1 in 34,480 people), seen in Iceland.

### **Containment and Testing Policies.**

For 161 countries, containment, testing, and health policies were scored by Oxford University.<sup>54</sup> The following countries with mask policies by April 16 were included in this analysis, but not in the previous multivariable model, for lack of data on numbers of tests performed: China, Macau, Cameroon, Sierra Leone, and Sudan. In univariate analysis, scores for school closing, cancelling public events, restrictions on gatherings, and international travel controls were significantly associated with lower per-capita mortality (all  $p \leq 0.03$ , Table 4). Policies regarding workplace closing, closing public transport, stay at home requirements, internal movement restrictions, public information campaigns, testing, and contact tracing were not significant predictors of mortality (all  $p > 0.05$ , Table 4). Likewise, overall indices of stringency, government response, and containment and health were not associated with mortality (all  $p > 0.05$ , Table 4).

**Table 4.** Government policies in 161 countries with low and high per-capita coronavirus mortality by May 9, 2020.

	Odds ratio (95% CI)	Mean (SD)		p value
		Low Mortality	High Mortality	
School closing (0-3)	0.46 (0.26 to 0.82)	2.09 (0.66)	1.83 (0.48)	0.006
Workplace closing (0-3)	1.62 (0.98 to 2.66)	1.14 (0.77)	1.33 (0.47)	0.07
Cancel public events (0-2)	0.32 (0.14 to 0.74)	1.38 (0.45)	1.20 (0.34)	0.008
Restrictions on gatherings (0-4)	0.76 (0.53 to 1.09)	1.99 (0.91)	1.78 (0.82)	0.03
Close public transport (0-2)	0.83 (0.43 to 1.59)	0.62 (0.50)	0.58 (0.45)	0.21
Stay at home requirements (0-3)	1.27 (0.73 to 2.21)	0.81 (0.66)	0.88 (0.44)	0.33
Internal movement restrictions (0-2)	0.71 (0.35 to 1.42)	0.92 (0.52)	0.85 (0.36)	0.06
International travel controls (0-4)	0.44 (0.28 to 0.68)	2.87 (0.78)	2.36 (0.82)	<0.001
Income support (0-2)	40.3 (11.1 to 147.0)	0.13 (0.21)	0.50 (0.40)	<0.001
Debt / contract relief (0-2)	4.44 (2.04 to 9.66)	0.28 (0.39)	0.57 (0.49)	<0.001
Public information campaigns (0-2)	0.63 (0.30 to 1.33)	1.68 (0.39)	1.60 (0.45)	0.31
Testing policy (0-3)	0.84 (0.48 to 1.49)	1.10 (0.60)	1.05 (0.49)	0.22
Contact tracing (0-2)	0.89 (0.54 to 1.47)	1.06 (0.63)	1.02 (0.60)	0.17
Stringency Index (0-100)	0.98 (0.96 to 1.01)	52.2 (15.6)	49.0 (12.4)	0.09
Government response index (0-100)	0.99 (0.97 to 1.03)	44.6 (12.3)	44.5 (11.1)	0.42
Containment & health index (0-100)	0.98 (0.96 to 1.01)	50.9 (14.3)	48.1 (11.9)	0.10
Economic support index (0-100)	1.07 (1.04 to 1.10)	9.8 (12.8)	24.7 (17.4)	<0.001

Government policies were scored by Oxford University.<sup>54</sup> Characterization as low or high mortality was defined by the median for all 198 countries.

A multivariable model in 161 countries found that duration of the infection, duration masks were recommended, prevalence of age at least 60 years, obesity, and international travel restrictions were independently predictive of per-capita mortality (Table 5). At baseline, each week of the infection in a country was associated with an increase in per-capita mortality of 26.68% (Table 5). In contrast, when masks were worn, the per-capita mortality increased by  $(1.2668)(0.7927) = 1.004$ , or just 0.4% each week (Table 5).

International travel restrictions were scored by Oxford as: (0) no measures, (1) screening, (2) quarantine arrivals from high-risk regions; and ban on arrivals from some (3) or all (4) regions. The regression analysis suggested that as compared with no border controls, a complete ban on entries from abroad was associated with a change in mortality of  $10^{4 \times (-0.207)} = 0.15$ , meaning an 85% reduction in per-capita mortality (Table 5). The international travel restrictions were scored as 4 in Greenland, 3.8 in Bermuda, 3.6 in Israel, 3.5 in Czechia and New Zealand, and 3.1 in Taiwan, and at the other extreme, were scored as 1.1 in Sweden, and as 0 in Iran, Luxembourg, and the UK.

**Table 5.** Predictors of (log) Country-wide Per-capita Coronavirus Mortality by May 9 by Multivariable Linear Regression in 161 Countries.

	10 <sup>coefficient</sup>	Coefficient (SE)	95% CI	P
Duration in country (wks)	1.2668	0.1027 (0.032)	0.040 to 0.166	0.002
Time wearing masks (wks)	0.7927	-0.1009 (0.024)	-0.148 to -0.053	<0.001
Time in lockdown (wks)	0.9438	-0.0251 (0.029)	-0.082 to 0.032	0.38
Population, age ≥ 60 (%)	1.1730	0.0693 (0.009)	0.051 to 0.087	<0.001
Urbanization (%)	1.0140	0.00606 (0.004)	-0.001 to 0.013	0.09
Obesity prevalence (%)	1.0621	0.0261 (0.009)	0.009 to 0.043	0.003
Temperature, ambient (C)	1.0181	0.00779 (0.008)	-0.007 to 0.023	0.31
Testing policy (0-3)	1.2012	0.0796 (0.109)	-0.136 to 0.295	0.47
Contact tracing (0-2)	0.7030	-0.153 (0.097)	-0.344 to 0.038	0.12
Internat. travel controls (0-4)	0.6211	-0.207 (0.077)	-0.359 to -0.054	0.008
Constant	--	-7.300 (0.412)	-8.12 to -6.49	<0.001

Duration of infection in country from estimated date of first infection until 23 days before May 9, 2020 (i.e. April 16). Mask and lockdown durations run from the stated event (mask recommendation or lockdown) or estimated date of first infection in the country (whichever was later) until 23 days before May 9, 2020 (i.e. April 16).

Per-capita mortality was not significantly associated with policies regarding either testing policy ( $p=0.47$ ), or contact tracing ( $p=0.12$ , Table 5). Testing policy was scored as: no policy (0), symptomatic with exposure, travel history, hospitalization, or key occupation (1), all symptomatic (2), or open to anyone (3). Testing policy tended to be positively associated with mortality. Contact tracing was scored as: none (0), some cases (1), or all cases (2), and tended to be inversely related with per-capita mortality (though not significantly). These countervailing associations meant that as compared with a country with no testing or tracing policy, a country which opened testing to the entire public with comprehensive contact tracing might be associated with a reported change in mortality of  $10^{(3 \times 0.0796 + 2 \times (-0.153))} = 0.857$ , i.e. a 14.3% reduction in per-capita mortality (though statistical significance was not demonstrated). Thus, testing and tracing seem unlikely to account for the almost 100-fold variation in per-capita mortality between low and high mortality countries.

## Discussion.

These results confirm that over 4 months since the appearance of COVID-19 in late 2019, there remains a great deal of variation between countries in related mortality. Countries in the lower half of mortality have experienced an average COVID-19-related per-capita mortality of 1.1 deaths per million population, in contrast with an average of 94.2 deaths per million in the remaining countries. Depending on the model and dataset evaluated, independent predictors of per-capita mortality included urbanization, fraction of the population age 60 years or over, prevalence of obesity, duration of the outbreak in the country, international travel restrictions, and the period of the outbreak subject to cultural norms or government policies favoring mask-wearing by the public.

These results support the universal wearing of masks by the public to suppress the spread of the coronavirus.<sup>1</sup> Given the low levels of coronavirus mortality seen in the Asian countries which adopted widespread public mask usage early in the outbreak, it seems highly unlikely that masks are harmful. One major limitation is that evidence concerning the actual levels of mask-wearing by the public are not available for most countries. Particularly in Western countries which only recommended (rather than mandated) mask-wearing by the public, such as the United States, the practice has been steadily increasing, but change has not occurred overnight. Our analysis based on norms and policies can be compared with “intention-to-treat” analysis in a clinical trial.

Much of the randomized controlled data on the effect of mask-wearing on the spread of respiratory viruses relates to influenza. One recent meta-analysis of 10 trials in families, students, or religious pilgrims found that the relative risk for influenza with the use of face masks was 0.78, a 22% reduction, though the findings were not statistically significant.<sup>48</sup> Combining all the trials, there were 29 cases in groups assigned to wear masks, compared with 51 cases in control groups.<sup>48</sup> The direct applicability of these results to mask-wearing at the population level is uncertain. For instance, there was some heterogeneity in methods of the component trials, with one trial assigning mask wearing to the person with a respiratory illness, another to his close contacts, and the remainder to both the ill and their contacts.<sup>48</sup> Mask-wearing was inconsistent. The groups living together could not wear a mask when bathing, sleeping, eating, or brushing teeth.<sup>49-51</sup> In one of the studies reviewed, parents wore a mask during the day, but not at night when sleeping next to their sick child.<sup>51</sup> In a different trial, students were asked to wear a mask in their residence hall for at least 6 hours daily (rather than all the time).<sup>49</sup> The bottom line is that it is nearly impossible for people to constantly maintain mask wear around the people with whom they live. In contrast, wearing a mask when on public transit or shopping is quite feasible. In addition, as an infection propagates through multiple generations in the population, the benefits multiply exponentially. Even if one accepts that masks would only reduce transmissions by 22%, then after 10 cycles of the infection, mask-wearing would reduce the level of infection in the population by 91.7%, as compared with a non-mask wearing population (because  $0.78^{10} = 0.083$ ). It is highly unlikely that entire countries or populations will ever be randomized to either wear, or not wear, masks. Public policies can only be formulated based on the best evidence available.

Limits on international travel were significantly associated with lower per-capita mortality from coronavirus. As compared with no restrictions, complete shutdown of the border throughout the outbreak was associated with 85% lower per-capita mortality.

Nationwide policies to ban large gatherings and to close schools or businesses, tended to be associated with lower mortality, though not in a statistically significant fashion. However, businesses, schools, and individuals made decisions to limit contact, independent of any government policies. The adoption of numerous public health policies at the same time can make it difficult to tease out the relative importance of each.

Colder average monthly temperature was associated with higher levels of COVID-19 mortality in univariate analysis, but not when accounting for other independent variables. Nonetheless, environmental factors which could influence either human behavior or the stability and spread of virus particles are worthy of further study.

Presumably, high levels of testing might identify essentially all coronavirus-related deaths, and still higher levels of testing, combined with contact tracing, might lower mortality. However, statistical support for the benefit of testing and tracing on mortality could not be demonstrated. Policies on testing and tracing were not significantly associated with mortality. In addition, per-capita testing both early (April 4, 16) and later (May 9) were positively associated with reported coronavirus-related mortality. It seems likely that countries which test at a low level are missing many cases. We previously identified just 3 countries (Iceland, the Faeroe Islands, and the UAE) which had performed over 75,000 tests per million population by April 16, and all 3 had mortality below 1 in 46,000 at that point.<sup>53</sup> By May 9, we could add to this “high-testing” group, the Falkland Islands and Bahrain, as all 5 countries had tested over one tenth of their population. All 5 countries had a mortality of 29 per million (1 in 34,480 people) or less. The degree to which these results would apply to larger, less isolated, or less wealthy countries is unknown. Statistical support for benefit of high levels of testing might be demonstrated if additional and more diverse countries are able to test at this level.

One limitation of our study is that the ultimate source of mortality data is often from governments which may not have the resources to provide a full accounting of their public health crises, or an interest in doing so. It should be noted that the benefit of wearing masks persisted in a model which excluded data from China (because no testing data were available, Appendix Table A2). We also acknowledge that country-wide analyses are subject to the ecologic fallacy.

In summary, older age of the population, urbanization, obesity, and longer duration of the outbreak in a country were independently associated with higher country-wide per-capita coronavirus mortality. International travel restrictions were associated with lower per-capita mortality. However, other containment measures, testing and tracing policies, and the amount of viral testing were not statistically significant predictors of country-wide coronavirus mortality, after controlling for other variables. The use of masks in public is an important and readily modifiable public health measure. Societal norms and government policies supporting mask-wearing by the public were independently associated with lower per-capita mortality from COVID-19.

**Funding:** None.

**Disclosures:** The authors have no conflicts of interest.

## References.

1. Leffler CT, Ing E, McKeown CA, Pratt D, Grzybowski A. Final Country-wide Mortality from the Novel Coronavirus (COVID-19) Pandemic and Notes Regarding Mask Usage by the Public. April 4, 2020. Available from: [https://www.researchgate.net/publication/340438732\\_Country-wide\\_Mortality\\_from\\_the\\_Novel\\_Coronavirus\\_COVID-19\\_Pandemic\\_and\\_Notes\\_Regarding\\_Mask\\_Usage\\_by\\_the\\_Public](https://www.researchgate.net/publication/340438732_Country-wide_Mortality_from_the_Novel_Coronavirus_COVID-19_Pandemic_and_Notes_Regarding_Mask_Usage_by_the_Public) Accessed: May 20, 2020.
2. Leffler CT, Hogan MC. Age-dependence of mortality from novel coronavirus disease (COVID-19) in highly exposed populations: New York transit workers and residents and Diamond Princess passengers. MedRxiv. Available from: <https://doi.org/10.1101/2020.05.14.20094847> Accessed May 20, 2020.
3. Jordan RE, Adab P, Cheng KK. Covid-19: risk factors for severe disease and death. BMJ. 2020;368:m1198. Available from: <https://www.bmj.com/content/368/bmj.m1198.long> Accessed May 20, 2020.
4. Zhu Y, Xie J. Association between ambient temperature and COVID-19 infection in 122 cities from China. Science of The Total Environment. 2020 Mar 30:138201. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7142675/> Accessed May 20, 2020.
5. Squalli J. Evaluating the determinants of COVID-19 mortality: A cross-country study. medRxiv. 2020 Jan 1. Available from: <https://www.medrxiv.org/content/10.1101/2020.05.12.20099093v1> Accessed May 20, 2020.
6. Syed Q, Sopwith W, Regan M, Bellis MA. Behind the mask. Journey through an epidemic: some observations of contrasting public health responses to SARS. Journal of Epidemiology & Community Health. 2003 Nov 1;57(11):855-6.
7. Worldometers. COVID-19 Coronavirus Pandemic. Available from: [https://www.worldometers.info/coronavirus/?utm\\_campaign=homeAdUOA?Si](https://www.worldometers.info/coronavirus/?utm_campaign=homeAdUOA?Si) Accessed May 9, 2020.
8. Worldometers. COVID-19 Coronavirus Pandemic. April 5, 2020. Available from: <https://web.archive.org/web/20200405020252/https://www.worldometers.info/coronavirus/> Accessed May 17, 2020.
9. European Centre for Disease Prevention and Control. Download today's data on the geographic distribution of COVID-19 cases worldwide. Available from: <https://www.ecdc.europa.eu/en/publications-data/download-todays-data-geographic-distribution-covid-19-cases-worldwide> Accessed April 16, 2020.
10. No author listed. World Climate Guide. Available from: <https://www.climatestotravel.com/> Accessed May 14, 2020.

11. No author listed. List of Cities by Average Temperature. Available from: [https://en.wikipedia.org/wiki/List\\_of\\_cities\\_by\\_average\\_temperature](https://en.wikipedia.org/wiki/List_of_cities_by_average_temperature) Accessed May 14, 2020.
12. Countries in the world by population (2020). Available from: <https://www.worldometers.info/world-population/population-by-country/> Accessed May 16, 2020.
13. United Nations Population Division, New York. World Population Prospects: The 2019 Revision. Population, surface area, and density. Available from: [http://data.un.org/Docs/SYB/PDFs/SYB61\\_T02\\_Population,%20Surface%20Area%20and%20Density.pdf](http://data.un.org/Docs/SYB/PDFs/SYB61_T02_Population,%20Surface%20Area%20and%20Density.pdf) Accessed May 16, 2020.
14. International Monetary Fund. World Economic Outlook Database, October 2019. Report for Selected Countries and Subjects. Available from: <https://www.imf.org/en/Publications/SPROLLs/world-economic-outlook-databases#sort=%40imfdate%20descending> Accessed April 11, 2020.
15. World Bank. Urban population (% of total population). United Nations Population Division. World Urbanization Prospects: 2018 Revision. Available from: <https://data.worldbank.org/indicator/SP.URB.TOTL.in.zs> Accessed May 16, 2020.
16. The American Cancer Society. The Tobacco Atlas. 2020 Available from: <https://tobaccoatlas.org/topic/prevalence/> Accessed April 11, 2020.
17. WHO Report on the Global Tobacco Epidemic, 2009. Surveys of adult tobacco use in WHO Member States. Available from: [https://www.who.int/tobacco/mpower/2009/Appendix\\_VIII-table\\_1.pdf](https://www.who.int/tobacco/mpower/2009/Appendix_VIII-table_1.pdf) Accessed May 16, 2020.
18. World Bank. Smoking prevalence, males (% of adults). <https://data.worldbank.org/indicator/SH.PR.V.SMOK.MA?locations=ME> Accessed May 16, 2020.
19. World Bank. Smoking prevalence, females (% of adults). Available from: <https://data.worldbank.org/indicator/SH.PR.V.SMOK.FE?locations=ME> Accessed May 16, 2020.
20. Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, Meredith HR, Azman AS, Reich NG, Lessler J. The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and application. *Annals of Internal Medicine*. Mar 10, 2020.
21. Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, Cuomo-Dannenburg G, Thompson H, Walker P, Fu H, Dighe A. Estimates of the severity of COVID-19 disease. *Lancet Infectious Disease*. 2020: 1-9.
22. Leffler CT, Ing E, Lykins JD, McKeown CA, Grzybowski A. Prevention of the spread of coronavirus using masks. *Annals of Internal Medicine*. April 30, 2020. Available from: <https://www.acpjournals.org/doi/10.7326/M20-1342> Accessed May 20, 2020.
23. European Centre for Disease Prevention and Control. COVID-19 Coronavirus data. Available from: <https://data.europa.eu/euodp/en/data/dataset/covid-19-coronavirus-data> Accessed April 16, 2020.

24. Feng S, Shen C, Xia N, Song W, Fan M, Cowling BJ. Rational use of face masks in the COVID-19 pandemic. *Lancet Respir Med*. 2020 Mar 20. pii: S2213-2600(20)30134-X. doi: 10.1016/S2213-2600(20) 30134-X.
25. Thongthab S. Ministry of Public Health Enhances Disease Prevention and Control. The Bangkok Insight. January 31, 2020. Available from: <https://www.thebangkokinsight.com/282702/> Accessed May 9, 2020.
26. Takahashi. Amid virus outbreak, Japan stores scramble to meet demand for face masks. *Japan Times*. Available from: <https://www.japantimes.co.jp/news/2020/01/31/national/coronavirus-japan-surgical-masks/#.Xrc8Z2hKhPY> Accessed May 9, 2020.
27. Kim ET. How South Korea Solved Its Face Mask Shortage: Neighborhood pharmacists and government intervention were the secret weapons. *New York Times*. April 1, 2020. Available from: <https://www.nytimes.com/2020/04/01/opinion/covid-face-maskshortage.html> Accessed April 17, 2020.
28. Blanchard B. Taiwan ups Chinese visitor curbs, to stop mask exports. *Reuters*. January 27, 2020. Available from: <https://www.reuters.com/article/us-china-health-taiwan/taiwan-ups-chinese-visitor-curbs-to-stop-mask-exports-idUSKBN1ZQ1C6> Accessed May 9, 2020.
29. Cheung E. China coronavirus: death toll almost doubles in one day as Hong Kong reports its first two cases. *South China Morning Post*. January 22, 2020. Available from: <https://www.scmp.com/news/hong-kong/health-environment/article/3047193/china-coronavirus-first-case-confirmed-hong-kong> Accessed May 9, 2020.
30. Department of Health, Hong Kong. Latest recommendations by Scientific Committee on Emerging and Zoonotic Diseases and Scientific Committee on Infection Control after reviewing cases of novel coronavirus infection. January 24, 2020. Available from: <https://www.info.gov.hk/gia/general/202001/24/P2020012400762.htm> Accessed May 9, 2020.
31. Taylor K. I've been traveling in Asia for 3 weeks amid the deadly coronavirus outbreak, and actually catching the virus is far from my biggest fear. *Business Insider*. February 18, 2020. Available from: <https://www.businessinsider.com/travel-in-asia-during-coronavirus-outbreak-2020-2#so-for-now-my-travels-continue-17> Accessed May 9, 2020.
32. Harun HN, Yusof TA, Solhi F. Demand for face masks, hand sanitisers soars. *New Straits Times*. January 30, 2020. Available from: <https://www.nst.com.my/news/nation/2020/01/561250/demand-face-masks-hand-sanitisers-soars> Accessed May 9, 2020.
33. Khan S. Cambodia Confirms First Coronavirus Case. *VOA Khmer*. January 27, 2020. Available from: <https://www.voanews.com/science-health/coronavirus-outbreak/cambodia-confirms-first-coronavirus-case> Accessed May 9, 2020.
34. Khan S. Cambodian Businesses Embrace Protective Measures Against Coronavirus. *VOA Khmer*. March 30, 2020. Available from: <https://www.voacambodia.com/a/cambodian-businesses-embrace-protective-measures-against-coronavirus/5352023.html>

Accessed May 9, 2020.

35. Lasco G. Why Face Masks Are Going Viral. Sapiens. February 7, 2020. Available from: <https://www.sapiens.org/culture/coronavirus-mask/> Accessed May 9, 2020.
36. Baljmaa.T. SEC: MNT 150,000 fine for not wearing face masks. April 14, 2020. Available from: <https://www.montsame.mn/en/read/222259> Accessed May 9, 2020.
37. No author listed. Laos reports no case of COVID-19, to import face masks from Vietnam. Vietnam Times. March 20, 2020. Available from: <https://vietnamtimes.org.vn/laos-reports-no-case-of-covid-19-to-import-face-masks-from-vietnam-18566.html> Accessed May 9, 2020.
38. AFP. Coronavirus: Maduro tells Venezuelans to make their own masks. March 17, 2020. Available from: <https://www.youtube.com/watch?v=EjTsjJNCAuc> Accessed May 15, 2020.
39. Berwick A, Nava M, Kinosian S. Venezuela confirms coronavirus cases amid public health concerns. Reuters. March 14, 2020. Available from: <https://www.physiciansweekly.com/venezuela-confirms-coronavirus-cases/> Accessed May 15, 2020.
40. Štefúnová I. "Slovensko pritvrdilo v boji s vírusom. Nosenie rúšok je povinné". Pravda. March 15, 2020. Available from: <https://spravy.pravda.sk/domace/clanok/545657-slovensko-pritvrdilo-v-boji-s-virusom-nosenie-rusok-je-povinne/> Accessed May 15, 2020.
41. Public Health Authority, Slovakia. "Opatrenie Úradu verejného zdravotníctva Slovenskej republiky pri ohrození verejného zdravia" March 24, 2020. Available from: [http://www.uvzsr.sk/docs/info/covid19/Opatrenie\\_UVZSR\\_povinnost\\_nosit\\_ruska\\_2403\\_2020.pdf](http://www.uvzsr.sk/docs/info/covid19/Opatrenie_UVZSR_povinnost_nosit_ruska_2403_2020.pdf) Accessed May 15, 2020.
42. Veronika B. Could Czech's Measure to Fight Coronavirus Save Thousands of Lives? Prague Morning. April 4, 2020. Available from: <https://www.praguemorning.cz/could-czechs-measure-to-fight-coronavirus-save-thousands-of-lives-2/> Accessed May 7, 2020.
43. COVID-19 Information. US Embassy in Uzbekistan. May 1, 2020. Available from: <https://uz.usembassy.gov/covid-19-information/> Accessed May 9, 2020.
44. STA. COVID-19 & Slovenia, Night 29 March: Movement Restrictions, Mandatory Masks, More Aid for Individuals. Total Slovenia News. March 29, 2020. Available from: <https://www.total-slovenia-news.com/politics/5951-covid-19-slovenia> Accessed May 8, 2020.
45. ORF. Regierung verschärft Maßnahmen. March 30, 2020. Available from: <https://orf.at/stories/3159909/> Accessed May 15, 2020.
46. Reuters. Austria widening face-mask requirement while loosening lockdown. April 6, 2020. Available from: <https://www.reuters.com/article/health-coronavirus-austria-masks/austria-widening-face-mask-requirement-while-loosening-lockdown-idUSV9N28601G> Accessed May 15, 2020.
47. Borissov B. "Здравният министър даде заден за задължителните маски". March 31, 2020. Available from:

<https://www.segabg.com/hot/category-bulgaria/gledayte-na-zhivo-vutreshniyat-i-zdravniyat-ministur-razyasnyavat-merkite/> Accessed May 15, 2020.

48. Xiao J, Shiu EY, Gao H, Wong JY, Fong MW, Ryu S, Cowling BJ.

Nonpharmaceutical Measures for Pandemic Influenza in Nonhealthcare Settings- Personal Protective and Environmental Measures. *Emerging Infectious Diseases*. 2020; 26(5): 967-75.

49. Aiello AE, Perez V, Coulborn RM, Davis BM, Uddin M, Monto AS. Facemasks, hand hygiene, and influenza among young adults: a randomized intervention trial. *PLoS One*. 2012;7:e29744.

50. Cowling BJ, Fung RO, Cheng CK, Fang VJ, Chan KH, Seto WH, et al. Preliminary findings of a randomized trial of non-pharmaceutical interventions to prevent influenza transmission in households. *PLoS One*. 2008;3:e2101.

51. MacIntyre CR, Cauchemez S, Dwyer DE, Seale H, Cheung P, Browne G, et al. Face mask use and control of respiratory virus transmission in households. *Emerg Infect Dis*. 2009;15:233–41.

53. Leffler CT, Zhan S. Suppression of the Covid-19 Outbreak by Mass Testing and Tracing, and Other Measures: Real-World Data. April 17, 2020. Available from:

[https://www.researchgate.net/publication/340720271\\_Suppression\\_of\\_the\\_Covid-19\\_Outbreak\\_by\\_Mass\\_Testing\\_and\\_Tracing\\_and\\_Other\\_Measures\\_Real-World\\_Data?\\_sg=qmL3zeijOGrKwY3pKS5BGQ\\_-H9KzOsEYC2mqy\\_71nifHMme737SciFndah2rLwHEKXAIQQN\\_-JNV5\\_rCOLawtoo36uhRDVW7cXDeCEId.MivUKSPrk2FO3h6BPweYc9NwomngmcwdCrIxxVDJWBozq01jbOPpwb5y4G1IVs3Gsgu2QZcmND\\_XnYcIY\\_j1g](https://www.researchgate.net/publication/340720271_Suppression_of_the_Covid-19_Outbreak_by_Mass_Testing_and_Tracing_and_Other_Measures_Real-World_Data?_sg=qmL3zeijOGrKwY3pKS5BGQ_-H9KzOsEYC2mqy_71nifHMme737SciFndah2rLwHEKXAIQQN_-JNV5_rCOLawtoo36uhRDVW7cXDeCEId.MivUKSPrk2FO3h6BPweYc9NwomngmcwdCrIxxVDJWBozq01jbOPpwb5y4G1IVs3Gsgu2QZcmND_XnYcIY_j1g) Accessed May 17, 2020.

54. Oxford Coronavirus Government Response Tracker. University of Oxford. Available from: <https://www.bsg.ox.ac.uk/research/research-projects/coronavirus-government-response-tracker> Accessed May 24, 2020.

55. Hale T, Angrist N, Kira B, Phillips T, Webster S. Variation in government responses to COVID-19. BSG-WP-2020/032. Version 6.0. University of Oxford. Blavatnik School of Government. May 2020 Available from:

<https://www.bsg.ox.ac.uk/sites/default/files/2020-05/BSG-WP-2020-032-v6.0.pdf> Accessed May 31, 2020.

56. Howard J. What Countries Require Masks in Public or Recommend Masks? #Masks4all Available from: <https://masks4all.co/what-countries-require-masks-in-public/> Accessed May 30, 2020.

57. Central Intelligence Agency. The World Factbook. Country Comparison. Obesity. Adult Prevalence Rate. Available from: <https://www.cia.gov/library/publications/the-world-factbook/rankorder/2228rank.html> Accessed May 29, 2020.

58. Parris D. The Alarming Increase of Diabetes in Bermuda. *West Indian Med J*. 2014; 63(7): 685-686. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4668975/> Accessed May 29, 2020.

59. Fu FH. A comparison of lifestyle management practices of residents of three Chinese cities: Hong Kong, Macau and Weihai (Shandong). *International Journal of Sports*. 2013;3:115-26.

60. Ramachandran A, Snehalatha C. Rising burden of obesity in Asia. *Journal of obesity*. 2010 Aug 30;2010.
61. Chobanyan N, Allison Kruger K, Nebb S, Jackson G, Asin V. Evaluation of Environmental Risk Factors for Type 2 Diabetes in Sint Maarten. *J Environ Anal Toxicol*. 2016;6(386):2161-0525.
62. Bjerregaard P, Jørgensen ME, Greenland Population Study Group. Prevalence of obesity among Inuit in Greenland and temporal trend by social position. *American Journal of Human Biology*. 2013 May;25(3):335-40.
63. Veyhe AS, Andreassen J, Halling J, Grandjean P, Petersen MS, Weihe P. Prevalence of type 2 diabetes and prediabetes in the Faroe Islands. *Diabetes research and clinical practice*. 2018 Jun 1;140:162-73.
64. Abdeen Z, Jildeh C, Dkeideek S, Qasrawi R, Ghannam I, Al Sabbah H. Overweight and obesity among Palestinian adults: analyses of the anthropometric data from the first national health and nutrition survey (1999-2000). *Journal of obesity*. 2012 Feb 21;2012.
65. Grol ME, Eimers JM, Alberts JF, Bouter LM, Gerstenbluth I, Halabi Y, Van Sonderen E, Van den Heuvel WJ. Alarming high prevalence of obesity in Curacao: data from an interview survey stratified for socioeconomic status. *International journal of obesity*. 1997 Nov;21(11):1002-9.
66. Corsenac P, Annesi-Maesano I, Hoy D, Roth A, Rouchon B, Capart I, Taylor R. Overweight and obesity in New Caledonian adults: Results from measured and adjusted self-reported anthropometric data. *Diabetes research and clinical practice*. 2017 Nov 1;133:193-203.
67. Favier F, Jaussent I, Le Moullec N, Debussche X, Boyer MC, Schwager JC, Papoz L, REDIA Study Group. Prevalence of Type 2 diabetes and central adiposity in La Reunion Island, the REDIA Study. *Diabetes research and clinical practice*. 2005 Mar 1;67(3):234-42.
68. Grievink L, Alberts JF, O'niel J, Gerstenbluth I. Waist circumference as a measurement of obesity in the Netherlands Antilles; associations with hypertension and diabetes mellitus. *European journal of clinical nutrition*. 2004 Aug;58(8):1159-65.
69. Solet JL, Baroux N, Pochet M, Benoit-Cattin T, De Montera AM, Sissoko D, Favier F, Fagot-Campagna A. Prevalence of type 2 diabetes and other cardiovascular risk factors in Mayotte in 2008: the MAYDIA study. *Diabetes & metabolism*. 2011 Jun 1;37(3):201-7.
70. Daigne JL, Atallah A, Boissin JL, Jean-Baptiste G, Kangambega P, Chevalier H, Balkau B, Smadja D, Inamo J. The prevalence of overweight and obesity, and distribution of waist circumference, in adults and children in the French Overseas Territories: the PODIUM survey. *Diabetes & metabolism*. 2012 Nov 1;38(5):404-11.
71. Guariguata L, Brown C, Sobers N, Hambleton I, Samuels TA, Unwin N. An updated systematic review and meta-analysis on the social determinants of diabetes and related risk factors in the Caribbean. *Revista Panamericana de Salud Pública*. 2018;42: e171.
72. World Health Organization. Nutrition, Physical Activity and Obesity San Marino. Available from: [http://www.euro.who.int/data/assets/pdf\\_file/0016/243322/San-Marino-WHO-Country-Profile.pdf?ua=1](http://www.euro.who.int/data/assets/pdf_file/0016/243322/San-Marino-WHO-Country-Profile.pdf?ua=1) Accessed May 29, 2020.

73. States of Jersey. Health Profile for Jersey 2014...with comparisons of Guernsey, English regions and Europe. Available from: <https://www.gov.je/SiteCollectionDocuments/Government%20and%20administration/R%20Health%20Profile%20Jersey%202014%2020140410%20MM%20v1.pdf> Accessed May 29, 2020.
74. Isle of Man Government. Island's plan to tackle childhood obesity launched. January 22, 2013. Available from: <https://www.gov.im/news/2013/jan/22/islands-plan-to-tackle-childhood-obesity-launched/> Accessed May 29, 2020.
75. Gibraltar Health Authority. Health and Lifestyle. Survey Report of the Adult Population of Gibraltar. 2015. Available from: [https://www.gha.gi/wp-content/uploads/2016/12/GHA\\_Lifestyle\\_Report2015LR.pdf](https://www.gha.gi/wp-content/uploads/2016/12/GHA_Lifestyle_Report2015LR.pdf) Accessed May 29, 2020.
76. James J, Soyibo AK, Hurlock L, Gordon-Strachan G, Barton EN. Cardiovascular Risk Factors in an Eastern Caribbean Island: Prevalence of Non-Communicable Chronic Diseases and Associated Lifestyle Risk Factors for Cardiovascular Morbidity and Mortality in the British Virgin Islands. *West Indian Med J* 2012; 61(4):429-36.
77. Chung K. Macau confirms second patient infected with Chinese coronavirus. *South China Morning Post*. Available from: <https://www.scmp.com/news/china/article/3047337/macau-confirms-second-patient-infected-chinese-coronavirus> Accessed May 26, 2020.
78. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497-506.
79. Wu H, Yu S. Masks on, Chinese start holiday travels as alarm mounts over mystery virus. *Reuters*. Jan. 20, 2020. Available from: <https://www.reuters.com/article/us-china-health-pneumonia-masks/masks-on-chinese-start-holiday-travels-as-alarm-mounts-over-mystery-virus-idUSKBN1ZJ0VU> Accessed May 26, 2020.
80. No author listed. All people in public places in Wuhan required to wear masks, local government says. *China Daily*. Jan. 22, 2020. Available from: <http://www.chinadaily.com.cn/a/202001/22/WS5e285cada310128217272d84.html> Accessed May 26, 2020.
81. Epidemic Prevention and Control Group. CDC. Notice regarding the issuance of guidelines for the protection of people with different risks of new coronavirus infection and guidelines for the use of pneumonia masks for the prevention of new coronavirus infection. Jan. 31, 2020. Available from: <http://www.nhc.gov.cn/jkj/s7916/202001/a3a261dabfcf4c3fa365d4eb07ddab34.shtml> Accessed May 26, 2020.
82. No author listed. Face mask and fruit prices soar in Sudan. *Dabanga*. March 16, 2020. Available from: <https://www.dabangasudan.org/en/all-news/article/face-mask-and-fruit-prices-soar-in-sudan>
83. No author listed. Sudan Flash Update, 16 Mar 2020: Sudan creates two COVID-19 isolation centres in Khartoum State. Available from: <https://reliefweb.int/report/sudan/sudan-flash-update-16-mar-2020-sudan-creates-two-covid-19-isolation-centres-khartoum>

Accessed May 26, 2020.

84. Health Alert: U.S. Embassy Sarajevo, Bosnia and Herzegovina. Available from: <https://ba.usembassy.gov/health-alert-u-s-embassy-sarajevo-bosnia-and-herzegovina-march-30-2020/> Accessed May 6, 2020.

85. Cuba Suspends International Passenger Flights. Xinhua in Telesur. Available from: <https://www.telesurenglish.net/news/Cuba-Suspends-International-Passenger-Flights-20200402-0012.html> Accessed May 29, 2020.

86. Ministry of Health, Singapore. Updates on COVID-19 (Coronavirus Disease 2019) Local

Situation. <https://www.moh.gov.sg/covid-19> Accessed April 4, 2020.

87. Ministry of Health, Singapore. Official Update of COVID -19 Situation in Singapore. <https://experience.arcgis.com/experience/7e30edc490a5441a874f9efe67bd8b89>

Accessed April 4, 2020.

88. Sim D. Coronavirus: what's behind Singapore's U-turn on wearing masks? South China Morning Post. April 3, 2020. Available from: <https://www.scmp.com/week-asia/health-environment/article/3078399/coronavirus-whats-behind-singapores-u-turn-wearing> Accessed May 30, 2020.

89. Thacker T. Amid outbreak, Health Ministry recommends homemade face masks. Economic times. Apr 4, 2020. Available from:

<https://economictimes.indiatimes.com/industry/healthcare/biotech/healthcare/amid-outbreak-health-ministry-recommends-homemade-face-masks/articleshow/74978175.cms> Accessed May 29, 2020.

90. Muraya J. Kenya: Masks Now Mandatory in Public Places, Kenya Declares. All Africa. April 5, 2020. Available from: <https://allafrica.com/stories/202004060049.html> Accessed May 29, 2020.

91. Chilunga Z. Malawi: Mutharika Urges Malawi Unity and 'Steadfast' in COVID-19 Fight - Announce New Measures to Stop Spread of Outbreak. Nyasa Times. April 4, 2020. Available from: <https://allafrica.com/stories/202004060182.html> Accessed May 29, 2020.

92. Yulisman L. Coronavirus: Indonesia makes face masks compulsory as death toll nears 200. Straits Times. Available from:

<https://www.straitstimes.com/asia/se-asia/coronavirus-indonesia-orders-citizens-to-wear-masks-as-infections-rise> Accessed May 29, 2020.

93. No author listed. Mandatory wearing of face masks will soon be enforceable. Antigua Nice. Available from:

<http://www.antiguanice.com/v2/client.php?id=943&news=12425>

Accessed May 29, 2020.

94. No author listed. La Tunisie décide l'obligation du port du masque pour toute la population. GNet News. April 7, 2020. Available from: <https://news.gnet.tn/la-tunisie->

[decide-lobligation-du-port-du-masque-pour-toute-la-population/](#) Accessed May 30, 2020.

95. Benin Police Enforce Mask Wearing In Bid To Stop Virus. Agence France Presse. April 8, 2020. Available from: <https://www.barrons.com/news/benin-police-enforce-mask-wearing-in-bid-to-stop-virus-01586363106> Accessed May 30, 2020.

96. No author listed. Mozambique: Government Orders Wearing of Masks. allAfrica. April 9, 2020. Available from: <https://allafrica.com/stories/202004091000.html> Accessed May 30, 2020.

97. No author listed. Bahrain makes face masks compulsory in public, allows shops to re-open. Arab News. April 9, 2020. Available from: <https://www.arabnews.com/node/1656026/middle-east>

Accessed May 30, 2020.

98. U.S. Embassy in Gabon. COVID-19 Information for Gabon and São Tomé and Príncipe. April 23, 2020. Available from: <https://web.archive.org/web/20200426105023/https://ga.usembassy.gov/u-s-citizen-services/coronavirus-update/> Accessed May 30, 2020.

99. Samuel G. Ethiopia Outlaws Handshakes, Obliges Masks in Public Places. Addis Fortune. April 12, 2020. Available from: <https://addisfortune.news/ethiopia-outlaws-handshakes-obliges-masks-in-public-places/> Accessed May 30, 2020.

100. Unah L, Mussa C. Masks, bans and questions: Inside Cameroon's COVID-19 response. April 23, 2020. Available from: <https://www.aljazeera.com/news/2020/04/masks-bans-questions-cameroon-covid-19-response-200422134140013.html> Accessed May 26, 2020.

## Appendix. Supplemental Multivariable Analyses.

**Table A1.** Predictors of (log) Country-wide Per-capita Coronavirus Mortality by May 9 by Multivariable Linear Regression in 198 Countries.

	$10^{\text{coefficient}}$	Coefficient (SE)	95% CI	P
Duration in country (weeks)	1.455	0.163 (0.032)	0.099 to 0.226	<0.001
Time wearing masks (weeks)	0.708	-0.150 (0.027)	-0.203 to -0.098	<0.001
Time in lockdown (weeks)	0.979	-0.009 (0.035)	-0.078 to 0.059	0.79
Population, age $\geq$ 60 (%)	1.147	0.059 (0.009)	0.042 to 0.077	<0.001
Urbanization (%)	1.019	0.00836 (0.003)	0.002 to 0.014	0.007
Constant	--	-7.774 (0.240)	-8.247 to -7.301	<0.001

Duration of infection in country from estimated date of first infection until 23 days before May 9, 2020 (i.e. April 16). Mask and lockdown durations run from the stated event (mask recommendation or lockdown) or estimated date of first infection in the country (whichever was later) until 23 days before May 9, 2020 (i.e. April 16).

**Table A2.** Predictors of (log) Country-wide Per-capita Coronavirus Mortality by May 9 by Multivariable Linear Regression in 179 Countries.

	10 <sup>coefficient</sup>	Coefficient (SE)	95% CI	P
Duration in country (weeks)	1.4907	0.173 (0.034)	0.106 to 0.241	<0.001
Time wearing masks (weeks)	0.7589	-0.120 (0.029)	-0.178 to -0.062	<0.001
Time in lockdown (weeks)	0.9667	-0.015 (0.040)	-0.094 to 0.064	0.71
Population, % age 60 or over	1.1010	0.042 (0.012)	0.019 to 0.065	0.001
Urbanization (%)	1.0140	0.00606 (0.004)	-0.002 to 0.014	0.12
Obesity prevalence (%)	1.0503	0.021 (0.010)	0.002 to 0.041	0.03
Temperature (C)	0.9772	-0.00999 (0.009)	-0.028 to 0.008	0.28
Testing (log per cap., by May 9)	1.1016	0.042 (0.116)	-0.187 to 0.271	0.72
Constant	--	-7.637 (0.600)	-8.82 to -6.45	<0.001

Based on 179 countries with both obesity and testing data by May 9. Duration of infection in country from estimated date of first infection until 23 days before May 9, 2020 (i.e. April 16). Mask and lockdown durations run from the stated event (mask recommendation or lockdown) or estimated date of first infection in the country (whichever was later) until 23 days before May 9, 2020 (i.e. April 16).

S