Estimation and monitoring of COVID-19's transmissibility from publicly available data

Antonio Silveira*, **Antonio Pereira**

Department of Electrical and Biomedical Engineering, Federal University of Pará, 66075-110, Belém, PA, Brazil

Correspondence*:
Antonio Silveira
asilveira@ufpa.br

2 ABSTRACT

The COVID-19 pandemic began in the city of Wuhan, China, at the end of 2019 and quickly
spread worldwide. The disease is caused by contact with the SARS-CoV-2 virus, which probably
jumped from an animal host to humans. SARS-CoV-2 infects various tissues in the body, notably
the lungs, and patients usually die from respiratory complications. Mathematical models of the
disease have been instrumental to guide the implementation of mitigation strategies aimed at
slowing the spread of the disease. One of the key parameters of mathematical models is the
basic reproduction ratio R_0, which measures the degree of infectivity of affected individuals. The
goal of mitigation is to reduce R_0 as close or below 1 as possible, as it means that new infections
are in decline. In the present work, we use the recursive least-squares algorithm to establish the
stochastic variability of a time-varying $R_0(t)$ from eight different countries: Argentina, Belgium,
Brazil, Germany, Italy, New Zealand, Spain, and the United States of America. The proposed
system can be implemented as an online tracking application providing information about the
dynamics of the pandemic to health officials and the public at large.

Keywords: COVID-19, Epidemic Spreading, Pattern Recognition, Mathematical modelling, Transmission dynamics, Disease Prediction.

1 INTRODUCTION

On the 11th of March 2020, the World Health Organization (WHO) declared the 2019 coronavirus disease
(COVID-19) a global pandemic [1]. COVID-19 is caused by the SARS-CoV-2 coronavirus and was first
reported in Wuhan, China, in December 2019 [2]. Since then, COVID-19 has spread globally with millions
of laboratory-confirmed cases and hundreds of thousands of deaths [3]. The median incubation period of
COVID-19 is 5.1 days and nearly all infected persons who have symptoms will do so within 12 days of
infection [4]. However, an unprecedented characteristic of COVID-19 is its capacity for asymptomatic
transmission [5], which contributes to increase the probability of transmission [6]. So far, there is no
specific treatment for the disease and many research teams are currently working on a vaccine that,
optimistically, will only be available in 2021. Even then, it will take some time to inoculate a significant
share of the population. Meanwhile, hospital structures around the globe (e.g. intensive care units' (ICU)
beds, ventilators, etc.) are becoming overwhelmed with new patients and the increasing caseload will
prove most catastrophic for poor countries, which lack the adequate health-care capacity to deal with the
unparalleled demand posed by COVID-19 [7, 8].

So far, efforts to contain the spread of the disease have focused on the adoption of non-pharmaceutical
interventions (NPI) based on population behavioral change and social distancing, such as banning large
gatherings, enforcing the use of face masks, washing hands, or imposing severe lockdowns [9, 10]. Due to

significant uncertainties regarding the transmissibility of SARS-CoV-2 as well as other political, social,
and economic considerations, it is necessary to delineate effective social distancing policies which are
able to alleviate COVID-19's burden on healthcare structure and borrow time for the development of a
vaccine or drug candidates while also simultaneously reducing the socioeconomic strain of living in a
locked-down, confined society [11]. The effective monitoring of the epidemic's dynamics plays a crucial
role in the ongoing containment efforts, but will also continue to do so for some time ahead when social
distancing control measures are eventually relaxed and the first wave of the pandemic is followed some
months later by second or third waves of infection that may be more severe than the first [12].

Mathematical models, by providing a quantitative framework for hypothesis evaluation and the estimation
of changes in transmission of infectious diseases over time and space, can indicate whether containment
measures are having a measurable effect while guiding the design of alternative interventions [13].
Mathematical models vary in many aspects, including complexity in terms of the number of variables
and parameters used, spatial and temporal resolution (e.g., discrete vs. continuous-time), and design
(e.g., deterministic or stochastic) [14]. Mechanistic models of the susceptible-infected-recovered SIR type
[15, 16] are the standard framework for a wide array of infectious diseases, including COVID-19 (see,
for example, Li et al. [17], Weitz et al. [18]). However, parameter estimates for a given model are subject
to two major sources of uncertainty: the noise intrinsic to the data and the ad-hoc assumptions used for
ascertaining parameter estimates [14].

The basic reproduction number [19], R_0, or the average number of new infections caused by an infectious
individual [20], is widely used to characterize the dynamics of infectious outbreaks and guide policy-
planning. For instance, the average R_0 at the start of the SARS pandemic in 2003 was estimated to be around
2.75 and was later reduced to 1 due to intervention strategies, including isolation and quarantine activities
[21]. R_0 is an imprecise estimate that is rarely measured directly (it is the product of disease parameters:
duration, opportunity, transmission probability and susceptibility - DOTS) and rests on particular model
structures and assumptions [22]. Modellers face many challenges when trying to provide robust estimations
of R_0 in the current pandemics, such as the existence of superspreaders, the fact that SARS-CoV-2 can
also be spread by asymptomatic individuals, and the scarce availability of testing supplies [17]. Several
methods have been proposed to track trends in R_0 during the course of an epidemic [23, 24, 25, 26, 27].
The access to reliable estimates of R_0 could provide useful information about the efficacy of containment
measures and allow their effective management in order to keep hospitalization rates within a desired
approximate range [28].

In the present study, we model the transmission dynamic of COVID-19 in eight countries (Fig. 1) using
least-squares algorithm (LSA) techniques. The criteria used to select the eight countries were due to their
geographical location in either the northern/southern hemisphere or due to specificities regarding their first
response to the pandemic threat as reported in the news media. Some (e.g. New Zealand, Argentina) were
very effective in responding rapidly to the pandemic by implementing lock down measures, while others (e.g.
USA and Brazil) initially downplayed the virus' threat and took longer to initiate containment protocols.
Our goal is to contribute to understanding the spread of SARS-CoV-2 and compare R_0 uncertainty arising
from noise in the time series data gathered from public online sources. We used machine learning algorithms
to optimally estimate a time-varying $R_0(t)$, which allows the monitoring of the ongoing pandemic in almost
real-time. We compared the daily country reports for R_0 to those we estimated in the present work in order
to assess the reliability of official data. Besides, LSA-based techniques were used to reveal clues regarding
the dynamics of $R_0(t)$ and its stochasticity in terms of the linear power of the estimation error and provide
information on how such stochastic behavior is correlated to the outcome of the ongoing pandemics.

The LSA is one of the most popular estimation methods in machine-learning and has been used in many
scientific and engineering applications [29, 30, 31], including epidemiology, for calibrating mathematical
models' parameters based on time series data while also generating disease forecasts in the near or long
terms [14, 32, 33, 34]. While it has been used for centuries as a classic curve-fitting technique [30, 31], it
is still a basic tool in modern data science because of its least-squares Euclidean ℓ_2-norm minimization that
is advantageous over other norms and metrics, such as the ℓ_∞ and ℓ_1 norms, granting reduced sensibility
to outliers due to the squared error [35, 31]. Higher-order norms and the use of more generalized cost
functions, both linear and non-linear, often require gradient descent solutions which are the foundation of
deep learning algorithms with none, many, or infinite solutions. In comparison, LSA is computationally
inexpensive and can even be solved analytically.

LSA estimation is based on the least mean-squares algorithm (LMS), a special case of Bayesian estimation
and the foundation of classical optimal estimation theory, where its applicability is commonly attributed to
offline *batch processing*, i.e., the whole data set must be *a priori* available and be processed at once "in one
single step into the estimate" [35], involving complex algebraic procedures such as the inversion of high
order matrices, with dimension equal to the length of the data set. Its offshoot, the recursive least-squares
algorithm (RLS), has been used for real-time estimation applications in diverse areas such as signal and
data processing, communications, and control systems [36, 37, 38], since it benefits from the recursive
method and avoids matrix inversion by working one sample at a time, both speeding up processing and
avoiding a possible ill conditioned (non-invertible) information matrix formation.

Some of the advantages of RLS over LMS, and other more complex gradient descent-based estimation
methods, are: its recursive or sequential processing, which requires less memory over a single iteration step;
the possibility to capture the dynamics of non-linear and time-varying systems; its native discrete-time
synthesis to deal with discrete-time signals; and its dead-beat convergence characteristics, i.e., convergence
in minimal time [30]. However, one important drawback is the possibility of RLS getting stuck in local
minima and becoming unstable when tracking certain classes of signals, as remarked in [30]. Thankfully,
the exponential functions used in epidemic modeling are within the stable scope of convergence of the
RLS method.

In the present work, we compare the performance of both RLS and LMS on estimating $R_0(t)$, in terms
of processing speed and accuracy. While both are LSA-based methods apparently differing just by the
sequential-recursive and the batch-processing forms of implementation, we are dealing with a general case
problem of estimating a random variable $R_0(t)$ given random variables as well in a maximum-likelihood
estimation problem that "implies ignorance of any statistics of the estimated variable" [35], $R_0(t)$. Thus, if
the available measurements are independent, RLS and LMS should achieve the same accuracy performance.
Otherwise, the RLS should perform better.

Our modeling approach is based on a discrete-time MIMO setup which uses a well established concept in
the design of aerospace navigation systems called sensor fusion and which is summarized by the following
statement "one always gains by adding a new measurement in terms of navigation error, and this no matter
how bad the additional measurement is" [39, Remark 4.8]. In the present work, we combine data from
the number of susceptibles, infections, recoveries, deaths, and individual parameters of three coupled
differential equations in order to improve the estimation of $R_0(t)$.

Country	Population	GDP	HDI	Covid-19		Date of 100th case	Day control started (in days since 100th case)			
				Cases	Deaths		L1	L2	L3	L4
Argentina	44,938,712	9,887	0.830	9,918	416	03/19/20	prior	prior	prior	prior
Belgium	11,589,623	46,724	0.919	56,511	9,212	03/06/20	4	6	11	-
Brazil	210,147,125	8,955	0.761	312,000	20,112	03/13/20	6	-	-	-
Germany	83,149,300	46,653	0.939	179,000	8,320	03/01/20	10	12	13	21
Italy	60,317,116	32,947	0.883	228,000	32,486	02/23/20	7	10	17	27
New Zealand	4,984,340	41,616	0.921	1,154	21	03/19/20	prior	2	4	6
Spain	47,007,367	30,774	0.893	235,000	28,628	03/2/20	8	13	13	13
USA	328,239,523	67,426	0.920	1,610,000	92,513	03/8/20	4	12	12	12

Figure 1. Sociodemographic data for the eight countries targeted in this study and associated COVID-19 data. COVID-19 data accessed on 22/05/20 on `https://coronavirus.jhu.edu/map.html`. Intervention data according to [40]. GDP per capita in US$. Interventions are categorised into Alert Levels 1 to 4 according to the New Zealand framework (L1 = Level 1, Prepare; L2 = Level 2, Reduce; L3 = Level 3, Restrict; L4 =Level 4, Eliminate) [40].

2 METHODS

2.1 Data sources

119   The COVID-19 data used in this report is publicly available from The Center for Systems Science and
120   Engineering of the Johns Hopkins University (JSU CCSE) [41], which maintains a Repository on Github
121   [42].

2.2 Procedures

123   The transmission dynamics of the COVID-19 outbreak is usually described by a compartmental model
124   (SEIR), where (S) susceptible – (E) exposed – (I) infected – (R) removed [43, 44]. In a closed population
125   of P_n individuals, the transitions between the compartments (cf. Fig. 2) are described through the following
126   set of differential equations:

Figure 2. SEIR Model Structure

$$\frac{d}{dt}s(t) = -\bar{\lambda}s(t)i(t) \tag{1}$$

$$\frac{d}{dt}e(t) = \bar{\lambda}s(t)i(t) - \kappa e(t) \tag{2}$$

$$\frac{d}{dt}i(t) = \kappa e(t) - \gamma i(t) \tag{3}$$

$$\frac{d}{dt}r(t) = \gamma i(t) \tag{4}$$

where $\bar{\lambda} = \lambda/P_n$, λ is the infection rate, γ is the remove rate, and κ is the incubation rate. From these parameters, it is possible to calculate the basic reproduction ratio, $R_0 = \lambda/\gamma$. Thus, R_0 is not solely dependent on the infection rate, but also on the frequency of removals due to death or recoveries.

Assuming that the incubation period of the disease is instantaneous and the duration of infectivity is the same as the length of the disease, we can consider both groups E and I as contagious and $E(t) := I(t)$. Also, according to the Akaike information criterion (AIC), the simpler SIR model performs much better than an SEIR model in representing the information contained in the confirmed-case data available for COVID-19 [45]. The basic SIR model is described by the set of Kermack–McKendrick equations [46]:

$$\frac{d}{dt}s(t) = -\frac{\lambda}{P_n}s(t)i(t) \tag{5}$$

$$\frac{d}{dt}i(t) = \frac{\lambda}{P_n}s(t)i(t) - \gamma i(t) \tag{6}$$

$$\frac{d}{dt}r(t) = \gamma i(t) \tag{7}$$

2.3 Discrete-time SIR system parametric estimation in real-time

Traditionally, mathematical epidemiology models have been approached with a continuous-time perspective, due in part to the fact that these are more tractable mathematically [47]. However, in order

138 to use machine learning techniques there is a need for a discrete-time equivalent realization to cope with
139 daily-sampled data [48]. Due to the slow dynamics of the pandemics, a first order continuous to discrete
140 Euler approximation can be applied to the Kermack–McKendrick equations.

141 For a general $f(t)$ function, a Backward discrete-time derivative approximation is:

$$\frac{d}{dt}f(t) := \frac{f(k+1) - f(k)}{T_s} = \Delta f(k+1)/T_s \tag{8}$$

142 where $T_s = 1$ is the sampling interval in days and $\Delta = 1 - q^{-1}$ is the discrete difference operator, defined
143 in q^{-1}, the backward shift operator domain. The discrete-time approximations of equations (5) to (7) are
144 given by the following difference equations, respectively:

$$s(k) = s(k-1) - \lambda(k-1)\frac{s(k-1)i(k-1)}{P_n} \tag{9}$$

$$i(k) = i(k-1) + \lambda(k-1)\frac{s(k-1)i(k-1)}{P_n} - \gamma(k-1)i(k-1) \tag{10}$$

$$r(k) = r(k-1) + \gamma(k-1)i(k-1) \tag{11}$$

145 The discrete-time SIR system described above considers time-varying parameters in order to continuously
146 adapt the model as new data becomes available. Using the time-series of infections and removals (due to
147 death or recovery), equations (9) to (11) can be used to estimate the model parameters.

148 Since Eq. (11) has an exclusive dependence with $\gamma(k)$, this poses a direct estimation problem which can
149 be stated as "for N registered samples, minimize the following quadratic cost function:"

$$J_r = \frac{1}{2}\sum_0^N e_r^2(k) = \frac{1}{2}\sum_{k=0}^N [r(k) - \hat{r}(k)]^2 \tag{12}$$

150 Eq. (12) is based on the estimation error of $r(k)$. By applying the recursive least-squares method to
151 minimize J_r, it is possible to optimally estimate $\hat{\gamma}(k)$ using the following equation:

$$\hat{r}(k) = r(k-1) + \hat{\gamma}(k-1)i(k-1) \tag{13}$$

152 We assume that the estimation error is Gaussian, $e_r(k) \sim (0, \sigma_{e_r}^2)$, with zero mean and variance $\sigma_{e_r}^2$, such
153 that $r(k) = \hat{r}(k) + e_r(k)$. The estimator gain, the parametric estimation and error covariance minimization
154 are solved recursively, as follows:

$$L_r(k) = \frac{p_r(k-1)i(k-1)}{1 + i(k-1)p_r(k-1)i(k-1)} \tag{14}$$

$$\hat{\gamma}(k) = \hat{\gamma}(k-1) + L_r(k)[r(k) - \hat{r}(k)] \tag{15}$$

　　　　　　　

$$p_r(k) = [1 - L_r(k)i(k-1)]p_r(k-1) \tag{16}$$

where the error covariance matrix $p_r(k)$ can be reset periodically to prioritize more recent data. Specifically in this work, $p_r(k)$ is a scalar, since only a single parameter is being estimated. Choices to initialize the covariance matrix may vary, depending on prior available covariance information or other positive definite matrix. The higher its magnitude the higher the estimator gain in the transitory dynamical stage of the estimation procedure.

In regard to Eq. (11) and the estimation of $\hat{\gamma}(k)$, since the infection numbers increase before any removal report is available during the first stages of the pandemic, during this period $\hat{\gamma}(k)$ tends to zero and eventually makes the time-varying estimated reproduction number tend to infinity:

$$\hat{R}_0(k) = \frac{\hat{\lambda}(k)}{\hat{\gamma}(k) \to 0} = \infty \tag{17}$$

As a consequence of (17), the estimation method proposed in this work cannot be applied when the number of recovered is not available.

Since Eq. (10) depends on both SIR parameters, due to its stronger dependence of $\hat{\gamma}(k)$ and Eq. (11), its estimated value is substituted into Eq. (10) and another recursive least-squares problem is constructed to estimate $\hat{\lambda}(k)$, using

$$\hat{i}(k) = i(k-1) + \hat{\lambda}(k-1)\frac{\hat{s}(k-1)i(k-1)}{P_n} - \hat{\gamma}(k)i(k-1) \tag{18}$$

The solution is akin to minimizing the estimation error $e_i(k) = i(k) - \hat{i}(k)$, using the following equations for estimator gain, parametric estimation update, and error covariance minimization, respectively:

$$L_i(k) = \frac{p_i(k-1)\hat{s}(k-1)i(k-1)/P_n}{1 + [\hat{s}(k-1)i(k-1)p_i(k-1)\hat{s}(k-1)i(k-1)]/P_n^2} \tag{19}$$

$$\hat{\lambda}(k) = \hat{\lambda}(k-1) + L_i(k)\left[i(k) - \hat{i}(k)\right] \tag{20}$$

$$p_i(k) = \left\{1 - L_i(k)\left[\frac{\hat{s}(k-1)i(k-1)}{P_n}\right]\right\}p_i(k-1) \tag{21}$$

Time-series data for Eq. (9) is not available and the evolution of the Susceptible compartment in time is in fact estimated based on the known initial condition (i.e., the population P_n) and on the estimated $\hat{\lambda}(k)$. Thus, it is always estimated and fed back to (18), such that the correct form to represent it, within this time-varying SIR model, is by rewriting Eq. (9) based on the estimated Susceptible:

$$\hat{s}(k) = \hat{s}(k-1) - \hat{\lambda}(k)\frac{\hat{s}(k-1)i(k-1)}{P_n} \tag{22}$$

174   The estimation of the time-varying reproduction number, based on the derived discrete-time SIR model,
175 is given by:

$$\hat{R}_0(k) = \frac{\hat{\lambda}(k)}{\hat{\gamma}(k)} \tag{23}$$

176   We also adopted two modifications to the nominal $\hat{R}_0(k)$ equation: a moving 4-day average to compensate
177 for the randomness of daily updates on incidence data, as seen in the German Daily Situation Report of the
178 Robert Koch Institute on COVID-19 [49], and the proportion of susceptible individuals in the population,
179 known as the effective reproduction number [14]:

$$\bar{R}_0(k) = \frac{\hat{s}(k)}{P_n} \left[\frac{\hat{R}_0(k) + \hat{R}_0(k-1) + \hat{R}_0(k-2) + \hat{R}_0(k-3)}{4} \right] \tag{24}$$

180   Figure 3 shows a block diagram of the proposed $\bar{R}_0(k)$ estimator. This diagram presents a clearer view of
181 the coupled multivariate dynamics and closed-loop characteristics of the proposed estimation approach.

Figure 3. Block diagram of the estimation system. Notice how its subsystems are interconnected using closed-loop strategies and data fusion.

With this formulation, it is possible to analyze the transmission ratio of the pandemics on a daily basis, as
with a sensor. Besides, estimations of the transmission ratio produce a dynamic representation from the
perspective of the time-series of $\bar{R}_0(t)$ (henceforth designated simply as $\hat{R}_0(t)$), which allows the modeling
of its dynamics and its randomness in order to assess stochastic properties correlated to the time-varying
reproduction number, which might reflect how health authorities have been handling the challenges posed
by the pandemics in each country considered in this work.

2.4 Modeling of R_0

Henceforth we assume that we are able to estimate the reproduction number on a daily basis and it is thus
possible to consider it as another output of the proposed pandemic model. Thus, relying on the time-series
of $\hat{R}_0(k)$ and knowing it is correlated with the number of infected and removed individuals, we deploy
machine learning techniques to identify a dynamic system that fits the data.

Differently from the real-time monitoring/sensing procedure adopted to estimate $\hat{R}_0(k)$, now we are
interested in obtaining a general model that can describe the dynamics of a system, $\hat{R}_0(q^{-1})$, for a certain
period of interest. In order to model such a system, we used the LMS, which is a non-recursive least-squares
estimation technique [31, 30], and propose a black-box linearized polynomial model:

$$\hat{R}_0(q^{-1}) = \frac{\hat{B}(q^{-1})I(q^{-1}) + \hat{C}(q^{-1})R(q^{-1}) + e_{R_0}(q^{-1})}{\hat{A}(q^{-1})} \tag{25}$$

$$\begin{aligned}
\hat{A}(q^{-1}) &= 1 + \hat{a}_1 q^{-1} + \hat{a}_2 q^{-2} \\
\hat{B}(q^{-1}) &= \hat{b}_0 q^{-1} + \hat{b}_1 q^{-2} \\
\hat{C}(q^{-1}) &= \hat{c}_0 q^{-1} + \hat{c}_1 q^{-2}
\end{aligned} \tag{26}$$

where $e_{R_0}(q^{-1})$ is the Gaussian process based on the estimation error $e_{R_0}(k)$ of estimated polynomials
shown in (26).

This second-order autoregressive with exogenous inputs (ARX) based model structure is assumed
considering the fundamental simplicity of the SIR model, in which the Infected and Removed systems
together form a second-order system.

The non-recursive least-squares estimator is a batch processing technique, used to optimally estimate
the set of parameters that minimizes a quadratic performance index as the one shown in (12), but using a
vector-matrix form of error, $\mathbf{e}_{R_0} = \hat{\mathbf{R}}_0 - \mathbf{\Phi}\hat{\theta}$. This vector-matrix system is defined as:

$$\mathbf{e}_{R_0} = \begin{bmatrix} e(0) & \cdots & e(N) \end{bmatrix}^T \tag{27}$$

$$\hat{\mathbf{R}}_0 = \begin{bmatrix} \hat{R}_0(0) & \cdots & \hat{R}_0(N) \end{bmatrix}^T \tag{28}$$

$$\hat{\theta}^{\mathbf{T}} = \begin{bmatrix} \hat{a}_1 & \hat{a}_2 & \hat{b}_0 & \hat{b}_1 & \hat{c}_0 & \hat{c}_1 \end{bmatrix} \tag{29}$$

$$\mathbf{\Phi}^{\mathbf{T}} = \begin{bmatrix} \phi^T(0) & \cdots & \phi^T(N) \end{bmatrix} \tag{30}$$

205   The above equations represent, respectively, the vector of errors, the vector of observed outputs, the
206 estimated parameters vector and the matrix of regressors. The latter is based on the vectors of regressors
207 formed up to N registered samples, with such vectors defined as:

$$\phi^T(k) = \begin{bmatrix} \hat{r}_0(k-1) & \hat{r}_0(k-2) & i(k-1) & i(k-2) & r(k-1) & r(k-2) \end{bmatrix} \tag{31}$$

208   The solution to obtain the estimated parameters is straightforward and given by

$$\hat{\theta} = \left(\mathbf{\Phi^T \Phi}\right)^{-1} \mathbf{\Phi^T \hat{R}_0} \tag{32}$$

By assuming an ARX linear model it is possible to evaluate the pandemics from the perspective of
linear stochastic systems theory, assessing how the reproduction number decays linearly in time in
different countries and how the random nature of events associated with the pandemics affects the model's
uncertainties, i.e., $e_{R_0}(k) \sim (0, \sigma^2_{e_{R_0}})$. This calculated uncertainty can give us some clues regarding the
effectiveness of pandemic control measures.

We propose that by analyzing the linear power of the $\hat{R}_0(q^{-1})$ estimation error, i.e. $\sigma^2_{e_{R_0}}$, we can use this
stochastic property to generate *stochastic ratio* curves based on the estimated linearized model of Eq. (25).
The higher the variability associated with the stochastic ratio the higher we expect the variance or linear
power of the error to be and, consequently, the more uncertain is the official COVID-19 data reported by
health authorities.

3   RESULTS

We used publicly available data to validate the algorithms and the estimated time-varying SIR model
parameters. This section is organized in the following way: we first present the estimated results based on
the number of infectious and removals available from different countries. Then, we present time-series of
daily estimates on $R_0(k)$ based on the RLS-estimated $\hat{\lambda}(k)$ and $\hat{\gamma}(k)$ and compare both processing time
and accuracy results to the classical LMS method to verify the efficiency of the proposed technique.

These results are followed by the presentation of linearized estimated outputs generated by the ARX-
based $\hat{R}(q^{-1})$ models (shown in (25)) using non-recursive or batch processing of 30-days of $R_0(k)$
estimates. The modeling residuals were assumed to be Gaussian (zero mean). Estimated error variances
were used to produce 200 discrete Gaussian sequences as surrogates to additive white noises, depicted
by $e_{R_0}(k)$ in (25). These 200 additive noise sequences were used to generate, for every analyzed country,
200 stochastic reproduction ratio trajectories, shown together with the linearized ratio and the real-time
estimated ratio. It must be remarked, however, that this value of 200 Monte Carlo simulations were selected
heuristically in order to give the necessary visual information in our figures, such that the readers could
verify by themselves how the modeling uncertainties of the pandemics could mislead our judgment of the
probable effective reproduction number of COVID-19 during the studied period. The number of Monte
Carlo simulations thus affects only a post-modeling phase and do not interfere with the estimation of the
reproduction number.

3.1   Discrete-time SIR model estimation results

The Achilles' heel of the RLS algorithm for parametric estimation may be the setup of initial conditions
and whether they are optimal or not. However, this is a major problem only if the estimated parameters are
to be used in a real-time adaptive control system where the closed-loop stability must be guaranteed [30].
This is not the case in the present work which is interested only in the modeling question itself. Thus, the
initial RLS parameters can be either arbitrarily set or set at zero since, theoretically, the RLS estimator is
dead-beat and converges in the minimum possible number of iterations [31, 35].

The discrete SIR model proposed in this work is described by three coupled differential equations, each based on a single recursive regression and thus forming a third order system. Then, theoretically, as a dead-beat estimator, the RLS would take three iterations to converge and estimate optimal parameters. However, we used plenty more iterations than the theoretical requirements, by commencing to process data from 22-March-2020 onwards but evaluating the results just after 23-April-2020, thus giving more than 30 days/iterations for the RLS to converge to an optimal set of parameters by 23-April-2020, when we started our analysis.

The estimators were implemented with arbitrary initial parameters of $\hat{\gamma}(0) = \hat{\lambda}(0) = 1$ and the initial guess for $\hat{R}_0(0) = 1$. The magnitudes of the estimation error covariance matrices were initialized as $p_r(0) = p_i(0) = 1$, considering that the initial error is large. Both $p_r(k)$ and $p_i(k)$ were reset to 1 every 7 days to prioritize more recent data [50]. The selection of the reset period considered not only the number of days required to wash-out outliers but also to allow weekly changes in the parameters' dynamical behavior, such as due to lockdowns or quarantine relaxation. we also adopted a moving 4-day average for $\hat{R}_0(k)$, shown in (24), to compensate for daily random effects [49]. We only used data from 22 March 2020 onwards, when all eight countries already had more than 100 infections reported.

Figures 4 and 5 show the dynamics of both Infected and Removed cases using the SIR model. It must be remarked that the Infected curve in the United States was downscaled by a factor of 4 in order to fit in the graph along with the other studied countries. It is evident from Fig. 4 that the RLS provide parametric estimates that fit the reported data. The same goodness of fit cannot be observed in Fig. 5, as the number of removals for the U.S., for example, poses some difficulties for the RLS estimator, as depicted by the estimated removals (dotted lines) showing a certain dispersion from the real data (continuous lines). However, the inclusion of the Removed data set, even with bad measurements, leads to a better estimate of $\hat{R}_0(k)$.

Figure 4. Estimation (dotted lines) of Infected individuals (continuous lines) using the RLS technique (The USA data was downscaled 4 times to fit in the graph).

Figure 5. Estimation (dotted lines) of Removed individuals (continuous lines) using the RLS technique.

Fig. 6 presents the $\hat{R}_0(k)$ of the eight different countries during a period of two weeks. Despite the large
variability of both Argentina and the United States, there are other countries that are already reducing the
number of new infections and where the frequency of removals has increased, such as in Germany, Italy,
and New Zealand (see Table 1).

The trajectory of the curves shown in Fig. 6 can also be associated with some extraordinary events that
occurred during the same period. For example, after 3 May 2020, when some U.S. states had relaxed social
distancing guidelines, it is possible to observe a corresponding phasic increase in the basic reproduction
ratio of the U.S., which was also reported in the Washington Post on 9 May 2020 [51].

One interesting trajectory in Fig. 6 regards Argentina. For most of the time, the estimated reproduction
ratio of Argentina was one of the highest and comparable only to the U.S., despite its low number of
infected individuals (cf. Fig. 4). This apparent contradiction is related to the low number of removals at the
beginning of the pandemic that tends to raise the $\hat{R}_0(k)$ (see Eq. 17). Recoveries in Argentina, based on
data of May 16 (cf. Table 1), are approximately 30.5% of its total infected, close to Belgium with 26.6%
and whose transmission ratio is below Argentina's, reinforcing the notion that the transmission ratio is not
a static parameter and is best approached by dynamical systems theory.

The results displayed in Figure 4 seem at odds with a recent report that claimed that Brazil's R_0 had
recently dropped to 1.4 [52]. However, on the same day a Brazilian newspaper quoted this report, the
country had a record number of new infections and deaths [53]. This reinforces the notion that machine
learning methods might give better and faster clues regarding the severity of the pandemic in terms
of $\hat{R}_0(k)$.

We compared the performance of the proposed RLS estimator with its most common counterpart, the
LMS algorithm. In this comparison, we considered the average processing time and the normalized relative

Figure 6. Daily monitoring of the COVID-19 pandemic: dynamics of the estimated reproduction number for two weeks in May, according to available official data and the SIR model.

Table 1. Recoveries (May 21, 2020).

Country	Recoveries
Argentina	30.5%
Belgium	26.6%
Brazil	40.5%
Germany	88.2%
Italy	59.0%
New Zealand	96.6%
Spain	70.3%
United States	23.5%

estimation error based on the mean-square-based cost function shown in Eq. (12). The average processing
time results for the RLS and LMS were, respectively, 14 ms and 20 ms. We used a computer with a 4th
Generation Intel Core i5-4200U CPU at 1.6 GHz, 4 GB of RAM, running Ubuntu 18.04 LTS and Matlab
R2018a (Mathworks, Inc.). The accuracy results expressed by normalized relative errors are summarized in
Fig. 7. Notice that the RLS outperformed the LMS in all cases, except for Argentina.

3.2 Assessing the pandemics through R_0 dynamics

Figures 8 to 11 show $\hat{R}_0(k)$ for a 30-day period for the eight investigated countries. These figures
show linearized R_0 together with the respective real-time estimated data that originated this second stage
estimation. The variance of the estimation error is then used to generate Gaussian sequences which
are superimposed on the linearized R_0 estimate, the stochastic ratio which synthetically reproduces the
stochasticity and uncertainties of R_0 estimates. Both Germany and Italy (Fig. 8) are through a period of

Figure 7. Comparison of normalized relative estimation error costs between the proposed RLS-based technique and the traditional LMS estimation algorithm. The average processing time of RLS and LMS was, respectively, 14ms and 20ms.

consistently decaying $R_0(t)$. Despite the hard way COVID-19 hit Italy before, its stochastic ratio currently
is among the lowest.

Brazil and the United States (Fig. 9) are the two most populous countries of our sample and the most
hard-hit by COVID-19 among them. Both countries also have been struggling with their uneven response
to the pandemics [54]. This outcome is captured by our R_0 sensors, with the Brazilian stochastic ratio
being in decrease, as Recovered data becomes more available (see Fig. 9).

In Fig. 10 we present linearized R_0 estimates for Spain and Belgium. Spain's estimates have suffered
the influence of annotation errors (by subtracting infected individuals in 24 Apr. 2020) which eventually
provoked oscillations in the real-time ratio estimate, making it zero-cross on April 27th. Such an error was
washed out by the linearized estimate and has certainly contributed to its increased degree of uncertainty
(see Table 2).

Belgium's degree of uncertainty was the worst among all the countries, at least during the period we
analyzed. One possible clue to understand why Belgium's stochastic ratio became so variable is to consider
the impossibility to linearize its dynamics and the associated increase in error. However, our estimation
procedure considers the error as Gaussian, and its mean value in fact approaches zero. Thus, the linearized
estimate based on 30-days of data has a high probability to be close to the values shown.

Belgium has also shown an increased R_0 during the analyzed period. The low number of recoveries
(see Table 1) influenced the frequency of removals and the basic reproduction number, which is the ratio
between the frequency of infections and the frequency of removals (see Eq. (23)). Even though a recent
report [55] proposed that Belgium's R_0 at the beginning of May 2020 was 0.8, according to our adaptive

Figure 8. 30-days of linearized R0 dynamics for Germany and Italy. Both countries have low variability in the stochastic ratio curve.

Figure 9. Brazil and the US display large variability in the stochastic ratio curve, which reflects data uncertainty and difficulties to control COVID-19's spread.

319 SIR-based R_0 estimations, the current situation in Belgium is uncertain with a probable R_0 close to 3.0,
320 which is the mean value of 30-days linearized dynamics (see Table 3).

Figure 10. 30-days of linearized R0 dynamics for Spain and Belgium. Both countries exhibit high variability of the stochastic ratio, which is suggestive of problems with either data compilation or difficulties to stabilize the pandemics.

Table 2.
Pandemic's uncertainty (up to 23 May 2020).

Country	Power	Uncertainty
Argentina	0.3099	30.9%
Belgium	0.2426	24.2%
Brazil	0.0546	5.4%
Germany	0.0220	2.2%
Italy	0.0113	1.1%
New Zealand	0.0349	3.4%
Spain	0.0521	5.2%
United States	0.1835	18.3%

Figure 11 shows the trends for Argentina and New Zealand. The R_0 estimates for the two countries, which have the lowest number of infections of the group, suggests that several recent R_0 reports are considering a ratio solely based on the number of infections and population, without estimating and taking into account the frequency of removals, which makes R_0 increase.

We compiled a table of the uncertainty of the estimated stochastic ratios by country (see Table 2), where the percentage of uncertainty is proportional to the linear power of the Gaussian estimation error. Table 3 shows the estimated R_0 estimates by country and demonstrate the usefulness of our approach to provide a real-time picture of the pandemic that can be used to support decision-making.

Figure 11. 30-days of linearized R0 dynamics for Argentina and New Zealand. New Zealand has been able to control the infection rates, which is reflected on the low variability stochastic ratio curve.

Table 3.
R0 estimates (on/up to 23 May 2020).

Country	Real-time	Linearized	30-days Mean
Argentina	4.6	5.3	3.5
Belgium	2.7	3.1	3.0
Brazil	2.9	2.8	3.0
Germany	1.6	1.4	1.5
Italy	1.2	1.1	1.5
New Zealand	1.1	1.2	1.1
Spain	2.6	2.1	1.4
United States	2.9	3.1	4.0

4 DISCUSSION

Almost two months after the WHO declared COVID-19 a global pandemic, health complications due
to SARS-CoV-2 have caused many deaths and upended the routines of billions of people around the
world. Research efforts for the development of a vaccine are being accelerated but it is still a distant target
[56]. At this moment, however, the most effective interventions are NPIs aimed at reducing SARS-CoV-2
transmission rates in order to increase the fraction of severe cases having access to scarce medical resources,
such as mechanical ventilation [57].

Mathematical modeling is a valuable instrument to gauge the epidemics' dynamics and evaluate the
effects of interventions aimed to control its spread. A crucial parameter is R_0, the basic reproduction
number, which is closely followed by health officials and the public alike. As the world hopefully transitions
to a gradual release from social distancing measures, many questions still remain about the SARS-CoV-2

virus and there is all but the inevitability of secondary waves of infection ahead. Thus, the continuing
use of mathematical models to track the disease will remain a necessity. However, the utility of models
depends on the quality of the data they are fed and there are many uncertainties regarding publicly available
data on COVID-19 cases. For instance, due to the lack of widespread testing and sub notifications on
the cause of deaths due to the disease, it is almost impossible to have a definitive picture of transitions
between the compartments used to model the disease. Thus, in this work, we provide a system that takes
into consideration the degree of uncertainty of the results presented by epidemiological SIR-type models.

The NPIs aimed to control the spread of the the COVID-19 pandemic are implemented in a way similar
to feedback control systems, with health authorities implementing social restriction measures in response
to real-time evaluation of the number of infected and removed cases. Therefore, as in control systems, we
are interfering with COVID-19's dynamics, affecting its behavior and parameters, and even discussing
the development of real-time monitoring techniques using automation and control systems technologies.
However, such closed-loop control systems still need humans in the loop and some data annotation mistakes
may occur such as: seasonal dynamic changes related to workers shifts, weekend reduced reports and even
possible data manipulation of the data log may compromise the stability of estimators.

We showed that RLS performed a better estimation job than LMS in the present work, both in speed and
accuracy. Real-time parametric estimation techniques, such as RLS, are widely used in the automation and
aerospace industries to support adaptive control systems and state estimators. In automation applications,
the RLS and LMS estimators are in fact divided into two different technologies, respectively, self-tuning
and auto-tuning [30], where the first refers to real-time online adaptive process and the second involves
the following steps: on-demand user request, acquiring the data set with sufficient samples, run offline
processing without time constraints, present the results to the user decide whether to deploy or not the
parametric update. Auto-tuning is less costlier to implement and thus more common, whereas the self-
tuning technology requires online real-time processing with time constraints and is more commonly applied
in the aerospace industry [58, 30].

LSA-based estimation is becoming more popular in industrial settings, such as: in single-input, single-
output systems (SISO) and in multiple-inputs, multiple-outputs systems (MIMO); in linear and non-linear
system identification, and in polynomial and state-space MIMO system realizations [59, 60]. Such an
increase in the use of RLS for real-time applications is justified by the advances of microprocessors to
cope with the time constraints of fast dynamical systems, where the sampling interval can go as short
as a few nanoseconds. Modern engineering applications with great impact in our society are associated
with discrete-time sampled systems and signals, thus requiring adequate and reliable digital estimation
algorithms such as LSA. At least from an engineering viewpoint, the COVID-19's pandemic is also a
discrete-time system, since the publicly available data for infections, deaths and recoveries are updated on
a daily basis, thus generating discrete-time sequences with the sampling period of 24h.

During COVID-19's pandemic the scientific community is struggling to study, design and deploy
engineering applications to better assist society with information that could somehow quantify the
pandemic's degree of severity based on $R_0(t)$ estimates [61]. Different from traditional modelling
approaches, we propose a discrete-time MIMO system realization instead of the continuous-time estimation
using ordinary differential equations; the MIMO-based parametric estimation of $R_0(t)$ is obtained through
data fusion in a closed-loop estimation arrangement (see Fig. 3).

The innovation of our modeling approach is the use of a discrete-time MIMO setup based on sensor
fusion, a well established strategy employed in the design of aerospace navigation systems under the

assumption that "one always gains by adding a new measurement in terms of navigation error, and this no
matter how bad the additional measurement is" [39, Remark 4.8]. Thus, by fusing data from the number
of susceptibles, infections, recoveries, deaths, and individual parameters of the three coupled dynamical
equations instead of a single one based solely on infections, the chance of having a better estimate of
$R_0(t)$ in the maximum-likelihood sense is higher. Of course, the drawback of our proposed approach is the
impossibility to apply it to nations lacking publicly reports on death and recovery numbers.

Our comparative approach shows that the strict measures adopted by some countries managed to stabilize
the epidemics, such as Germany, Italy, Spain, and New Zealand. In others, such as the U.S. and Brazil, the
delay in adopting such measures and lack of coordination proved decisive to keep R_0 values high and with
a high degree of uncertainty. The real-time estimation of model parameters such as R_0 provides important
insight into the underlying epidemic process and provides robustness in the face of imperfect data. This
strategy can be eventually implemented as an online tracking application providing information about the
dynamics of the pandemic to health officials and the public at large.

CONFLICT OF INTEREST STATEMENT

395   The authors declare that the research was conducted in the absence of any commercial or financial
396   relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

397   A.S. and A.P. conceived and wrote the manuscript.

FUNDING

398   Antonio Silveira acknowledges the financial support of the Brazilian Research Agency, the CNPq (*Conselho*
399   *Nacional de Desenvolvimento Científico e Tecnológico*), under grant 408559/2016-0.

DATA AVAILABILITY STATEMENT

400   The datasets analyzed for this study can be found in the COVID-19 data repository from The Center for
401   Systems Science and Engineering of the Johns Hopkins University (JSU CCSE) [41, 42].

REFERENCES

[1] WHO announces COVID-19 outbreak a pandemic. URL `http://www.euro.who.int/en/`
`health-topics/health-emergencies/coronavirus-covid-19/news/news/`
`2020/3/who-announces-covid-19-outbreak-a-pandemic`. Accessed on Mon, May
18, 2020.
[2] Na Zhu, Dingyu Zhang, Wenling Wang, Xingwang Li, Bo Yang, Jingdong Song, Xiang Zhao,
Baoying Huang, Weifeng Shi, Roujian Lu, Peihua Niu, Faxian Zhan, Xuejun Ma, Dayan Wang,
Wenbo Xu, Guizhen Wu, George F. Gao, and Wenjie Tan. A Novel Coronavirus from Patients with
Pneumonia in China 2019. *New England Journal of Medicine*, 382(8):727–733, feb 2020. doi:
10.1056/nejmoa2001017.

[3] Global situation report No. 118. URL `https://www.who.int/docs/default-source/coronavirus/situation-reports/20200517-covid-19-sitrep-118.pdf?sfvrsn=21c0dafe_6`. Accessed on Mon, May 18, 2020.

[4] Stephen A. Lauer, Kyra H. Grantz, Qifang Bi, Forrest K. Jones, Qulu Zheng, Hannah R. Meredith, Andrew S. Azman, Nicholas G. Reich, and Justin Lessler. The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Annals of Internal Medicine*, 172(9):577, mar 2020. doi: 10.7326/m20-0504.

[5] Melissa M. Arons, Kelly M. Hatfield, Sujan C. Reddy, Anne Kimball, Allison James, Jesica R. Jacobs, Joanne Taylor, Kevin Spicer, Ana C. Bardossy, Lisa P. Oakley, Sukarma Tanwar, Jonathan W. Dyal, Josh Harney, Zeshan Chisty, Jeneita M. Bell, Mark Methner, Prabasaj Paul, Christina M. Carlson, Heather P. McLaughlin, Natalie Thornburg, Suxiang Tong, Azaibi Tamin, Ying Tao, Anna Uehara, Jennifer Harcourt, Shauna Clark, Claire Brostrom-Smith, Libby C. Page, Meagan Kay, James Lewis, Patty Montgomery, Nimalie D. Stone, Thomas A. Clark, Margaret A. Honein, Jeffrey S. Duchin, and John A. Jernigan. Presymptomatic SARS-CoV-2 Infections and Transmission in a Skilled Nursing Facility. *New England Journal of Medicine*, apr 2020. doi: 10.1056/nejmoa2008457.

[6] Monica Gandhi, Deborah S. Yokoe, and Diane V. Havlir. Asymptomatic Transmission the Achilles' Heel of Current Strategies to Control Covid-19. *New England Journal of Medicine*, apr 2020. doi: 10.1056/nejme2009758.

[7] Wafaa M. El-Sadr and Jessica Justman. Africa in the Path of Covid-19. *New England Journal of Medicine*, apr 2020. doi: 10.1056/nejmp2008193.

[8] Tony Kirby. South America prepares for the impact of COVID-19. *The Lancet Respiratory Medicine*, apr 2020. doi: 10.1016/s2213-2600(20)30218-6.

[9] Roy M Anderson, Hans Heesterbeek, Don Klinkenberg, and T Déirdre Hollingsworth. How will country-based mitigation measures influence the course of the COVID-19 epidemic? *The Lancet*, 395 (10228):931–934, mar 2020. doi: 10.1016/s0140-6736(20)30567-5.

[10] Robert West, Susan Michie, G. James Rubin, and Richard Amlôt. Applying principles of behaviour change to reduce SARS-CoV-2 transmission. *Nature Human Behaviour*, may 2020. doi: 10.1038/s41562-020-0887-9.

[11] Julio Torales, Marcelo O'Higgins, João Mauricio Castaldelli-Maia, and Antonio Ventriglio. The outbreak of COVID-19 coronavirus and its impact on global mental health. *International Journal of Social Psychiatry*, page 002076402091521, mar 2020. doi: 10.1177/0020764020915212.

[12] Shunqing Xu and Yuanyuan Li. Beware of the second wave of COVID-19. *The Lancet*, 395(10233): 1321–1322, apr 2020. doi: 10.1016/s0140-6736(20)30845-x.

[13] Neil M. Ferguson, Derek A. T. Cummings, Christophe Fraser, James C. Cajka, Philip C. Cooley, and Donald S. Burke. Strategies for mitigating an influenza pandemic. *Nature*, 442(7101):448–452, apr 2006. doi: 10.1038/nature04795.

[14] Gerardo Chowell. Fitting dynamic models to epidemic outbreaks with quantified uncertainty: A primer for parameter uncertainty identifiability, and forecasts. *Infectious Disease Modelling*, 2(3): 379–398, aug 2017. doi: 10.1016/j.idm.2017.08.001.

[15] A. G. M'Kendrick. Applications of Mathematics to Medical Problems. *Proceedings of the Edinburgh Mathematical Society*, 44:98–130, feb 1925. doi: 10.1017/s0013091500034428.

[16] A contribution to the mathematical theory of epidemics. *Proceedings of the Royal Society of London. Series A Containing Papers of a Mathematical and Physical Character*, 115(772):700–721, aug 1927. doi: 10.1098/rspa.1927.0118.

[17] Ruiyun Li, Sen Pei, Bin Chen, Yimeng Song, Tao Zhang, Wan Yang, and Jeffrey Shaman. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). *Science*, 368(6490):489–493, mar 2020. doi: 10.1126/science.abb3221.

[18] Joshua S Weitz, Stephen J Beckett, Ashley R Coenen, David Demory, Marian Dominguez-Mirazo, Jonathan Dushoff, Chung-Yin Leung, Guanlin Li, Andreea Magalie, Sang Woo Park, Rogelio Rodriguez-Gonzalez, Shashwat Shivam, and Conan Zhao. Modeling shield immunity to reduce COVID-19 epidemic spread. *Nat Med*, 26:849 – 854, 2020. doi: 10.1038/s41591-020-0895-3.

[19] Giulio Viceconte and Nicola Petrosillo. COVID-19 R0: Magic number or conundrum? *Infectious Disease Reports*, 12(1), feb 2020. doi: 10.4081/idr.2020.8516.

[20] Benjamin Ridenhour, Jessica M. Kowalik, and David K. Shay. Unraveling R0: Considerations for Public Health Applications. *American Journal of Public Health*, 104(2):e32–e41, feb 2014. doi: 10.2105/ajph.2013.301704.

[21] S. Riley. Transmission Dynamics of the Etiological Agent of SARS in Hong Kong: Impact of Public Health Interventions. *Science*, 300(5627):1961–1966, jun 2003. doi: 10.1126/science.1086478.

[22] Paul L. Delamater, Erica J. Street, Timothy F. Leslie, Y. Tony Yang, and Kathryn H. Jacobsen. Complexity of the Basic Reproduction Number (R0). *Emerging Infectious Diseases*, 25(1):1–4, jan 2019. doi: 10.3201/eid2501.171901.

[23] Thomas Obadia, Romana Haneef, and Pierre-Yves Boëlle. The R0 package: a toolbox to estimate reproduction numbers for epidemic outbreaks. *BMC Medical Informatics and Decision Making*, 12 (1), dec 2012. doi: 10.1186/1472-6947-12-147.

[24] Luís M. A. Bettencourt and Ruy M. Ribeiro. Real Time Bayesian Estimation of the Epidemic Potential of Emerging Infectious Diseases. *PLoS ONE*, 3(5):e2185, may 2008. doi: 10.1371/journal.pone. 0002185.

[25] J. Wallinga. Different Epidemic Curves for Severe Acute Respiratory Syndrome Reveal Similar Impacts of Control Measures. *American Journal of Epidemiology*, 160(6):509–516, sep 2004. doi: 10.1093/aje/kwh255.

[26] Simon Cauchemez, Pierre-Yves Boëlle, Christl A. Donnelly, Neil M Ferguson, Guy Thomas, Gabriel M. Leung, Anthony J Hedley, Roy M. Anderson, and Alain-Jacques Valleron. Real-time Estimates in Early Detection of SARS. *Emerging Infectious Diseases*, 12(1):110–113, jan 2012. doi: 10.3201/eid1201.050593.

[27] S Cauchemez, PY Boëlle, G Thomas, and AJ Valleron. Estimating in real time the efficacy of measures to control emerging communicable diseases. *Am J Epidemiol*, 164:591–7, Sep 2006.

[28] Seyed M. Moghadas, Affan Shoukat, Meagan C. Fitzpatrick, Chad R. Wells, Pratha Sah, Abhishek Pandey, Jeffrey D. Sachs, Zheng Wang, Lauren A. Meyers, Burton H. Singer, and Alison P. Galvani. Projecting hospital utilization during the COVID-19 outbreaks in the United States. *Proceedings of the National Academy of Sciences*, 117(16):9122–9126, apr 2020. doi: 10.1073/pnas.2004064117.

[29] Yves Nievergelt. A tutorial history of least squares with applications to astronomy and geodesy. *Journal of Computational and Applied Mathematics*, 121(1-2):37–72, sep 2000. doi: 10.1016/ s0377-0427(00)00343-5.

[30] K. J. Astrom and B. Wittenmark. *Adaptive Control*. Dover Publications, Mineola, NY, USA, 2nd edition, 2008.

[31] Steven L. Brunton and J. Nathan Kutz. *Data-Driven Science and Engineering: Machine Learning, Dynamical Systems, and Control*. Cambridge University Press, 2019. ISBN 1108422098,9781108422093.

[32] Alexandra Smirnova and Gerardo Chowell. A primer on stable parameter estimation and forecasting in epidemiology by a problem-oriented regularized least squares algorithm. *Infectious Disease Modelling*, 2(2):268–275, may 2017. doi: 10.1016/j.idm.2017.05.004.

[33] Harvey Thomas Banks, Shuhua Hu, and W Clayton Thompson. *Modeling and inverse problems in the presence of uncertainty*. CRC Press, 2014.

[34] Julia Shen. A Recursive Bifurcation Model for Predicting the Peak of COVID-19 Virus Spread in United States and Germany. apr 2020. doi: 10.1101/2020.04.09.20059329.

[35] Frank L. Lewis, Lihua Xie, and Dan Popa. *Optimal and Robust Estimation*. CRC Press. Taylor & Francis Group, 2nd edition, 2008.

[36] Giseo Park and Seibum B. Choi. An Integrated Observer for Real-Time Estimation of Vehicle Center of Gravity Height. *IEEE Transactions on Intelligent Transportation Systems*, pages 1–12, 2020. doi: 10.1109/tits.2020.2988508.

[37] Vasundhara Puttagunta and Konstantinos Kalpakis. Adaptive Methods for Activity Monitoring of Streaming Data. In *ICMLA, vol. 2*, 2002.

[38] Rene Jaros, Radek Martinek, Radana Kahankova, and Jiri Koziorek. Novel hybrid extraction systems for fetal heart rate variability monitoring based on non-invasive fetal electrocardiogram. *IEEE Access*, 7:131758–131784, 2019.

[39] Pierre T. Kabamba and Anouck R. Girard. *Fundamentals of Aerospace Navigation and Guidance*. Cambridge Aerospace Series. Cambridge University Press, 2014. doi: 10.1017/CBO9781107741751.

[40] Rachelle N Binny, Shaun C Hendy, Alex James, Audrey Lustig, Michael J Plank, and Nicholas Steyn. Effect of Alert Level 4 on effective reproduction number: review of international COVID-19 cases. may 2020. doi: 10.1101/2020.04.30.20086934.

[41] Novel Coronavirus (COVID-19) Cases Data - Humanitarian Data Exchange. https://data.humdata.org/dataset/novel-coronavirus-2019-ncov-cases. URL `https://data.humdata.org/dataset/novel-coronavirus-2019-ncov-cases`. Accessed on Fri, May 08, 2020.

[42] CSSEGISandData/COVID-19. https://github.com/CSSEGISandData/COVID-19. URL `https://github.com/CSSEGISandData/COVID-19`. Accessed on Fri, May 08, 2020.

[43] Roy M Anderson and Robert M May. *Infectious diseases of humans: dynamics and control*. Oxford university press, 1992.

[44] Fred Brauer and Carlos Castillo-Chavez. Epidemic Models. In *Texts in Applied Mathematics*, pages 345–409. Springer New York, nov 2011. doi: 10.1007/978-1-4614-1686-9_9.

[45] Weston C. Roda, Marie B. Varughese, Donglin Han, and Michael Y. Li. Why is it difficult to accurately predict the COVID-19 epidemic? *Infectious Disease Modelling*, 5:271–281, 2020. doi: 10.1016/j.idm.2020.03.001.

[46] Roy M. Anderson and Robert M. May. Population biology of infectious diseases: Part I. *Nature*, 280 (5721):361–367, aug 1979. doi: 10.1038/280361a0.

[47] Fred Brauer, Zhilan Feng, and Carlos Castillo-Chavez. Discrete epidemic models. *Mathematical Biosciences & Engineering*, 7(1):1.

[48] S. Gómez, A. Arenas, J. Borge-Holthoefer, S. Meloni, and Y. Moreno. Discrete-time Markov chain approach to contact-based disease spreading in complex networks. *EPL (Europhysics Letters)*, 89(3): 38009, feb 2010. doi: 10.1209/0295-5075/89/38009.

[49] RKI - Coronavirus SARS-CoV-2 - Aktueller Lage-/Situationsbericht des RKI zu COVID-19. URL `https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Situationsberichte/Gesamt.html`. Accessed on Wed, May 06, 2020.

[50] Kevin Systrom. http://systrom.com/blog/author/ksys1983/. URL `http://systrom.com/blog/author/ksys1983/`. Accessed on Mon, May 04, 2020.

[51] Coronavirus flares as states and countries ease social distancing guidelines. URL `https://www.washingtonpost.com/national-security/coronavirus-flares-as-states-and-countries-ease-social-distancing-guideli 2020/05/09/cccb3c0c-9219-11ea-9e23-6914ee410a5f_story.html`. Accessed on Sun, May 17, 2020.

[52] Rubens Lichtenthaler Filho and Daniel Gomes Lichtenthaler. A dynamic model for Covid-19 in Brazil. may 2020. doi: 10.1101/2020.05.10.20097550.

[53] Brazil Registers 888 New Deaths from Coronavirus; Total Number of Deaths Exceeds 18 thousand. https://www1.folha.uol.com.br/internacional/en/scienceandhealth/2020/05/brazil-registers-888-new-deaths-from-coronavirus-total-number-of-deaths-exceeds-18-thousand.shtml. URL `https://www1.folha.uol.com.br/internacional/en/scienceandhealth/2020/05/brazil-registers-888-new-deaths-from-coronavirus-total-number-of-deaths-e shtml`. Accessed on Thu, May 21, 2020.

[54] Brazil, Once a Leader, Struggles to Contain Virus Amid Political Turmoil. https://www.nytimes.com/2020/05/16/world/americas/virus-brazil-deaths.html. URL `https://www.nytimes.com/2020/05/16/world/americas/virus-brazil-deaths.html`. Accessed on Wed, May 20, 2020.

[55] Basic reproduction number of novel coronavirus in Belgium falls to 0.6. https://www.vrt.be/vrtnws/en/2020/05/04/basic-reproduction-number-of-novel-coronavirus-in-belgium-falls/. URL `https://www.vrt.be/vrtnws/en/2020/05/04/basic-reproduction-number-of-novel-coronavirus-in-belgium-falls/`. Accessed on Fri, May 22, 2020.

[56] Barney S. Graham. Rapid COVID-19 vaccine development. *Science*, page eabb8923, may 2020. doi: 10.1126/science.abb8923.

[57] Ezekiel J. Emanuel, Govind Persad, Ross Upshur, Beatriz Thome, Michael Parker, Aaron Glickman, Cathy Zhang, Connor Boyle, Maxwell Smith, and James P. Phillips. Fair Allocation of Scarce Medical Resources in the Time of Covid-19. *New England Journal of Medicine*, mar 2020. doi: 10.1056/nejmsb2005114.

[58] NASA Administrator. Simplify, Simplify - Streamlined Adaptive System Works - Capabilities Verified, June 2013. URL `http://www.nasa.gov/centers/dryden/news/X-Press/mrac.html`.

[59] Antonio Silveira, Anderson Silva, Antonio Coelho, José Real, and Orlando Silva. Design and real-time implementation of a wireless autopilot using multivariable predictive generalized minimum variance control in the state-space. *Aerospace Science and Technology*, 105:106053, 2020. ISSN 1270-9638. doi: 10.1016/j.ast.2020.106053.

[60] Francesco Vicario. *OKID as a general approach to linear and bilinear system identification*. PhD thesis, Graduate School of Arts and Sciences, Columbia University, 2014.

[61] David Adam. A guide to r – the pandemic's misunderstood metric. *Nature*, 5883:346 – 348, 2020. doi: 10.1038/d41586-020-02009-w.