Prioritizing COVID-19 tests based in Participatory Surveillance and Spatial Scanning.

Leal-Neto, OB[1,2]*; Santos, FAS[3]; Lee, JY[4]; Albuquerque, JO[2,5]; Souza, WV[6]

[1]Department of Economics, University of Zurich, Zurich, Switzerland
[2]Epitrack, Recife, Brazil
[3]Agreste Academic Center, Federal University of Pernambuco, Caruaru, Brazil
[4]Colab, São Paulo, Brazil
[5]Immunopathology Laboratory Keizo Asami, Federal University of Pernambuco, Recife, Brazil
[6]Aggeu Magalhães Research Center, Oswaldo Cruz Foundation, Recife, Brazil

*Corresponding author at: Department of Economics, University of Zurich, Schöenberggasse 1. 8001, Zurich, Switzerland.
E-mail address: onicio.leal@econ.uzh.ch

Abstract:

Objectives

This study aimed to identify, describe and analyze priority areas for Covid-19 testing combining participatory surveillance and traditional surveillance.

Design

It was carried out a descriptive transversal study in the city of Caruaru, Pernambuco state, Brazil, within the period of 20/02/2020 to 05/05/2020. Data included all official reports for influenza-like illness notified by the municipality health department and the self-reports collected through the participatory surveillance platform Brasil Sem Corona.

Methods

We used linear regression and loess regression to verify a correlation between Participatory Surveillance (PS) and Traditional Surveillance (TS) . Also a spatial scanning approach was deployed in order to identify risk clusters for Covid-19.

Results

In Caruaru, the PS had 861 active users, presenting an average of 1.2 reports per user per week. The platform Brasil Sem Corona started on March 20[th] and since then, has been officially used by the Caruaru health authority to improve the quality of information from the traditional surveillance system. Regarding the respiratory syndrome cases from TS, 1,588 individuals were positive for this clinical outcome. The spatial scanning analysis detected 18 clusters and 6 of them presented statistical significance (p-value < 0.1). Clusters 3 and 4 presented an overlapping area that was chosen by the local authority to deploy the Covid-19 serology, where 50 individuals were tested. From there, 32% (n=16) presented reagent results for antibodies related to Covid-19.

Conclusion

Participatory surveillance is an effective epidemiological method to complement the traditional surveillance system in response to the COVID-19 pandemic by adding real-time spatial data to detect priority areas for Covid-19 testing.

Keywords: Participatory Surveillance, Epidemiology, Spatial scanning, Covid-19

Introduction

Participatory surveillance has shown promising results from its conception to its application in several public health events [1-6]. The use of a collaborative information pathway provides a rapid way for the data collection on symptomatic individuals in the territory, to complement traditional health surveillance systems [7, 8]. In Brazil, this methodology has been used at the national level since 2014 during mass gatherings events since they have great importance for monitoring public health emergencies [9, 10]. With the occurrence of the Covid-19 pandemic, and the limitation of the main non-pharmaceutical interventions for epidemic

control - in this case, testing and social isolation - added to the challenge of existing underreporting of cases and delay of notifications, there is a demand on alternative sources of up to date information to complement the current system for disease surveillance. Several studies [11-14] have demonstrated the benefits of participatory surveillance in coping with Covid-19, reinforcing the opportunity to modernize the way health surveillance has been carried out. Additionally, spatial scanning techniques have been used to understand syndromic scenarios, investigate outbreaks, and analyze epidemiological risk, constituting relevant tools for health management [15-18]. While there are limitations in the quality of traditional health systems, the data generated by participatory surveillance reveals an interesting application combining traditional techniques to clarify epidemiological risks that demand urgency in decision-making. Moreover, with the limitations of testing available, the identification of priority areas for intervention is an important activity in the early response to public health emergencies. This study aimed to describe and analyze priority areas for Covid-19 testing combining data from participatory surveillance and traditional surveillance for respiratory syndromes.

Methods

The study's method is a descriptive transversal, performed in the city of Caruaru, Pernambuco state, within the period of 02/20/2020 to 05/05/2020. Data included all official reports for influenza-like illness notified by municipality health department and the self-reports collected through the participatory surveillance platform Brasil Sem Corona[19].

To volunteer for the Brasil Sem Corona participatory surveillance platform, the individual should download an app called Colab. Through that, the volunteer has the option to agree with its terms of use that describe the purpose of this platform and ethical aspects, including the use of data for scientific studies that will help local public health authorities to improve the understanding of the epidemiological setting. After accepting the terms and conditions, participants can fill out a self-report questionnaire on symptoms and exposure in order to actively inform their health status. Daily reports are permitted, and a participant can report a maximum of two reports a day. Each time a participant completes and submits a report, it

provides the respective latitude and longitude, which are anonymized in the server within a radius of 2 kilometers. Personal identification is also anonymized, and participants can drop out their enrollment in the platform at any time, and the data is completely deleted from the servers, following national privacy laws.

Data from traditional disease surveillance system was accessed by the epidemiological bulletins, publicly available at local government communication channels.

To verify a correlation between Participatory Surveillance (PS) and Traditional Surveillance (TS), it was carried out a linear regression [20] extracting summary of fit and parameter estimates:

$$\gamma = \alpha + \beta\chi + \varepsilon_i$$

A local polynomial regression (LOESS) [21] was also performed to generalize a moving average to generate a scatterplot smoothing among the data points, where its function can be shown as:

$$\omega\left(\chi\right) = (1 - |d|^3)^3,$$

where d is the distance of data point from the point on the curve being fitted, scaled to lie in the range from 0 to 1.

On the spatial scanning method, it was performed a purely spatial analysis scanning for clusters with high rates using Bernoulli model assuming that in the studied area there are cases and non-cases represented by a dummy variable, acting as a spatial case (respiratory syndrome cases)/control (users that responded no symptoms) approach. The likelihood function for the Bernoulli [22] model is expressed as

$$L(Z, p, q) = p^{nZ}(1-p)^{\mu(Z)-nZ}q^{nG-nZ}(1-q)^{(\mu(G)-\mu(Z))-(nG-nZ)}$$

In order to detect the zone that is most likely to be a cluster, it was found the zone $\check{Z}$ that maximized the likelihood function, where $\check{Z}$ is the maximum likelihood estimator of the parameter Z. To achieve this, it needs to have two-stage actions. First, maximize the likelihood function conditioned on Z.

$$L(Z) \overset{def}{=} \underset{p > q}{\sup}(Z, p, q) = \left(\frac{n_Z}{\mu(Z)}\right)^{n_Z}\left(1 - \frac{n_Z}{\mu(Z)}\right)^{\mu(Z)-n_Z}$$

$$\times$$

$$\left(\frac{(n_G - n_Z)}{\mu(G) - \mu(Z)}\right)^{n_G - n_Z} \left(1 - \frac{n_G - n_{GZ}}{\mu(G) - \mu(Z)}\right)^{(\mu(G) - \mu(Z)) - (n_G - n_Z)}$$

$$\text{when } \frac{n_Z}{\mu(Z)} > \frac{(n_G - n_Z)}{(\mu(G) - \mu(Z))},$$

and otherwise

$$L(Z) = \left(\frac{n_G}{\mu(G)}\right)^{n_G} \left(\frac{\mu(G) - n_G}{\mu(G)}\right)^{\mu(G) - n_G}$$

Second, to find the solution $\hat{Z} = \{Z : L(Z) \geq L(Z') \forall Z' \in \mathcal{Z}\}$.

The most likely cluster is of interest in itself, but it is important also to have a statistical inference. Let $L_0 \stackrel{\text{def}}{=} \sup\limits_{p=q} L(Z, p, q) = \left(\frac{n_G}{\mu(G)}\right)^{n_G} \left(\frac{\mu(G) - \mu_G}{\mu(G)}\right)^{\mu(G) - n_G}$

The likelihood ratio, λ, can be written as

$$\lambda = \frac{\sup_{Z \in Z, p > q} L(Z, p, q)}{\sup_{p=q} L(Z, p, q)} = \frac{L(\hat{Z})}{L_0}$$

In addition, the SatScan runs a Monte Carlo simulation based on pseudo-random number generators. This allows to obtain the p-value for hypothesis testing, comparing the rank of the maximum likelihood from the real data set with the maximum likelihood from the random data sets.

In this case of spatial scanning clusters, statistical significance was defined as p-value < 0.1, bringing not only the clusters with a confidence interval of 95% and a maximum error of 5% but also considering a credible interval of 90% and a maximum error of 10%.

For the clusters that presented statistical significance [23], an overlaying in a Geographic Information System was done using QGIS 3.12 [24]. This map was used by the local authority to choose the area to deploy 50 serology tests for Covid-19. Other 50 tests were deployed in areas outside those clusters. For data wrangling, data tidying, and data visualization, an R language framework was used (Exploratory.io) [25]. For spatial scanning analysis, SatScan v9.6[26] was adopted.

The study had an exemption from the oversight body (Caruaru Health Department and Brasil Sem Corona scientific committee) since only public domain data available from secondary sources were used.

Results

In Caruaru, the PS had 861 active users, presenting an average of 1.2 reports per user per week. The platform Brasil Sem Corona started on March 20th and since then, has been officially used by the Caruaru health authority to improve the quality of information from the traditional surveillance system. Regarding the respiratory syndrome cases from TS, 1,588 individuals were positive for this clinical outcome.

When performed the signal (PS or TS) over time, it was found a linear correlation among these outputs and predictors where PS showed a correlation of 0.7242 (R squared 0.5245) and a coefficient of 1.1700 (p-value < 0.000). When verified the results for TS, correlation showed an output of 0.6198 (R squared 0.3841) and a coefficient of 1.12916 (p-value < 0.000). Regarding the LOESS, PS has shown an equivalent number of parameters of 4.374, with a residual standard error of 17.003 and a trace of smoother matrix of 4.7794. Finally, TS has shown as an equivalent number of parameters 4.628, residual standard error of 18.013, and a trace of smoother matrix of 5.072.

The spatial scanning analysis has detected 18 clusters and 6 of them have presented statistical significance (p-value < 0.1). Clusters 3 and 4 presented an overlapping and this area was chosen by the local authority to deploy the Covid-19 serology, where 50 individuals were tested. From there, 32% (*n*=16) presented reagent results for antibodies related to Covid-19, where the spatial scanning showed a hit rate between 84% and 94%. Another group of 50 individuals were tested outside identified clusters and from there, only 3% were positive for Covid-19 serology.

Figure 1. Time series distribution with linear regression for each source.

Figure 2. Time series distribution with LOESS regression for each source.

Cluster	Relative risk	Log likelihood ratio	P-value
1	1.54	44.504679	0.000
2	1.53	26.568310	0.000
3	1.34	13.654698	0.003
4	1.38	12.143518	0.013
5	1.45	11.350351	0.021
6	1.46	9.770844	0.094
7	1.51	8.619205	0.274
8	1.37	8.025044	0.404
9	1.31	7.421699	0.544
10	1.51	7.382095	0.616
11	1.51	6.970157	0.728
12	1.51	6.970157	0.728
13	1.51	6.558434	0.848
14	1.35	6.492181	0.854
15	1.51	5.735634	0.975
16	1.29	5.413135	0.982
17	1.51	5.324557	0.994
18	1.51	5.324557	0.994

Table 1. Summary of clusters including relative risk, log likelihood ratio and p-value.

Figure 3. Map showing the significant clusters, including an overlapping area between clusters 3 and 4 and testing targeted area.

Discussion

There was a strong correlation between the participation of individuals in the PS strategy over time, whereas there was a moderate association in the same period with the increase in notifications related to influenza syndrome from TS.

Regarding spatial scanning, the use of clusters to target priority regions for testing was of relevant importance in the adequacy and rationalization of the use of diagnostic resources, which are sometimes scarce in different contexts. The finding of 32% people reagent for the Covid-19 antibody in areas that the analysis showed a variation between 34% and 38% demonstrates an excellent result for the method to be replicated not only in this city, but also in other places that need tools to prioritization of tests. The remaining clusters that did not overlap but were statistically significant will be implemented in the coming weeks.

The challenge of coping with COVID-19 is even greater in settings where there is a lack of access for testing, low schooling of the population, and minor investment in health policies. In addition, there are a large number of cases that are not registered due to the small number of symptoms [27] or by complete absence [28] of any symptoms that may guide accurate health care, at the right time and in the right place.

Brazil is one of the countries that have the lowest number of tests among those that have the highest number of confirmed cases for COVID-19, and one of the epicenters is the state of Pernambuco, where the number of cases can be up to 10 times greater than those officially presented in epidemiological reports [29]. The study region became known as one of the main areas affected by the Zika Virus syndrome in 2015 and 2016 [30], where there is a population that is affected by several problems such as measles, leprosy, and other diseases overcome elsewhere of the world. This increases the need for rapid response during the SARS-CoV-2 pandemic, since the health system is already overloaded with basal demands.

Most of the documented cases of COVID-19, in studies carried out in China, are in the age group between 30 to 79 years old, a fact that can be observed as convergent [31]. In the current study, another relevant finding is that detected clusters were present in low-income areas that corroborate with other finds in Brazil [32].

In a study carried out by the Imperial College of London comparing the evolution of COVID-19 in 54 countries on several continents, it showed that Brazil

is the country most threatened to be one of the main epicenters of the disease in the world, mainly due to the great possibility of spreading the disease. Disease, with an R_0 greater than 2 and an increasing number of deaths [33].

Knowledge of risk areas is essential for appropriate non-pharmacological interventions [34], as well as for the definition of priorities regarding medical care, for example. The best time to start treatment is another big challenge because as the disease has a rapid and almost silent evolution, knowing where the probable cases are can save lives, expand the possibility of differential diagnosis and favor an effective treatment.

The use of alternative methods as participatory surveillance showed a relevant role by taking advantage on the insertion at community levels, generating real-time spatial information not only for self-reported symptoms individuals but also for participants that informed to have no symptoms. Although the presented approach demands other sorts of validation and diagnostic methods (PCR) and it also needs to be carried out in different cities to address potential biases, our application shows an alternative to rapid mass screening in areas to prioritize tests and supporting local public health authorities to implement public policies based in evidence.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

1. Guerrisi, C., Turbelin, C., Blanchon, T., Hanslik, T., Bonmarin, I., Levy-Bruhl, D., ... & Franco, A. O. Participatory syndromic surveillance of influenza in Europe. The Journal of infectious diseases, 214(suppl_4), (2016) S386-S392.

2. Meyers, D. J., Ozonoff, A., Baruwal, A., Pande, S., Harsha, A., Sharma, R., ... & Maru, D. S. Combining healthcare-based and participatory approaches to surveillance: trends in diarrheal and respiratory conditions collected by a mobile phone system by community health workers in rural Nepal. PloS one, 11(4), (2016).

3. Geneviève, L. D., Martani, A., Wangmo, T., Paolotti, D., Koppeschaar, C., Kjelsø, C., ... & Flahault, A. Participatory disease surveillance systems: ethical framework. Journal of medical Internet research, 21(5), (2019) e12273.

4. Wójcik, O. P., Brownstein, J. S., Chunara, R., & Johansson, M. A. Public health for the people: participatory infectious disease surveillance in the digital age. Emerging themes in epidemiology, 11(1),(2014) p. 7.

5. Smolinski, M. S., Crawley, A. W., Baltrusaitis, K., Chunara, R., Olsen, J. M., Wójcik, O., ... & Brownstein, J. S. Flu near you: crowdsourced symptom reporting spanning 2 influenza seasons. American journal of public health, 105(10)(2015) 2124-2130.

6. Santillana, M., Nguyen, A. T., Dredze, M., Paul, M. J., Nsoesie, E. O., & Brownstein, J. S. Combining search, social media, and traditional data sources to improve influenza surveillance. PLoS computational biology, 11(10) (2015).

7. Pini, A., Merk, H., Carnahan, A., Galanis, I., Van Straten, E., Danis, K., ... & Wallensten, A. High added value of a population-based participatory surveillance system for community acute gastrointestinal, respiratory and influenza-like illnesses in Sweden, 2013–2014 using the web. Epidemiology & Infection, 145(6) (2017) 1193-1202.

8. Leal-Neto, O. B., Dimech, G. S., Libel, M., Oliveira, W., & Ferreira, J. P. Digital disease detection and participatory surveillance: overview and perspectives for Brazil. Revista de saude publica, 50, 17 (2016).

9. Neto, O. L., Dimech, G. S., Libel, M., de Souza, W. V., Cesse, E., Smolinski, M., ... & Albuquerque, J. Saúde na Copa: the world's first application of participatory surveillance for a mass gathering at FIFA World Cup 2014, Brazil. JMIR public health and surveillance, 3(2) (2017) e26.

10. Neto, O. L., Cruz, O., Albuquerque, J., de Sousa, M. N., Smolinski, M., Cesse, E. Â. P., ... & de Souza, W. V. Participatory surveillance based on crowdsourcing during the rio 2016 olympic games using the guardians of health platform: Descriptive study. JMIR Public Health and Surveillance, 6(2) (2020) e16119.

11. Drew, D. A., Nguyen, L. H., Steves, C. J., Menni, C., Freydin, M., Varsavsky, T., ... & Spector, T. D. . Rapid implementation of mobile technology for real-time epidemiology of COVID-19. Science. (2020)

12. Menni, C., Valdes, A. M., Freidin, M. B., Sudre, C. H., Nguyen, L. H., Drew, D. A., ... & Visconti, A. Real-time tracking of self-reported symptoms to predict potential COVID-19. Nature Medicine,(2020) 1-4.

13. Garg, S., Bhatnagar, N., & Gangadharan, N. A Case for Participatory Disease Surveillance of the COVID-19 Pandemic in India. JMIR Public Health and Surveillance, 6(2) (2020) e18795.

14. Luo, H., Lie, Y., & Prinzen, F. W. Surveillance of COVID-19 in the General Population Using an Online Questionnaire: Report From 18,161 Respondents in China. JMIR Public Health and Surveillance, 6(2) (2020) e18576.

15. Greene, S. K., Peterson, E. R., Kapell, D., Fine, A. D., & Kulldorff, M. Daily reportable disease spatiotemporal cluster detection, New York City, New York, USA, 2014–2015. Emerging infectious diseases, 22(10) (2016) p.1808.

16. Van Den Wijngaard, C. C., Van Asten, L., Van Pelt, W., Doornbos, G., Nagelkerke, N. J., Donker, G. A., ... & Koopmans, M. P Syndromic surveillance for local outbreaks of lower-respiratory infections: would it work?. PLoS One, 5(4) (2010).

17. Chen, J., Roth, R. E., Naito, A. T., Lengerich, E. J., & MacEachren, A. M. Geovisual analytics to enhance spatial scan statistic interpretation: an analysis of US cervical cancer mortality. International journal of health geographics, 7(1) (2008) p.57.

18. de Souza Gomes, E. C., Leal-Neto, O. B., de Oliveira, F. J. M., Campos, J. V., Souza-Santos, R., & Barbosa, C. S. Risk analysis for occurrences of schistosomiasis in the coastal area of Porto de Galinhas, Pernambuco, Brazil. BMC infectious diseases, 14(1), (2014) p.101.

19. Brasil Sem Corona. Available at www.brasilsemcorona.com.br. (Accessed in 05/01/2020)

20. Freedman, D. A. Statistical models: theory and practice. Cambridge university press. (2009).

21. Garimella, R. V. A Simple Introduction to Moving Least Squares and Local Regression Estimation (No. LA-UR-17-24975). Los Alamos National Lab.(LANL), Los Alamos, NM (United States) (2017).

22. Kulldorff, M. A spatial scan statistic. Communications in Statistics-Theory and methods, 26(6) (1997) 1481-1496.

23. Kulldorff, M., & Nagarwalla, N. Spatial disease clusters: detection and inference. Statistics in medicine, 14(8) (1995) 799-810.

24. Team, Q. D. QGIS geographic information system. Open source geospatial foundation project. (2016)

25. Exploratory.io v5, computer program. (2020)

26. Coleman, M., Coleman, M., Mabuza, A. M., Kok, G., Coetzee, M., & Durrheim, D. N. Using the SaTScan method to detect local malaria clusters for guiding malaria control programmes. Malaria Journal, 8(1) (2009) p.68.

27. Bai, Y., Yao, L., Wei, T., Tian, F., Jin, D. Y., Chen, L., & Wang, M. Presumed asymptomatic carrier transmission of COVID-19. Jama, 323(14), (2020) 1406-1407.

28. Li, R., Pei, S., Chen, B., Song, Y., Zhang, T., Yang, W., & Shaman, J. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). Science, 368(6490), (2020) 489-493.

29. As Brazil's COVID-19 testing lags, available labs go unused. https://www.reuters.com/article/us-health-coronavirus-brazil-testing/as-brazils-covid-19-testing-lags-available-labs-go-unused-idUSKCN21X36V (accessed in 20th May 2020).

30. de Oliveira, W. K., de França, G. V. A., Carmo, E. H., Duncan, B. B., de Souza Kuchenbecker, R., & Schmidt, M. I. Infection-related microcephaly after the 2015 and 2016 Zika virus outbreaks in Brazil: a surveillance-based analysis. The Lancet, 390(10097), (2017) 861-870.

31. Surveillances, V. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19)—China, 2020. China CDC Weekly, 2(8) (2020) 113-122.

32. Lichand, G. Prokisch, G. Leal-Neto, O. A incidência econômica do coronavírus. Pre-print 2020. DOI: 10.17605/OSF.IO/HF27W

33. Cintra, P. H. P., & Nunes, F. F. Estimative of real number of infections by COVID-19 on Brazil and possible scenarios. medRxiv. (2020)

34. Garcia, L. P., & Duarte, E. Nonpharmaceutical interventions for tackling the COVID-19 epidemic in Brazil. Epidemiol. Serv. Saúde 29 (2) (2020).

Testing targeted area		

Relative risk

	1.34 – 1.34
	1.34 – 1.38
	1.38 – 1.47
	1.47 – 1.54
	Neighborhood boundaries
	Streets