Title: Risk for COVID-19 among Migrants from different areas of the World in Spain: A population-based cohort study in a country with universal health coverage.

Authors

[1]Carlos Guijarro, [2]Elia Pérez-Fernández, [,3,]Beatriz González-Piñeiro, [4]Victoria Meléndez, [5]Maria José Goyanes, [6] Mª Esther Renilla, [7]Maria Luisa Casas, [3]Isabel Sastre, BCE, [8]María Velasco, Alcorcon COVID Investigators .

[1]Internal Medicine Unit Hospital Universitario Fundación Alcorcón-Universidad Rey Juan Carlos, 28922 Alcorcón, Spain, [2] Research Unit, Hospital Universitario Fundación Alcorcón, 28922 Alcorcón, Spain, [,3,] Technology and Information Systems, Hospital Universitario Fundación Alcorcón, 28922 Alcorcón, Spain, [4]Department of Social, Senior and Public Health Services. City Council. 28922 Alcorcón, Spain, [5]Microbiology Unit , Hospital Universitario Fundación Alcorcón. Alcorcón. 20922, Spain, [6] Emergency Unit, , Hospital Universitario Fundación Alcorcón. Alcorcón. 20922, Spain, [7] Laboratory Unit. Hospital Universitario Fundación Alcorcón. Alcorcón. 20922, Spain, [8]Infectious Diseases Section, Internal Medicine Unit, Hospital Universitario Fundación Alcorcón. Alcorcón. 20922, Spain

Correspondence to. Carlos Guijarro, Internal Medicine Unit. Hospital Universitario Fundación Alcorcon - Universidad Rey Juan Carlos. Budapest 1, 28922 Alcorcón, Madrid (Spain)

Tel + 34 600 400 404,

Fax + 34 91 621 9905

e-mail: cguijarro@fhalcorcon.es ; carlos.guijarro@urjc.es

Abstract

Objective. To evaluate the COVID-19 incidence among migrants from different areas of the world as compared to Spaniards living in Alcorcón

Design. Population-based cohort analysis of the cumulative incidence of PCR-confirmed COVID-19 cases until April 25 (2020) among adult residents at Alcorcón (Spain) attended at the only public hospital serving this city. Crude incident rates for Spaniards and migrants from different areas of the world were estimated. Age and sex-adjusted relative risks for COVID19 were estimated by negative bomial regression.

Setting. University public Hospital at Alcorcón, Madrid, Spain

Participants. All adult residents living in Alcorcon classified by their country and region of the world of origin.

Main outcome. PCR confirmed COVID-19.

Results. PCR confirmed COVID-19 cumulative incidence was 6.81 cases per 1000 inhabitants among residents of Alcorcón. The crude incidence among migrants (n=20419) was higher than among Spaniards (n=131599): 8.81 and 6.51 and per 1000 inhabitants respectively (p<0.001).

By regions of the world, crude cumulative COVID-19 incidence rates were: European Union 2.38, Asia 2.01, , Northern Africa 3.59, East ern Europe 4.37, Sub-Saharan Africa 11.24, Caribbean 18.26 and Latin-America 20.77 8 per 1000 inhabitants. Migrant residents were markedly younger than Spaniards (median age 52 vs 73 years, p<0.001). By negative binomial regression, adjusted for age and sex, relative risks (RR) for COVID-19 were not significantly different from Spaniards for individuals from Europe, Asia or Northern Africa. In contrast, there was an increased risk for Sub-Saharan Africa (RR 3.66, 95% confidence interval (CI) 1.42-9.41, p=0.007), Caribbean (RR 6.35, 95% CI 3.83-10.55, p<0.001) and Latin-America (RR 6.92, 95% CI 4.49-10.67, p <0.001).

Conclusions. There was a marked increased risk for COVID-19 among migrants from Sub-Saharan Africa, Caribbean and Latin-America residing in Spain. The reasons underlying this increased risk and health and social implications deserve further attention.

What is known about the topic

Recent reports suggest an increased burden of COVID-19 among migrants or ethnic minorities in the United Kingdom and the USA, particularly regarding mortality. Reports have failed to dissociate clinical outcomes from differences in access to medical care or pre-existing medical conditions. There is no information regarding COVID risk for latinos in countries with universal health coverage

What this study add

Migrants from subsaharian Africa resident in Spain exhibit an increased risk for COVID-19. This risk is further increased for migrants from the Caribbean and Latin-America and cannot be attributed to unequal access to medical care. Studies in countries with universal health coverage may help to dissociate COVID burden in migrants and ethnic populations from access to health care.

Main text.

Introduction.

COVID-19 pandemic is imposing a tremendous challenge to the humankind with dramatic direct health consequences and dire secondary economic perspectives [1,2]. As the epidemic spreads out to different countries, with different social, economic and health systems, as well as different societal and political responses, the spectrum of COVID-19 is acquiring a variety of shapes [3]. Nonetheless, little is known regarding potential different effects in people from different countries and diverse ethnic background[4–7].

Recent reports have described a disproportionate burden of COVID-19 among migrants as well as ethnical minorities in Western countries [7–12]. However, the clinical outcomes (in particular mortality) associated to ethnic minorities are inextricably connected to different socioeconomic backgrounds, pre-existing medical conditions, and unequal access to medical services[10,11,13–15].

Studies in countries with a universal health care coverage may help to attenuate the contribution of unequal access to medical services from other factors associated to differential COVID-19 incidence and clinical outcomes in migrants.

We sought to describe the incidence of COVID-19 among migrants from different areas of the world as compared to Spaniards, both of them living in Alcorcón, a city in the suburbs of Madrid (Spain) with a substantial proportion of foreign residents. The high proportion of migrants into Spain from Latin America may provide a particular opportunity to evaluate COVID risk in this population.

Methods.

Case-population cohort study conducted at Alcorcón (Madrid), with a total population of about 17000 inhabitants. The population at risk was defined as all adults included in the official municipal live registry of population of the City Council of Alcorcón (last updated March 14, 2020).[16] Distribution of the population by nationality and country of origin stratified by age and sex was obtained from this registry.

Case was defined as a patient with a COVID-19 diagnosis at Hospital Universitario Fundación Alcorcón confirmed by Polymerase Chain Reaction (PCR) for SARS-CoV2 for adults residing in Alcorcón. Incident cases of COVID-19 were obtained from the Electronic Patient Record (Selene ©) that is used for all medical and administrative interactions with patients. COVID-19 clinical diagnosis was established by the attending clinicians at the Emergency Room according to European Center for Disease Control and Prevention (ECDCP)-World Health Organization Criteria (WHO) [17].

For molecular diagnosis of SARS-CoV-2 infection, nasopharyngeal swabs, sputum, or bronchopulmonary aspirates were processed by automatized extraction using the MagNaPureLc instrument (Roche Applied Science, Mannheim, Germany) and real time reverse-transcription PCR using the SARS-Cov-2 nucleic acid detection Viasure kit (CerTestBiotec S.L.), following the manufacturer's instructions. For this rRT-PCR, we used Bio-Rad CFX96™ Real-Time PCR Detection System. We amplified two different viral regions: ORF1ab gene (FAM channel), N gene (ROX channel), and the internal (HEX channel). Cycle threshold values ≤40 were considered positive. Positive and negative controls were included in each run for every assay.

An anonymized set of data was extracted on April 25 from the electronic patient record containing the following parameters: age, sex, nationality, country of birth, city of residence, date of COVID-19 diagnosis (clinical), date and results of SARS-CoV2 PCR, clinical evolution / outcomes (hospital admission, critical care admission, hospital discharge, length of hospital stay, and in-hospital death). For the present study patients whose residence was outside the city of Alcorcón were excluded.

Patients were classified according to their country of origin into one the following groups: a) Spain; b) European Union (including Switzerland and Norway); c). Eastern Europe (including Russia); d) Asia, e). Australia & New Zealand; f) Northern Africa; g) Sub Saharan Africa; h) Latin America (Mexico and continental Latin-American countries); i) Caribbean; j) United States of America & Canada.

Statistics. Results are described as means (± standard deviation), medians (interquartile range) or percentages as appropriate. Quantitative variables were compared by the student's t test,

ANOVA, or U Mann-Whitney 's test, as appropriate. Qualitative variables were compared by the Chi^2 test or the Fisher's exact test as appropriate.

Aggregated data by world zone (or country), sex and age group (<30, 30-39, 40-49, 50-59, 60-69, 70-79, 80-89 and >=90) were analyzed. Incidence rates and 95% exact Poisson confidence intervals were calculated by world zone of origin. Multivariate negative binomial regression model with robust variance was used to estimate the incidence rate by world zone adjusted for sex and age. These models are appropriated to analyze over-dispersed count data [18,19]. Statistical significance was assumed for p values <0.05. All statistical analyses were conducted with Stata 14 (StataCorp LLC, Texas, USA).

The protocol received the approval of the local Institutional Review Board / CEIm (Medicines Research Ethical Committee).

Results

Alcorcon city registry (last updated on March 14) comprised 152,018 individuals 17 years of age or older. Among them, 20,419 (13.4%) residents had a nationality different from Spain. Their countries of origin (and associated areas of the world) are summarized in Table 1. Different areas of the world were chosen because of geographical, socioeconomic and ethnic background reasons provided there was a minimum of 500 individuals per area. In this regard, Switzerland and Norway were included within the European Union group. The highest proportion of migrants from EU were from Romania (61.4%), followed by Poland (10.1%). Eastern Europe was vastly dominated by Ukraine (90.4%). China was predominant in Asia (60.4%), followed by Pakistan (15.4%). Equatorial Guinea (39,1%) and Nigeria (21.5%) were predominant among sub-Saharan Africa, while Northern Africa was essentially represented by Morocco (96.5%). Major contributors for continental Latin America were Colombia (20.2%), Peru (18.0%), and Venezuela (17.3%). Predominant Caribbean countries were Dominican Republic (71.3%) and Cuba (26.8%). Finally, migrants from the the USA, Canada, Australia and New Zealand) added up less than 200 residents. Given its residual nature, this group was removed for further analysis.

Male sex was more frequent in Spaniards (58.6%) than migrants (52.5%), although the difference did not reach statistical significance (p=0.15). In contrast, Spaniards were

substantially older than migrants (median age 73 vs 52 years, p<0.001) (Supplemental Figure 1). Similarly, age distribution pyramids in the reference population were strikingly different for Spaniards and migrants (Supplemental Figure 2).

A total of 1691 residents in Alcorcon were diagnosed as having COVID-19 following ECDCP-WHO clinical criteria[17] from February 1 until April 25 at our hospital (Figure 1). The number of cases increased steadily until the end of March and declined thereafter (figure 1), roughly 2 weeks after the lockout Decree of the Spanish Government (March 14, 2020)[20]. In order to avoid potential misdiagnosis, we restricted our analysis to the 1036 cases with a positive CRP for SARS-CoV2 sample (hereafter COVID-A-PCR+ cases). Median age of COVID-A-PCR+ patients was 71 years (IQ range 54-79), and 57.2 % were male. The global accumulated COVID-A-PCR+ incidence was 6.81 per 1000 inhabitants. By origin, the global accumulated COVID-A-PCR+ incidence in Spaniards was 6.50 cases per 1000 inhabitants as compared to 8.82 per 1000 for non-Spaniards (p<0.001; Table 2, Figure 1B). The global cumulative incidence increased dramatically with age for both Spaniards and migrants (Supplemental Figure 3A).

We next evaluated cumulative incidence rates of COVID-A-PCR+ for natives from different regions of the world as compared to Spaniards (Table2, Figure 1B,). Unadjusted incident rates for individuals from Europe, Asia and North Africa and Spain were lower than average, while for Sub-Saharan Africa, Caribbean and Latin-America were higher.

In order to have a better understanding of risks for migrants as compared to the local Spanish population we conducted a multivariate negative binomial regression analysis [18,19] adjusting for age and sex, as they are proven determinants of COVID-19 risk. The adjusted relative risks for COVID-PCR+ among migrants from Asia, European Union and Eastern Europe did not significantly differ from Spain (table 3). In contrast, the relative risk for Latin-America migrants was about 7-fold higher than that for Spaniards (RR 6.92; 95% CI 4.49-10.67, p <0.001). In addition, the adjusted risk was also increased for Sub-Saharan Africa (RR 3.66 95% 1.42-9.41, p=0.007) as well as Caribbean (RR 6.35 95% CI 3.83-10.55, p<0.001) in a highly clinical and statistically significant manner.

As a sensitive analysis (Table 3), we repeated our estimation with wider criteria (clinical

diagnosis regardless of PCR results; n=1,691) and stricter criteria (patients requiring hospital admission because of moderate-to-severe COVID-19, n=877). In both cases results remained essentially unchanged. Adjusted relative risks for the spectrum of COVID-19 among migrants from Europe, Asia or Northern African (essentially Morocco) did not differ from Spaniards.

In addition, adjusted risk for Sub-Saharan Africans, Caribbeans and Latinos was about 3, 6 and 7 fold higher in all clinical scenarios (Table 3). As an exploratory evaluation, we performed a similar analysis for countries that had at least 800 inhabitants in Alcorcón (Table 1). Citizens from Romania, Ukraine, China, or Morocco did not exhibit increased rates for COVID-19 cumulative incidence (Supplemental Figure 4). In agreement with regional data, migrants from all Latin American countries showed a consistent increased risk for COVID-19.

Severe COVID-19, defined as death, critical care admission or hospital stay longer than 7 days occurred in 63% of the Spaniards as compared to 33% for migrants. Similarly, unadjusted mortality was higher for Spaniards (25% of admissions) than for migrants (6%). Most likely this mirrors the advanced age of patients with severe disease (median 76 years IQR 69-82 years) or death (median 79 years IQR 74-85), an age range virtually absent in migrants (Supplemental Figures 1, 2) . Our data do not support a more severe disease among migrants residing in Spain.

Discussion.

Our data highlights that the incidence of COVID-19 varies for citizens of national versus migrants with a population-based approach in a Western European country, with particularly striking results for migrants from Latin America. The major finding of our work is an apparent higher risk for COVID-19 for individuals from Sub-Saharan, Caribbean or Latin-American origin. All of them have equal access to the virtually universal health coverage available for Spaniards or migrants from other areas of the world. Migrants from these three areas of the world are younger than Spaniards, and as opposed to *Latinos* or African Americans in the USA, the do not exhibit higher prevalence of obesity, hypertension, diabetes or cardiovascular disorders[10,21–23].. A higher risk is consistently detected when it is estimated globally for their regions of origin or for particular countries. Furthermore, the risk is clearly enhanced when adjusted for age and gender.

In contrast to the previous findings, we found a lower unadjusted risk for COVID-19 risk for other European citizens as compared to Spaniards. This may be related to a younger age connected to the so called 'healthy migrant effect' [21–24]. Indeed, when adjusted for age and gender the relative risk for COVID-19 for migrants from Europe, Asia or Northern Africa was not different from Spaniards. These migrants are usually employed for low-income jobs and pertain to a low socioeconomic status that may not be very different from migrants from Latin America [25,26]. The fact that no increased risk is found for other European, Asian or Northern Africa migrants is found underscores the relevance of the increased risk for Latin America, Caribbean and Sub-Saharan migrants

One strength of our study is the population-based approach. We used the best available updated registry of citizens residing in the reference population. This is also likely the case for migrants, since a number of social services rely upon the actual place of residence, even in the case of citizens in an irregular administrative position for the immigration authorities [16] Our estimation of the incidence of COVID-19 is a reasonable approach since the population primarily attended by our hospital is precisely the city of Alcorcón and there is no other public hospital in the city[27].

Another critical issue relates to the actual access to medical care for migrants. Since July 2018 all individuals resident in Spain have access to free medical care in conditions virtually identical to Spaniards [28]. In addition, the hospital is close to the city centre and serves a single urban area, with an easy access either by foot or public transportation. It might be argued that some migrants may rely less on primary care physicians and thus show a disproportionate use of hospital services [22,29]. However, our sensitive analysis, restricted to more severe COVID-19 requiring hospital admission does not support this potential overuse of hospital resources.

Some limitations of our study must be addressed. First, we did not have local data regarding socioeconomic status, education or health conditions for migrants. However, on aggregate, migrants from Romania, Ukraine or Morocco residing in Spain belong to a relatively lower socioeconomic stratum all along Spain[26] and do not exhibit an increased risk for COVID-19 as opposed to migrants from Sub-Saharan Africa or Latin-America.

Once the above mentioned factors do not seem to adequately explain this increased incidence, we should pay attention to the ethnic background of the populations. Our results suggest a gradual increased risk for Sub-Saharan and Latin-American with an intermediate value for Caribbeans consistent with their mixed ethnic background. There are limited data regarding different susceptibility to respiratory viruses from different ethnic backgrounds. Some studies have suggested a greater susceptibility of racial minorities to influenza[30,31]. However, many studies have failed to dissociate clinical outcomes from socioeconomic issues, particularly unequal access to health care [4,5,32]. Different susceptibilities to COVID-19 with a genetic background with unequal world distribution are just beginning to be explored[33,34].

In summary, we report a selective increased risk for COVID-19 among certain migrant populations in Spain: Sub-Saharan, Caribbean and Latin-America that is not related to unequal access to health care. These groups may deserve a particular attention, particularly when our country, as well as others, is beginning a de-escalation of social distancing measures. In addition, our data are compatible with different genetic susceptibility to COVID-19 among diverse ethnic backgrounds that warrants further studies. Those studies are of particular relevance since the focus of COVID-19 has shifted from China to Europe, then to the United States and now is unfortunately emerging in Africa and Latin America.

Acknowledgements. There was no funding for the present work
We are grateful to Dr F. Rodriguez-Artalejo for critically reviewing the manuscript.

References

Coronavirus Disease (COVID-19) Situation Reports. https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports (accessed June 21, 2020).

Peckham R. COVID-19 and the anti-lessons of history. *Lancet Lond Engl* 2020; **395**: 850–2.

Wynants L, Van Calster B, Collins GS, *et al.* Prediction models for diagnosis and prognosis of covid-19 infection: systematic review and critical appraisal. *BMJ* 2020; **369**: m1328.

Khunti K, Singh AK, Pareek M, Hanif W. Is ethnicity linked to incidence or outcomes of covid-19? *BMJ* 2020; **369**. DOI:10.1136/bmj.m1548.

de Lusignan S, Dorward J, Correa A, *et al.* Risk factors for SARS-CoV-2 among patients in the Oxford Royal College of General Practitioners Research and Surveillance Centre primary care network: a cross-sectional study. *Lancet Infect Dis* 2020; published online May 15. DOI:10.1016/S1473-3099(20)30371-6.

Orcutt M, Patel P, Burns R, *et al.* Global call to action for inclusion of migrants and refugees in the COVID-19 response. *Lancet Lond Engl* 2020; published online April 23. DOI:10.1016/S0140-6736(20)30971-5.

Office for National Statistics. Coronavirus (COVID-19) related deaths by ethnic group, England and Wales. https://backup.ons.gov.uk/wp-content/uploads/sites/3/2020/06/Coronavirus-COVID-19-related-deaths-by-ethnic-group-England-and-Wales-2-March-2020-to-15-May-2020.pdf.

Aldridge RW, Lewer D, Katikireddi SV, *et al.* Black, Asian and Minority Ethnic groups in England are at increased risk of death from COVID-19: indirect standardisation of NHS mortality data. *Wellcome Open Res* 2020; **5**: 88.

Kmietowicz Z, Ladher N, Rao M, Salway S, Abbasi K, Adebowale V. Ethnic minority staff and patients: a health service failure. *BMJ* 2019; **365**. DOI:10.1136/bmj.l2226.

Hooper MW, Nápoles AM, Pérez-Stable EJ. COVID-19 and Racial/Ethnic Disparities. *JAMA* 2020; published online May 11. DOI:10.1001/jama.2020.8598.

Price-Haywood EG, Burton J, Fort D, Seoane L. Hospitalization and Mortality among Black Patients and White Patients with Covid-19. *N Engl J Med* 2020; **0**: null.

CDC. Coronavirus Disease 2019 (COVID-19) in Racial and Ethnic Minority Groups. Cent. Dis. Control Prev. 2020; published online Feb 11. https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/racial-ethnic-minorities.html (accessed June 21, 2020).

Ji Y, Ma Z, Peppelenbosch MP, Pan Q. Potential association between COVID-19 mortality and health-care resource availability. *Lancet Glob Health* 2020; **8**: e480.

Dodds C, Fakoya I. Covid-19: ensuring equality of access to testing for ethnic minorities. *BMJ* 2020; **369**. DOI:10.1136/bmj.m2122.

Martinez DA, Hinson JS, Klein EY, *et al.* SARS-CoV-2 Positivity Rate for Latinos in the Baltimore-Washington, DC Region. *JAMA* 2020; published online June 18. DOI:10.1001/jama.2020.11374.

Ayuntamiento de Alcorcón. Portal Ciudadano. https://portalciudadano.ayto-alcorcon.es/portal/literal.do?opc_id=10000&pes_cod=-2&ent_id=1 (accessed May 10, 2020).

European Centre for Disease Prevention and Control. Case definition and European surveillance for COVID-19, as of 2 March 2020. Eur. Cent. Dis. Prev. Control.

https://www.ecdc.europa.eu/en/case-definition-and-european-surveillance-human-infection-novel-coronavirus-2019-ncov (accessed May 9, 2020).

Sroka CJ, Nagaraja HN. Odds ratios from logistic, geometric, Poisson, and negative binomial regression models. *BMC Med Res Methodol* 2018; **18**: 112.

Lloyd-Smith JO. Maximum likelihood estimation of the negative binomial dispersion parameter for highly overdispersed data, with applications to infectious diseases. *PloS One* 2007; **2**: e180.

Boletin Oficial del Estado. Real Decreto 463/2020, de 14 de marzo, por el que se declara el estado de alarma para la gestión de la situación de crisis sanitaria ocasionada por el COVID-19. https://www.boe.es/eli/es/rd/2020/03/14/463/con (accessed May 24, 2020).

Regidor E, Astasio P, Calle ME, Martínez D, Ortega P, Domínguez V. The association between birthplace in different regions of the world and cardiovascular mortality among residents of Spain. *Eur J Epidemiol* 2009; **24**: 503–12.

Ministerio de Sanidad, Consumo y Bienestar Social. Portal Estadístico del SNS - Encuesta Nacional de Salud de España 2017. https://www.mscbs.gob.es/estadEstudios/estadisticas/encuestaNacional/encuesta2017.htm (accessed May 10, 2020).

Cainzos-Achirica M, Vela E, Cleries M, *et al.* Cardiovascular risk factors and disease among non-European immigrants living in Catalonia. *Heart Br Card Soc* 2019; **105**: 1168–74.

Helgesson M, Johansson B, Nordquist T, Vingård E, Svartengren M. Healthy migrant effect in the Swedish context: a register-based, longitudinal cohort study. *BMJ Open* 2019; **9**: e026972.

World Bank Country and Lending Groups – World Bank Data Help Desk. https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups (accessed May 10, 2020).

Instituto Nacional de Estadística. (National Statistics Institute). Situación de los inmigrantes y de sus hijos en el mercado laboral. https://www.ine.es/dynt3/inebase/index.htm?path=/t22/e308/meto_05/modulo/base_2011/2014/ (accessed May 10, 2020).

Servicio Madrileño de Salud. Consejería de Sanidad de la Comunidad de Madrid. Memoria anual 2018. http://www.madrid.org/bvirtual/BVCM020283.pdf (accessed May 10, 2020).

Boletin Oficial del Estado. Real Decreto-ley 7/2018, de 27 de julio, sobre el acceso universal al Sistema Nacional de Salud. https://www.boe.es/eli/es/rdl/2018/07/27/7/con.

Velasco M, Castilla V, Guijarro C, Moreno L, Barba R, Losa JE. Differences in the use of health resources by Spanish and immigrant HIV-infected patients. *Enferm Infecc Microbiol Clin* 2012; **30**: 458–62.

Doxey M, Chrzaszcz L, Dominguez A, James RD. A Forgotten Danger: Burden of Influenza Mortality Among American Indians and Alaska Natives, 1999-2016. *J Public Health Manag Pract JPHMP* 2019; **25 Suppl 5, Tribal Epidemiology Centers: Advancing Public Health in Indian Country for Over 20 Years**: S7–10.

Centers for Disease Control and Prevention (CDC). Deaths related to 2009 pandemic influenza A (H1N1) among American Indian/Alaska Natives - 12 states, 2009. *MMWR Morb Mortal Wkly Rep* 2009; **58**: 1341–4.

CDC. Coronavirus Disease 2019 (COVID-19). Cent. Dis. Control Prev. 2020; published online Feb 11. https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/racial-ethnic-minorities.html (accessed May 10, 2020).

Nguyen A, David JK, Maden SK, *et al.* Human leukocyte antigen susceptibility map for SARS-CoV-2. *J Virol* 2020; published online April 17. DOI:10.1128/JVI.00510-20.

Devaux CA, Rolain J-M, Raoult D. ACE2 receptor polymorphism: Susceptibility to SARS-CoV-2, hypertension, multi-organ failure, and COVID-19 disease outcome. *J Microbiol Immunol Infect* 2020; published online May 6. DOI:10.1016/j.jmii.2020.04.015.

Table1. World distribution of migrants residing in Alcorcon by world region, country and sex.

WORLD REGION	Men	Women	Both	COUNTRIES	Men	Women	Both
1. SPAIN	62588	69011	131599	Spain	62588	69011	131599
2. EUROPEAN UNION[*]	3819	3749	7568	Romania	2281	2368	4649
				Poland	407	355	762
				Bulgaria	305	299	604
				Italy	330	260	590
				Portugal	182	177	359
				Others	314	290	604
3. EASTERN EUROPE & RUSSIA	921	1137	2058	Ukraine	850	1015	1865
				Others	71	122	193
4. ASIA	758	681	1439	China	418	451	869
				Others	340	230	570
5. NORTHERN AFRICA	1174	778	1952	Morocco	1129	755	1884
				Others	45	23	68
6. SUB-SAHARAN AFRICA	409	392	801	Equatorial Guinea	101	212	313
				Nigeria	103	69	172
				Others	205	111	316
7. LATIN AMERICA	2496	3330	5826	Colombia	537	641	1178
				Peru	484	570	1054
				Venezuela	412	595	1007
				Ecuador	357	304	661
				Honduras	148	380	528
				Others	558	840	1398
8. CARIBBEAN	319	338	657	Dominican Republic	226	247	473
				Cuba	90	88	178
				Others	3	3	6
9. MISCELLANEA[#]	55	63	118	United States	38	54	92
				Others	17	9	26
Total general	72539	79479	152018				

[*]-Excluding Spain; including Norway & Switzerland

[#]United States, Canada, Australia and New Zealand were merged because of their low number.

Table 2. COVID-19 PCR+ patients at Alcorcón. Accumulated incidence at Alcorcón by world region (A) and representative countries (B)

A

	All Regions	Spain	European Union	Eastern Europe	Asia	Northern Africa	Sub-Saharan Africa	Caribbean	Latin America
Age (median)	71.0	73.0	54.5	57.0	49.0	72.0	53.0	45.5	51.0
Interquartile range	54-79	62-79	45.75-64	45.5-61.5	42-49	34-76	49.5-66.5	36.25-57	39-55
Male (%)	57.5	58.6	54.9	52.3	47.0	61.9	57.2	48.0	48.4
Accumulated Incidence	6.81	6.50	2.38	4.37	2.08	3.59	11.24	18.26	20.77
Poisson Exact 95% confidence interval	6.41-7.24	6.08-6.96	1.41-3.76	2.00-8.30	0.43-6.09	1.44-7.39	5.14-21.33	9.44-31.91	17.23-24.82

B

			Europe		Asia	Northern Africa	Latin-America		
	All Countries	Spain	Romania	Ukraine	China	Morocco	Venezuela	Colombia	Peru
Age (median)	72.0	73.0	42.0	58.0	46.0	70.5	55.0	46.0	52.0
Interquartile range	59-79	62-80	38-55	57-62	44-48	44,7-75	53-67	38-52	38-58
Male (%)	55.0	58.6	57,1	0.0	50.0	66.7	62,5	47.4	47.0
Accumulated Incidence	6.54	6.08	1.51	3.75	2.30	3.18	7.94	16.13	36.05
Poisson Exact 95% confidence interval	6.13-6.98	6.08-6.96	0.61-3.10	1.51-7.73	0.28-8.31	1.17-6.93	3.43-15.65	9.71-25.19	25.51-49.49

Table 3. Negative Binomial Regression Models for COVID-19 Incidence by World Region (A) and Sensitive Analysis (B and C).

	Negative binomial regression model (adjusted for age and sex)			
	Zone of the World	Relative Risk	95% Cofidence.Interval	p value
A. PCR confirmed cases (n=1036)	Spain	1 (Reference)		
	Asia	0.64	0.27-1.52	0.32
	European Union	0.66	0.43-1.02	0.062
	Eastern Europe	1.23	0.49-3.06	0.646
	Northern Africa	1.11	0.37-3.31	0.84
	Sub-Saharan Africa	**3.66**	1.42-9.41	0.007
	Caribbean	**6.35**	3.83-10.55	<0.001
	Latin-America	**6.92**	4.49-10.67	<0.001
B. Clinical diagnosis (n=1691)	Spain	1 (Reference)		
	Asia	0.77	0.44-1.34	0.36
	European Union	0.56	0.4-0.78	0.001
	Eastern Europe	1.28	0.63-2.57	0.486
	Northern Africa	0.99	0.51-1.88	0.964
	Sub-Saharan Africa	**3.73**	2.11-6.57	<0.001
	Caribbean	**5.22**	3.37-8.09	<0.001
	Latin-America	**6.50**	5.11-8.27	<0.001
C. C. PCR confirmed cases & hospital admission (n=877)	Spain	1 (Reference)		
	Asia	0.57	0.2-1.66	0.311
	European Union	0.64	0.33-1.24	0.188
	Eastern Europe	0.95	0.35-2.54	0.921
	Northern Africa	1.46	0.5-4.24	0.48
	Sub-Saharan Africa	**3.32**	1.29-8.53	0.013
	Caribbean	**6.39**	3.79-10.76	<0.001
	Latin-America	**7.65**	4.99-11.72	<0.001

Figure Legends

Figure 1. Evolution of clinical diagnosis of COVID-19 at Alcorcón. Panel A: daily diagnosis (graph represents for everyday the average number of daily diagnosis of the preceding week). Panel B Cumulative incidence rate of new diagnosis per region of the world.

Figure 2. Relative risks for age intervals (reference age < 30 years), sex (Panel A), as well as areas of the world (reference Spain) were calculated by negative binomial regression analysis (Panel B).