

TITLE PAGE

Title: Differential risk for COVID-19 in the first wave of the disease (February-April 2020) among migrants from several areas of the world living in Alcorcón (Spain): A population-based cohort study.

Authors

^{1,3,7}Carlos Guijarro MD, PhD

⁴Elia Pérez-Fernández, M Stat

⁴Beatriz González-Piñeiro, BCE

⁹Victoria Meléndez, PhD

⁵Maria José Goyanes, MD

⁶ M^a Esther Renilla, MD

⁷Maria Luisa Casas, MD

⁴Isabel Sastre, BCE

^{1,2,3,8}María Velasco, MD, PhD, DTMH

& Alcorcon COVID Investigators .

¹Internal Medicine Unit (²Infectious Diseases Section), ³Research Unit, ⁴Technology and Information Systems, ⁴Microbiology Unit, ⁵Emergency Unit, ⁶Laboratory Unit. Hospital Universitario Fundación Alcorcón. Alcorcón. Madrid. Spain

⁷Department of Medical Specialties and Preventive Medicine. Universidad Rey Juan Carlos. Alcorcón. Madrid. Spain

⁸Department of Social, Senior and Public Health Services. City Council. Alcorcón

Correspondence to. Carlos Guijarro, Internal Medicine Unit. Hospital Universitario Fundación Alcorcon - Universidad Rey Juan Carlos. Budapest 1, 28922 Alcorcón, Madrid (Spain)

Tel + 34 600 400 404,

Fax + 34 91 621 9905

e-mail: cguijarro@fhalcorcon.es ; carlos.guijarro@urjc.es

***Alcorcón COVID Investigators: (Collaborators):**

Alejandro Algora, Juan Carlos Alonso-Punter, Maria Teresa Alonso-Salazar, Gregorio Bonilla Zafra, Mercedes Bueno-Campaña, Virgilio Castilla, Ana Isabel Diaz-Cuasante, Aurora Fabero, Rosa Maria Fariña, Isabel M^a Galán, Isabel González-Anglada, M^a Mercedes Izquierdo, Susana Lorenzo, Juan Emilio Losa, Margarita Mosquera, Carmen Noguera, Montserrat Pérez-Encinas, Gil Rodríguez-Caravaca, José Francisco Valverde

MANUSCRIPT

Title: Differential risk for COVID-19 in the first wave of the disease (February-April 2020) among migrants from several areas of the world living in Alcorcón (Spain): A population-based cohort study.

Abstract

Word count 176

Little is known regarding the relevance of racial / ethnic background on the risk of COVID-10 infection, particularly in Europe. We evaluated the risk for COVID-19 among migrants from different areas of the world within a context of universal free access to medical care. We conducted a population-based cohort analysis of the cumulative incidence of PCR-confirmed COVID-19 among adult residents at Alcorcón (Spain) in the first wave of the disease. The crude cumulative incidence among migrants (n=20419) was higher than among Spaniards (n=131599): 8.81 and 6.51 and per 1000 inhabitants respectively ($p < 0.001$) but differed by world region of origin. By negative binomial regression, adjusted for age and sex, relative risks (RR) for COVID-19 were not significantly different from Spaniards for individuals from Europe, Asia or Northern Africa. In contrast, there was a marked increased risk for Sub-Saharan Africa (RR 3.66, 95% confidence interval (CI) 1.42-9.41, $p = 0.007$), Caribbean (RR 6.35, 95% CI 3.83-10.55, $p < 0.001$) and Latin-America (RR 6.92, 95% CI 4.49-10.67, $p < 0.001$). Migrants from these areas of the world may deserve a closer attention both for clinical and epidemiological reasons.

Main text.

Word Count 2670

Introduction.

COVID-19 pandemic is imposing a tremendous challenge to the humankind with dramatic direct health consequences and dire secondary economic perspectives (1). Recent reports have highlighted an increased COVID-19 related mortality among migrants and ethnical minorities in Western countries (2–8). However, the clinical outcomes (in particular mortality) associated to ethnic minorities are inextricably connected to different socioeconomic backgrounds, pre-existing medical conditions, and unequal access to medical services (9–12). There have been a number of mass-media reports and calls to action but very few sounded scientific publications addressing this issue, most of them from the USA or United Kingdom, and are virtually absent from other European countries (13–15). In addition, the pattern of migration into different countries is quite variable: migrants from Latin America are quite common in Spain but are scarce in the United Kingdom.

We sought to describe the incidence of COVID-19 in the first wave of the disease among migrants from different areas of the world as compared to Spaniards, both of them living in Alcorcón, a city in the suburbs of Madrid (Spain) with a substantial proportion of foreign residents. The differential proportion of migrants into Spain from several areas of the world may provide a particular opportunity to evaluate COVID risk in these populations.

Methods.

Case-population cohort study conducted at Alcorcón (Madrid), with a total population of about 170,000 inhabitants. The population at risk was defined as all adults included in the official municipal live registry of population of the City Council of Alcorcón (last updated March 14, 2020). (16) Distribution of the population by nationality and country of origin stratified by age and sex was obtained from this registry.

Case was defined as a patient with a COVID-19 diagnosis at Hospital Universitario Fundación Alcorcón confirmed by Reverse Transcription (RT)-Polymerase Chain Reaction (PCR) for SARS-CoV2 for adults residing in Alcorcón. Incident cases of COVID-19 were obtained from the Electronic Patient Record (Selene ©) that is used for all medical and administrative interactions with patients. COVID-19 clinical diagnosis was established by the attending clinicians at the Emergency Room according to European Center for Disease Control and Prevention (ECDC)-World Health Organization Criteria (WHO) (17).

For molecular diagnosis of SARS-CoV-2 infection, nasopharyngeal swabs, sputum, or bronchopulmonary aspirates from symptomatic patients were processed by automatized extraction using the MagNaPureLc instrument (Roche Applied Science, Mannheim, Germany) and real time reverse-transcription PCR using the SARS-Cov-2 nucleic acid detection Viasure kit (CerTestBiotec S.L.), following the manufacturer's instructions. For this rRT-PCR, we used Bio-Rad CFX96™ Real-Time PCR Detection System. We amplified two different viral regions: ORF1ab gene (FAM channel), N gene (ROX channel), and the internal (HEX channel). Cycle threshold values ≤ 40 were considered positive. Positive and negative controls were included in each run for every assay.

An anonymized set of data was extracted from the electronic patient record (from February 1st to April 25th) containing the following parameters: age, sex, nationality, country of birth, city of residence, date of COVID-19 diagnosis (clinical), date and results of SARS-CoV2 PCR, clinical evolution / outcomes (hospital admission, critical care admission, hospital discharge, length of hospital stay, and in-hospital death). For the present study patients whose residence was outside the city of Alcorcón were excluded.

Patients were classified according to their country of origin into one the following groups: a) Spain; b) European Union (including Switzerland and Norway); c). Eastern Europe (including Russia); d) Asia, e). Australia & New Zealand; f) Northern Africa; g) Sub Saharan Africa; h) Latin America (Mexico and continental Latin-American countries); i) Caribbean; j) United States of America & Canada.

Statistics. Results are described as means (\pm standard deviation), medians (interquartile range) or percentages as appropriate. Quantitative variables were compared by the student's t test, ANOVA, or U Mann-Whitney's test, as appropriate. Qualitative variables were compared by the Chi² test or the Fisher's exact test as appropriate.

Aggregated data by world zone (or country), sex and age group (<30, 30-39, 40-49, 50-59, 60-69, 70-79, 80-89 and >=90 years) were analyzed. Incidence rates and 95% exact Poisson confidence intervals were calculated by world zone of origin. Multivariate negative binomial regression model with robust variance was used to estimate the incidence rate by world zone adjusted for sex and age. These models are appropriated to analyze over-dispersed count data (18). Statistical significance was assumed for p values <0.05. All statistical analyses were conducted with Stata 14 (StataCorp LLC, Texas, USA).

The protocol received the approval of the local Independent Review Board / CEIm (Medicines Research Ethical Committee).

Results

Alcorcon city registry (last updated on March 14) comprised 152,018 individuals 17 years of age or older. Among them, 20,419 (13.4%) residents had a nationality different from Spain. Their countries of origin (and associated areas of the world) are summarized in Table 1. Different areas of the world were chosen because of geographical, socioeconomic and ethnic background reasons provided there was a minimum of 500 individuals per area. In this regard, Switzerland and Norway were included within the European Union group. The highest proportion of migrants from EU were from Romania (61.4%), followed by Poland (10.1%). Eastern Europe was vastly dominated by Ukraine (90.4%). China was predominant in Asia (60.4%), followed by Pakistan (15.4%). Equatorial Guinea (39,1%) and Nigeria (21.5%) were predominant among sub-Saharan Africa, while Northern Africa was essentially represented by Morocco (96.5%). Major contributors for continental Latin America were Colombia (20.2%), Peru (18.0%), and Venezuela (17.3%). Predominant Caribbean countries were Dominican Republic (71.3%) and Cuba (26.8%). Finally, migrants from the the USA, Canada, Australia and New Zealand) added up less than 200 residents. Given its residual nature, this group was removed for further analysis.

Table1. World distribution of migrants residing in Alcorcón (Spain) on by world region, country and sex on April 14 2020.							
WORLD REGION	Men	Women	Both	COUNTRIES	Men	Women	Both
1. SPAIN	62588	69011	131599	Spain	62588	69011	131599
2. EUROPEAN UNION^a	3819	3749	7568	Romania	2281	2368	4649
				Poland	407	355	762
				Bulgaria	305	299	604
				Italy	330	260	590
				Portugal	182	177	359
				Others	314	290	604
3. EASTERN EUROPE & RUSSIA	921	1137	2058	Ukraine	850	1015	1865
				Others	71	122	193
4. ASIA	758	681	1439	China	418	451	869
				Others	340	230	570
5. NORTHERN AFRICA	1174	778	1952	Morocco	1129	755	1884
				Others	45	23	68
6. SUB-SAHARAN AFRICA	409	392	801	Equatorial Guinea	101	212	313
				Nigeria	103	69	172
				Others	205	111	316
7. LATIN AMERICA	2496	3330	5826	Colombia	537	641	1178
				Peru	484	570	1054
				Venezuela	412	595	1007
				Ecuador	357	304	661
				Honduras	148	380	528
				Others	558	840	1398
8. CARIBBEAN	319	338	657	Dominican Republic	226	247	473
				Cuba	90	88	178
				Others	3	3	6
9. MISCELLANEA^b	55	63	118	United States	38	54	92
				Others	17	9	26
Total general	72539	79479	152018				

^a-Excluding Spain; including Norway & Switzerland

^bUnited States, Canada, Australia and New Zealand were merged because of their low number.

Migrant residents were substantially younger than Spaniards (median age 40 years IQR 30-50 vs 49,5 years IQR 38-67 respectively, Supplemental Figure 1), with a slightly higher proportion of males (48,7 vs 47,6% p=0,002). Indeed, age distribution pyramids in the reference population were strikingly different for Spaniards and migrants (Supplemental Figure 2).

A total of 1691 residents in Alcorcon were diagnosed as having COVID-19 following ECDCP-WHO clinical criteria (17) from February 1 until April 25 at our hospital (Figure 1A). The number of cases increased steadily until the end of March and declined thereafter (Figure 1B), roughly 2 weeks after the lockout Decree of the Spanish Government (March 14, 2020)(19). In order to avoid potential misdiagnosis, we restricted our analysis to the 1036 cases with a positive PCR for SARS-CoV2 sample (hereafter COVID-19-PCR+ cases). Median age of COVID-19-PCR+ patients was 71 years (IQ range 54-79), and 57.2 % were male.

The global accumulated COVID-19-PCR+ incidence was 6.81 per 1000 inhabitants. By origin, the global accumulated COVID-19-PCR+ incidence in Spaniards was 6.50 cases per 1000 inhabitants as compared to 8.82 per 1000 for non-Spaniards (p<0.001; Table 2, Figure 1B). Spanish COVID-19-PCR+ cases were

substantially older than migrants (median age 73 years, IQR 62-80 vs 51,5, IQR 41,7-58, $p < 0,001$), with a non-significant higher proportion of males (58.6% vs 52.5%; $p = 0.15$) (Supplemental figure 3). The global cumulative incidence increased dramatically with age for both Spaniards and migrants (Supplemental Figure 3A).

Table 2. COVID-19 PCR+ patients at Alorcón (Spain). Accumulated incidence by world region of origin (February-April 2020)

	All	Spain	European Union	Eastern Europe	Asia	Northern Africa	Sub-Saharan Africa	Caribbean	Latin America
Cases (no)	1036	856	18	9	3	7	9	12	121
Age (median)	71.0	73.0	54.5	57.0	49.0	72.0	53.0	45.5	51.0
Interquartile range	54-79	62-79	45.75-64	45.5-61.5	42-49	34-76	49.5-66.5	36.25-57	39-55
Male (%)	57.5	58.6	54.9	52.3	47.0	61.9	57.2	48.0	48.4
Accumulated Incidence per 1000 inhabitants	6.81	6.50	2.38	4.37	2.08	3.59	11.24	18.26	20.77
Poisson Exact 95% confidence interval	6.41-7.24	6.08-6.96	1.41-3.76	2.00-8.30	0.43-6.09	1.44-7.39	5.14-21.33	9.44-31.91	17.23-24.82

Figure 1. Evolution of clinical diagnosis of COVID-19 at Alorcón Spain.

February- April 2020. Panel A: daily diagnosis (graph represents for every date the 7 day moving average = number of daily diagnosis of the preceding week) at Hospital Universitario Fundación Alcorcón. Panel B Cumulative incidence rate of new COVID-19 clinical diagnosis per region of the world at Hospital Universitario Fundación Alcorcón. (See image)

We next evaluated cumulative incidence rates of COVID-19-PCR+ for natives from different regions of the world as compared to Spaniards (Table2, Figure 1B). Unadjusted incident rates for individuals from Europe, Asia and North Africa were lower than average, while for Sub-Saharan Africa, Caribbean and Latin-America were higher. Cumulative incident rates for countries with at least 800 Alcorcón residents are described in Table 3.

Table 3. COVID-19 PCR+ patients at Alorcón. Accumulated incidence at Alorcón by representative countries (February-April 2020)

	All	Spain	Europe		Asia	Northern Africa	Latin-America		
			Romania	Ukraine	China	Morocco	Venezuela	Colombia	Peru
Cases (no)	943	856	7	7	2	6	8	19	38
Age (median)	72.0	73.0	42.0	58.0	46.0	70.5	55.0	46.0	52.0
Interquartile range	59-79	62-80	38-55	57-62	44-48	44,7-75	53-67	38-52	38-58
Male (%)	55.0	58.6	57,1	0.0	50.0	66.7	62,5	47.4	47.0
Accumulat ed Incidence	6.54	6.08	1.51	3.75	2.30	3.18	7.94	16.13	36.05
Poisson Exact 95% confidence interval	6.13-6.98	6.08-6.96	0.61-3.10	1.51-7.73	0.28-8.31	1.17-6.93	3.43-15.65	9.71-25.19	25.51-49.49

In order to have a better understanding of risks for migrants as compared to the local Spanish population we conducted a multivariate negative binomial regression analysis (18) adjusting for age and sex, as they are proven determinants of COVID-19 risk (Figure 2). The adjusted relative risks for COVID-PCR+ among migrants from Asia, European Union and Eastern Europe did not significantly differ from Spain (Table 4, Figure 2). In contrast, the relative risk for Latin-America migrants was about 7-fold higher than that for Spaniards (RR 6.92; 95% CI 4.49-10.67, $p < 0.001$). In addition, the adjusted risk was also increased for Sub-Saharan Africa (RR 3.66 95% 1.42-9.41, $p=0.007$) as well as Caribbean (RR 6.35 95% CI 3.83-10.55, $p < 0.001$) in a highly clinical and statistically significant manner.

Figure 2. Relative risks for COVID-19 for age intervals (reference age < 30 years), sex (Panel A), as well as areas of the world (reference Spain) calculated by negative binomial regression analysis (Panel B). Alorcón (Spain), February-April 2020.

Table 4. Negative Binomial Regression Models for COVID-19 Incidence by World Region of origin (A) and Sensitive Analysis (B and C). Alorcón (Spain) February-April 2020				
Negative binomial regression model (adjusted for age and sex)				
	Zone of the World	Relative Risk	95% Confidence Interval	p value
A. PCR confirmed cases (n=1036)	Spain	1 (Reference)		
	Asia	0.64	0.27-1.52	0.32
	European Union	0.66	0.43-1.02	0.062
	Eastern Europe	1.23	0.49-3.06	0.646
	Northern Africa	1.11	0.37-3.31	0.84
	Sub-Saharan Africa	3.66	1.42-9.41	0.007
	Caribbean	6.35	3.83-10.55	<0.001
	Latin-America	6.92	4.49-10.67	<0.001
B. Clinical diagnosis (n=1691)	Spain	1 (Reference)		
	Asia	0.77	0.44-1.34	0.36
	European Union	0.56	0.4-0.78	0.001
	Eastern Europe	1.28	0.63-2.57	0.486
	Northern Africa	0.99	0.51-1.88	0.964
	Sub-Saharan Africa	3.73	2.11-6.57	<0.001
	Caribbean	5.22	3.37-8.09	<0.001
	Latin-America	6.50	5.11-8.27	<0.001
C. PCR confirmed cases & hospital admission (n=877)	Spain	1 (Reference)		
	Asia	0.57	0.2-1.66	0.311
	European Union	0.64	0.33-1.24	0.188
	Eastern Europe	0.95	0.35-2.54	0.921
	Northern Africa	1.46	0.5-4.24	0.48
	Sub-Saharan Africa	3.32	1.29-8.53	0.013
	Caribbean	6.39	3.79-10.76	<0.001
	Latin-America	7.65	4.99-11.72	<0.001

Relative risks in **bold** indicate statistical significance ($p < 0,05$)

As a sensitive analysis (Table 4), we repeated our estimation with wider criteria (clinical diagnosis regardless of PCR results; n=1,691) and stricter criteria (patients requiring hospital admission because of moderate-to-severe COVID-19, n=877). In both cases results remained essentially unchanged. Adjusted relative risks for the spectrum of COVID-19 among migrants from Europe, Asia or Northern African (essentially Morocco) did not differ from Spaniards.

Adjusted risks for Sub-Saharan Africans, Caribbeans and Latinos were about 3, 6 and 7 fold higher in all clinical scenarios (Table 3). As an exploratory evaluation, we performed a similar analysis for countries that had at least 800 inhabitants in Alcorcón (Table 1). Citizens from Romania, Ukraine, China, or Morocco did not exhibit increased rates for COVID-19 cumulative incidence (Supplemental Figure 4). In agreement with regional data, migrants from all Latin American countries showed a consistent increased risk for COVID-19. Severe COVID-19, defined as death, critical care admission or hospital stay longer than 7 days occurred in 63% of the Spaniards as compared to 33% for migrants. Similarly, unadjusted mortality was higher for Spaniards (25% of admissions) than for migrants (6%). Most likely this mirrors the advanced age of patients with severe disease (median 76 years IQR 69-82 years) or death (median 79 years IQR 74-85), an age range virtually absent in migrants (Supplemental Figures 1, 2). Our data do not support a more severe disease among migrants residing in Spain.

Discussion.

On a population-based approach, our data highlight that the incidence of COVID-19 varies among migrants from different areas of the world as well as compared to the Spanish reference population.

A major finding of our work is an apparent higher risk for COVID-19 for individuals from Sub-Saharan, Caribbean or Latin-American origin, as compared to migrants from Europe, Asia or Northern Africa or Spaniards. Preliminary data from other area in Madrid are consistent with our findings (8). The higher risk is consistently detected when it is estimated globally for their regions of origin or for particular countries. Furthermore, the risk is clearly enhanced when adjusted for age and gender. In addition, we found a lower unadjusted relative risk for COVID-19 risk for other European citizens as compared to Spaniards. However, migrants represent a younger population when compared to Spaniards (20,21), as it is the case in our setting. Indeed, when adjusted for age and gender the relative risk for COVID-19 for migrants from Europe (both for European Union and Eastern Europe) did not differ from Spaniards. Similarly, the adjusted incidence rate of COVID-19 in migrants from Asia or Northern Africa was not different from the reference population.

The reasons for this apparent risk for COVID-19 among migrants from different areas of the world are not apparent. First, all migrants have equal access to the virtually universal health coverage available for Spaniards. Second, most migrants have low-wage jobs and pertain to a low socioeconomic status (22)(23). This is also the case for 'European' migrants, since this population is vastly represented by Romania and Ukraine in our city. An increased prevalence of hypertension or obesity may underlie a higher risk for migrants from Sub-Saharan Africa or Latin America, as it has been described in blacks or Hispanics in the USA (24). However, in our setting migrants from these areas do not seem to exhibit higher prevalence of obesity, hypertension, diabetes or cardiovascular disorders. This may be related to a younger age and the so called 'healthy migrant effect' (20,21,25).

One strength of our study is the population-based approach. We used the best available updated registry of citizens residing in the reference population. This is also likely the case for migrants, since a number of social services rely upon the actual place of residence, even in the case of citizens in an irregular administrative position for the immigration authorities (16). Our estimation of the incidence of COVID-19 is a reasonable approach since the population primarily attended by our hospital is precisely the city of Alcorcón and there is no other public hospital in the city. In case of an emergency such as COVID-19, it is likely that citizens rely even more into their local facilities. In any event, it is almost certain that a number, albeit small, of Alcorcón residents have been attended because of COVID-19 in hospitals different from ours and thus our data are likely an underestimation of the real incidence. We have excluded from our analysis 16% of COVID-19 cases attended in our hospital whose residence was not Alcorcón. Official data from the Community of Madrid indicates that our hospital receives a slightly higher number of patients from other areas than viceversa (26). Accordingly, our underestimation of COVID-19 incidence in Alcorcón may be somewhere between 10-15%. We do not believe that these data modify substantially the relative incidence for migrants, since the proportion of COVID-19 migrants excluded from our study because of residence outside Alcorcon was essentially identical (17%) as that for Spaniards (16%).

Another critical issue relates to the actual access to medical care for migrants. Since July 2018 all individuals resident in Spain have access to free medical care in conditions virtually identical to Spaniards (27). In addition, the hospital is close to the city center and serves a single urban area, with an easy access either by foot or public transportation. It might be argued that some migrants may rely less on primary care physicians and thus show a disproportionate use of hospital services (21,28). However, our sensitive analysis, restricted to more severe COVID-19 requiring hospital admission does not support this potential overuse of hospital resources.

Some limitations of our study must be addressed. First, we did not have local data regarding socioeconomic status, education or other health conditions for migrants. However, on aggregate, migrants from Romania, Ukraine or Morocco residing in Spain belong to a relatively lower socioeconomic stratum all along Spain (23) and do not exhibit an increased risk for COVID-19 as opposed to migrants from Sub-Saharan Africa or Latin-America, suggesting that socioeconomic status may not be a major reason for the differences in COVID-19 incidence.

Once the above mentioned factors do not seem to adequately explain this increased incidence, we should pay attention to the ethnic background of the populations. Interestingly, a recent report from a large database from the Veteran administration suggests that Black and Hispanic individuals in the USA are experiencing an excess burden of Covid-19 not entirely explained by underlying medical conditions (24), in agreement with our results. Interestingly, recent reports from the USA suggest that the increased risk for black minorities does not occur in Asian minorities, also in agreement with our data (29). These data point to a different ethnic background, in addition to other socio-economical reasons, as a potential risk factor for COVID-19.

There are limited data regarding different susceptibility to respiratory viruses from different ethnic backgrounds. Some studies have suggested a greater susceptibility of racial minorities to influenza (30). However, many studies have failed to dissociate clinical outcomes from socioeconomic issues, particularly unequal access to health care. Different susceptibilities to COVID-19 with a genetic background with unequal world distribution are just beginning to be explored (31).

Finally, our data do not suggest an increased mortality risk for migrants. Of note, black and Hispanic individuals did not exhibit increased mortality in the Veteran Administration database (24), in contrast with other reports in the USA (4). These data suggest that health care access, rather than ethnicity may be the major determinant for survival in COVID-19.

In summary, we report a selective increased risk for COVID-19 among certain migrant populations in Spain: Sub-Saharan, Caribbean and Latin-America that is not related to unequal access to health care. These groups may deserve a particular attention, particularly when our country, as well as others, is beginning a de-escalation of social distancing measures. In addition, our data are compatible with different genetic susceptibility to COVID-19 among diverse ethnic backgrounds that warrants further studies. Regardless of the reasons underlying the differential risk of COVID-19 in different migrant populations, our data can help to identify individuals or groups at particularly high risk who may merit special attention for both clinical and epidemiological reasons.

Acknowledgements. There was no funding for the present work

References

1. Coronavirus Disease (COVID-19) Situation Reports. [Internet]. [cited 2020 Aug 30]. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>
2. CDC. Coronavirus Disease 2019 (COVID-19) in Racial and Ethnic Minority Groups [Internet]. Centers for Disease Control and Prevention. 2020 [cited 2020 Aug 30]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/racial-ethnic-minorities.html>
3. Aldridge RW, Lewer D, Katikireddi SV, Mathur R, Pathak N, Burns R, et al. Black, Asian and Minority Ethnic groups in England are at increased risk of death from COVID-19: indirect standardisation of NHS mortality data. *Wellcome Open Res.* 2020;5:88.
4. Price-Haywood EG, Burton J, Fort D, Seoane L. Hospitalization and Mortality among Black Patients and White Patients with Covid-19. *N Engl J Med.* 2020 May 27;
5. Holtgrave DR, Barranco MA, Tesoriero JM, Blog DS, Rosenberg ES. Assessing racial and ethnic disparities using a COVID-19 outcomes continuum for New York State. *Ann Epidemiol.* 2020;48:9–14.
6. The COVID Racial Data Tracker [Internet]. The COVID Tracking Project. [cited 2020 Aug 30]. Available from: <https://covidtracking.com/race>
7. Williamson EJ, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature.* 2020;584(7821):430–6.
8. Jaqueti Aroca J, Molina Esteban LM, García-Arata I, García-Martínez J. COVID-19 en pacientes españoles e inmigrantes en un área sanitaria de Madrid. *Rev Esp Quimioter.* 2020;33(4):289–91.
9. Hooper MW, Nápoles AM, Pérez-Stable EJ. COVID-19 and Racial/Ethnic Disparities. *JAMA.* 2020 May 11;
10. Ji Y, Ma Z, Peppelenbosch MP, Pan Q. Potential association between COVID-19 mortality and health-care resource availability. *Lancet Glob Health.* 2020;8(4):e480.
11. Martinez DA, Hinson JS, Klein EY, Irvin NA, Saheed M, Page KR, et al. SARS-CoV-2 Positivity Rate for Latinos in the Baltimore–Washington, DC Region. *JAMA [Internet].* 2020 Jul 28 [cited 2020 Aug 30];324(4):392–5. Available from: <https://doi.org/10.1001/jama.2020.11374>
12. Chen JT, Krieger N. Revealing the unequal burden of COVID-19 by income, race/ethnicity, and household crowding: US county vs ZIP code analyses. *Harv Cent Popul Dev Stud Work Pap Ser.* 2020;19(1).
13. Chowkwanyun M, Reed AL. Racial Health Disparities and Covid-19 - Caution and Context. *N Engl J Med.* 2020 May 6;
14. Pareek M, Bangash MN, Pareek N, Pan D, Sze S, Minhas JS, et al. Ethnicity and COVID-19: an urgent public health research priority. *Lancet Lond Engl.* 2020 02;395(10234):1421–2.
15. Cowling BJ, Leung GM. Epidemiological research priorities for public health control of the ongoing global novel coronavirus (2019-nCoV) outbreak. *Eurosurveillance [Internet].* 2020 Feb 13 [cited 2020 Sep 2];25(6):2000110. Available from:

https://www.eurosurveillance.org/content/10.2807/1560-7917.ES.2020.25.6.2000110#html_fulltext

16. Ayuntamiento de Alcorcón. Portal Ciudadano [Internet]. [cited 2020 May 10]. Available from: https://portalciudadano.ayto-alcorcon.es/portal/literal.do?opc_id=10000&pes_cod=-2&ent_id=1
17. European Centre for Disease Prevention and Control. Case definition and European surveillance for COVID-19, as of 2 March 2020 [Internet]. European Centre for Disease Prevention and Control. [cited 2020 May 9]. Available from: <https://www.ecdc.europa.eu/en/case-definition-and-european-surveillance-human-infection-novel-coronavirus-2019-ncov>
18. Lloyd-Smith JO. Maximum likelihood estimation of the negative binomial dispersion parameter for highly overdispersed data, with applications to infectious diseases. *PloS One*. 2007 Feb 14;2(2):e180.
19. Boletín Oficial del Estado. Real Decreto 463/2020, de 14 de marzo, por el que se declara el estado de alarma para la gestión de la situación de crisis sanitaria ocasionada por el COVID-19. [Internet]. [cited 2020 May 24]. Available from: <https://www.boe.es/eli/es/rd/2020/03/14/463/con>
20. Regidor E, Astasio P, Calle ME, Martínez D, Ortega P, Domínguez V. The association between birthplace in different regions of the world and cardiovascular mortality among residents of Spain. *Eur J Epidemiol*. 2009;24(9):503–12.
21. Ministerio de Sanidad, Consumo y Bienestar Social. Portal Estadístico del SNS - Encuesta Nacional de Salud de España 2017 [Internet]. [cited 2020 May 10]. Available from: <https://www.mscbs.gob.es/estadEstudios/estadisticas/encuestaNacional/encuesta2017.htm>
22. World Bank Country and Lending Groups – World Bank Data Help Desk [Internet]. [cited 2020 Aug 30]. Available from: <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-grou>
23. Instituto Nacional de Estadística. (National Statistics Institute). Situación de los inmigrantes y de sus hijos en el mercado laboral. [Internet]. [cited 2020 Aug 30]. Available from: https://www.ine.es/dynt3/inebase/index.htm?path=/t22/e308/meto_05/modulo/base_2011/2014/
24. Rentsch CT, Kidwai-Khan F, Tate JP, Park LS, King JT, Skanderson M, et al. Covid-19 by Race and Ethnicity: A National Cohort Study of 6 Million United States Veterans. medRxiv [Internet]. 2020 May 12;2020.05.12.20099135. Available from: <http://medrxiv.org/content/early/2020/05/18/2020.05.12.20099135.abstract>
25. Helgesson M, Johansson B, Nordquist T, Vingård E, Svartengren M. Healthy migrant effect in the Swedish context: a register-based, longitudinal cohort study. *BMJ Open*. 2019 15;9(3):e026972.
26. Servicio Madrileño de Salud. Consejería de Sanidad de la Comunidad de Madrid. Memoria anual 2018 [Internet]. [cited 2020 Aug 30]. Available from: <http://www.madrid.org/bvirtual/BVCM020283.pdf>

27. Boletín Oficial del Estado. Real Decreto-ley 7/2018, de 27 de julio, sobre el acceso universal al Sistema Nacional de Salud. [Internet]. Available from: <https://www.boe.es/eli/es/rdl/2018/07/27/7/con>
28. Velasco M, Castilla V, Guijarro C, Moreno L, Barba R, Losa JE. Differences in the use of health resources by Spanish and immigrant HIV-infected patients. *Enferm Infecc Microbiol Clin*. 2012 Oct;30(8):458–62.
29. Moore JT. Disparities in Incidence of COVID-19 Among Underrepresented Racial/Ethnic Groups in Counties Identified as Hotspots During June 5–18, 2020 — 22 States, February–June 2020. *MMWR Morb Mortal Wkly Rep* [Internet]. 2020 [cited 2020 Aug 27];69. Available from: <https://www.cdc.gov/mmwr/volumes/69/wr/mm6933e1.htm>
30. Doxey M, Chrzaszcz L, Dominguez A, James RD. A Forgotten Danger: Burden of Influenza Mortality Among American Indians and Alaska Natives, 1999-2016. *J Public Health Manag Pract JPHMP*. 2019 Oct;25 Suppl 5, Tribal Epidemiology Centers: Advancing Public Health in Indian Country for Over 20 Years:S7–10.
31. Devaux CA, Rolain J-M, Raoult D. ACE2 receptor polymorphism: Susceptibility to SARS-CoV-2, hypertension, multi-organ failure, and COVID-19 disease outcome. *J Microbiol Immunol Infect*. 2020 Jun;53(3):425–35.



