COVID-19 Trend and Forecast in India:
A Joinpoint Regression Analysis

Aalok Ranjan Chaurasia

Abstract

This paper analyses the trend in daily reported confirmed cases of COVID-19 in India using joinpoint regression analysis. The analysis reveals that there has been little impact of the nation-wide lockdown and subsequent extension on the progress of the COVID-19 epidemic in the country and there is no empirical evidence to suggest that relaxations under the third and the fourth phase of the lockdown has resulted in spiking the reported confirmed cases of COVID-19. The analysis also suggests that if the current trend continues, in the immediate future, then the daily reported confirmed cases of COVID-19 in the country is likely to increase to 21 thousand by 15 June 2020 whereas the total number of confirmed cases of COVID-19 will increase to around 422 thousand.

Address for Correspondence

Aalok Ranjan Chaurasia
MLC Foundation
51, Lake City Farms (Ganesh Puri),
Kalkheda Road, Neelbad,
Bhopal, MP-462044, India
aalok@mlcfoundation.org.in
aranjan53@gmail.com

COVID-19 Trend and Forecast in India:
A Joinpoint Regression Analysis

Background

Total number of reported confirmed cases of COVID-19 in India crossed 100 thousand mark on 19 May 2010 according to the database maintained by the World Health Organization. The first confirmed COVID-19 case in India was reported on 30 January 2020 but no confirmed COVID-19 case was reported during 4 February 2020 through 1 March 2020. By 15 March 2020, more than 100 confirmed cases of COVID-19 were reported which increased to 500 by 24 March 2020 when the nation-wide lockdown was announced in the country. Since then, the number of daily reported confirmed cases crossed the 10000 mark by 14 April 2020 and the 50000 mark by 7 May 2020. An analysis of the trend in the daily reported confirmed cases of COVID-19 may provide an idea about how the COVID-19 epidemic has progressed in the country. The trend analysis also permit forecasting the likely trend in the reported confirmed cases of COVID-19 in the immediate future.

A trend analysis of daily reported confirmed cases of COVID-19 is also needed as it is widely claimed that the imposition of the nation-wide lockdown on 25 March 2020 has significantly decelerated the progress of COVID-19 epidemic in the country. It has also been claimed that loosening the restrictions under the nation-wide lockdown during its third and the fourth phase has primarily been responsible for the recently witnessed spike in the number of daily reported confirmed cases of COVID-19 in the country. It has even been argued that reimposing the harsh restrictions as part of the nation-wide lockdown is the only way of stopping or decelerating the progress of COVID-19 epidemic despite the fact that the social and economic cost of nation-wide lockdown has been found to be quite complex and exorbitant. It has repeatedly been stressed that because of serious social and economic implications of the nation-wide lockdown, it cannot be prolonged.

One way of empirically examining these and many other claims regarding the progress of the epidemic is whether the trend in the daily reported confirmed cases of COVID-19 has changed after the imposition of the nation-wide lockdown or after loosening the restrictions under the nation-wide lockdown. If the trend in the daily reported confirmed cases of COVID-19 has changed after the imposition of the nation-wide lockdown, then it can be concluded that the nation-wide lockdown indeed has an impact on the progress of the epidemic. Similarly, if it is found that the trend in the daily reported confirmed cases of COVID-19 has changed after loosening the restrictions under the nation-wide lockdown, then it can be concluded that loosening of the restrictions has been responsible for the spike in reported

confirmed COVID-19 cases in the country. However, if there is no change in the trend, then there is little empirical evidence to suggest that either the nation-wide lockdown or loosening of restrictions under the nation-wide lockdown has any telling impact on the progress of the COVID-19 epidemic.

A review of the daily reporting of the confirmed COVID-19 cases in India reveals that during the 28 days from 4 February 2020 through 1 March 2020, no confirmed case of COVID-19 case was reported in the country (Table 1). Moreover, during the period 2 March 2020 through 31 March 2020, daily reporting of confirmed COVID-19 cases has been highly inconsistent. For example, no confirmed case of COVID-19 was reported on 3 March, 20 March and 28 March 2020 whereas on 29 March 2020 alone, 255 confirmed cases of COVID-19 were reported. These inconsistencies in the reporting of daily confirmed cases of COVID-19 may bias any analysis of the trend in the daily reported confirmed cases of COVID-19. It is therefore necessary that these irregular fluctuations in the daily reporting of confirmed cases of COVID-19 are first ironed out before any analysis of the trend in the reported confirmed cases of COVID-19 is carried out.

One approach to minimise the impact of reporting inconsistencies in the analysis of the trend in daily reporting of confirmed COVID-19 cases is to use moving average instead of actual daily reported confirmed cases of COVID-19. The same approach has been followed in the present analysis. To minimise irregular fluctuations in the reporting of COVID-19 cases, five-day moving average has been used for the trend analysis. In other words, the reported confirmed cased of COVID-19 in a day used in the present analysis are actually the average of the reported confirmed cases of COVID-19 two days prior to the day in question; two days after the day in question and the reported confirmed cases of COVID-19 on the day itself. For example, the reported confirmed cases of COVID-19 on 3 March 2020 used in the present analysis are actually the simple average of reported confirmed cases of COVID-19 on 1 March through 5 March 2020, etc.

This paper analyses the trend in daily reported confirmed cases of COVID-19 in the country using joinpoint regression (Kim et al, 2000). Joinpoint regression is used to study the trend that varies over time. It first identifies the time point(s) at which the trend in the reported confirmed cases of COVID-19 has changed or the joinpoint(s). Once the joinpoint(s) are identified, then the average per cent change between two joinpoints is calculated to reflect how the trend in the reported confirmed cases of COVID-19 has varied over time. The goal of the joinpoint regression analysis is not to provide the statistical model that best fits the time series data. Rather, the purpose of the joinpoint regression analysis is to provide the model that best summarises the trend in the data (Marrot, 2010). The underlying assumption of joinpoint regression is that trend in the data is not the same throughout the period under reference.

Joinpoint Regression Model

The joinpoint regression model is essentially different from the conventional piecewise or segmented regression model in the sense that the identification of joinpoint(s) and their location(s) is estimated within the model and are not set arbitrarily as is the case with the piecewise or segmented regression analysis. The minimum and the maximum number of joinpoint(s) are, however, set in advance but the final number of joinpoint(s) or the time point(s) when the trend changes is determined statistically. The model first identifies the time point(s) when there is a change in the trend and calculates the average percentage change (APC) which reflects the rate of change between two joinpoint(s). When the number of joinpoint(s) is zero, the model reduces to simple linear regression model.

Let y_i denotes the infant mortality rate for the year t_i such that $t_1 < t_2 < ... < t_n$. Then the joinpoint regression model is defined as

$$\ln y_i = \alpha + \beta_1 t_1 + \delta_1 u_1 + \delta_2 u_2 + ... + \delta_j u_j + \varepsilon_i \tag{1}$$

where

$$u_j = \begin{cases} (t_j - k_j) & \text{if } t_j > k_j \\ 0 & \text{otherwise} \end{cases}$$

and $k_1 < k_2 .... < k_j$ are joinpoints. The details of joinpoint regression analysis are given elsewhere (Kim et al, 2000; Kim et al, 2004).

Joinpoint regression analysis has commonly been used when the temporal trend of a given quantity, like incidence, prevalence and mortality is of interest (Tyczynski and Berkel, 2005; Doucet, Rochette and Hamel, 2016; John and Hanke, 2015; Chaurasia, 2020). However, this method has generally been applied with the calendar year as the time scale (Akinyede and Soyemi, 2016; Mogos et al, 2016; Chatenoud et al, 2015; Missikpode at al, 2015). The joinpoint regression analysis can also be applied in epidemiological studies in which the starting date can be easily established such as the day when the disease is detected for the first time as is the case in the present analysis. Joinpoint regression analysis can, therefore, be applied in the public health framework also and can explore whether the number of reported cases of a specific disease has decreased after the introduction of an intervention to check the disease. In the present context, the application of the joinpoint regression analysis can answer the question whether the imposition of the nation-wide lockdown has resulted in the decrease in the reported confirmed cases of COVID-19 in the country or not. If the day of introducing the intervention turns out to be a joinpoint, then, a comparison of daily per cent change in the reported confirmed cases of COVID-19 before and after the joinpoint can tell whether the intervention has been able to bring down the reported confirmed cases of COVID-19 or not.

Actual calculations in the present analysis were carried out using the Joinpoint Regression Program developed by the Statistical Research and Application Branch of the National Cancer Institute of the United States of America (NIC, 2013). The software requires specification of minimum (0) and maximum number of joinpoints (>0) in advance. In the present analysis, the minimum number of joinpoints is specified as 0 while the maximum number of joinpoints have been specified as 5. The programme starts with the minimum number of joinpoints (0, which is actually a straight line and the model is simple linear regression model) and tests whether more joinpoints are statistically significant and must be added to the model (up to the pre-specified maximum number of joinpoints). The tests of significance is based on a Monte Carlo Permutation method (Kim et al, 2000).

The Bayesian Information Criterion (BIC) was used to identify the number of joinpoints in the model. There are other methods also for the purpose. These include the permutation test and the data driven BIC methods. Relative merits and demerits of different methods are discusses elsewhere (NIC, 2013). The permutation method is regarded as the best method but it is very highly computationally intensive. The BIC method, on the other hand, is less computationally intensive. This method selects that model for which the object function, which is either the sum of the model fit error or the penalty term is minimised.

Trend in Daily Reported Confirmed Cases of COVID-19

Results of the joinpoint regression analysis of the five-days moving average of the daily reported confirmed cases of COVID-19 in India for the period 1 March 2020 through 23 May 2020 are summarised in table 2 and figure 1. The five-days moving average is centred at the mid-point of the five-day interval. For example, the five-day moving average of the period 1 March through 5 March 2020 is centred on 3 March 2020. In other words the joinpoint regression analysis is carried out for the period 3 March 2020 through 21 May 2020, although, it covers the data on daily reported confirmed cases of COVID-19 1 March 2020 through 23 May 2020. The period prior to 1 March 2020 has not been included in the analysis as the daily reported confirmed cases of COVID-19 during the period 30 January 2020 through 1 March 2020 have mostly been found to be zero.

The application of the joinpoint regression analysis divides the duration 1 March 2020 through 23 May 2020 or a period of 84 days into five time segments and the trend in the daily reported confirmed cases of COVID-19 cases is found to be different in different time segments. During the first five days of the period under reference - 3 March 2020 (day 1) through 7 March 2020 (day 5), the trend in the daily reported confirmed cases of COVID-19 in the country has been found to be negative which means that daily reported

confirmed cases of COVID-19 in the country actually decreased, instead increased during this period, on average, at a rate of around 8 per cent per day. This decrease in the reported confirmed cases of COVID-19 may be attributed to reporting inconsistencies. The daily per cent change during this period has, however, been found to be statistically significant. On the other hand during the next 10 days - from 7 March 2020 (day 5) through 16 March 2020 (day 14) - the daily reported confirmed cases of COVID-19 increased, on average, at a rate of almost 15 per cent per day. The increase in the daily reported confirmed cases of COVID-19 accelerated further during the next eight days - from 16 March 2020 (day 14) through 23 March 2020 (day 21) - when the daily reported confirmed cases of COVID-19 in the country increased at a rate of more than 28 per cent per day. However, the increase in the daily reported confirmed cases of COVID-19 decelerated during the period 23 March 2020 (day 21) through 26 March 2020 (day 24) at a rate of almost 9 per cent per day, although the daily per cent change was not found to be statistically significant. The daily reported confirmed cases of COVID-19 increased again at a rate of 27 per cent per day, on average, during the next 10 days - from 26 March 2020 (day 24) through 4 April 2020 (day 33). After 4 April 2020, however, there has been no change in the trend in the daily reported number of confirmed cases of COVID-19 till 21 May 2020. During 4 April 2020 through 21 May 2020, the daily reported confirmed cases of COVID-19 increased, on average, at a rate of almost 5.2 per cent per day.

The joinpoint regression analysis suggests that the trend in the daily reported confirmed cases of COVID-19 changed statistically significantly at the 5[th] day (7 March 2020); 14[th] day (16 March 2020); 24[th] day (26 March 2020); and 33[rd] day (4 May 2020) of the 80 days period beginning 3 March 2020. The nation-wide lockdown in the country was imposed on 25 March 2020 initially for a period of 21 days which was then extended to 3 May 2020 on 15 April 2020. On 4 May 2020, the lockdown was again extended up to 17 May 2020 but with a relaxed set of restrictions which, on 18 May 2020, was further extended to 31 May 2020 with even more relaxed set of restrictions. The analysis suggests that the day of the fifth change in the trend in daily reported confirmed cases of COVID-19 only matched with the date of third extension of the nation-wide lockdown. The day of the change in the trend in daily reported confirmed cases of COVID-19 has not been found to be linked with the imposition of the nation-wide lockdown on 25 March 2020 as well as its extension on 15 April 2020. There has also been no change in the trend in the daily reported confirmed cases of COVID-19 after the fourth extension of the nation-wide lockdown on 18 May 2020 when restrictions under the nation-wide lockdown were significantly loosened.

After 4 April 2020, the daily reported confirmed cases of COVID-19 in the country have been found to have increased, on average, at almost 5.2 per cent per day. In other words, the joinpoint regression analysis of the trend in daily reported confirmed cases of COVID19 does not support the claim that the imposition of the nation-wide lockdown on 25 March 2020 had resulted in a deceleration in the increase in daily reported confirmed cases of COVID-19. At the same time, the trend analysis also does not support the claim that the relaxations in the restrictions under the nation-wide lockdown has resulted in the spiking of daily reported confirmed cases of COVID-19.

Forecasting Number of COVID-19 Cases

The average daily percent change in the reported confirmed cases of COVID-19 during the period 4 April 2020 through 21 May 2020 permit forecasting the daily reported confirmed cases of COVID-19 under the assumption that there is no change in the trend. This exercise suggests that by 15 June 2020, the daily reported confirmed cases of COVID-19 will increase to 21243 with a 95 per cent confidence interval of 18246 - 24725 (Table 3 and Figure 2). This increase in the reported confirmed cases of COVID-19 may change only when there is a significant change in the trend in the reported confirmed cases of COVID-19. A significant change in the trend is possible only when a new set of interventions are introduced to combat COVID-19 epidemic in the country. It is already being emphasised that the nation-wide lockdown imposed on 25 March 2020 is now getting increasingly irrelevant in checking the progress of the epidemic because of a host of factors, the most important

of which is that the national-wide lockdown could not prevent large scale movement, especially of migrant workers from urban areas to rural hinterland. It is, therefore, being stressed that population-wide testing for COVID-19 followed by active contact tracing and isolation of the positive cases and their contacts is necessary to stop the increase and even decrease the reported confirmed cases of COVID-19 in the country. The need for such a strategy stems from the fact that almost 40 per cent of the individuals tested positive for COVID-19 are found to be asymptomatic. Chaurasia (2020a) has suggested a cluster-based approach of population-wide testing for COVID-19 which significantly reduces the number of tests to be done.

The forecast of the daily reported confirmed cases of COVID-19, on the basis of the joinpoint regression analysis present here also suggests that the total number of confirmed COVID-19 cases in the country are likely to increase to almost 422 thousand by 15 June 2020 with a 95 per cent confidence interval of around 376 thousand to around 473 thousand. According to the latest information available from the database maintained by the World Health Organization, the total number of confirmed COVID-19 cases in the country has crossed the 138 thousand mark by 25 May 2020. This implies that during the next 20 days, there will be most probably around 284 thousand additional COVID-19 cases in the country. This scenario can be change through introducing appropriate interventions to halt or to even reverse the progress of the epidemic. The good sign, however, is that recovery rate of the disease in the country is quite encouraging while the risk of death from the disease is quite low by international standards.

Conclusions

The present analysis, based on the daily reported confirmed cases of COVID-19, suggests that there has virtually been little impact of the nation-wide lockdown and subsequent extension and relaxation in restrictions on the progress of the COVID-19 epidemic in India. There has also been little empirical evidence to suggest that relaxation in the restrictions under the third and the fourth phase of the nation-wide lockdown has resulted in spiking the reported confirmed cases of COVID-19 in the country. The analysis also suggests that if the trend in the reported confirmed cases of COVID-19 during the period 4 April through 21 May 2020 continues in the immediate future, then the daily reported confirmed cases of COVID-19 is likely to increase to around 21 thousand by 15 June 2020 whereas the total number of confirmed cases of COVID-19 will increase to around 422 thousand. This trend can be changed or reverted by introducing appropriate interventions that may help in containing the spread of the disease. In this context, population-wide testing for COVID-19 along with isolation of positive cases and contacts to the positive cases appears to be the need of the time.

References

Akinyede O, Soyemi K (2016) Joinpoint regression analysis of pertussis crude incidence rates, Illinois, 1990-2014. *American Journal of Infection Control* 44(12):1732–3.

Chatenoud L, Garavello W, Pagan E, Bertuccio P, Gallus S, La Vecchia C, Negri E, Bosetti C (2015) Laryngeal cancer mortality trends in European countries. *International Journal of Cancer* 842: 833–42.

Chaurasia AR (2020) Long-term trend in infant mortality in India: a joinpoint regression analysis for 1981-2018. Bhopal, MLC Foundation.

Chaurasia AR (2020a) Cluster approach to population-wide testing for COVID-19. Bhopal, MLC Foundation.

Doucet M, Rochette, Hamel D (2016) Prevalence and mortality trends in Chronic Obstructive Pulmonary Disease over 2001 to 2011: a public health point of view of the burden. *Canadian Respiratory Journal* 2016:1–10.

John U, Hanke M (2015) Liver cirrhosis mortality, alcohol consumption and tobacco consumption over a 62 year period in a high alcohol consumption country: a trend analysis. *BMC Research Notes* 8(1):822.

Kim HJ, Fay MP, Feuer EJ, Midthune DN (2000) Permutation tests for joinpoint regression with applications to cancer rates. *Statistics in Medicine* 19: 335-351.

Kim HJ, Fay MP, Yu B, Barrett MJ, Feuer EJ (2004) Comparability of segmented line regression models. *Biometrics* 60(4): 1005-1014.

Marrot LD (2010) Colorectal cancer network (CRCNet) user documentation for surveillance analytic software: Joinpoint. *Cancer Care Ontario*: 1-28.

Missikpode C, Peek-Asa C, Young T, Swanton A, Leinenkugel K, Torner J (2015) Trends in non-fatal agricultural injuries requiring trauma care. *Injury Epidemiology* 2(1): 30.

Mogos MF, Salemi JL, Spooner KK, McFarlin BL, Salihu HM (2016) Differences in mortality between pregnant and nonpregnant women after cardiopulmonary resuscitation. *Obstetrics and Gynecology* 128(4): 880–8.

National Cancer Institute (2013) *Joinpoint Regression Program*. Bethesda, MD: National Institutes of Health, United States Department of Health and Human Services.

Rea F, Pagan E, Compagnoni MM, Cantarutti A, Pigni P, Bagnardi V, Cprrap G (2017) Joinpoint regression analysis with time-on-study as time-scale. Application to three Italian population-based cohort studies. *Epidemiology, Biostatistics and Public Health* 14(3): e12616.

Tyczynski JE, Berkel HJ (2005) Mortality from lung cancer and tobacco smoking in Ohio (U.S.): will increasing smoking prevalence reverse current decreases in mortality? *Cancer Epidemiol Biomarkers Preview. United States* 14(5):1182–7.

Table 1
Reported confirmed cases of COVID-19 in India, 1 March 2020 - 23 May 2020.

Date	Reported confirmed cases	Cumulative confirmed cases	Five days moving average of reported confirmed cases
January 30, 2020	1	1	
January 31, 2020	0	1	
February 1, 2020	0	1	1
February 2, 2020	1	2	0
February 3, 2020	1	3	0
February 4, 2020	0	3	0
February 5, 2020	0	3	0
February 6, 2020	0	3	0
February 7, 2020	0	3	0
February 8, 2020	0	3	0
February 9, 2020	0	3	0
February 10, 2020	0	3	0
February 11, 2020	0	3	0
February 12, 2020	0	3	0
February 13, 2020	0	3	0
February 14, 2020	0	3	0
February 15, 2020	0	3	0
February 16, 2020	0	3	0
February 17, 2020	0	3	0
February 18, 2020	0	3	0
February 19, 2020	0	3	0
February 20, 2020	0	3	0
February 21, 2020	0	3	0
February 22, 2020	0	3	0
February 23, 2020	0	3	0
February 24, 2020	0	3	0
February 25, 2020	0	3	0
February 26, 2020	0	3	0
February 27, 2020	0	3	0
February 28, 2020	0	3	0
February 29, 2020	0	3	0
March 1, 2020	0	3	0
March 2, 2020	2	5	5
March 3, 2020	0	5	5
March 4, 2020	22	27	6

Date	Reported confirmed cases	Cumulative confirmed cases	Five days moving average of reported confirmed cases
March 5, 2020	2	29	6
March 6, 2020	2	31	7
March 7, 2020	3	34	3
March 8, 2020	5	39	4
March 9, 2020	5	44	6
March 10, 2020	6	50	8
March 11, 2020	10	60	8
March 12, 2020	14	74	8
March 13, 2020	7	81	11
March 14, 2020	3	84	11
March 15, 2020	23	107	13
March 16, 2020	7	114	14
March 17, 2020	23	137	22
March 18, 2020	14	151	18
March 19, 2020	44	195	34
March 20, 2020	0	195	41
March 21, 2020	88	283	57
March 22, 2020	58	341	65
March 23, 2020	93	434	82
March 24, 2020	85	519	73
March 25, 2020	87	606	77
March 26, 2020	43	649	58
March 27, 2020	75	724	92
March 28, 2020	0	724	93
March 29, 2020	255	979	120
March 30, 2020	92	1071	182
March 31, 2020	180	1251	248
April 1, 2020	385	1636	264
April 2, 2020	329	1965	366
April 3, 2020	336	2301	425
April 4, 2020	601	2902	486
April 5, 2020	472	3374	491
April 6, 2020	693	4067	595
April 7, 2020	354	4421	593
April 8, 2020	853	5274	608
April 9, 2020	591	5865	676
April 10, 2020	547	6412	787

Date	Reported confirmed cases	Cumulative confirmed cases	Five days moving average of reported confirmed cases
April 11, 2020	1035	7447	776
April 12, 2020	909	8356	900
April 13, 2020	796	9152	1005
April 14, 2020	1211	10363	987
April 15, 2020	1076	11439	1006
April 16, 2020	941	12380	1045
April 17, 2020	1007	13387	1070
April 18, 2020	991	14378	1165
April 19, 2020	1334	15712	1244
April 20, 2020	1553	17265	1319
April 21, 2020	1336	18601	1403
April 22, 2020	1383	19984	1473
April 23, 2020	1409	21393	1448
April 24, 2020	1684	23077	1579
April 25, 2020	1429	24506	1582
April 26, 2020	1990	26496	1608
April 27, 2020	1396	27892	1651
April 28, 2020	1543	29435	1709
April 29, 2020	1897	31332	1709
April 30, 2020	1718	33050	1889
May 1, 2020	1993	35043	2109
May 2, 2020	2293	37336	2240
May 3, 2020	2644	39980	2677
May 4, 2020	2553	42533	2870
May 5, 2020	3900	46433	3123
May 6, 2020	2958	49391	3272
May 7, 2020	3561	52952	3426
May 8, 2020	3390	56342	3301
May 9, 2020	3320	59662	3552
May 10, 2020	3277	62939	3561
May 11, 2020	4213	67152	3588
May 12, 2020	3604	70756	3668
May 13, 2020	3525	74281	3806
May 14, 2020	3722	78003	3758
May 15, 2020	3967	81970	4034
May 16, 2020	3970	85940	4378
May 17, 2020	4987	90927	4627

Date	Reported confirmed cases	Cumulative confirmed cases	Five days moving average of reported confirmed cases
May 18, 2020	5242	96169	4956
May 19, 2020	4970	101139	5284
May 20, 2020	5611	106750	5504
May 21, 2020	5609	112359	5768
May 22, 2020	6088	118447	
May 23, 2020	6564	125011	

Source: WHO COVID-19 database

Table 2
Results of the joinpoint regression analysis

Segment	Lower Endpoint	Upper Endpoint	Number of days	Average daily per cent change	95 per cent confidence interval		't'	P>|t|
					Lower	Upper		
1	3 March 2020	7 March 2020	5	-8.363	-16.063	0.044	-1.988	0.051
2	7 March 2020	16 March 2020	10	14.648*	11.228	18.173	9.021	0.000
3	16 March 2020	23 March 2020	8	28.113*	22.242	34.267	10.553	0.000
4	23 March 2020	26 March 2020	4	-8.612	-30.761	20.622	-0.648	0.519
5	26 March 2020	4 April 2020	10	27.008*	23.219	30.913	15.777	0.000
6	4 April 2020	21 May 2020	43	5.194*	4.972	5.416	47.944	0.000
All	3 March 2020	21 May 2020	80	9.094	7.694	10.513	13.206	0.000

Source: Author's calculations
Remarks: * Statistically significant at p=0.05.

Table 3

Forecast of daily reported confirmed cases of COVID-19 till 15 June 2020.

Date	Estimate	95% Confidence interval	
		Lower	Upper
May 22, 2020	6301	5694	6972
May 23, 2020	6629	5977	7350
May 24, 2020	6973	6274	7748
May 25, 2020	7335	6586	8168
May 26, 2020	7716	6913	8610
May 27, 2020	8117	7257	9076
May 28, 2020	8539	7618	9568
May 29, 2020	8982	7997	10086
May 30, 2020	9449	8394	10633
May 31, 2020	9939	8812	11208
June 1, 2020	10456	9250	11815
June 2, 2020	10999	9710	12455
June 3, 2020	11570	10192	13130
June 4, 2020	12171	10699	13841
June 5, 2020	12803	11231	14591
June 6, 2020	13468	11790	15381
June 7, 2020	14167	12376	16214
June 8, 2020	14903	12991	17092
June 9, 2020	15677	13637	18018
June 10, 2020	16492	14315	18994
June 11, 2020	17348	15027	20022
June 12, 2020	18249	15774	21107
June 13, 2020	19197	16558	22250
June 14, 2020	20194	17381	23455
June 15, 2020	21243	18246	24725

Source: Author's calculations