Reliability of P3 Event-Related Potential during Working Memory Across the Spectrum of Cognitive Aging

Hannes Devos[1*], Jeffrey Burns[2,3], Ke Liao[4], Pedram Ahmadnezhad,[1] Jonathan D. Mahnken,[3,4]

William M. Brooks,[2,3,4] Kathleen Gustafson[3,4]

[1]Laboratory for Advanced Rehabilitation Research in Simulation, Department of Physical Therapy

and Rehabilitation Science, University of Kansas Medical Center, Kansas City, KS, USA

[2]Department of Neurology, University of Kansas Medical Center, Kansas City, KS

[3]University of Kansas Alzheimer's Disease Center, University of Kansas Medical Center, Kansas

City, KS

[4]Hoglund Brain Imaging Center, University of Kansas Medical Center, Kansas City KS

[5]Department of Biostatistics & Data Science, University of Kansas Medical Center, Kansas City, KS

* Correspondence:

Hannes Devos

Department of Physical Therapy and Rehabilitation Science, University of Kansas Medical Center,

MS 2002 3901 Rainbow Blvd., Kansas City KS

E-mail: hdevos@kumc.edu

Keywords: event-related potentials, electro-encephalography, reliability, working memory,

older adults, mild cognitive impairment, Alzheimer disease, preclinical

Abstract

Event-related potentials (ERPs) offer unparalleled temporal resolution in tracing distinct

electrophysiological processes related to normal and pathological cognitive aging. The stability of

ERPs in older individuals, who inherently show more intraindividual variability in cognitive

functions, has not been established. In this test-retest reliability study, 39 older individuals (age 74.10

(5.4) years; 23 (59%) women; 15 non β-amyloid elevated, 16 β-amyloid elevated, 8 cognitively

impaired) with scores on the Montreal Cognitive Assessment (MOCA) ranging between 3 and 30

completed a working memory (n-back) test with three levels of difficulty at baseline and two-week

follow-up. Stability of the ERP was evaluated on grand averaged task effects for both visits. P3 peak

amplitude and latency were measured in frontal channels. P3 peak amplitude at Fz, our main outcome

variable, showed excellent reliability in 0-back (intraclass correlation coefficient (ICC), 95%

confidence interval = 0.82 (0.67 – 0.90) and 1-back (ICC = 0.87 (0.76 – 0.93), however, only fair

reliability in 2-back (ICC = 0.53 (0.09 – 0.75). Reliability of P3 peak latencies was substantially

lower, with ICCs ranging between 0.17 for 2-back and 0.54 for 0-back. Generalized linear mixed

models showed no confounding effect of age, group, or task difficulty on stability of P3 amplitude

and latency of Fz. By contrast, MOCA scores tended to negatively correlate with P3 amplitude of Fz

(p=0.07). We conclude that P3 peak amplitude and latency provide a stable measure of

electrophysiological processes in older individuals. However, impaired cognition may affect the

stability of the ERP response.

1   Introduction

The aging process is characterized by gradual decline in physical, neurobiological and cognitive

functions that may impact instrumental activities of daily living (iADL) such as driving, doing

household chores, managing finances, medication adherence, or grocery shopping.(1, 2)

Deterioration in these iADL becomes more apparent with age-related neurodegeneration such as mild

cognitive impairment (MCI) and Alzheimer's disease (AD).(3) Executive functions in particular are

paramount in carrying out numerous iADL, but are also vulnerable to the effects of normal and

pathological cognitive aging.(4, 5) Working memory is one core executive function that relates to the

ability to temporarily store, process, and manipulate the information necessary for higher order

cognitive tasks such as decision making, learning, and reasoning.(6) Working memory stems from

the interaction between attention, short-term retention and manipulation of information, carried out

by the coordinated activation of many brain regions.(7)

The prefrontal cortex has particularly been associated with working memory.(8) Consequently, the

prefrontal cortex is highly susceptible to the effects of aging and early neurodegeneration.(9, 10) A

recent meta-analysis pooling functional magnetic resonance imaging studies suggested a gradual and

linear decline in prefrontal cortex engagement in older individuals.(11) Similarly,

electrophysiological processes also decline with age. The P3, a positive peak that appears with a

latency between 250 to 500ms in the event-related potential (ERP), has been implicated in attention

and working memory processes across the lifespan.(12) A previous study showed reduced positivity

in P3 central-frontal and parietal ERPs in older adults,(13) whereas others demonstrated frontal

hyperactivity in P3 coupled with parietal or posterior hypoactivity.(14, 15). Despite the ambiguity in

ERP findings, most studies conclude that the abnormal ERP response in older individuals reflects

inefficient or compensatory use of neural resources due to frontal cortex dysfunction.(13, 14)

Therefore, electrophysiological responses to working memory tasks are convenient measures to test

hypotheses related to frontal cortex function, normal cognitive aging, and early neurodegeneration.

The ability to distinguish natural variability and measurement error from biologically relevant

cognitive changes due to aging or early neurodegeneration is valuable to provide informed decisions

on diagnosis, monitoring, and treatment of cognitive impairments.(16) However, older adults show

more intraindividual variability in performance measures of working memory compared to younger

adults. The age-related changes in intraindividual variability of performance measures become even

more apparent with increasing cognitive demand.(9) This increased intraindividual may also stem

from the heterogeneity of cognitive profiles in older individuals, especially when patients with MCI

and AD are included.(17) The intraindividual variability observed in performance measures is

believed to be linked to frontal cortex dysfunction,(9) which may therefore also affect intraindividual

variability of the ERP response in older adults.(18) To date, few studies have investigated test-retest

reliability of P3 ERP in healthy older adults(19-21) The test-retest reliability of the P3 ERP in older

individuals with a heterogeneous cognitive profile has yet to be established.

The aims were to (1) characterize test-retest reliability of P3 amplitude and latency, particularly in

the frontal-midline channels; and (2) investigate the impact of age, disease groups, cognitive status,

and task difficulty P3 amplitude and latency.

2   Materials and Methods

2.1   Participants

This test-retest reliability study included 39 right-handed participants recruited from the KU

Alzheimer's Disease Center between 05/03/2018 and 03/10/2020. Inclusion criteria were (1) age

older than 65; (2) having undergone a previous amyloid PET scan of the brain; (3) ability to

understand the instructions in English; and (4) informed consent. Exclusion criteria were: (1)

currently taking steroids, benzodiazepines, or neuroleptics; (2) history of any substance abuse; and

(3) history of a neurological disorder other than MCI or AD. Sixteen were cognitively normal older

adults with no elevated amyloid PET scans (Aβ-), 15 were cognitive normal with elevated amyloid

PET scans (Aβ+), and eight had a clinical diagnosis of mild cognitive impairment with positive

amyloid PET scans. Participants completed their two-week follow-up session 16 ± 8 days after the

first session. Each session lasted about 60 minutes including rest breaks.

2.2   Procedure

2.2.1 Demographic and Clinical Information

92   Age, sex, and education were recorded. General cognitive functions were evaluated with the

93   Montreal Cognitive Assessment (MOCA).(22) Scores on the MOCA range between 0 and 30.

2.2.2 N-back Test

In the n-back test, participants are shown a series of letters and are instructed to press a button when

the current stimulus is the same as the item presented n-positions back. The cognitive demand of the

n-back task increases with each number, while the perceptual and motor demands remain constant.

Whereas in the 0-back and 1-back the stimulus on screen is held in the focus of attention, the more

difficult versions of the n-back tests require constant switching from the focus of attention to short-

term memory.(23) Higher levels of difficulty require continuous mental effort to update information

of new stimuli and maintain representations of recently presented stimuli.(24)

In this study, the 0-back, 1-back, and 2-back tests were administered. Participants sat in a

comfortable chair at 26 inches in front of the computer screen with the center of the screen at eye

level. White letters appeared on a black screen. Prior to each test, participants were given a practice

trial consisting of 7 non-targets and 3 targets. The practice trials were repeated until the participant

felt comfortable with the instructions. Each test comprised 180 trials, including 60 trials that needed a

response (target, 33.3%) and 120 trials for which a response was not required (nontarget, 66.7%).

Each letter was presented for 500 ms on the computer screen followed by a blank interstimulus

interval for 1700 ms, with a random jitter of +/-50 ms. The maximum time to accept the response was

2150 ms. The total task time was ~7 minutes. In the 0-back test, participants were instructed to press

the left mouse button as soon as the letter "X" (target) appeared on the screen while ignoring the

other letters (nontarget). In the 1-back test, participants were instructed to press the button if the

current letter on the screen was the same as the letter previously shown (target). In the 2-back test,

participants were instructed to press the button when the current letter was the same as the one

presented two places before (target). The number of hits (accuracy) and response times to the hits

were the main behavioral performance outcome measures.

2.2.3 P3 ERP

Continuous electro-encephalogram (EEG) was acquired using a Philips EGI high-density system

from 256 scalp electrodes, digitized at 1,000 Hz. Data were filtered from 0.50 to 30 Hz using EGI

software. Data were online referenced to Cz and offline rereferenced to the left and right mastoid. All

other EEG processing was done in EEGLab(25) and in ERPLab.(26) Various artifacts unrelated to

cognitive functions, including ocular and muscular movement, cardiovascular signal were identified

and removed using independent component analysis (ICA). Signal from bad electrodes were

interpolated using surrounding electrode data. Stimulus-locked ERPs were extracted from the n-back

tests and segmented into epochs of 100 ms before to 1000ms after stimulus onset, and baseline

corrected using the prestimulus interval. Scalp locations and measurement windows for the P3

component were based on their spatial extent and latency after inspection of grand average

waveforms (collapsed across the two sessions). P3 peak amplitude and peak latency of the task effect

were considered the main electrophysiological outcome measures. The difference waveform was

calculated by subtracting the average ERP elicited from the targets from the average ERP elicited by

non-targets for each participant. The P3 component time window was established between 200 ms

and 400 ms for all three tests. Because of the prefrontal cortex involvement in working memory, we

identified a priori the Fz channel as the main channel, but also calculated reliability of other pre-

identified electrode locations, i.e., Cz, Pz, F3, and F4. Cz was interpolated using the surrounding five

channels. No participants were removed from the analyses because of artifacts. However, one

participant disengaged during the 2-back test and was therefore excluded from the analyses.

2.3   Data Analysis

Descriptive analysis including mean (standard deviation) and frequency count of participants'

general, performance measures, and ERP data were performed as appropriate. Intra-class correlation

coefficients (ICC) were used to calculate test-retest reliability of performance measures and P3

amplitude and latency. ICCs reflect the consistency of a measure taking into account variance related

to the time of testing.(27) ICC values less than 0.40 were considered poor; values between 0.40 and

0.59 fair, values between 0.60 and 0.74 good, and values between 0.75 and 1.00 excellent.(28).

Bland-Altman plots were used to visualize the measurement precision of amplitude and latency

across the test moments.(29) Intersubject stability according to subject rankings was calculated using

the Pearson r correlation coefficient. Generalized linear mixed models were employed to evaluate the

effect of age, diagnosis (Aβ-; Aβ+; MCI/AD), MOCA scores, and task difficulty on stability of the

P3 amplitude and latency. Stability of P3 amplitude (latency) was calculated as the squared

difference of P3 amplitude (latency) at follow-up and baseline. The Kolmogorov–Smirnov test was

employed to test the normality of our data distribution in addition to visualization of Q-Q plots. All

analyses were done using SAS 9.4 software. The threshold of significance was set at $p = 0.05$.

3   Results

3.1   Participant Characteristics

Participants (n = 39) were on average 74.05 (5.37) years old and scored 26.44 (4.76) on the MOCA

scale. However, MOCA scores ranged from 3 to 30, indicating a wide range of severe to normal

cognitive functions. About 60% (n = 23) were women.

3.2   Test-Retest Reliability of Performance Measures

All ICC values of hits (accuracy) and response times of each n-back test demonstrated excellent

reliability (Supplementary Table 1). ICCs of hits ranged between 0.92 (1- and 2-back) and 0.99 (0-

back) and were slightly higher than the ICCs of response times, ranging between 0.76 (2-back) and

0.89 (1-back). Pearson r correlations ranged from 0.65 (0-back response time) to 0.99 (0-back hits).

3.3   Test-Retest Reliability of ERP Measures

Grand average waveforms of the task effect from all channels at baseline and follow-up are displayed

in Figure 1. The 3D scalp map is embedded in the figure to demonstrate the task effect at P3.

Considerable overlap in ERP response within the P3 time window (200 – 400 ms post-stimulus) was

observed at baseline and two-week follow-up.

The ICC values of P3 peak amplitude and peak latency of the key electrode locations are displayed in

Table 1. Overall, P3 amplitude showed greater reliability compared to P3 latency across channels and

task difficulty levels. Also, ICCs of the 0-back and 1-back were consistently higher than those

calculated for the 2-back.

For the main channel location Fz, excellent reliability was found in P3 amplitude for 0-back (ICC =

0.82) and 1-back (ICC = 0.87). P3 amplitude of Fz for 2-back only showed fair reliability. Reliability

scores of P3 latency at Fz were fair for 0-back (ICC = 0.54) and 1-back (ICC = 0.47), but poor for 2-

back (ICC = 0.17). Figure 2 shows the Bland-Altman plots for P3 peak amplitude and peak latency at

the Fz channel. All plots demonstrated equal distribution of the data around zero, indicating no bias

in the results and no heteroscedasticity within the data.

Finally, generalized linear mixed models were employed to evaluate the effect of age, disease

diagnosis (Aβ-; Aβ+; MCI/AD), cognitive status, and task difficulty on stability of squared P3 peak

amplitude and latency at the Fz channel. Age (p=0.74), disease diagnosis (p = 0.67), and task

difficulty (p = 0.70) did not affect the stability of the P3 amplitude response, although individuals

with lower MOCA cognitive scores tended to show more variability in P3 amplitude (p = 0.07).

Age (p = 0.60), disease diagnosis (p = 0.55), MOCA (p = 0.52), or task difficulty (p = 0.95) did not

affect the stability of the P3 latency response.

We recalculated ICC for 0-back, 1-back, and 2-back in participants who scored 26 or higher on

MOCA (n = 32) and those scoring lower than 26 (n = 7). ICC values showed more variance in 0-back

and in 2-back in the group with lower MOCA scores, but ICC values were not worse across the n-

back tests in this group (Supplementary Tables 2 and 3). Whereas ICCs were similar in the Aβ- and

Aβ+ groups, lower ICCs were found for the MCI/AD group (Supplementary Tables 2 and 3).

4   Discussion

This test-retest reliability study provides critical information on the stability of electrophysiological

measures related to working memory in healthy older adults, older adults with increased risk of

dementia, and those with MCI or AD. Our results showed that most P3 ERPs in the frontal channels

provide fair to excellent reliability to measure electrophysiological processes of cognitive aging in

older adults with and without cognitive impairments. Similar to previous studies, the reliability is

superior in measures of amplitude compared to latency.(20, 21, 30, 31) The robustness of P3 stability

is not affected by age, disease diagnosis, or task difficulty, however, there is a trend that lower

MOCA scores may affect the stability of the P3 amplitude response.

The body of evidence related to reliability of P3 ERPs is sparse, and typically restricted to healthy

young,(30-34) middle-aged,(31) and older individuals.(19-21). Few studies have reported reliability

measures in neurological conditions.(35) The reliability analyses in our study produced fair to

excellent ICC values across the n-back tests. Whereas ICC values provide a single measure of the

magnitude of agreement, Bland-Altman plots depict a graphical display of bias across the two test

moments.(36) Visual inspection of the Bland-Altman plots showed average difference in ERP

responses between first and second testing close to 0, with equal spread of data points around the

average difference line. These findings suggest that 2 weeks follow-up is sufficient to wash out any

potential adaptation, test, or practice effect of the n-back on ERPs in older individuals.

Comparison of our results with other test-retest studies of ERPs in older adults is complicated by lack

of consistency in terms of the ERP components that are investigated, the tests of working memory,

the choice of channel locations, the extracted P3 metric, the P3 window measurement, and the test-

retest reliability intervals.(19-21). Our research design most closely aligns with a study that

compared ERPs to novel stimuli collected at baseline and 7-week follow-up in healthy older

individuals.(21) Similar to our study, this study also found excellent reliability for P3 mean

amplitude (ICC =0.86, 95% CI, 0.78 – 0.92), and poorer reliability for P3 mean latency (ICC = 0.56,

0.30 – 0.73). Our study demonstrated larger confidence intervals in some of the amplitude and

latency measures, which might have been due to the greater cognitive heterogeneity of our sample.

Another reliability study also reported considerably lower reliability in P3 amplitude (ICC = -0.02)

and latency (ICC = -0.17) in seven individuals experiencing cognitive difficulties following traumatic

brain injury compared to healthy peers (ICC = 0.84 for amplitude and 0.64 for latency).(35)

Combined, these findings point towards a potential confounding effect of cognitive impairment on

stability of ERPs in neurological conditions.

No effect of age, task difficulty, or disease diagnosis was found on stability of the P3 ERP in the n-

back task. Most participants in our study were cognitively normal, either without (n = 15) or with (n

= 16) elevated beta amyloid. Accumulation of beta amyloid deposits in the brain is known to increase

the risk of developing AD.(37) The fair to excellent reliability of P3 amplitude and latency in our

group of older adults provides opportunities for studying the effect of amyloid deposits on neural

transmission in preclinical AD using ERP. P3 amplitudes are smaller in AD compared to

controls.(38) ERPs also show useful in predicting conversion to AD, with accuracy rates ranging

between 70% and 94%.(39) Patients with AD exhibit prolonged latency in P3 ERP compared to age-

matched controls(40). These prolonged latencies observed in patients with AD become particularly

apparent in the cognitive domains of executive function, memory, and language.(41) The ability of

P3 ERP to discriminate between mild cognitive impairment (MCI) and AD(42) opens avenues for

investigation of ERP in detecting preclinical AD.(43)

Although this study included participants with a wide range of cognitive impairment, most were

cognitively normal. Future studies should include a larger sample of participants with MCI and AD

to confirm the confounding effect of impaired cognition on the stability of the P3 response. Although

the n-back is arguably the most ubiquitous working memory test used in ERP studies across the age

spectrum,(23) we did not establish reliability of ERP in other cognitive domains known to deteriorate

in older age, such as memory and language, and this remains an opportunity for further investigation.

Our results are therefore only generalizable to the specific subdomain of working memory. Future

research should also include multiple testing sessions over extended periods of time to evaluate the

sensitivity of ERP to detect subtle neurobiological changes due to normal and pathological aging.

5   Conclusion

We set out to assess the test-retest reliability of ERP response in older adults with a heterogeneous

cognitive profile. Consistent with other studies, P3 amplitude and latency show fair to excellent

reliability across different levels of task difficulty. However, impaired cognition may potentially

affect the stability of the P3 ERP response.

References

1.   Carmona-Torres JM, Rodríguez-Borrego MA, Laredo-Aguilera JA, López-Soto PJ,
Santacruz-Salas E, Cobo-Cuenca AI. Disability for basic and instrumental activities of daily living in
older individuals. *PLOS ONE* (2019) 14(7):e0220157. doi: 10.1371/journal.pone.0220157.

2.   Moon S, Ranchet M, Abiodun, Tant M, David, Mukaila, et al. The Impact of Advanced Age
on Driving Safety in Adults with Medical Conditions. *Gerontology* (2018) 64(3):291-9. doi:
10.1159/000486511.

3.   Jekel K, Damian M, Wattmo C, Hausner L, Bullock R, Connelly PJ, et al. Mild cognitive
impairment and deficits in instrumental activities of daily living: a systematic review. *Alzheimer's
Research & Therapy* (2015) 7(1). doi: 10.1186/s13195-015-0099-0.

4.   Overdorp EJ, Kessels RPC, Claassen JA, Oosterman JM. The Combined Effect of
Neuropsychological and Neuropathological Deficits on Instrumental Activities of Daily Living in
Older Adults: a Systematic Review. *Neuropsychology Review* (2016) 26(1):92-106. doi:
10.1007/s11065-015-9312-y.

5.   Tabira T, Hotta M, Murata M, Yoshiura K, Han G, Ishikawa T, et al. Age-Related Changes in
Instrumental and Basic Activities of Daily Living Impairment in Older Adults with Very Mild
Alzheimer's Disease. *Dementia and Geriatric Cognitive Disorders Extra* (2020):27-37. doi:
10.1159/000506281.

6.   Baddeley A. Working memory. *Science* (1992) 255(5044):556-9. doi:
10.1126/science.1736359.

7.   Eriksson J, Edward, Lansner A, Bergström F, Nyberg L. Neurocognitive Architecture of
Working Memory. *Neuron* (2015) 88(1):33-46. doi: 10.1016/j.neuron.2015.09.020.

8.   Bahmani Z, Clark K, Merrikhi Y, Mueller A, Pettine W, Isabel Vanegas M, et al. Prefrontal
Contributions to Attention and Working Memory. *Curr Top Behav Neurosci* (2019) 41:129-53. Epub
2019/02/11. doi: 10.1007/7854_2018_74. PubMed PMID: 30739308; PubMed Central PMCID:
PMCPMC6689265.

9.   West R, Murphy KJ, Armilio ML, Craik FIM, Stuss DT. Lapses of Intention and Performance
Variability Reveal Age-Related Increases in Fluctuations of Executive Control. (2002) 49(3):402-19.
doi: 10.1006/brcg.2001.1507.

10.   Ranchet M, Morgan JC, Akinwuntan AE, Devos H. Cognitive workload across the spectrum
of cognitive impairments: A systematic review of physiological measures. *Neuroscience &
Biobehavioral Reviews* (2017) 80:516-37. doi: 10.1016/j.neubiorev.2017.07.001.

11.   Yaple ZA, Stevens WD, Arsalidou M. Meta-analyses of the n-back working memory task:
fMRI evidence of age-related changes in prefrontal cortex involvement across the adult lifespan.
NeuroImage (2019) 196:16-31. doi: 10.1016/j.neuroimage.2019.03.074.

12.   Van Dinteren R, Arns M, Jongsma MLA, Kessels RPC. P300 Development across the
Lifespan: A Systematic Review and Meta-Analysis. *PLoS ONE* (2014) 9(2):e87347. doi:
10.1371/journal.pone.0087347.

13.   Lubitz AF, Niedeggen M, Feser M. Aging and working memory performance:
Electrophysiological correlates of high and low performing elderly. *Neuropsychologia* (2017)
106:42-51. Epub 2017/09/12. doi: 10.1016/j.neuropsychologia.2017.09.002. PubMed PMID:
28889995.

14.   Saliasi E, Geerligs L, Lorist MM, Maurits NM. The Relationship between P3 Amplitude and
Working Memory Performance Differs in Young and Older Adults. (2013) 8(5):e63701. doi:
10.1371/journal.pone.0063701.

15.   Fjell AM, Walhovd KB. P300 and neuropsychological tests as measures of aging: Scalp
topography and cognitive changes. *Brain Topography* (2001) 14(1):25-40. doi:
10.1023/a:1012563605837.

16.   Feinkohl I, Borchers F, Burkhardt S, Krampe H, Kraft A, Speidel S, et al. Stability of
neuropsychological test performance in older adults serving as normative controls for a study on
postoperative cognitive dysfunction. *BMC Research Notes* (2020) 13(1). doi: 10.1186/s13104-020-
4919-3.

17.   Troyer AK, Vandermorris S, Murphy KJ. Intraindividual variability in performance on
associative memory tasks is elevated in amnestic mild cognitive impairment. *Neuropsychologia*
(2016) 90:110-6. Epub 2016/06/15. doi: 10.1016/j.neuropsychologia.2016.06.011. PubMed PMID:
27297728.

18.   Robertson S, Myerson J, Hale S. Are There Age Differences in Intraindividual Variability in
Working Memory Performance? *The Journals of Gerontology Series B: Psychological Sciences and
Social Sciences* (2006) 61(1):P18-P24. doi: 10.1093/geronb/61.1.p18.

19.   Sandman CA, Patterson JV. The auditory event-related potential is a stable and reliable
measure in elderly subjects over a 3 year period. *Clinical Neurophysiology* (2000) 111(8):1427-37.
doi: 10.1016/s1388-2457(00)00320-5.

20.   Walhovd KB, Fjell AM. One-year test-retest reliability of auditory ERPs in young and old
adults. *Int J Psychophysiol* (2002) 46(1):29-40. Epub 2002/10/11. doi: 10.1016/s0167-
8760(02)00039-9. PubMed PMID: 12374644.

21.   Behforuzi H, Feng NC, Billig AR, Ryan E, Tusch ES, Holcomb PJ, et al. Markers of Novelty
Processing in Older Adults Are Stable and Reliable. *Frontiers in Aging Neuroscience* (2019) 11. doi:
10.3389/fnagi.2019.00165.

22.   Nasreddine ZS, Phillips NA, Bedirian V, Charbonneau S, Whitehead V, Collin I, et al. The
Montreal Cognitive Assessment, MoCA: A brief screening tool for mild cognitive impairment. *J Am
Geriatr Soc* (2005) 53(4):695-9. doi: 10.1111/j.1532-5415.2005.53221.x. PubMed PMID:
WOS:000227899200021.

23.   Bopp KL, Verhaeghen P. Aging and n-Back Performance: A Meta-Analysis. *The Journals of
Gerontology: Series B* (2018). doi: 10.1093/geronb/gby024.

24.   Gevins A, Smith ME, McEvoy LK, Ilan AB, Chan CS, Jiang A, et al. A cognitive and
neurophysiological test of change from an individual's baseline. *Clinical Neurophysiology* (2011)
122(1):114-20. doi: 10.1016/j.clinph.2010.06.010.

25.   Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis. *J Neurosci Methods* (2004) 134(1):9-21. Epub
2004/04/23. doi: 10.1016/j.jneumeth.2003.10.009. PubMed PMID: 15102499.

26.   Lopez-Calderon J, Luck SJ. ERPLAB: an open-source toolbox for the analysis of event-
related potentials. *Frontiers in Human Neuroscience* (2014) 8(213). doi: 10.3389/fnhum.2014.00213.

27.   Shrout PE, Fleiss JL. Intraclass correlations: uses in assessing rater reliability. *Psychol Bull*
(1979) 86(2):420-8. Epub 1979/03/01. doi: 10.1037//0033-2909.86.2.420. PubMed PMID: 18839484.

28.   Cicchetti DV. Guidelines, criteria, and rules of thumb for evaluating normed and standardized
assessment instruments in psychology. *Psychological assessment* (1994) 6(4):284.

29.   Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of
clinical measurement. *Lancet (London, England)* (1986) 1(8476):307-10. Epub 1986/02/08. PubMed
PMID: 2868172.

30.   Cassidy SM, Robertson IH, O'Connell RG. Retest reliability of event-related potentials:
Evidence from a variety of paradigms. *Psychophysiology* (2012) 49(5):659-64. doi: 10.1111/j.1469-
8986.2011.01349.x.

31.   Kinoshita S, Inoue M, Maeda H, Nakamura J, Morita K. Long-term patterns of change in
ERPs across repeated measurements. *Physiol Behav* (1996) 60(4):1087-92. Epub 1996/10/01. doi:
10.1016/0031-9384(96)00130-8. PubMed PMID: 8884937.

32.   Segalowitz SJ, Barnes KL. The reliability of ERP components in the auditory oddball
paradigm. *Psychophysiology* (1993) 30(5):451-9. Epub 1993/09/01. doi: 10.1111/j.1469-
8986.1993.tb02068.x. PubMed PMID: 8416071.

33.   Brunner JF, Hansen TI, Olsen A, Skandsen T, Haberg A, Kropotov J. Long-term test-retest
reliability of the P3 NoGo wave and two independent components decomposed from the P3 NoGo
wave in a visual Go/NoGo task. *Int J Psychophysiol* (2013) 89(1):106-14. Epub 2013/06/19. doi:
10.1016/j.ijpsycho.2013.06.005. PubMed PMID: 23770084.

34.   Huffmeijer R, Bakermans-Kranenburg MJ, Alink LR, van Ijzendoorn MH. Reliability of
event-related potentials: the influence of number of trials and electrodes. *Physiol Behav* (2014)
130:13-22. Epub 2014/03/20. doi: 10.1016/j.physbeh.2014.03.008. PubMed PMID: 24642000.

35.   Lew HL, Gray M, Poole JH. Temporal Stability of Auditory Event-Related Potentials in
Healthy Individuals and Patients With Traumatic Brain Injury. *Journal of Clinical Neurophysiology*
(2007) 24(5):392-7. doi: 10.1097/wnp.0b013e31814a56e3.

36.   Ranganathan P, Pramesh CS, Aggarwal R. Common pitfalls in statistical analysis: Measures
of agreement. *Perspect Clin Res* (2017) 8(4):187-91. doi: 10.4103/picr.PICR_123_17. PubMed
PMID: 29109937.

37.   Klunk WE, Engler H, Nordberg A, Wang Y, Blomqvist G, Holt DP, et al. Imaging brain
amyloid in Alzheimer's disease with Pittsburgh Compound-B. *Ann Neurol* (2004) 55(3):306-19.
Epub 2004/03/03. doi: 10.1002/ana.20009. PubMed PMID: 14991808.

38.   Hedges D, Janis R, Mickelson S, Keith C, Bennett D, Brown BL. P300 Amplitude in
Alzheimer's Disease. *Clinical EEG and Neuroscience* (2016) 47(1):48-55. doi:
10.1177/1550059414550567.

39.   Chapman RM, McCrary JW, Gardner MN, Sandoval TC, Guillily MD, Reilly LA, et al. Brain
ERP components predict which individuals progress to Alzheimer's disease and which do not.
(2011) 32(10):1742-55. doi: 10.1016/j.neurobiolaging.2009.11.010.

40.   Pedroso RV, Fraga FJ, Corazza DI, Andreatto CAA, Coelho FGDM, Costa JLR, et al.
Latência e amplitude do P300 auditivo na doença de Alzheimer: uma revisão sistemática. *Brazilian
Journal of Otorhinolaryngology* (2012) 78(4):126-32. doi: 10.1590/s1808-86942012000400023.

41.   Lee M-S, Lee S-H, Moon E-O, Moon Y-J, Kim S, Kim S-H, et al. Neuropsychological
correlates of the P300 in patients with Alzheimer's disease. *Progress in Neuro-Psychopharmacology
and Biological Psychiatry* (2013) 40:62-9. doi: 10.1016/j.pnpbp.2012.08.009.

42.   Bennys K, Portet F, Touchon J, Rondouin G. Diagnostic value of event-related evoked
potentials N200 and P300 subcomponents in early diagnosis of Alzheimer's disease and mild
cognitive impairment. *J Clin Neurophysiol* (2007) 24(5):405-12. Epub 2007/10/04. doi:
10.1097/WNP.0b013e31815068d5. PubMed PMID: 17912065.

43.   Rossini PM, Di Iorio R, Vecchio F, Anfossi M, Babiloni C, Bozzali M, et al. Early diagnosis
of Alzheimer's disease: the role of biomarkers including advanced EEG signal analysis. Report from
the IFCN-sponsored panel of experts. *Clinical Neurophysiology* (2020) 131(6):1287-310. doi:
10.1016/j.clinph.2020.03.003.

Tables

Table 1. Comparison of task effect (target – nontarget) P3 response at baseline and two-week follow-up.

Variable	Baseline	Follow-up	Pearson r	ICC, (95% CI)
0-back, Fz amplitude (µV)	4.87 (3.23)	5.02 (3.13)	0.63[a]	0.82 (0.67 – 0.90)[a]
0-back, Fz latency (ms)	291.49 (49.98)	293.31 (40.26)	0.38[b]	0.54 (0.13 – 0.76)[b]
0-back, Cz amplitude (µV)	2.71 (2.46)	2.59 (2.20)	0.56[a]	0.73 (0.48 – 0.85)[a]
0-back, Cz latency (ms)	289.49 (55.06)	290.03 (46.57)	0.30[b]	0.51 (0.07 – 0.74)[a]
0-back, Pz amplitude (µV)	1.94 (1.47)	2.01 (2.00)	0.38[b]	0.54 (0.11 – 0.76)[b]
0-back, Pz latency (ms)	309.74 (62.36)	315.92 (63.96)	0.34[b]	0.52 (0.08 – 0.75)[b]
0-back, F3 amplitude (µV)	3.99 (2.80)	4.29 (2.41)	0.58[a]	0.74 (0.49 – 0.86)[a]
0-back, F3 latency (ms)	297.33 (45.51)	295.51 (38.37)	0.31[b]	0.47 (-0.02 – 0.72)[b]
0-back, F4 amplitude (µV)	4.29 (2.93)	4.44 (2.72)	0.62[a]	0.77 (0.56 – 0.88)[a]
0-back, F4 latency (ms)	300.54 (52.41)	297.95 (35.36)	0.31	0.45 (-0.05 – 0.71)[b]
1-back, Fz amplitude (µV)	3.97 (3.27)	4.08 (3.76)	0.78[a]	0.87 (0.76 – 0.93)[a]
1-back, Fz latency (ms)	300.90 (38.45)	300.26 (45.77)	0.31	0.47 (-0.02 – 0.72)[a]
1-back, Cz amplitude (µV)	3.48 (3.05)	3.56 (3.02)	0.79[a]	0.86 (0.74 – 0.92)[a]
1-back, Cz latency (ms)	302.11 (37.55)	300.03 (42.39)	0.33	0.44 (-0.04 – 0.70)[a]
1-back, Pz amplitude (µV)	1.64 (1.51)	1.54 (1.70)	0.58[a]	0.73 (0.48 – 0.86)[a]
1-back, Pz latency (ms)	318.44 (66.66)	309.97 (64.17)	0.26	0.42 (-0.11 – 0.69)[b]
1-back, F3 amplitude (µV)	3.30 (3.43)	3.42 (3.47)	0.76[a]	0.87 (0.74 – 0.93)[a]
1-back, F3 latency (ms)	301.92 (44.80)	300.67 (49.86)	0.38[b]	0.55 (0.14 – 0.77)[b]
1-back, F4 amplitude (µV)	3.51 (2.95)	3.46 (3.37)	0.68[a]	0.81 (0.64 – 0.90)[a]
1-back, F4 latency (ms)	304.44 (43.16)	311.03 (45.42)	0.59[a]	0.75 (0.51 – 0.87)[a]
2-back, Fz amplitude (µV)	3.39 (2.29)	3.21 (2.15)	0.36[b]	0.53 (0.09 – 0.75)[b]
2-back, Fz latency (ms)	300.97 (39.67)	303.68 (45.65)	0.09	0.17 (-0.60 – 0.49)
2-back, Cz amplitude (µV)	2.28 (1.26)	2.08 (1.65)	0.39[b]	0.51 (0.12 – 0.78)[b]
2-back, Cz latency (ms)	300.63 (49.62)	302.21 (45.84)	0.31	0.46 (-0.04 – 0.70)
2-back, Pz amplitude (µV)	1.24 (1.20)	1.38 (1.83)	-0.03	-0.06 (-1.04 – 0.44)

2-back, Pz latency (ms)	318.74 (66.70)	321.63 (63.18)	0.08	0.15 (-0.63 – 0.56)
2-back, F3 amplitude (µV)	2.89 (2.03)	2.30 (1.65)	0.38[b]	0.54 (0.11 – 0.76)[b]
2-back, F3 latency (ms)	297.29 (47.56)	292.89 (46.84)	0.30	0.47 (-0.03 – 0.72)[b]
2-back, F4 amplitude (µV)	3.37 (2.32)	3.23 (1.89)	0.46[b]	0.63 (0.28 – 0.81)[a]
2-back, F4 latency (ms)	306.26 (44.99)	308.92 (43.12)	0.09	0.16 (-0.61 – 0.56)

Abbreviations: ICC, Intraclass correlation coefficient; CI, confidence interval

[a]$p < 0.0001$

[b]$p < 0.05$

Ethics Statement

The study was carried out in accordance with the Declaration of Helsinki. The study was approved by the University of Kansas Medical Center Internal Review Board. Informed consent was obtained from each participant. Each participant received $100 for participating.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

HD, JB, JM, WMB, and KG conceptualized the study. HD, KL, and KG worked out the EEG data processing steps. HD, PA, and KL administered the tests. HD and JM analysed the data. HD wrote the initial manuscript. JB, KL, PA, JM, WMB, and KG reviewed the manuscript and provided valuable comments.

Funding

Research reported in this publication was supported by the National Institute on Aging of the National Institutes of Health under Award Number K01 AG058785. This study was supported in part by a pilot grant of the KU Alzheimer Disease Center (P30 AG035982). The Hoglund Biomedical

Imaging Center is supported in part by S10 RR29577 and generous gifts from Forrest and Sally

Hoglund. The content is solely the responsibility of the authors and does not necessarily represent the

official views of the National Institutes of Health.

Acknowledgments

The authors thank the volunteers for their time and willingness to participate in this research. We are

also grateful for the staff at the KU Alzheimer Disease Center.

Figure Legends

Figure 1. Grand Average Event-related Potential Waveform of (a) 0-back, (b) 1-back and (c) 2-back.

Figure 2. Bland Altman Plotsof (a) 0-back Fz Peak Amplitude (b) 1-back Fz Peak Amplitude; (c) 2-

back Fz Peak Amplitude; (d) 0-back Fz Peak Latency; (e) 1-back Peak Latency; (f) 2-back Peak

Latency.