Title

Lessons from movement ecology for the return to work: modeling contacts and the spread of COVID-19

Authors

Allison K. Shaw[1*], Lauren A. White[2], Matthew Michalska-Smith[3,4], Elizabeth T. Borer[1], Meggan E. Craft[3], Eric W. Seabloom[1], Emilie Snell-Rood[1], Michael Travisano[1,5]

1. Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, Minnesota, USA
2. National Socio-Environmental Synthesis Center, Annapolis, Maryland, USA
3. Department of Veterinary Population Medicine, University of Minnesota, St. Paul, Minnesota, USA
4. Department of Plant Pathology, University of Minnesota, St. Paul, Minnesota, USA
5. BioTechnology Institute, University of Minnesota, St. Paul, Minnesota, USA

* corresponding author: ashaw@umn.edu

Abstract

Human behavior (movement, social contacts) plays a central role in the spread of pathogens like SARS-CoV-2. The rapid spread of SARS-CoV-2 was driven by global human movement, and initial lockdown measures aimed to localize movement and contact in order to slow spread. Thus, movement and contact patterns need to be explicitly considered when making reopening decisions, especially regarding return to work. Here, as a case study, we consider the initial stages of resuming research at a large research university, using approaches from movement ecology and contact network epidemiology. First, we develop a dynamical pathogen model describing movement between home and work; we show that limiting social contact, via reduced people or reduced time in the workplace are fairly equivalent strategies to slow pathogen spread. Second, we develop a model based on spatial contact patterns within a specific office and lab building on campus; we show that restricting on-campus activities to labs (rather than labs and offices) could dramatically alter (modularize) contact network structure and thus, potentially reduce pathogen spread by providing a workplace mechanism to reduce contact. Here we argue that explicitly accounting for human movement and contact behavior in the workplace can provide additional strategies to slow pathogen spread that can be used in conjunction with ongoing public health efforts.

Keywords

contact network epidemiology; disease ecology; infection; mathematical model; network; SARS-CoV-2; theoretical ecology; theory; transmission

Introduction

The cosmopolitan connectivity of modern society facilitated the rapid spread of SARS-CoV-2 around the globe in early 2020 [1]. The rate at which any pathogen spreads depends critically on host

movement behavior [2]. Indeed, estimates of key epidemiological parameters like the basic reproduction number (R_0) are highly variable in part because they are context-specific and are a function of behaviors

like movement and heterogenous contact structure [3,4]. Although most cases of COVID-19 (the disease caused by SARS-CoV-2) seem to be mild or even asymptomatic [5,6], the sheer number of cases to date

means that limited personnel, hospital beds, and ICU equipment can be rapidly overwhelmed, increasing mortality [7,8]. Thus, continuing our normal movement patterns, unmitigated, is not a viable strategy.

Without a vaccine or widespread immunity to SARS-CoV-2, our best defense to slow pathogen spread has been restricting movement and contacts through physical distancing [9], testing for SARS-CoV-2

when available [10] and contact tracing [11]. Lockdown measures have drastically reduced human movement [12,13] and consequently have reduced the effective reproduction number, R_e [14]. However,

such measures are affecting mental health [15] and have had a devastating impact on the economy, so individual regions are considering best practices for the reopening of businesses, schools, and other places

where people gather (e.g.,[16]). Decisions regarding next steps can be informed by recognizing that not all movement patterns nor all contact behaviors are equal in terms of pathogen spread.

Concepts from movement ecology and contact network epidemiology can provide helpful frameworks for understanding the nuanced interactions between movement, contacts and infection.

Increased movement does not always mean increased transmission risk [17]; for example, movement that either takes individuals away from infected areas or reduces contact with infected conspecifics can reduce

transmission risk (migratory escape; [18,19]). Increased movement can even increase some aspects of infection risk while decreasing others, simultaneously [20]. Thus, explicitly considering how movement

relates to transmission can help us understand what effect different movement patterns have on infection dynamics [21,22]. Similarly, from disease ecology and contact network epidemiology, we know that

structured contacts among individuals in a population have different effects on disease spread than

random contacts. For example, long-range connections in otherwise locally-connected small world

68 networks can have dramatic effects on disease spread at a population level [23].

Individual movement across multiple scales — from occasional global movements to smaller-

70 scale daily patterns — is critical for shaping contact and thus the spread of pathogens. To date, models of

SARS-CoV-2/COVID-19 spread have focused on comparing patterns of spread across countries, states,

72 and counties [24,25]. Indeed, a plethora of epidemiological models have proven useful in generating

recommendations for reducing the virus spread rate, from understanding the role of contact-tracing and

74 society-wide physical distancing [11,26], to travel restrictions, and region-wide lockdowns (e.g., [27–

30]). However, few if any offer guidance at scales as fine as individual workplaces, despite the fact that

76 this local scale is where individual decisions are made and where most transmission occurs. Furthermore,

apart from time at home, the most predictable component of many people's days is time spent in the

78 workplace. Thus, knowledge of work commute patterns, contact networks of individuals in the

workplace, and related workplace-specific factors could help mitigate pathogen spread during the period

80 that total population immunity remains low. In many cases, commute trajectories are not random but

involve regularity in timing, location, and encounters with other individuals along the way (e.g., on public

82 transport). Here, we consider the implications for mitigating COVID-19 transmission using a case study

of the initial stages of resuming research at a large research university.

84 Implicit in this analysis is that COVID-19 is currently spreading in local communities around the

world, and every individual in a workplace is part of a home community. Even under many weeks of

86 extreme restrictions with only society's most essential employees present in workplaces (i.e., Stay at

Home orders), the number of new cases have continued to rise in most locations. For example, in late

88 April 2020, even after three weeks of a Stay at Home order and extreme physical distancing in Minnesota,

a state with moderate spread and commendable compliance with the order, the number of new cases

90 confirmed each day had tripled [31]. With community spread of this pathogen, it is unrealistic to expect

zero workplace infection or widespread virus containment primarily through workplace practices. Any

92 return-to-work plan, therefore, must include the explicit expectation that new infections may arise while

concurrently prioritizing worker safety and optimizing the work that can be done. Thus, reopening

94 businesses requires an evidence-based plan to reduce contacts through time to minimize new infections at

the workplace, when an infected individual, presumably pre-symptomatic [32], brings the virus to a

96 workplace.

Here we develop a pair of models to understand how movement and contact structure shape

98 infection spread. As a case study, we consider the context of moving from full-time work at home to part-

time resumption of research at a university; however, results from this model are general to many other

100 settings as well. We take a dual modeling approach by developing a general movement model and a

network case study of one academic laboratory and office building. We explore tradeoffs between

102 limiting contact, people, or time on campus. We find that moving back to work on campus does not

necessarily speed up infection spread, and depends particularly on the infection risk associated with

104 commutes and how well physical distancing can be maintained on campus. Thus our findings allow us to

set evidence-based expectations and generate specific behavioral recommendations for a safer return to

106 work.

Materials and Methods

We develop two models: a movement model to explore movement between home and work

110 environments, and a network model to explore contact patterns within the work environment. Both

models are SEIR, tracking susceptible (S), exposed (E), infected (I), and removed (recovered and

112 immune, or deceased; R) individuals. We assume there is no loss of immunity (removed individuals never

move back to the susceptible class) over the short time scales we consider, and we assume a closed

114 population (no births, deaths, immigration, or emigration).

Methods: movement model

Setup

118   Our first model explores infection dynamics as individuals move between home, commuting, and
120 work environments. Workplaces, including universities, face a number of different decisions about how
to slowly ramp up work following easing of lockdown. Here, we simulate three potential strategies for
122 returning to work: (i) allowing people to return while maintaining physical distancing, (ii) limiting the
number of people returning to campus during the work day, and (iii) limiting the time each person spends
124 on campus. For each strategy combination, we simulate infection dynamics and quantify two output
metrics: (1) the 'final epidemic size' (cumulative fraction of the population infected, at equilibrium), and
126 (2) the 'epidemic peak size' (maximum fraction of the population infected during the outbreak). The aim
of this type of conceptual model is to clarify the connections between assumptions and outcomes, unlike
128 predictive models which would contain an abundance of empirical data and aim to generate forecasts for
a specific system [33].

Daily cycle

132   Our model dynamics have a combination of continuous and discrete time (e.g., [34]), where each
day is broken into discrete phases (T_h spent at home, T_w spent at work, and T_c spent commuting each way,
134 with $T_h + T_w + 2T_c = 1$) and infection dynamics occur continuously during each phase (Fig 1, see Tables
1-2 for model variables and parameters). All individuals start at home and spend a fraction of their day (of
136 length T_h) there and not working. During this time, the infection dynamics are given by

$$dS/dt = -\beta_h\, S\,(I/N)$$

138   $$dE/dt = \beta_h\, S\,(I/N) - \sigma\, E \hspace{3cm} \text{[eqn. 1]}$$

$$dI/dt = \sigma\, E - \gamma\, I$$

140   $$dR/dt = \gamma\, I$$

where S is the number of susceptible individuals, E is the number of exposed individuals, I is the number

142   of infected individuals, R is the number of removed (recovered and immune, or deceased) individuals, N

is the total number of individuals in the population ($N = S + E + I + R$), β_h is the rate of transmission while

144   at home, σ is the rate of moving from exposed to infected (inverse of the latent period) and γ is the rate of

recovery from infection.

[Fig 1. Movement model schematic, showing a daily cycle.

148   All individuals — Susceptible (S), Exposed (E), Infected (I) or Removed (R) — spend part of their day

(T_h) at home. A proportion θ of individuals move to campus, spending T_c time commuting in each

150   direction, and work from campus during the workday (time T_w), while the other fraction (1 - θ) works

from home. A total 24 hour cycle is then represented by: $T_h + T_w + 2\,T_c = 1$. Transmission rates can vary

152   among home (β_h; this includes transmission during essential trips e.g., to the grocery store), commute (β_c;

traveling between home and work), and work (β_w; campus-based interactions) environments, while the

154   rate of moving from exposed to infected (σ) and recovery rate (γ) are the same regardless of where

individuals are located.]

Here, the rate at which new susceptible individuals (S) become exposed (E) depends on three

158   components [35]. First is the rate of contact between two individuals in a location. Here we assume this

contact rate is constant (does not change with population density) but can differ across environments

160   (home vs. work vs. commuting). Critically, we assume that β_h accounts for transmission not just in an

individual's actual home, but transmission that occurs during other essential activities during lockdown

162   (e.g., grocery store trips). Second is the probability that the contact for each susceptible individual is with

an infected individual; this is given by the proportion of infected individuals in the local population (I/N).

164   Third is the probability that contact with an infectious individual results in transmission. In equation 1

above (and the other equations below), we have combined the first and third factors into a single term, β,

166   while the second factor is given by I/N. Overall, this gives us frequency-dependent transmission

(transmission rate depends on the frequency — not density — of infected individuals in the population);

168 an appropriate assumption for spatially structured environments [2,35].

After the period of time at home (T_h), a fraction, θ, of all individuals commute to work while the

170 remaining (1- θ) stay to work from home. At this point we subdivide the population based on the number

of individuals of each type and fraction commuting. We denote location by subscripts (*h* for home, *c* for

172 commute), so the number of individuals of each type are

$$S_c(T_h) = \theta\, S(T_h)$$

174 $$E_c(T_h) = \theta\, E(T_h)$$

$$I_c(T_h) = \theta\, I(T_h)$$

176 $$R_c(T_h) = \theta\, R(T_h) \qquad\qquad\qquad \text{[eqn. 2]}$$

$$S_h(T_h) = (1 - \theta)\, S(T_h)$$

178 $$E_h(T_h) = (1 - \theta)\, E(T_h)$$

$$I_h(T_h) = (1 - \theta)\, I(T_h)$$

180 $$R_h(T_h) = (1 - \theta)\, R(T_h) \ .$$

During the commute phase, the infection dynamics for those commuting are given by

182 $$dS_c/dt = -\,\beta_c\, S_c\, (I_c/N_c)$$

$$dE_c/dt = \beta_c\, S_c\, (I_c/N_c) - \sigma\, E_c \qquad\qquad \text{[eqn. 3]}$$

184 $$dI_c/dt = \sigma\, E_c - \gamma\, I_c$$

$$dR_c/dt = \gamma\, I_c$$

186 where N_c is the total number of individuals commuting ($N_c = S_c + E_c + I_c + R_c$), and β_c is the rate of

transmission while commuting. Similarly, during the commute phase, the infection dynamics for those

188 still at home are given by

$$dS_h/dt = -\,\beta_h\, S_h\, (I_h/N_h)$$

190 $$dE_h/dt = \beta_h\, S_h\, (I_h/N_h) - \sigma\, E_h \qquad\qquad \text{[eqn. 4]}$$

$$dI_h/dt = \sigma\, E_h - \gamma\, I_h$$

192 $$dR_h/dt = \gamma\, I_h$$

where N_h is the total number of individuals at home ($N_h = S_h + E_h + I_h + R_h$).

16   8

194   After the commute phase (of length T_c), comes a work phase. Here, the population continues to be

subdivided into eight types, where the number of individuals of each type are

196 $$S_w(T_h+T_c) = S_c(T_h+T_c)$$

$$E_w(T_h+T_c) = E_c(T_h+T_c)$$

198 $$I_w(T_h+T_c) = I_c(T_h+T_c)$$

$$R_w(T_h+T_c) = R_c(T_h+T_c) \hspace{3cm} \text{[eqn. 5]}$$

200 $$S_h(T_h+T_c) = S_h(T_h+T_c)$$

$$E_h(T_h+T_c) = E_h(T_h+T_c)$$

202 $$I_h(T_h+T_c) = I_h(T_h+T_c)$$

$$R_h(T_h+T_c) = R_h(T_h+T_c)$$

204 where the subscript w denotes work. Note that individuals that work on campus switch from a commute

(c) subscript to a work (w) one here, while individuals that work at home continue with the same

206 subscript (h). Both groups are still experiencing infection dynamics. During the work phase, the infection

dynamics in the workplace are given by

208 $$dS_w/dt = -\beta_w\, S_w\, (I_w/N_w)$$

$$dE_w/dt = \beta_w\, S_w\, (I_w/N_w) - \sigma\, E_w \hspace{2cm} \text{[eqn. 6]}$$

210 $$dI_w/dt = \sigma\, E_w - \gamma\, I_w$$

$$dR_w/dt = \gamma\, I_w$$

212 where N_w is the total number of individuals at work ($N_w = S_w + E_w + I_w + R_w$), and β_w is the rate of

transmission while at work. During the work phase, the infection dynamics for those working at home are

214 given by [eqn. 4] above.

        After the work phase (of length T_w), we describe a second commute phase. The population

216 continues to be subdivided into eight types, where the number of individuals of each type are

$$S_c(T_h+T_c+T_w) = S_w(T_h+T_c+T_w)$$

218 $$E_c(T_h+T_c+T_w) = E_w(T_h+T_c+T_w)$$

$$I_c(T_h+T_c+T_w) = I_w(T_h+T_c+T_w)$$

220 $$R_c(T_h+T_c+T_w) = R_w(T_h+T_c+T_w) \qquad \text{[eqn. 7]}$$

$$S_h(T_h+T_c+T_w) = S_h(T_h+T_c+T_w)$$

222 $$E_h(T_h+T_c+T_w) = E_h(T_h+T_c+T_w)$$

$$I_h(T_h+T_c+T_w) = I_h(T_h+T_c+T_w)$$

224 $$R_h(T_h+T_c+T_w) = R_h(T_h+T_c+T_w) \; .$$

During this second commute phase (also of length T_c), the infection dynamics for those commuting are

226 given by [eqn. 3] above, and the infection dynamics for those still at home are given by [eqn. 4] above. At

the end of the second commute phase, all individuals are back in the home environment (no longer

228 subdivided) and the number of individuals of each type are

$$S(T_h+2T_c+T_w) = S_w(T_h+2T_c+T_w) + S_h(T_h+2T_c+T_w)$$

230 $$E(T_h+2T_c+T_w) = E_w(T_h+2T_c+T_w) + E_h(T_h+2T_c+T_w) \qquad \text{[eqn. 8]}$$

$$I(T_h+2T_c+T_w) = I_w(T_h+2T_c+T_w) + I_h(T_h+2T_c+T_w)$$

232 $$R(T_h+2T_c+T_w) = R_w(T_h+2T_c+T_w) + R_h(T_h+2T_c+T_w) \; .$$

This ends the cycle for a single day; the next day starts the cycle again.

Model Parameters

236   We used a fixed population size (N) of 3,000 individuals. We did not include births or deaths, or

movement in and out of the population. These are reasonable assumptions given the scope of our

238 simulations: a work population that is not hiring new employees and has few retirements or actual deaths

over a few months. Because we assumed frequency-dependent transmission, the relative fraction of the

240 population infected is the same regardless of population size.

      Infection parameters were calculated as follows. We used 4.2 days as the latent period [36] and

242 calculated the rate of moving from exposed to infected (σ) as the inverse of this: $\sigma = 1/4.2 = 0.238$ per

day. We used 9.5 days as the infectious period (the estimated length of viral shedding for SARS-coV-2;

244 [37]), and calculated the recovery rate (γ) as the inverse of this: $\gamma = 1/9.5 = 0.105$ per day. Transmission

rate (β) was calculated based on the basic reproductive number, R_0. We assumed a 'baseline' R_0

246   (unmitigated; no behavioral changes like physical distancing) of 2.5 based on current estimates for SARS-

coV-2 [38], although some estimates put R_0 as high as 5.7 [36]. To quantify how behavioral changes to

248   movement and contact affect transmission we defined effective reproduction numbers (R_e) for each of the

environments (home, work, commute). We assumed that stay-at-home measures to reduce pathogen

250   spread in the community halved the rate of contacts at home (e.g., [39]), that is $R_{e-h} = 0.5R_0$. We assumed

that infection at work could be anywhere between current infection rates at home ($R_{e-w} = 0.5R_0$) and

252   unmitigated rates ($R_{e-w} = R_0$). To facilitate interpretation of our results, we also describe infection at work

in terms of the fraction increase in transmission compared to home, where 0 indicates transmission is the

254   same at work and home, 0.5 indicates transmission at work is 50% higher than at home and 1 indicates

transmission at work is 100% higher than at home (i.e., double). Finally, we assumed that infection while

256   commuting spanned a broader range of possible values than either home or work. At one extreme,

commuting by private transport effectively has no risk of transmission from others ($R_{e-c} = 0$). At the other

258   extreme, commuting by crowded public transport can reduce feasible physical distancing ($R_{e-c} = 2R_0$),

both because individuals have a greater number of contacts while commuting and because these contacts

260   potentially last for longer than normal. Transmission rates (β) were back-calculated from R_e values, based

on rearranging the expression $R_e = \beta/\gamma$ to $\beta = \gamma R_e$. We assume that R_0 and R_e values estimated for the

262   general public apply to our population of University workers. If instead our population had lower

transmission rates than the general public under stay-at-home measures, going back to work could lead to

264   faster pathogen spread than we predict here.

266   **Simulations**

Since our aim was to understand the relative importance of model parameters on infection

268   dynamics (rather than try to forecast outcomes), we started each simulation with one individual infected

($I(t{=}0){=}1$), zero exposed ($E(t{=}0){=}0$), zero removed ($R(t{=}0){=}0$), and the rest susceptible ($S(t{=}0){=}2{,}999$).

270   Each simulation was run until it reached equilibrium (where the fraction of the population in the R class

did not change from one day to the next). We defined a baseline set of values for each parameter (see

272   Table 2). Then we ran the following simulations that varied some parameters while holding others

constant:

274   (i)   Varying transmission while at work (β_w) and during the commutes (β_c). We considered three

scenarios that differed in the degree of risk of a commute to work and back. For low risk, we assumed

276   low contact both during commutes and on campus ($R_{e\text{-}w} = 0.5R_0 = 1.25$, equivalent to at home). For

moderate risk, we assumed unmitigated contact during commute ($R_{e\text{-}c} = R_0 = 2.5$, shared transport)

278   and partial physical distancing at work ($R_{e\text{-}w} = 0.75R_0 = 1.875$, intermediate between home and

unmitigated). For high risk, we assumed elevated contact during commute ($R_{e\text{-}c} = 2R_0 = 5$, crowded

280   shared transport), and unmitigated contact at work ($R_{e\text{-}w} = R_0 = 2.5$). These results are presented in Fig

2.

282   (ii) Varying the fraction of the population commuting (θ) and fraction of the day spent on campus

(T_w). We considered eleven values of the fraction of the population commuting ($\theta = 0,0.1,...,0.9,1$)

284   and eleven values of the fraction of an 8-hour workday spent on campus ($T_w = x(8/24)$ where x =

0,0.1,...,0.9,1). These results are presented in Fig 3a.

286   (iii) Varying the fraction of the population commuting (θ) and fraction increase in transmission at

work compared to home ($R_{e\text{-}w}$). We considered eleven values of the fraction of the population

288   commuting ($\theta = 0,0.1,...,0.9,1$) and eleven values of the fraction increase in transmission at work

compared to at home ($R_{e\text{-}w} = (1+x)R_{e\text{-}h}$ where x = 0,0.1,...,0.9,1). These results are presented in Fig 3b.

290   (iv) Varying the fraction of the day spent on campus (T_w) and fraction increase in transmission at

work compared to home ($R_{e\text{-}w}$). We considered eleven values of the fraction of an 8-hour workday

292   spent on campus ($T_w = y(8/24)$ where y = 0,0.1,...,0.9,1), and eleven values of the fraction increase in

transmission at work compared to at home ($R_{e\text{-}w} = (1+x)R_{e\text{-}h}$ where x = 0,0.1,...,0.9,1). These results

294   are presented in Fig 3c.

Movement model simulations were conducted in Matlab 2018b.

24   12

[**Fig 2. Movement model: varying the degree of physical distancing on campus and during**

commutes.

The fraction of the population that is Susceptible (S), Exposed (E), Infected (I), and Removed (R), when

all individuals either work from home (solid lines) or commute to work on campus (dashed lines), for

different degrees of physical distancing both on campus and during the commute: (a) low risk: low

contact during commute and on campus ($R_{e-c} = R_{e-w} = 0.5R_0 = 1.25$, equivalent to at home), dashed and

solid lines are identical, (b) moderate risk: unmitigated contact during commute ($R_{e-c} = R_0 = 2.5$, shared

transport) and partial physical distancing at work ($R_{e-w} = 0.75R_0 = 1.875$, intermediate between home and

unmitigated), (c) high risk: elevated contact during commute ($R_{e-c} = 2R_0 = 5$, crowded shared transport),

and unmitigated contact at work ($R_{e-w} = R_0 = 2.5$).]

[**Fig 3. Movement model: limiting people, time and contact on campus.**

The final epidemic size (cumulative fraction of the population infected) as a function of (a) the fraction of

an 8-hour workday spent on campus (x-axis) and the fraction of the population working on campus (y-

axis) with no physical distancing, (b) the fraction increase in transmission while at work compared to at

home (x-axis) and the fraction of the population working on campus (y-axis) with an 8-hour work day, (c)

the fraction increase in transmission while at work compared to at home (x-axis) and the fraction of an 8-

hour workday spent on campus (y-axis) with 100% of people on campus.]

Sensitivity Analysis

Finally, we performed a sensitivity analysis to determine how sensitive the two model output

metrics (final epidemic size, epidemic peak size) were to each of the model parameters, using a

combination of Latin Hypercube Sampling (LHS) and Partial Rank Correlation Coefficients (PRCC). The

LHS/PRCC sensitivity analysis is appropriate when the relationship between model output and each

model parameter is monotonic and nonlinear [40]. For our model, this relationship was monotonic for all

322 nine parameters considered (N, T_c, T_w, θ, σ, γ, $R_{e\text{-}c}$, $R_{e\text{-}h}$, and $R_{e\text{-}w}$; S1-S2 Figs). The LHS/PRCC sensitivity

analysis has two steps.

324   First, we used Latin Hypercube Sampling (LHS; [41]), a Monte Carlo approach, to generate sets

of parameter value combinations from preset ranges of parameter values. LHS has a minimum required

326 sample size (n) which is given by: $n \geq k+1$ or $n \geq k(4/3)$ where k is the number of parameters included in

the LHS [42], nine for our analysis. We chose the number of samples (see below) to meet these criteria.

328 Each of the nine model parameters considered was sampled from a uniform probability density function

based on the ranges given in Table 2. The model was run for each parameter value set, and the final

330 epidemic size (cumulative fraction of the population infected, in the long-term) and epidemic peak size

(maximum fraction of the population infected at any time) were both saved as output metrics.

332   Second, we measured the sensitivity of the output metrics to each parameter using Spearman

Partial Rank Correlation Coefficients (PRCC). To determine how many samples of each parameter was

334 needed to generate stable PRCC value, we calculated PRCC value for an increasing number of samples

(S3 Fig) and noted that the PRCC values were relatively stable past 1000 samples. Thus, we used 1000

336 samples of each parameter value for our final PRCC analysis. A positive PRCC value indicates that

increasing the value of that parameter increases the output metric while a negative PRCC value indicates

338 that increasing the value of that parameter decreases the output metric. PRCC values that were not

significant at the 0.05 level are marked with 'ns' in Fig 4 (not corrected for multiple comparisons).

340 Finally, we used a z-test to rank significant model parameters in terms of their relative importance, since

larger PRCC values do not always indicate more important parameters [40]. For our results (Fig 4), model

342 output sensitivity was indeed given by the size of PRCC values.

344 [**Fig 4. Movement model: sensitivity analysis.**

The partial rank correlation coefficient (PRCC) values for each model parameter (Table 2) for the final

346 epidemic size metric (blue bars) and the epidemic peak size metric (orange bars). Positive values indicate

parameters that increase epidemic size as they are increased (negative values indicate parameters that

348   decrease epidemic size as they are increased). Cases where the relationship between the parameter and

model output metric was not significant are indicated with 'ns'.]

Methods: network model

352   Our second model explores infection dynamics as individuals work on campus either in both

office and lab spaces or just in lab spaces. We created a network map of all the individuals housed in the

354   Ecology building on the St. Paul campus of the University of Minnesota. We created our dataset by

merging information on the office and lab room assignments for each individual with an office or lab in

356   the building. (The methods for collection and analyses of these data were reviewed by the University of

Minnesota's Institutional Review Board and were determined not to be human subjects research.) Work

358   in the Ecology building is structured by two primary space types, laboratories that can include one to

three research groups, each associated with a single faculty member, and offices which can be single-

360   occupancy or shared. Office space is generally shared by groups of graduate students and postdoctoral

scholars, often from different lab groups. Because undergraduates are generally not permitted to work on

362   campus during the resumption of research, we included faculty, staff, postdocs, and graduate students, but

excluded all undergraduates from this visualization.

364   We considered two types of bipartite networks: shared office space and shared lab space.

Individuals sharing an office or a lab all had an edge with that location node. We then consider the one-

366   mode projection of each network, creating a weighted unipartite network connecting individuals

according to their shared spaces. The binary representation of these networks was used to create static

368   network visualizations of connections among individuals using the igraph, tidygraph, and ggraph libraries

in R [43–45], shown in Fig 5a and b. Animations of disease progression through the networks were

370   produced using the gganimate library in R (S5-S6 Figs; [46]). For each network, we computed the

distribution of (finite) shortest paths between each pair of nodes (Fig 5c) and for each distinct component

372 of the networks, we noted its size (number of nodes), diameter (longest shortest path), and mean path

length (average shortest path length; S7 Fig).

[**Fig 5. Network model structure.**

376 Space-sharing, or 'contacts' (edges) are shown among all individuals (nodes) for two scenarios: (a) when

individuals at work share either office or lab spaces, or (b) when individuals only used shared lab space

378 and not shared offices (e.g., bench work is done on campus while office work is done at home). (c)

Histograms showing the distribution of shortest paths between all connected pairs of individuals.

380 Importantly, though all shortest paths between nodes in the network containing only links of shared lab

spaces are less than or equal to three, the vast majority (approximately 95%) of pairwise combinations of

382 individuals actually have no chain of interactions connecting them. In contrast, the combined network

contains a component consisting of almost 90% of individuals in the network, corresponding to nearly

384 80% of all pairs of individuals having a chain of interactions connecting them.]

386   For the network simulations, we used an SEIR model framework, starting with a randomly

selected index case to serve as the first infected individual in an entirely susceptible population (Fig 6).

388 Simulations proceeded in discrete time. At each time step, individuals who had been exposed to the virus

transitioned into the infectious class (E→I) based on the result of a Bernouli trial using the disease

390 progression rate as the probability of success. Likewise, currently infectious individuals were removed

(i.e., either recovered and immune or deceased; I→R) based on the result of a Bernouli trial using the

392 recovery rate as the probability of success. Finally, one Bernouli trial using the transmission rate as

probability of success was conducted for each edge connecting a susceptible individual to an infectious

394 one. Susceptible individuals became exposed (S→E) if at least one such trial resulted in success. At the

end of each simulation, we took note of the epidemic peak size, the final epidemic size, and the time

396 needed to read the epidemic peak (Fig 7). Network analysis and simulations were conducted in R

(Version 3.6.3).

[**Fig 6. Network model simulations.**

400   Final disease status of members of networks based on use of (a) both shared lab and office space and (c)

only shared lab space, following a simulated epidemic with susceptible individuals in blue, exposed

402   individuals in green, infectious individuals in orange, and removed individuals in red. (b, d) the

cumulative number of susceptible, infectious, and removed individuals over time for each network

404   simulation.]

406   [**Fig 7. Network model simulations.**

Outcome of 100 infection simulations on networks: the maximum peak number of individuals infected at

408   any one time (epidemic peak size), total number of individuals infected (final epidemic size), and time

until peak number of individuals infected for simulations of pathogen spread on networks based on use of

410   both shared lab and office space (blue) and only shared lab space (orange).]

Results

Movement model

416   Whether returning to work on campus affects the epidemic outcomes (measured as final epidemic

size and epidemic peak size) depends critically on the degree of physical distancing maintained both on

418   campus and during the commute between home and campus (Fig 2). If the current degree of physical

distancing that is achieved while working from home can be maintained while on campus, then working

420   from campus will not speed up infection dynamics compared to working from home (Fig 2a). However, if

physical distancing on campus or during the commute is less successful than current physical distancing

422   at home, then returning to work on campus will both increase the epidemic peak size in the short-term and

increase the final epidemic size in the long term (Fig 2b-c). When physical distancing cannot be

424   maintained on campus or during the commute, then infection dynamics can be kept slower by limiting the

fraction of workers on campus and the amount of time workers are on campus (Fig 3a).

426   Intriguingly the three strategies we considered (limiting contact, people, or time on campus) are

interchangeable with approximately equivalent effects on both the long-term metric, final epidemic size

428   (Fig 3) and the short-term metric, epidemic peak size (S4 Fig). That is, in situations where one of these

strategies cannot be fully implemented, a different strategy can be used in its stead. For example, if

430   individuals need to be on campus for an extended period of time to run an experiment (thus limiting time

on campus is not a feasible strategy), this can be compensated for by limiting the number of other

432   individuals on campus at the same time. However, of the three strategies, reducing the fraction of the

population on campus had a bigger impact than reducing either time or contact on campus, due to the

434   effect of commuting to and from campus. Regardless of time or physical distancing on campus, more

people working on campus requires that more people commute. Thus, if commuting substantially

436   increases transmission risk compared to staying at home (i.e., any form of shared transport vs. commuting

alone), reducing the number of people commuting will be a more effective strategy than reducing either

438   time or contact while on campus.

The sensitivity analysis revealed that both model metrics (final epidemic size, epidemic peak

440   size) were most sensitive to transmission at home ($R_{\text{e-h}}$), since most of the day is spent in that

environment, as well as the fraction of the population commuting (θ) to campus (Fig 4). Transmission on

442   campus ($R_{\text{e-w}}$) and transmission during commutes ($R_{\text{e-c}}$) were the next most influential; the first because

most time during the workday is spent on campus and the second because we allowed transmission to

444   vary across a wider range during commuting than on campus. The time spent on campus (T_{w}) and time

commuting (T_{c}) were somewhat influential. For each of these parameters, increasing the parameter value

446   increased the final and peak epidemic sizes. Finally, population size (N) did not significantly affect either

metric (but would be critical for the total number of individuals infected). The rate of moving from

448   exposed to infected (σ) and the recovery rate (γ) did not significantly affect final epidemic size, but both

had a minor effect on epidemic peak size: increasing σ (i.e., a shorter latent period) increased epidemic

450   peak size, while increasing γ (i.e., a shorter infectious period) decreased epidemic peak size.

Network model

The mixing of researchers from different labs in shared office spaces had a substantial impact on

454   the modularity of the resulting network. In particular, when people do not use shared office spaces (i.e.,

work from home if they share an office), but work on campus only in labs, the network is far more

456   modular, with smaller, more densely connected groups and few connections among groups (Fig 5b, S7

Fig). In this case, most individuals are directly connected to all other members of their group (i.e.,

458   "shortest path" of one, Fig 5c); however the absence of connections between groups means that, on

average, an infected individual lacks a path of connections to 95% of the rest of the population. In

460   contrast, when individuals share both lab and office space, the connectedness of the network is relatively

high because students, staff, and postdocs that share offices are often from different labs. For this

462   combined case, most individuals are four or fewer connections from one another (Fig 5c) and the largest

component contains nearly 90% of all individuals in the network (Fig 5a, S7 Fig). Thus, in the case where

464   an infected individual (presumably pre-symptomatic or asymptomatic) came to work, the combined lab

and office network has the potential for greater disease incidence than in the lab-only network, where the

466   infection could be constrained to a single lab (Figs 6-7, S5-S6 Figs). In general, when compared to the

combined network, the lab-only network had outbreaks that were less explosive (i.e., had less variance

468   and a lower mean number of individuals infected at any one time), fewer individuals infected overall, and

a shorter time until the peak number of infectious individuals (Fig 7).

Discussion

472   Movement and contact behaviors are key drivers of the spread of pathogens like SARS-CoV-2,

and not all movement and contacts have the same impact on pathogen spread. However, basic

474 compartmental models used to describe SARS-CoV-2 dynamics assume all individuals move and contact

each other at random (i.e., populations are well-mixed). Our models show how explicitly accounting for

476 movement, space use in a building, and contact behaviors can provide a more nuanced understanding of

relative risk. Our movement model, capturing the predictable movement between home and work/campus

478 environments, shows that reducing the number of people, rate of contact, and amount of time spent on

campus are all equivalently effective strategies for slowing pathogen spread. However, if commutes

480 specifically increase transmission risk (i.e., shared transport), reducing the number of people on campus is

the most effective strategy to reduce the infection spread rate. We also considered heterogeneity in

482 contact behavior once at the workplace; our network model captures the regular interactions among

workers in shared workspaces on campus and shows that restricting building use to lab spaces (rather than

484 lab and office space) may reduce pathogen spread. Our results provide a number of tools to distinguish

among different movement and contact patterns at the scale of individuals and workplace communities.

486   A number of future directions could be explored, by changing some of our simplifying

assumptions. First, staying within the broad structure of our model, alternative spatiotemporal strategies

488 could be explored including: structured work weeks (e.g., four days on-campus and ten off; [47]), or

further compartmentalizing time (e.g., sequential work shifts) or space (e.g., different buildings on

490 campus). For instance, if evidence suggests that infection can occur through air circulation within

buildings [48], these models could be altered to account for connections arising from shared ventilation

492 systems. These models also could be modified to account for movement and contact behavior that

explicitly depends on infection status [49]; e.g., splitting infected individuals into infectious but

494 asymptomatic (who still potentially commute to work) and symptomatic individuals (who stay home) or

building dynamic networks where contact behavior can change in response to infectious status (e.g.

496 infectious and symptomatic individuals reducing their interpersonal contacts either through staying home

or altering their behavior at work). Second, one could expand the scale of the model. This could be done

498 foremost by combining the movement model (movement between work, commute, campus) with the

network model (movement while on campus). Further expansions could consider both larger scales

500   (linking in regional patterns) as well as smaller ones (allowing contacts within buildings to vary over

time). For instance, structuring interaction networks with respect to age can greatly reduce deaths [50]

502   and integrating local models such as this with regional variation in disease [51] could inform

recommendations as rates of active infection and immunity shift. Third, as contact-tracing data

504   accumulate that implicate the most common routes of transmission, we can alter specific players or

interactions in the model. For instance, while the virus can survive on surfaces and in aerosols [52], most

506   transmission appears to be mediated by extended person-to-person interactions in close spaces [53–57],

so minimizing temporal overlap of workers in shared spaces is key [58]. It may also be possible to further

508   refine risk for certain classes of risk based on age, immune status, and pre-existing conditions; for

example, current data suggests children can be carriers for COVID-19 [56,59], although the role of

510   children and school closures in mitigating COVID-19 spread is a topic of current debate [14,16,60]. As

such data emerge, the model could be updated with additional data for individual members that could

512   indirectly increase transmission, such as children or elderly family members at home [61].

Our findings mesh with concepts in the broader movement and disease ecology literature. Within

514   movement ecology, there has long been a distinction between random/undirected movement like dispersal

versus predictable movements like diel and seasonal migration [62]. Human movement between home

516   and work is often a predictable and daily occurrence and thus is better viewed from the lens of predictable

migratory movements (as we do here) rather than random dispersive ones (as implicit in basic

518   compartmental models). Moving predictably between two environments does not always increase

infection (either for individuals or at the population level) compared to remaining in a single location; the

520   relative transmission in each environment is critical [20]. We find that transmission risk during a

commute is key to infection dynamics when considering the impact of movement between home and

522   work, paralleling recent work calling for the explicit consideration of how transient phases of movement

affect infection dynamics [22] and theory showing that infection dynamics during transit can have a

524   similar impact to dynamics in the second environment [63].

There are important insights that emerge from our movement and contact-network models that

526   can guide policy. For example, basic disease models assume random movement and equal probability of

contact, whereas many hosts, including humans, move in directed ways and in very structured social

528   networks. For this reason, disease mitigation policies will likely be more effective when they consider

disease risk in a more holistic way that integrates risk across the various components of a person's daily

530   movement. For example, in settings where many people commute by mass transit (e.g., New York City),

the efficacy of workplace safety protocols may be overwhelmed by transmission during daily commutes

532   rather than contacts at work. Careful examination of social network patterns could also help guide policy

to provide intermediate scenarios between business as usual and complete lock down. For example, in our

534   case-study contact rates and potential disease spread were significantly reduced when people's contacts at

the workplace were restricted to single lab groups, as opposed to linking separate lab and office networks.

536   Even so, the protective effects of heterogeneity in contact structure should not be overemphasized

for decision making. First, outbreaks on more heterogeneous networks may be less likely, but they can

538   also be more explosive [23]. Second, because SARS-CoV-2 spread appears to be primarily by

aerosolized transmission, the potential contact behaviors needed for transmission are more ubiquitous

540   than for pathogens with more specific transmission modes (e.g., sexually transmitted diseases like

HIV/AIDS). Importantly, the networks presented here consider only the room in which an employee

542   works (their office or lab space), explicitly omitting broader workplace considerations like air flow,

shared surfaces, entry points, etc., these additional points must be addressed in conjunction with thinking

544   about explicit contact behavior when forming a public health strategy. Lastly, these static networks are a

simplification of an inherently dynamic process of movement, contact, and infection. Using a time-

546   ordered or dynamic network approach could provide better insights to actual duration of exposures and

sickness-induced behavioral changes [64].

Conclusions

550   Human movement and contact behaviors are critical for the spread of pathogens like SARS-

CoV-2, yet are rarely addressed explicitly in the current conversations about decision-making in the face

552 of relaxing Stay at Home orders. Here we have drawn on movement and network models to demonstrate

the effect of these behaviors. First, we have shown that regular movement between two 'environments'

554 (i.e., work and home) does not inherently increase infection spread the way random dispersive

movements might. Rather the outcome depends on the relative degree of transmission (e.g., degree of

556 physical distancing) in each environment. Second, we have shown that different contact patterns (e.g.,

space usage) within the work environment could lead to different outcomes in terms of SARS-CoV-2

558 spread. In sum, we advocate for using an understanding of movement and contact patterns as an

adjunctive approach (alongside widespread testing, contact tracing, vaccine development and other tools)

560 to mitigate the effects of SARS-CoV-2 and COVID-19, particularly when considering return to work

environments.

Funding

564   This material is based in part upon work supported by the University of Minnesota's Office of

Academic Clinical Affairs COVID-19 Rapid Response Grant (https://clinicalaffairs.umn.edu/umn-covid-

566 19-research) (to MEC, LAW, and MMS) and by the National Science Foundation (https://www.nsf.gov/)

under Grants DEB-2030509 (to MEC and MMS), DEB-1654609 (to AKS and MEC) and DEB-1556649

568 (to EWS and ETB). The funders had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript.

Acknowledgements

572   We thank William Harcombe for helpful feedback and discussion, and Valery Forbes, David

Greenstein and Daniel Stanton for encouragement.

Data Availability

576   Model code and simulation output for the movement model will be made available on Data

Dryad. All Code to generate the network model figures and animations will be made publicly available on

578   Github.

References

1. Kraemer MUG, Yang C-H, Gutierrez B, Wu C-H, Klein B, Pigott DM, et al. The effect of human mobility and control measures on the COVID-19 epidemic in China. Science. 2020;368: 493–497. doi:10.1126/science.abb4218

2. Keeling MJ, Rohani P. Modeling Infectious Diseases in Humans and Animals. Princeton, NJ: Princeton University Press; 2008.

3. Keeling MJ, Grenfell BT. Individual-based Perspectives on R0. J Theor Biol. 2000;203: 51–61. doi:10.1006/jtbi.1999.1064

4. Hébert-Dufresne L, Althouse BM, Scarpino SV, Allard A. Beyond R0: Heterogeneity in secondary infections and probabilistic epidemic forecasting. ArXiv200204004 Phys Q-Bio. 2020 [cited 7 May 2020]. Available: http://arxiv.org/abs/2002.04004

5. Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. Eurosurveillance. 2020;25. doi:10.2807/1560-7917.ES.2020.25.10.2000180

6. Wu JT, Leung K, Bushman M, Kishore N, Niehus R, de Salazar PM, et al. Estimating clinical severity of COVID-19 from the transmission dynamics in Wuhan, China. Nat Med. 2020;26: 506–510. doi:10.1038/s41591-020-0822-7

7. Moghadas SM, Shoukat A, Fitzpatrick MC, Wells CR, Sah P, Pandey A, et al. Projecting hospital utilization during the COVID-19 outbreaks in the United States. Proc Natl Acad Sci. 2020;117: 9122–9126. doi:10.1073/pnas.2004064117

8. IHME COVID-19 health service utilization forecasting team, Murray CJ. Forecasting COVID-19

impact on hospital bed-days, ICU-days, ventilator-days and deaths by US state in the next 4 months. medRxiv. 2020 [cited 7 May 2020]. doi:10.1101/2020.03.27.20043752

9.   Lewnard JA, Lo NC. Scientific and ethical basis for social-distancing interventions against COVID-19. Lancet Infect Dis. 2020; S1473309920301900. doi:10.1016/S1473-3099(20)30190-0

10. Balilla J. Assessment of COVID-19 Mass Testing: The Case of South Korea. SSRN Electron J. 2020 [cited 3 May 2020]. doi:10.2139/ssrn.3556346

11. Koo JR, Cook AR, Park M, Sun Y, Sun H, Lim JT, et al. Interventions to mitigate early spread of SARS-CoV-2 in Singapore: a modelling study. Lancet Infect Dis. 2020; S1473309920301626. doi:10.1016/S1473-3099(20)30162-6

12. Clarke S. The traffic data that shows the road into - and out of - Covid-19 lockdown. The Guardian. 27 Apr 2020: 1.

13. Klein B, LaRock T, McCabe S, Torres L, Privitera F, Lake B, et al. Assessing changes in commuting and individual mobility in major metropolitan areas in the United States during the COVID-19 outbreak. 2020;https://uploads-ssl.webflow.com/5c9104426f6f88ac129ef3d2/5e8374ee75221201609ab586_Assessing_mobility_changes_in_the_United_States_during_the_COVID_19_outbreak.pdf: 29.

14. Zhang J, Litvinova M, Liang Y, Wang Y, Wang W, Zhao S, et al. Changes in contact patterns shape the dynamics of the COVID-19 outbreak in China. Science. 2020; eabb8001. doi:10.1126/science.abb8001

15. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. The Lancet. 2020;395: 912–920. doi:10.1016/S0140-6736(20)30460-8

16. Viner RM, Russell SJ, Croker H, Packer J, Ward J, Stansfield C, et al. School closure and management practices during coronavirus outbreaks including COVID-19: a rapid systematic review. Lancet Child Adolesc Health. 2020;4: 397–404. doi:10.1016/S2352-4642(20)30095-X

17. Binning SA, Shaw AK, Roche DG. Parasites and host performance: Incorporating infection into our understanding of animal movement. Integr Comp Biol. 2017;57: 267–280. doi:10.1093/icb/icx024

18. Loehle C. Social barriers to pathogen transmission in wild animal populations. Ecology. 1995;76: 326–335. doi:10.2307/1941192

19. Shaw AK, Binning SA. Recovery from infection is more likely to favour the evolution of migration than social escape from infection. J Anim Ecol. 2020;89: 1448–1457. doi:10.1111/1365-2656.13195

20. Shaw AK, Sherman J, Barker FK, Zuk M. Metrics matter: the effect of parasite richness, intensity and prevalence on the evolution of host migration. Proc R Soc B Biol Sci. 2018;285: 20182147. doi:10.1098/rspb.2018.2147

21. Boulinier T, Kada S, Ponchon A, Dupraz M, Dietrich M, Gamble A, et al. Migration, prospecting, dispersal? What host movement matters for infectious agent circulation? Integr Comp Biol. 2016;56: 330–342. doi:10.1093/icb/icw015

22. Daversa DR, Fenton A, Dell AI, Garner TWJ, Manica A. Infections on the move: how transient phases of host movement influence disease spread. Proc R Soc B Biol Sci. 2017;284: 20171807. doi:10.1098/rspb.2017.1807

23. Keeling MJ, Eames KT. Networks and epidemic models. J R Soc Interface. 2005;2: 295–307. doi:10.1098/rsif.2005.0051

24. Altieri N, Barter R, Duncan J, Dwivedi R, Kumbier K, Li X, et al. Curating a COVID-19 data repository and forecasting county-level death counts in the United States.

2020;https://bids.berkeley.edu/publications/curating-covid-19-data-repository-and-forecasting-county-level-death-counts-united: 18.

25. Woody S, Garcia Tec M, Dahan M, Gaither K, Lachmann M, Fox SJ, et al. Projections for first-wave COVID-19 deaths across the US using social-distancing measures derived from mobile phones. Infectious Diseases (except HIV/AIDS); 2020 Apr. doi:10.1101/2020.04.16.20068163

26. Kissler SM, Tedijanto C, Goldstein E, Grad YH, Lipsitch M. Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period. Science. 2020; eabb5793. doi:10.1126/science.abb5793

27. Chinazzi M, Davis JT, Ajelli M, Gioannini C, Litvinova M, Merler S, et al. The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. Science. 2020; eaba9757. doi:10.1126/science.aba9757

28. Kucharski AJ, Russell TW, Diamond C, Liu Y, Edmunds J, Funk S, et al. Early dynamics of transmission and control of COVID-19: a mathematical modelling study. Lancet Infect Dis. 2020;20: 553–558. doi:10.1016/S1473-3099(20)30144-4

29. Prem K, Liu Y, Russell TW, Kucharski AJ, Eggo RM, Davies N, et al. The effect of control strategies to reduce social mixing on outcomes of the COVID-19 epidemic in Wuhan, China: a modelling study. Lancet Public Health. 2020;5: e261–e270. doi:10.1016/S2468-2667(20)30073-6

30. IHME COVID-19 health service utilization forecasting team, Murray CJ. Forecasting the impact of the first wave of the COVID-19 pandemic on hospital demand and deaths for the USA and European Economic Area countries. medRxiv. 2020 [cited 9 May 2020]. doi:10.1101/2020.04.21.20074732

31. MN Department of Health. Situation Update for Coronavirus Disease 2019 (COVID-19). 2020.

Available: https://www.health.state.mn.us/diseases/coronavirus/situation.html#cases1

32. Wei WE, Li Z, Chiew CJ, Yong SE, Toh MP, Lee VJ. Presymptomatic transmission of SARS-CoV-2 — Singapore, January 23–March 16, 2020. MMWR Morb Mortal Wkly Rep. 2020;69: 411–415. doi:10.15585/mmwr.mm6914e1

33. Servedio MR, Brandvain Y, Dhole S, Fitzpatrick CL, Goldberg EE, Stern CA, et al. Not just a theory—The utility of mathematical models in evolutionary biology. PLoS Biol. 2014;12: e1002017.

34. Johns S, Shaw AK. Theoretical insight into three disease-related benefits of migration. Popul Ecol. 2016;58: 213–221. doi:10.1007/s10144-015-0518-x

35. Begon M, Bennett M, Bowers RG, French NP, Hazel SM, Turner J. A clarification of transmission terms in host-microparasite models: numbers, densities and areas. Epidemiol Infect. 2002;129: 147–153. doi:10.1017/S0950268802007148

36. Sanche S, Lin YT, Xu C, Romero-Severson E, Hengartner N, Ke R. High Contagiousness and Rapid Spread of Severe Acute Respiratory Syndrome Coronavirus 2. Emerg Infect Dis. 2020;26. doi:10.3201/eid2607.200282

37. Ling Y, Xu S-B, Lin Y-X, Tian D, Zhu Z-Q, Dai F-H, et al. Persistence and clearance of viral RNA in 2019 novel coronavirus disease rehabilitation patients: Chin Med J (Engl). 2020;133: 1039–1043. doi:10.1097/CM9.0000000000000774

38. Zhang S, Diao M, Yu W, Pei L, Lin Z, Chen D. Estimation of the reproductive number of novel coronavirus (COVID-19) and the probable outbreak size on the Diamond Princess cruise ship: A data-driven analysis. Int J Infect Dis. 2020;93: 201–204. doi:10.1016/j.ijid.2020.02.033

39. Enns EA, Kirkeide M, Mehta A, MacLehose R, Knowlton GS, Smith MK, et al. Modeling the

impact of social distancing measures on the spread of SARS-CoV-2 in Minnesota. MN Department
of Health; 2020 Apr p. 16. Available: Enns EA, Kirkeide M, Mehta A, MacLehose R, Knowlton
GS, Smith https://mn.gov/covid19/assets/MNmodel_tech_doc_tcm1148-427724.pdf

40. Marino S, Hogue IB, Ray CJ, Kirschner DE. A methodology for performing global uncertainty and
     sensitivity analysis in systems biology. J Theor Biol. 2008;254: 178–196.
     doi:10.1016/j.jtbi.2008.04.011

41. Mckay MD, Beckman RJ, Conover WJ. A comparison of three methods for selecting values of
     input variables in the analysis of output from a computer code. Technometrics. 1979;42: 55–61.
     doi:10.1080/00401706.2000.10485979

42. Blower SM, Dowlatabadi H. Sensitivity and Uncertainty Analysis of Complex Models of Disease
     Transmission: An HIV Model, as an Example. Int Stat Rev Rev Int Stat. 1994;62: 229.
     doi:10.2307/1403510

43. Csárdi G, Nepusz T. The igraph software package for complex network research. InterJournal
     Complex Syst. 2006;1695: 9.

44. Pedersen TL. tidygraph: A Tidy API for Graph Manipulation. 2019. Available: https://CRAN.R-
     project.org/package=tidygraph

45. Pedersen TL. ggraph: An Implementation of Grammar of Graphics for Graphs and Networks. 2020.
     Available: https://CRAN.R-project.org/package=ggraph

46. Pedersen TL, Robinson D. gganimate: A Grammar of Animated Graphics. 2020. Available: https://
     CRAN.R-project.org/package=gganimate

47. Karin O, Bar-On YM, Milo T, Katzir I, Mayo A, Korem Y, et al. Cyclic exit strategies to suppress
     COVID-19 and allow economic activity. medRxiv. 2020 [cited 13 May 2020].

doi:10.1101/2020.04.04.20053579

48. Lu J, Gu J, Li K, Xu C, Su W, Lai Z, et al. COVID-19 Outbreak Associated with Air Conditioning in Restaurant, Guangzhou, China, 2020. Emerg Infect Dis. 2020;26. doi:10.3201/eid2607.200764

49. Narayanan N, Binning SA, Shaw AK. Infection state can affect host migratory decisions. Oikos. in press.

50. Acemoglu D, Chernozhukov V, Werning I, Whinston MD. A multi-risk SIR model with optimally targeted lockdown. National Bureau of Economic Research; 2020 May p. 38. Available: http://www.nber.org/papers/w27102

51. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. Lancet Infect Dis. 2020;20: 533–534. doi:10.1016/S1473-3099(20)30120-1

52. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. N Engl J Med. 2020;382: 1564–1567. doi:10.1056/NEJMc2004973

53. Ghinai I, Woods S, Ritger KA, McPherson TD, Black SR, Sparrow L, et al. Community Transmission of SARS-CoV-2 at Two Family Gatherings — Chicago, Illinois, February–March 2020. MMWR Morb Mortal Wkly Rep. 2020;69: 446–450. doi:10.15585/mmwr.mm6915e1

54. Ghinai I, McPherson TD, Hunter JC, Kirking HL, Christiansen D, Joshi K, et al. First known person-to-person transmission of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in the USA. The Lancet. 2020;395: 1137–1144. doi:10.1016/S0140-6736(20)30607-3

55. Li W, Zhang B, Lu J, Liu S, Chang Z, Cao P, et al. The characteristics of household transmission of COVID-19. Clin Infect Dis. 2020; ciaa450. doi:10.1093/cid/ciaa450

56. Bi Q, Wu Y, Mei S, Ye C, Zou X, Zhang Z, et al. Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study. Lancet Infect Dis. 2020; S1473309920302875. doi:10.1016/S1473-3099(20)30287-5

57. Park SY, Kim YM, Yi S, Lee S, Na BJ, Kim CB, et al. Coronavirus disease outbreak in call center, South Korea. Emerg Infect Dis. 2020;26. doi:10.3201/eid2608.201274

58. Qian H, Miao T, Liu L, Zheng X, Luo D, Li Y. Indoor transmission of SARS-CoV-2. medRxiv. 2020 [cited 11 May 2020]. doi:10.1101/2020.04.04.20053058

59. Jones TC, Mühlemann B, Veith T, Zuchowski M, Hofmann J, Stein A, et al. An analysis of SARS-CoV-2 viral load by patient age. 2020;https://zoonosen.charite.de/fileadmin/user_upload/microsites/m_cc05/virologie-ccm/dateien_upload/Weitere_Dateien/analysis-of-SARS-CoV-2-viral-load-by-patient-age.pdf: 19.

60. Zhu Y, Bloxham CJ, Hulme KD, Sinclair JE, Tong ZWM, Steele LE, et al. Children are unlikely to have been the primary source of household SARS-CoV-2 infections. medRxiv. 2020. doi:https://doi.org/10.1101/2020.03.26.20044826

61. Chen Y, Wang A, Yi B, Ding K, Wang H, Wang J, et al. The epidemiological characteristics of infection in close contacts of COVID-19 in Ningbo city. Chin J Epidemiol. 2020;41: 668–672. doi:10.3760/cma.j.cn112338-20200304-00251

62. Heape W. Emigration, Migration and Nomadism. Cambridge, UK: Heffer and Sons; 1931.

63. Shaw AK, Craft ME, Zuk M, Binning SA. Host migration strategy is shaped by forms of parasite transmission and infection cost. J Anim Ecol. 2019;88: 1601–1612. doi:10.1111/1365-2656.13050

64. Enright J, Kao RR. Epidemics on dynamic networks. Epidemics. 2018;24: 88–97. doi:10.1016/j.epidem.2018.04.003

	Table 1. Movement model state variables and their meaning.

Variable	Meaning
S	Total number of susceptible individuals
E	Total number of exposed individuals
I	Total number of infected individuals
R	Total number of removed individuals
S_c	Number of susceptible individuals commuting to campus (during the commute phases)
E_c	Number of exposed individuals commuting to campus (during the commute phases)
I_c	Number of infected individuals commuting to campus (during the commute phases)
R_c	Number of removed individuals commuting to campus (during the commute phases)
S_w	Number of susceptible individuals working from work (during the work phase)
E_w	Number of exposed individuals working from work (during the work phase)
I_w	Number of infected individuals working from work (during the work phase)
R_w	Number of removed individuals working from work (during the work phase)
S_h	Number of susceptible individuals working from home (during the commute and work phases)
E_h	Number of exposed individuals working from home (during the commute and work phases)
I_h	Number of infected individuals working from home (during the commute and work phases)
R_h	Number of recovered individuals working from home (during the commute and work phases)

582 Table 2. Movement model parameters, meaning, and default value (with units). References:

[a] https://en.wikipedia.org/wiki/University_of_Minnesota_College_of_Biological_Sciences

584 [b] Zhang et al. 2020 IJID

[c] https://mn.gov/covid19/assets/MNmodel_tech_doc_tcm1148-427724.pdf

586 [d] Sanche et al. 2020

[e] Ling et al. 2020

Parameter	Meaning	Default values [Units]	Sensitivity analysis range
N	Population size	3000 [people] [a]	(1500 to 6000)
R_0	Basic reproductive number (number of new infections that each infection generates)	2.5 [unitless] [b]	fixed
R_{e-c}	Effective reproductive number while commuting between work and campus	R_0 [unitless]	(1 to 4 R_0)
R_{e-h}	Effective reproductive number while at home	$0.5R_0$ [unitless] [c]	(0.25 R_0 to R_0)
R_{e-w}	Effective reproductive number while at work at campus	R_0 [unitless]	(0.5 R_0 to 2 R_0)
T_c	Fraction of a 24-hour day spent commuting each way for those that commute to campus	1/24 [unitless]	(0.5/24 to 2/24)
T_h	Fraction of a 24-hour day spent not working (everyone is off campus)	= 1 - 2 T_c - T_w [unitless]	= 1 - 2 T_c - T_w
T_w	Fraction of a 24-hour day spent at work on campus for those commuting (some individual are on campus)	8/24 [unitless]	(2/24 to 12/24)
β_c	Transmission rate while commuting	= γ R_{e-c} [day^{-1}]	= γ R_{e-c}
β_h	Transmission rate while at home	= γ R_{e-h} [day^{-1}]	= γ R_{e-h}

β_w	Transmission rate while at work	$= \gamma\, R_{e\text{-}w}$ [day^{-1}]	$= \gamma\, R_{e\text{-}w}$
σ	Rate of moving from exposed to infected (inverse of the latent period)	1/4.2 [day^{-1}] [d]	(1/5.1 to 1/3.5)
γ	Recovery rate	1/9.5 [day^{-1}] [e]	(1/11 to 1/6)
θ	Fraction of the campus population commuting to work on campus (instead of continuing to work at home)	1 [unitless]	(0.0001 to 1)

 SUPPORTING INFORMATION FIGURE CAPTIONS

592 **S1 Fig. Movement model monotonicity plots.**

The relationship between each of the nine model parameters (x-axis) and the model output, final epidemic

594 size (y-axis) for (a) population size (N); (b) fraction of a 24-hour day spent commuting each way for those

that commute to campus (T_c); (c) fraction of a 24-hour day spent on campus for those commuting (T_w); (d)

596 fraction of the campus population commuting to work on campus (θ); (e) recovery rate (γ); (f) effective

reproductive number while at home (R_{e-h}); (g) effective reproductive number while commuting between

598 work and campus (R_{e-c}); and (h) effective reproductive number while at work on campus (R_{e-w}).

600 **S2 Fig. Movement model monotonicity plots.**

The relationship between each of the nine model parameters (x-axis) and the model output, epidemic peak

602 size (y-axis) for (a) population size (N); (b) fraction of a 24-hour day spent commuting each way for those

that commute to campus (T_c); (c) fraction of a 24-hour day spent on campus for those commuting (T_w); (d)

604 fraction of the campus population commuting to work on campus (θ); (e) recovery rate (γ); (F) effective

reproductive number while at home (R_{e-h}); (g) effective reproductive number while commuting between

606 work and campus (R_{e-c}); and (h) effective reproductive number while at work on campus (R_{e-w}).

608 **S3 Fig. Movement model sample numbers**.

Absolute value of PRCC for the final epidemic size model output and each of the nine model parameters

610 (N, T_c, T_w, θ, σ, γ, R_{e-h}, R_{e-c}, R_{e-w}) as a function of different numbers of LHS samples generated. The results

seem to stabilize after about 1000 samples.

S4 Fig. Movement model: limiting people, time and contact on campus.

614 The epidemic peak size (maximum fraction of the population infected) as a function of (a) the fraction of

an 8-hour workday spent on campus (x-axis) and the fraction of the population working on campus (y-

616   axis) with no physical distancing, (b) the fraction increase in transmission while at work compared to at

home (x-axis) and the fraction of the population working on campus (y-axis) with an 8-hour work day, (c)

618   the fraction increase in transmission while at work compared to at home (x-axis) and the fraction of an 8-

hour workday spent on campus (y-axis) with 100% of people on campus.

S5 Fig. Network model simulations.

622   Simulations of pathogen spread across networks based on use of both shared office and lab space.

624   **S6 Fig. Network model simulations.**

Simulations of pathogen spread across networks based on use of only shared lab space.

S7 Fig. Component-wise network structural metrics.

628   Measures of the size (number of individuals), diameter (longest shortest path between two individuals),

and mean path length (average shortest path length between individuals) for each distinct component of

630   networks presented in Fig 5a,b. The combined lab and office network (blue points) has 8 distinct

components (8 points for each metric), while the shared lab space network contains 31 distinct

632   components (31 points for each metric).

a)

Final epidemic size

b)

Final epidemic size

c)

Final epidemic size

A) Combined Lab and Office
8 distinct components

B) Just Shared Labspace
31 distinct components

C) Distribution of shortest paths

Susceptible ■ Exposed ■ Infectious ■ Removed

Click here to access/download
Supporting Information
figS1.tiff

Click here to access/download

Supporting Information
figS2.tiff

Click here to access/download
Supporting Information
figS3.tiff

Click here to access/download

Supporting Information

figS4.tiff

Click here to access/download

Supporting Information

figS5.gif

Click here to access/download
Supporting Information
figS6.gif

Supporting Information Fig S7

Click here to access/download
Supporting Information
figS7.tiff