Title

2   Lessons from movement ecology for the return to work: modeling contacts and the spread of COVID-19

Authors

6   Allison K. Shaw[1*], Lauren A. White[2], Matthew Michalska-Smith[3,4], Elizabeth T. Borer[1], Meggan E.

Craft[3], Eric W. Seabloom[1], Emilie Snell-Rood[1], Michael Travisano[1,5]

1. Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, Minnesota, USA

10   2. National Socio-Environmental Synthesis Center, Annapolis, Maryland, USA

3. Department of Veterinary Population Medicine, University of Minnesota, St. Paul, Minnesota, USA

12   4. Department of Plant Pathology, University of Minnesota, St. Paul, Minnesota, USA

5. BioTechnology Institute, University of Minnesota, St. Paul, Minnesota, USA

* corresponding author: ashaw@umn.edu

Abstract

Human behavior (movement, social contacts) plays a central role in the spread of pathogens like SARS-CoV-2. The rapid spread of SARS-CoV-2 was driven by global human movement, and initial lockdown measures aimed to localize movement and contact in order to slow spread. Thus, movement and contact patterns need to be explicitly considered when making reopening decisions, especially regarding return to work. Here, as a case study, we consider the initial stages of resuming research at a large research university, using approaches from movement ecology and contact network epidemiology. First, we develop a dynamical pathogen model describing movement between home and work; we show that limiting social contact, via reduced people or reduced time in the workplace are fairly equivalent strategies to slow pathogen spread. Second, we develop a model based on spatial contact patterns within a specific office and lab building on campus; we show that restricting on-campus activities to labs (rather than labs and offices) could dramatically alter (modularize) contact network structure and thus, potentially reduce pathogen spread by providing a workplace mechanism to reduce contact. Here we argue that explicitly accounting for human movement and contact behavior in the workplace can provide additional strategies to slow pathogen spread that can be used in conjunction with ongoing public health efforts.

Keywords

contact network epidemiology; disease ecology; infection; mathematical model; network; SARS-CoV-2; theoretical ecology; theory; transmission

Introduction

The cosmopolitan connectivity of modern society facilitated the rapid spread of SARS-CoV-2 around the globe in early 2020 [1]. The rate at which any pathogen spreads depends critically on host movement behavior [2]. Indeed, estimates of key epidemiological parameters like the basic reproduction number (R_0) are highly variable in part because they are context-specific and are a function of behaviors like movement and heterogenous contact structure [3,4]. Although most cases of COVID-19 (the disease caused by SARS-CoV-2) seem to be mild or even asymptomatic [5,6], the sheer number of cases to date means that limited personnel, hospital beds, and ICU equipment can be rapidly overwhelmed, increasing mortality [7,8]. Thus, continuing normal movement patterns, unmitigated, is not a viable containment strategy. Without a vaccine or widespread immunity to SARS-CoV-2, our best defense to slow pathogen spread has been restricting movement and contacts through physical distancing [9], testing for SARS-CoV-2 when available [10] and contact tracing [11]. Lockdown measures have drastically reduced human movement [1,12] and consequently have reduced the effective reproduction number, R_e [4,13]. However, such measures are affecting mental health [14,15] and have had a devastating impact on the economy, so individual regions are considering best practices for the reopening of businesses, schools, and other places where people gather (e.g.,[16–18]). Decisions regarding next steps can be informed by recognizing that not all movement patterns nor all contact behaviors are equal in terms of pathogen spread.

Concepts from movement ecology and contact network epidemiology can provide helpful frameworks for understanding the nuanced interactions between movement, contacts and infection. Increased movement does not always mean increased transmission risk [19]; for example, movement that either takes individuals away from infected areas or reduces contact with infected conspecifics can reduce transmission risk (migratory escape; [20,21]). Increased movement can even increase some aspects of infection risk while decreasing others, simultaneously [22]. Thus, explicitly considering how movement relates to transmission can help us understand what effect different movement patterns have on infection

60 dynamics [23,24]. Similarly, from disease ecology and contact network epidemiology, we know that

structured contacts among individuals in a population have different effects on disease spread than

62 random contacts. For example, long-range connections in otherwise locally-connected small world

networks can have dramatic effects on disease spread at a population level [25].

64   Individual movement across multiple scales — from occasional global movements to smaller-

scale daily patterns — is critical for shaping contact and thus the spread of pathogens. To date, models of

66 SARS-CoV-2/COVID-19 spread have focused on comparing patterns of spread across countries, states,

and counties [26,27]. Indeed, a plethora of epidemiological models have proven useful in generating

68 recommendations for reducing the virus spread rate, from understanding the role of contact-tracing and

society-wide physical distancing [11,28], to travel restrictions and lockdowns (e.g., [4,29,30]), to mask-

70 wearing [31]. However, few models offer guidance at scales as fine as individual workplaces, despite the

fact that this local scale is where individual decisions are made and where most transmission occurs.

72 Furthermore, apart from time at home, the most predictable component of many people's days is time

spent in the workplace. Thus, knowledge of work commute patterns, contact networks of individuals in

74 the workplace, and related workplace-specific factors could help mitigate pathogen spread during the

period that total population immunity remains low. In many cases, commute trajectories are not random

76 but involve regularity in timing, location, and encounters with other individuals along the way (e.g., on

public transport). Here, we consider the implications for mitigating COVID-19 transmission using a case

78 study of the initial stages of resuming research at a large research university.

        Implicit in this analysis is that COVID-19 is currently spreading in local communities around the

80 world, and every individual in a workplace is part of a home community. Even under many weeks of

extreme restrictions with only society's most essential employees present in workplaces (i.e., Stay at

82 Home orders), the number of new cases have continued to rise in most locations. For example, in late

April 2020, even after three weeks of a Stay at Home order and extreme physical distancing in Minnesota,

84 a state with moderate spread and commendable compliance with the order, the number of new cases

confirmed each day had tripled [32]. With community spread of this pathogen, it is unrealistic to expect

86   zero workplace infection or widespread virus containment primarily through workplace practices. Any

return-to-work plan, therefore, must include the explicit expectation that new infections may arise while

88   concurrently prioritizing worker safety and optimizing the work that can be done. Thus, reopening

businesses requires an evidence-based plan to reduce contacts through time to minimize new infections at

90   the workplace, when an infected individual, presumably pre-symptomatic [33], brings the virus to a

workplace.

92   Here we develop a pair of models to understand how movement and contact structure shape

infection spread. As a case study, we consider the context of moving from full-time work at home to part-

94   time resumption of research at a university; however, results from this model are general to many other

settings as well. We take a dual modeling approach by developing a general movement model and a

96   network case study of one academic laboratory and office building. We explore tradeoffs between

limiting contact, people, or time on campus. We find that moving back to work on campus does not

98   necessarily speed up infection spread, and depends particularly on the infection risk associated with

commutes and how well physical distancing can be maintained on campus. Thus our findings allow us to

100 set evidence-based expectations and generate specific behavioral recommendations for a safer return to

work.

Materials and Methods

104   We develop two models: a movement model to explore movement between home and work

environments, and a network model to explore contact patterns within the work environment. Both

106 models are SEIR, tracking susceptible (S), exposed (E), infected (I), and removed (recovered and

immune, or deceased; R) individuals. We assume there is no loss of immunity (removed individuals never

108 move back to the susceptible class) over the short time scales we consider, and we assume a closed

population (no births, deaths, immigration, or emigration).

Methods: movement model

Setup

114   Our first model explores infection dynamics as individuals move between home, commuting, and

work environments. Workplaces, including universities, face a number of different decisions about how to

116 slowly ramp up work following easing of lockdown. Here, we simulate three potential strategies for

returning to work: (i) allowing people to return while maintaining physical distancing, (ii) limiting the

118 number of people returning to campus during the work day, and (iii) limiting the time each person spends

on campus. For each strategy combination, we simulate infection dynamics and quantify two output

120 metrics: (1) the 'final epidemic size' (cumulative fraction of the population infected, at equilibrium), and

(2) the 'epidemic peak size' (maximum fraction of the population infected during the outbreak). The aim

122 of this type of conceptual model is to clarify the connections between assumptions and outcomes, unlike

predictive models which would contain an abundance of empirical data and aim to generate forecasts for a

124 specific system [34].

Daily cycle

Our model dynamics have a combination of continuous and discrete time (e.g., [35]), where each

128 day is broken into discrete phases (T_h spent at home, T_w spent at work, and T_c spent commuting each way,

with $T_h + T_w + 2T_c = 1$) and infection dynamics occur continuously during each phase (Fig 1, see Tables

130 1-2 for model variables and parameters). All individuals start at home and spend a fraction of their day (of

length T_h) there and not working. During this time, the infection dynamics are given by

132                          $$dS/dt = - \beta_h \, S \, (I/N)$$

$$dE/dt = \beta_h \, S \, (I/N) - \sigma \, E \qquad\qquad \text{[eqn. 1]}$$

$$dI/dt = \sigma \, E - \gamma \, I$$

$$dR/dt = \gamma \, I$$

136 where S is the number of susceptible individuals, E is the number of exposed individuals, I is the number

of infected individuals, R is the number of removed (recovered and immune, or deceased) individuals, N

138 is the total number of individuals in the population ($N = S + E + I + R$), β_h is the rate of transmission while

at home, σ is the rate of moving from exposed to infected (inverse of the latent period) and γ is the rate of

140 recovery from infection.

142 [**Fig 1. Movement model schematic, showing a daily cycle.**

All individuals — Susceptible (S), Exposed (E), Infected (I) or Removed (R) — spend part of their day

144 (T_h) at home. A proportion θ of individuals move to campus, spending T_c time commuting in each

direction, and work from campus during the workday (time T_w), while the other fraction (1 - θ) works

146 from home. A total 24 hour cycle is then represented by: $T_h + T_w + 2\, T_c = 1$. Transmission rates can vary

among home (β_h; this includes transmission during essential trips e.g., to the grocery store), commute (β_c;

148 traveling between home and work), and work (β_w; campus-based interactions) environments, while the

rate of moving from exposed to infected (σ) and recovery rate (γ) are the same regardless of where

150 individuals are located.]

152 Here, the rate at which new susceptible individuals (S) become exposed (E) depends on three

components [36]. First is the rate of contact between two individuals in a location. Here we assume this

154 contact rate is constant (does not change with population density) but can differ across environments

(home vs. work vs. commuting). Critically, we assume that β_h accounts for transmission not just in an

156 individual's actual home, but transmission that occurs during other essential activities during lockdown

(e.g., grocery store trips). Second is the probability that the contact for each susceptible individual is with

158   an infected individual; this is given by the proportion of infected individuals in the local population (I/N).

Third is the probability that contact with an infectious individual results in transmission. In equation 1

160   above (and the other equations below), we have combined the first and third factors into a single term, β,

while the second factor is given by I/N. Overall, this gives us frequency-dependent transmission

162   (transmission rate depends on the frequency — not density — of infected individuals in the population);

an appropriate assumption for spatially structured environments [2,36].

164   After the period of time at home (T_h), a fraction, θ, of all individuals commute to work while the

remaining (1- θ) stay to work from home. At this point we subdivide the population based on the number

166   of individuals of each type and fraction commuting. We denote location by subscripts (h for home, c for

commute), so the number of individuals of each type are

168    $$S_c(T_h) = \theta\, S(T_h)$$

$$E_c(T_h) = \theta\, E(T_h)$$

170    $$I_c(T_h) = \theta\, I(T_h)$$

$$R_c(T_h) = \theta\, R(T_h) \qquad\qquad \text{[eqn. 2]}$$

172    $$S_h(T_h) = (1 - \theta)\, S(T_h)$$

$$E_h(T_h) = (1 - \theta)\, E(T_h)$$

174    $$I_h(T_h) = (1 - \theta)\, I(T_h)$$

$$R_h(T_h) = (1 - \theta)\, R(T_h) \ .$$

176   During the commute phase, the infection dynamics for those commuting are given by

$$dS_c/dt = -\, \beta_c\, S_c\, (I_c/N_c)$$

178    $$dE_c/dt = \beta_c\, S_c\, (I_c/N_c) - \sigma\, E_c \qquad\qquad \text{[eqn. 3]}$$

$$dI_c/dt = \sigma\, E_c - \gamma\, I_c$$

180    $$dR_c/dt = \gamma\, I_c$$

where N_c is the total number of individuals commuting ($N_c = S_c + E_c + I_c + R_c$), and β_c is the rate of

182   transmission while commuting. Similarly, during the commute phase, the infection dynamics for those

still at home are given by

184
$$dS_h/dt = -\beta_h \, S_h \, (I_h/N_h)$$

$$dE_h/dt = \beta_h \, S_h \, (I_h/N_h) - \sigma \, E_h \qquad \text{[eqn. 4]}$$

186
$$dI_h/dt = \sigma \, E_h - \gamma \, I_h$$

$$dR_h/dt = \gamma \, I_h$$

188   where N_h is the total number of individuals at home ($N_h = S_h + E_h + I_h + R_h$).

After the commute phase (of length T_c), comes a work phase. Here, the population continues to be

190   subdivided into eight types, where the number of individuals of each type are

$$S_w(T_h+T_c) = S_c(T_h+T_c)$$

192
$$E_w(T_h+T_c) = E_c(T_h+T_c)$$

$$I_w(T_h+T_c) = I_c(T_h+T_c)$$

194
$$R_w(T_h+T_c) = R_c(T_h+T_c) \qquad \text{[eqn. 5]}$$

$$S_h(T_h+T_c) = S_h(T_h+T_c)$$

196
$$E_h(T_h+T_c) = E_h(T_h+T_c)$$

$$I_h(T_h+T_c) = I_h(T_h+T_c)$$

198
$$R_h(T_h+T_c) = R_h(T_h+T_c)$$

where the subscript w denotes work. Note that individuals that work on campus switch from a commute

200   (c) subscript to a work (w) one here, while individuals that work at home continue with the same

subscript (h). Both groups are still experiencing infection dynamics. During the work phase, the infection

202   dynamics in the workplace are given by

$$dS_w/dt = -\beta_w \, S_w \, (I_w/N_w)$$

204
$$dE_w/dt = \beta_w \, S_w \, (I_w/N_w) - \sigma \, E_w \qquad \text{[eqn. 6]}$$

$$dI_w/dt = \sigma \, E_w - \gamma \, I_w$$

206
$$dR_w/dt = \gamma \, I_w$$

where N_w is the total number of individuals at work ($N_w = S_w + E_w + I_w + R_w$), and β_w is the rate of

208 transmission while at work. During the work phase, the infection dynamics for those working at home are given by [eqn. 4] above.

210 After the work phase (of length T_w), we describe a second commute phase. The population continues to be subdivided into eight types, where the number of individuals of each type are

212
$$S_c(T_h+T_c+T_w) = S_w(T_h+T_c+T_w)$$

$$E_c(T_h+T_c+T_w) = E_w(T_h+T_c+T_w)$$

214
$$I_c(T_h+T_c+T_w) = I_w(T_h+T_c+T_w)$$

$$R_c(T_h+T_c+T_w) = R_w(T_h+T_c+T_w) \qquad \text{[eqn. 7]}$$

216
$$S_h(T_h+T_c+T_w) = S_h(T_h+T_c+T_w)$$

$$E_h(T_h+T_c+T_w) = E_h(T_h+T_c+T_w)$$

218
$$I_h(T_h+T_c+T_w) = I_h(T_h+T_c+T_w)$$

$$R_h(T_h+T_c+T_w) = R_h(T_h+T_c+T_w) \ .$$

220 During this second commute phase (also of length T_c), the infection dynamics for those commuting are given by [eqn. 3] above, and the infection dynamics for those still at home are given by [eqn. 4] above. At

222 the end of the second commute phase, all individuals are back in the home environment (no longer subdivided) and the number of individuals of each type are

224
$$S(T_h+2T_c+T_w) = S_w(T_h+2T_c+T_w) + S_h(T_h+2T_c+T_w)$$

$$E(T_h+2T_c+T_w) = E_w(T_h+2T_c+T_w) + E_h(T_h+2T_c+T_w) \qquad \text{[eqn. 8]}$$

226
$$I(T_h+2T_c+T_w) = I_w(T_h+2T_c+T_w) + I_h(T_h+2T_c+T_w)$$

$$R(T_h+2T_c+T_w) = R_w(T_h+2T_c+T_w) + R_h(T_h+2T_c+T_w) \ .$$

228 This ends the cycle for a single day; the next day starts the cycle again.

230 **Model Parameters**

We used a fixed population size (N) of 3,000 individuals. We did not include births or deaths, or

232 movement in and out of the population. These are reasonable assumptions given the scope of our

simulations: a work population that is not hiring new employees and has few retirements or actual deaths

234 over a few months. Because we assumed frequency-dependent transmission, the relative fraction of the

population infected is the same regardless of population size.

236   Infection parameters were calculated as follows. We used 4.2 days as the latent period [37] and

calculated the rate of moving from exposed to infected (σ) as the inverse of this: $\sigma = 1/4.2 = 0.238$ per

238 day. We used 9.5 days as the infectious period (the estimated length of viral shedding for SARS-coV-2;

[38]), and calculated the recovery rate (γ) as the inverse of this: $\gamma = 1/9.5 = 0.105$ per day. Transmission

240 rate (β) was calculated based on the basic reproductive number, R_0. We assumed a 'baseline' R_0

(unmitigated; no behavioral changes like physical distancing) of 2.5 based on current estimates for SARS-

242 coV-2 [39], although some estimates put R_0 as high as 5.7 [37]. To quantify how behavioral changes to

movement and contact affect transmission we defined effective reproduction numbers (R_e) for each of the

244 environments (home, work, commute). We assumed that stay-at-home measures to reduce pathogen

spread in the community halved the rate of contacts at home (e.g., [40]), that is $R_{e\text{-}h} = 0.5R_0$. We assumed

246 that infection at work could be anywhere between current infection rates at home ($R_{e\text{-}w} = 0.5R_0$) and

unmitigated rates ($R_{e\text{-}w} = R_0$). To facilitate interpretation of our results, we also describe infection at work

248 in terms of the fraction increase in transmission compared to home, where 0 indicates transmission is the

same at work and home, 0.5 indicates transmission at work is 50% higher than at home and 1 indicates

250 transmission at work is 100% higher than at home (i.e., double). Finally, we assumed that infection while

commuting spanned a broader range of possible values than either home or work. At one extreme,

252 commuting by private transport effectively has no risk of transmission from others ($R_{e\text{-}c} = 0$). At the other

extreme, commuting by crowded public transport can reduce feasible physical distancing ($R_{e\text{-}c} = 2R_0$),

254 both because individuals have a greater number of contacts while commuting and because these contacts

potentially last for longer than normal. Transmission rates (β) were back-calculated from R_e values, based

256 on rearranging the expression $R_e = \beta/\gamma$ to $\beta = \gamma R_e$. We assume that R_0 and R_e values estimated for the

general public apply to our population of University workers. If instead our population had lower

258 transmission rates than the general public under stay-at-home measures, going back to work could lead to

faster pathogen spread than we predict here.

Simulations

262   Since our aim was to understand the relative importance of model parameters on infection

dynamics (rather than try to forecast outcomes), we started each simulation with one individual infected

264 ($I(t=0)=1$), zero exposed ($E(t=0)=0$), zero removed ($R(t=0)=0$), and the rest susceptible ($S(t=0)=2,999$).

Each simulation was run until it reached equilibrium (where the fraction of the population in the R class

266 did not change from one day to the next). We defined a baseline set of values for each parameter (see

Table 2). Then we ran the following simulations that varied some parameters while holding others

268 constant:

(i) Varying transmission while at work (β_w) and during the commutes (β_c). We considered three

270   scenarios that differed in the degree of risk of a commute to work and back. For low risk, we assumed

low contact both during commutes and on campus ($R_{e\text{-}w} = 0.5R_0 = 1.25$, equivalent to at home). For

272   moderate risk, we assumed unmitigated contact during commute ($R_{e\text{-}c} = R_0 = 2.5$, shared transport)

and partial physical distancing at work ($R_{e\text{-}w} = 0.75R_0 = 1.875$, intermediate between home and

274   unmitigated). For high risk, we assumed elevated contact during commute ($R_{e\text{-}c} = 2R_0 = 5$, crowded

shared transport), and unmitigated contact at work ($R_{e\text{-}w} = R_0 = 2.5$). These results are presented in Fig

276   2.

(ii) Varying the fraction of the population commuting (θ) and fraction of the day spent on campus

278   (T_w). We considered eleven values of the fraction of the population commuting ($\theta = 0,0.1,...,0.9,1$)

and eleven values of the fraction of an 8-hour workday spent on campus ($T_w = x(8/24)$ where x $=$

280      $0,0.1,...,0.9,1$). These results are presented in Fig 3a.

(iii) Varying the fraction of the population commuting (θ) and fraction increase in transmission at

282   work compared to home ($R_{\text{e-w}}$). We considered eleven values of the fraction of the population

commuting ($\theta = 0,0.1,...,0.9,1$) and eleven values of the fraction increase in transmission at work

284   compared to at home ($R_{\text{e-w}} = (1+x)R_{\text{e-h}}$ where x = 0,0.1,...,0.9,1). These results are presented in Fig 3b.

(iv) Varying the fraction of the day spent on campus (T_{w}) and fraction increase in transmission at

286   work compared to home ($R_{\text{e-w}}$). We considered eleven values of the fraction of an 8-hour workday

spent on campus ($T_{\text{w}} = y(8/24)$ where y = 0,0.1,...,0.9,1), and eleven values of the fraction increase in

288   transmission at work compared to at home ($R_{\text{e-w}} = (1+x)R_{\text{e-h}}$ where x = 0,0.1,...,0.9,1). These results

are presented in Fig 3c.

290   Movement model simulations were conducted in Matlab 2018b.

292   [**Fig 2. Movement model: varying the degree of physical distancing on campus and during**

commutes.

294   The fraction of the population that is Susceptible (S), Exposed (E), Infected (I), and Removed (R), when

all individuals either work from home (solid lines) or commute to work on campus (dashed lines), for

296   different degrees of physical distancing both on campus and during the commute: (a) low risk: low

contact during commute and on campus ($R_{\text{e-c}} = R_{\text{e-w}} = 0.5R_0 = 1.25$, equivalent to at home), dashed and

298   solid lines are identical, (b) moderate risk: unmitigated contact during commute ($R_{\text{e-c}} = R_0 = 2.5$, shared

transport) and partial physical distancing at work ($R_{\text{e-w}} = 0.75R_0 = 1.875$, intermediate between home and

300   unmitigated), (c) high risk: elevated contact during commute ($R_{\text{e-c}} = 2R_0 = 5$, crowded shared transport),

and unmitigated contact at work ($R_{\text{e-w}} = R_0 = 2.5$).]

[**Fig 3. Movement model: limiting people, time and contact on campus.**

304   The final epidemic size (cumulative fraction of the population infected) as a function of (a) the fraction of

an 8-hour workday spent on campus (x-axis) and the fraction of the population working on campus (y-

306   axis) with no physical distancing, (b) the fraction increase in transmission while at work compared to at

home (x-axis) and the fraction of the population working on campus (y-axis) with an 8-hour work day, (c)

308   the fraction increase in transmission while at work compared to at home (x-axis) and the fraction of an 8-

hour workday spent on campus (y-axis) with 100% of people on campus.]

Sensitivity Analysis

312   Finally, we performed a sensitivity analysis to determine how sensitive the two model output

metrics (final epidemic size, epidemic peak size) were to each of the model parameters, using a

314   combination of Latin Hypercube Sampling (LHS) and Partial Rank Correlation Coefficients (PRCC). The

LHS/PRCC sensitivity analysis is appropriate when the relationship between model output and each

316   model parameter is monotonic and nonlinear [41]. For our model, this relationship was monotonic for all

nine parameters considered (N, T_c, T_w, θ, σ, γ, $R_{e\text{-}c}$, $R_{e\text{-}h}$, and $R_{e\text{-}w}$; S1-S2 Figs). The LHS/PRCC sensitivity

318   analysis has two steps.

          First, we used Latin Hypercube Sampling (LHS; [42]), a Monte Carlo approach, to generate sets

320   of parameter value combinations from preset ranges of parameter values. LHS has a minimum required

sample size (n) which is given by: $n \geq k+1$ or $n \geq k(4/3)$ where k is the number of parameters included in

322   the LHS [43], nine for our analysis. We chose the number of samples (see below) to meet these criteria.

Each of the nine model parameters considered was sampled from a uniform probability density function

324   based on the ranges given in Table 2. The model was run for each parameter value set, and the final

epidemic size (cumulative fraction of the population infected, in the long-term) and epidemic peak size

326   (maximum fraction of the population infected at any time) were both saved as output metrics.

          Second, we measured the sensitivity of the output metrics to each parameter using Spearman

328   Partial Rank Correlation Coefficients (PRCC). To determine how many samples of each parameter was

needed to generate stable PRCC value, we calculated PRCC value for an increasing number of samples

330   (S3 Fig) and noted that the PRCC values were relatively stable past 1000 samples. Thus, we used 1000

samples of each parameter value for our final PRCC analysis. A positive PRCC value indicates that

332   increasing the value of that parameter increases the output metric while a negative PRCC value indicates

that increasing the value of that parameter decreases the output metric. PRCC values that were not

334   significant at the 0.05 level are marked with 'ns' in Fig 4 (not corrected for multiple comparisons).

Finally, we used a z-test to rank significant model parameters in terms of their relative importance, since

336   larger PRCC values do not always indicate more important parameters [41]. For our results (Fig 4), model

output sensitivity was indeed given by the size of PRCC values.

[**Fig 4. Movement model: sensitivity analysis.**

340   The partial rank correlation coefficient (PRCC) values for each model parameter (Table 2) for the final

epidemic size metric (blue bars) and the epidemic peak size metric (orange bars). Positive values indicate

342   parameters that increase epidemic size as they are increased (negative values indicate parameters that

decrease epidemic size as they are increased). Cases where the relationship between the parameter and

344   model output metric was not significant are indicated with 'ns'.]

Methods: network model

Our second model explores infection dynamics as individuals work on campus either in both

348   office and lab spaces or just in lab spaces. We created a network map of all the individuals housed in the

Ecology building on the St. Paul campus of the University of Minnesota. We created our dataset by

350   merging information on the office and lab room assignments for each individual with an office or lab in

the building. (The methods for collection and analyses of these data were reviewed by the University of

352   Minnesota's Institutional Review Board and were determined not to be human subjects research.) Work

in the Ecology building is structured by two primary space types, laboratories that can include one to

354 three research groups, each associated with a single faculty member, and offices which can be single-

occupancy or shared. Office space is generally shared by groups of graduate students and postdoctoral

356 scholars, often from different lab groups. Because undergraduates are generally not permitted to work on

campus during the resumption of research, we included faculty, staff, postdocs, and graduate students, but

358 excluded all undergraduates from this visualization.

We considered two types of bipartite networks: shared office space and shared lab space.

360 Individuals sharing an office or a lab all had an edge with that location node. We then consider the one-

mode projection of each network, creating a weighted unipartite network connecting individuals

362 according to their shared spaces. The binary representation of these networks was used to create static

network visualizations of connections among individuals using the igraph, tidygraph, and ggraph libraries

364 in R [44–46], shown in Fig 5a and b. Animations of disease progression through the networks were

produced using the gganimate library in R (S5-S6 Figs; [47]). For each network, we computed the

366 distribution of (finite) shortest paths between each pair of nodes (Fig 5c) and for each distinct component

of the networks, we noted its size (number of nodes), diameter (longest shortest path), and mean path

368 length (average shortest path length; S7 Fig).

370 [**Fig 5. Network model structure.**

Space-sharing, or 'contacts' (edges) are shown among all individuals (nodes) for two scenarios: (a) when

372 individuals at work share either office or lab spaces, or (b) when individuals only used shared lab space

and not shared offices (e.g., bench work is done on campus while office work is done at home). (c)

374 Histograms showing the distribution of shortest paths between all connected pairs of individuals.

Importantly, though all shortest paths between nodes in the network containing only links of shared lab

376 spaces are less than or equal to three, the vast majority (approximately 95%) of pairwise combinations of

individuals actually have no chain of interactions connecting them. In contrast, the combined network

378 contains a component consisting of almost 90% of individuals in the network, corresponding to nearly

80% of all pairs of individuals having a chain of interactions connecting them.]

For the network simulations, we used an SEIR model framework, starting with a randomly

382   selected index case to serve as the first infected individual in an entirely susceptible population (Fig 6).

Simulations proceeded in discrete time. At each time step, individuals who had been exposed to the virus

384   transitioned into the infectious class (E→I) based on the result of a Bernoulli trial using the disease

progression rate as the probability of success. Likewise, currently infectious individuals were removed

386   (i.e., either recovered and immune or deceased; I→R) based on the result of a Bernoulli trial using the

recovery rate as the probability of success. Finally, one Bernoulli trial using the transmission rate as

388   probability of success was conducted for each edge connecting a susceptible individual to an infectious

one. Susceptible individuals became exposed (S→E) if at least one such trial resulted in success. At the

390   end of each simulation, we took note of the epidemic peak size, the final epidemic size, and the time

needed to read the epidemic peak (Fig 7). We evaluated the sensitivity of these results by comparing them

392   to simulations run on randomized versions of these empirical networks (S8 Supporting Information).

Network analysis and simulations were conducted in R (Version 3.6.3).

[**Fig 6. Network model simulations.**

396   Final disease status of members of networks based on use of (a) both shared lab and office space and (c)

only shared lab space, following a simulated epidemic with susceptible individuals in blue, exposed

398   individuals in green, infectious individuals in orange, and removed individuals in red. (b, d) the

cumulative number of susceptible, infectious, and removed individuals over time for each network

400   simulation.]

402   [**Fig 7. Network model simulations.**

Outcome of 100 infection simulations on networks: the maximum peak number of individuals infected at

404 any one time (epidemic peak size), total number of individuals infected (final epidemic size), and time

until peak number of individuals infected for simulations of pathogen spread on networks based on use of

406 both shared lab and office space (blue) and only shared lab space (orange).]

Results

Movement model

412 Whether returning to work on campus affects the epidemic outcomes (measured as final epidemic

size and epidemic peak size) depends critically on the degree of physical distancing maintained both on

414 campus and during the commute between home and campus (Fig 2). If the current degree of physical

distancing that is achieved while working from home can be maintained while on campus, then working

416 from campus will not speed up infection dynamics compared to working from home (Fig 2a). However, if

physical distancing on campus or during the commute is less successful than current physical distancing

418 at home, then returning to work on campus will both increase the epidemic peak size in the short-term and

increase the final epidemic size in the long term (Fig 2b-c). When physical distancing cannot be

420 maintained on campus or during the commute, then infection dynamics can be kept slower by limiting the

fraction of workers on campus and the amount of time workers are on campus (Fig 3a).

422 Intriguingly the three strategies we considered (limiting contact, people, or time on campus) are

interchangeable with approximately equivalent effects on both the long-term metric, final epidemic size

424 (Fig 3) and the short-term metric, epidemic peak size (S4 Fig). That is, in situations where one of these

strategies cannot be fully implemented, a different strategy can be used in its stead. For example, if

426 individuals need to be on campus for an extended period of time to run an experiment (thus limiting time

on campus is not a feasible strategy), this can be compensated for by limiting the number of other

428   individuals on campus at the same time. However, of the three strategies, reducing the fraction of the

population on campus had a bigger impact than reducing either time or contact on campus, due to the

430   effect of commuting to and from campus. Regardless of time or physical distancing on campus, more

people working on campus requires that more people commute. Thus, if commuting substantially

432   increases transmission risk compared to staying at home (i.e., any form of shared transport vs. commuting

alone), reducing the number of people commuting will be a more effective strategy than reducing either

434   time or contact while on campus.

The sensitivity analysis revealed that both model metrics (final epidemic size, epidemic peak

436   size) were most sensitive to transmission at home ($R_{e\text{-}h}$), since most of the day is spent in that

environment, as well as the fraction of the population commuting (θ) to campus (Fig 4). Transmission on

438   campus ($R_{e\text{-}w}$) and transmission during commutes ($R_{e\text{-}c}$) were the next most influential; the first because

most time during the workday is spent on campus and the second because we allowed transmission to

440   vary across a wider range during commuting than on campus. The time spent on campus (T_w) and time

commuting (T_c) were somewhat influential. For each of these parameters, increasing the parameter value

442   increased the final and peak epidemic sizes. Finally, population size (N) did not significantly affect either

metric (but would be critical for the total number of individuals infected). The rate of moving from

444   exposed to infected (σ) and the recovery rate (γ) did not significantly affect final epidemic size, but both

had a minor effect on epidemic peak size: increasing σ (i.e., a shorter latent period) increased epidemic

446   peak size, while increasing γ (i.e., a shorter infectious period) decreased epidemic peak size.

448   # Network model

The mixing of researchers from different labs in shared office spaces had a substantial impact on

450   the modularity of the resulting network. In particular, when people do not use shared office spaces (i.e.,

work from home if they share an office), but work on campus only in labs, the network is far more

452 modular, with smaller, more densely connected groups and few connections among groups (Fig 5b, S7

Fig). In this case, most individuals are directly connected to all other members of their group (i.e.,

454 "shortest path" of one, Fig 5c); however the absence of connections between groups means that, on

average, an infected individual lacks a path of connections to 95% of the rest of the population. In

456 contrast, when individuals share both lab and office space, the connectedness of the network is relatively

high because students, staff, and postdocs that share offices are often from different labs. For this

458 combined case, most individuals are four or fewer connections from one another (Fig 5c) and the largest

component contains nearly 90% of all individuals in the network (Fig 5a, S7 Fig). Thus, in the case where

460 an infected individual (presumably pre-symptomatic or asymptomatic) came to work, the combined lab

and office network has the potential for greater disease incidence than in the lab-only network, where the

462 infection could be constrained to a single lab (Figs 6-7, S5-S6 Figs). In general, when compared to the

combined network, the lab-only network had outbreaks that were less explosive (i.e., had less variance

464 and a lower mean number of individuals infected at any one time), fewer individuals infected overall, and

a shorter time until the peak number of infectious individuals (Fig 7).

Discussion

468   Movement and contact behaviors are key drivers of the spread of pathogens like SARS-CoV-2,

and not all movement and contacts have the same impact on pathogen spread. However, basic

470 compartmental models used to describe SARS-CoV-2 dynamics assume all individuals move and contact

each other at random (i.e., populations are well-mixed). Our models show how explicitly accounting for

472 movement, space use in a building, and contact behaviors can provide a more nuanced understanding of

relative risk. Our movement model, capturing the predictable movement between home and work/campus

474 environments, shows that reducing the number of people, rate of contact, and amount of time spent on

campus are all equivalently effective strategies for slowing pathogen spread. However, if commutes

476   specifically increase transmission risk (i.e., shared transport), reducing the number of people on campus is

the most effective strategy to reduce the infection spread rate. We also considered heterogeneity in

478   contact behavior once at the workplace; our network model captures the regular interactions among

workers in shared workspaces on campus and shows that restricting building use to lab spaces (rather than

480   lab and office space) may reduce pathogen spread. Our results provide a number of tools to distinguish

among different movement and contact patterns at the scale of individuals and workplace communities.

482   A number of future directions could be explored, by changing some of our simplifying

assumptions. First, staying within the broad structure of our model, alternative spatiotemporal strategies

484   could be explored including: structured work weeks (e.g., four days on-campus and ten off; [48]), or

further compartmentalizing time (e.g., sequential work shifts) or space (e.g., different buildings on

486   campus). For instance, if evidence suggests that infection can occur through air circulation within

buildings [49], these models could be altered to account for connections arising from shared ventilation

488   systems. These models also could be modified to account for movement and contact behavior that

explicitly depends on infection status [50]; e.g., splitting infected individuals into infectious but

490   asymptomatic (who still potentially commute to work) and symptomatic individuals (who stay home) or

building dynamic networks where contact behavior can change in response to infectious status (e.g.

492   infectious and symptomatic individuals reducing their interpersonal contacts either through staying home

or altering their behavior at work). Second, one could expand the scale of the model. This could be done

494   foremost by combining the movement model (movement between work, commute, campus) with the

network model (movement while on campus). Further expansions could consider both larger scales

496   (linking in regional patterns) as well as smaller ones (allowing contacts within buildings to vary over

time). For instance, integrating local models such as ours with regional variation in infection rates and

498   degree of community social mixing [11,51,52] could further inform recommendations. Third, as data

accumulate on transmission dynamics and individual susceptibility, we can alter specific players or

500   interactions in the model. For instance, while the virus can survive on surfaces [53], most transmission

        appears to be aerosolized, mediated by extended person-to-person interactions in close spaces [54–58], so

502   masking and minimizing temporal and spatial overlap of workers in shared spaces is key [59–61]. In

        addition, susceptibility and thus local demographic data can provide additional layers of tailored

504   recommendations [62].

            Our findings mesh with concepts in the broader movement and disease ecology literature. Within

506   movement ecology, there has long been a distinction between random/undirected movement like dispersal

        versus predictable movements like diel and seasonal migration [63]. Human movement between home

508   and work is often a predictable and daily occurrence and thus is better viewed from the lens of predictable

        migratory movements (as we do here) rather than random dispersive ones (as implicit in basic

510   compartmental models). Moving predictably between two environments does not always increase

        infection (either for individuals or at the population level) compared to remaining in a single location; the

512   relative transmission in each environment is critical [22]. We find that transmission risk during a

        commute is key to infection dynamics when considering the impact of movement between home and

514   work, paralleling recent work calling for the explicit consideration of how transient phases of movement

        affect infection dynamics [24] and theory showing that infection dynamics during transit can have a

516   similar impact to dynamics in the second environment [64].

            There are important insights that emerge from our movement and contact-network models that

518   can guide policy. For example, basic disease models assume random movement and equal probability of

        contact, whereas many hosts, including humans, move in directed ways and in very structured social

520   networks. For this reason, disease mitigation policies will likely be more effective when they consider

        disease risk in a more holistic way that integrates risk across the various components of a person's daily

522   movement. For example, in settings where many people commute by mass transit (e.g., New York City),

        the efficacy of workplace safety protocols may be overwhelmed by transmission during daily commutes

524   rather than contacts at work. Careful examination of social network patterns could also help guide policy

to provide intermediate scenarios between business as usual and complete lock down. For example, in our

526   case-study contact rates and potential disease spread were significantly reduced when people's contacts at

the workplace were restricted to single lab groups, as opposed to linking separate lab and office networks.

528   These findings are consistent with emerging calls to reduce COVID-19 spread by creating "learning

pods" and "social bubbles" of interacting children and adults as schools and workplaces re-open [65,66].

530   Even so, the protective effects of heterogeneity in contact structure should not be overemphasized

for decision making. First, although the threshold for herd immunity can be lower in heterogeneous

532   networks [67], making outbreaks less likely, outbreaks that do occur can also be more explosive [25].

Second, because SARS-CoV-2 spread appears to be primarily by aerosolized transmission, the potential

534   contact behaviors needed for transmission are more ubiquitous than for pathogens with more specific

transmission modes (e.g., sexually transmitted diseases like HIV/AIDS). Importantly, the networks

536   presented here consider only the room in which an employee works (their office or lab space), explicitly

omitting broader workplace considerations like air flow, shared surfaces, entry points, etc., these

538   additional points must be addressed in conjunction with thinking about explicit contact behavior when

forming a public health strategy. Lastly, these static networks are a simplification of an inherently

540   dynamic process of movement, contact, and infection. Using a time-ordered or dynamic network

approach could provide better insights to actual duration of exposures and sickness-induced behavioral

542   changes [68].

Conclusions

Human movement and contact behaviors are critical for the spread of pathogens like SARS-

546   CoV-2, yet are rarely addressed explicitly in the current conversations about decision-making in the face

of relaxing Stay at Home orders. Here we have drawn on movement and network models to demonstrate

548   the effect of these behaviors. First, we have shown that regular movement between two 'environments'

(i.e., work and home) does not inherently increase infection spread the way random dispersive

550   movements might. Rather the outcome depends on the relative degree of transmission (e.g., degree of

physical distancing) in each environment. Second, we have shown that different contact patterns (e.g.,

552   space usage) within the work environment could lead to different outcomes in terms of SARS-CoV-2

spread. In sum, we advocate for using an understanding of movement and contact patterns as an

554   adjunctive approach (alongside widespread testing, contact tracing, vaccine development and other tools)

to mitigate the effects of SARS-CoV-2 and COVID-19, particularly when considering return to work

556   environments.

558   Funding

This material is based in part upon work supported by the University of Minnesota's Office of

560   Academic Clinical Affairs COVID-19 Rapid Response Grant (https://clinicalaffairs.umn.edu/umn-covid-

19-research) (to MEC, LAW, and MMS), by the National Science Foundation (https://www.nsf.gov/)

562   under Grants DEB-2030509 (to MEC and MMS), DEB-1654609 (to AKS and MEC) and DEB-1556649

(to EWS and ETB) and by the National Socio Environmental Synthesis Center (SESYNC) under funding

564   received from the NSF DBI 1639145. The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript. The funders had no role in study design,

566   data collection and analysis, decision to publish, or preparation of the manuscript.

568   Acknowledgements

We thank William Harcombe for helpful feedback and discussion, and Valery Forbes, David

570   Greenstein and Daniel Stanton for encouragement.

Data Availability

Model code and simulation output for the movement model is available on Data Dryad [69]. All Code to

574 generate the network model figures and animations is available on Github

(https://github.com/whit1951/EEBCovid).

References

1. Kraemer MUG, Yang C-H, Gutierrez B, Wu C-H, Klein B, Pigott DM, et al. The effect of human mobility and control measures on the COVID-19 epidemic in China. Science. 2020;368: 493–497. doi:10.1126/science.abb4218

2. Keeling MJ, Rohani P. Modeling Infectious Diseases in Humans and Animals. Princeton, NJ: Princeton University Press; 2008.

3. Keeling MJ, Grenfell BT. Individual-based perspectives on R0. J Theor Biol. 2000;203: 51–61. doi:10.1006/jtbi.1999.1064

4. Kucharski AJ, Russell TW, Diamond C, Liu Y, Edmunds J, Funk S, et al. Early dynamics of transmission and control of COVID-19: a mathematical modelling study. Lancet Infect Dis. 2020;20: 553–558. doi:10.1016/S1473-3099(20)30144-4

5. Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. Eurosurveillance. 2020;25. doi:10.2807/1560-7917.ES.2020.25.10.2000180

6. Wu JT, Leung K, Bushman M, Kishore N, Niehus R, de Salazar PM, et al. Estimating clinical severity of COVID-19 from the transmission dynamics in Wuhan, China. Nat Med. 2020;26: 506–510. doi:10.1038/s41591-020-0822-7

7. Moghadas SM, Shoukat A, Fitzpatrick MC, Wells CR, Sah P, Pandey A, et al. Projecting hospital utilization during the COVID-19 outbreaks in the United States. Proc Natl Acad Sci. 2020;117: 9122–9126. doi:10.1073/pnas.2004064117

8. Duque D, Morton DP, Singh B, Du Z, Pasco R, Meyers LA. Timing social distancing to avert unmanageable COVID-19 hospital surges. Proc Natl Acad Sci. 2020;117: 19873–19878. doi:10.1073/pnas.2009033117

600   9.   Lewnard JA, Lo NC. Scientific and ethical basis for social-distancing interventions against

COVID-19. Lancet Infect Dis. 2020; S1473309920301900. doi:10.1016/S1473-3099(20)30190-0

602   10.   Balilla J. Assessment of COVID-19 Mass Testing: The Case of South Korea. SSRN Electron J.

2020 [cited 3 May 2020]. doi:10.2139/ssrn.3556346

604   11.   Koo JR, Cook AR, Park M, Sun Y, Sun H, Lim JT, et al. Interventions to mitigate early spread of

SARS-CoV-2 in Singapore: a modelling study. Lancet Infect Dis. 2020; S1473309920301626.

606   doi:10.1016/S1473-3099(20)30162-6

     12.   Hadjidemetriou GM, Sasidharan M, Kouyialis G, Parlikad AK. The impact of government

608   measures and human mobility trend on COVID-19 related deaths in the UK. Transp Res

Interdiscip Perspect. 2020;6: 100167. doi:10.1016/j.trip.2020.100167

610   13.   Zhang J, Litvinova M, Liang Y, Wang Y, Wang W, Zhao S, et al. Changes in contact patterns

shape the dynamics of the COVID-19 outbreak in China. Science. 2020; eabb8001.

612   doi:10.1126/science.abb8001

     14.   Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The

614   psychological impact of quarantine and how to reduce it: rapid review of the evidence. The

Lancet. 2020;395: 912–920. doi:10.1016/S0140-6736(20)30460-8

616   15.   Sønderskov KM, Dinesen PT, Santini ZI, Østergaard SD. The depressive state of Denmark during

the COVID-19 pandemic. Acta Neuropsychiatr. 2020;32: 226–228. doi:10.1017/neu.2020.15

618   16.   Viner RM, Russell SJ, Croker H, Packer J, Ward J, Stansfield C, et al. School closure and

management practices during coronavirus outbreaks including COVID-19: a rapid systematic

620   review. Lancet Child Adolesc Health. 2020;4: 397–404. doi:10.1016/S2352-4642(20)30095-X

     17.   Levinson M, Cevik M, Lipsitch M. Reopening primary schools during the pandemic. N Engl J

622   Med. 2020; NEJMms2024920. doi:10.1056/NEJMms2024920

     18.   Paltiel AD, Zheng A, Walensky RP. Assessment of SARS-CoV-2 screening strategies to permit

624   the safe Reopening of college campuses in the United States. JAMA Netw Open. 2020;3:

e2016818. doi:10.1001/jamanetworkopen.2020.16818

626 19. Binning SA, Shaw AK, Roche DG. Parasites and host performance: Incorporating infection into

our understanding of animal movement. Integr Comp Biol. 2017;57: 267–280.

628 doi:10.1093/icb/icx024

20. Loehle C. Social barriers to pathogen transmission in wild animal populations. Ecology. 1995;76:

630 326–335. doi:10.2307/1941192

21. Shaw AK, Binning SA. Recovery from infection is more likely to favour the evolution of

632 migration than social escape from infection. J Anim Ecol. 2020;89: 1448–1457.

doi:10.1111/1365-2656.13195

634 22. Shaw AK, Sherman J, Barker FK, Zuk M. Metrics matter: the effect of parasite richness, intensity

and prevalence on the evolution of host migration. Proc R Soc B Biol Sci. 2018;285: 20182147.

636 doi:10.1098/rspb.2018.2147

23. Boulinier T, Kada S, Ponchon A, Dupraz M, Dietrich M, Gamble A, et al. Migration, prospecting,

638 dispersal? What host movement matters for infectious agent circulation? Integr Comp Biol.

2016;56: 330–342. doi:10.1093/icb/icw015

640 24. Daversa DR, Fenton A, Dell AI, Garner TWJ, Manica A. Infections on the move: how transient

phases of host movement influence disease spread. Proc R Soc B Biol Sci. 2017;284: 20171807.

642 doi:10.1098/rspb.2017.1807

25. Keeling MJ, Eames KT. Networks and epidemic models. J R Soc Interface. 2005;2: 295–307.

644 doi:10.1098/rsif.2005.0051

26. Altieri N, Barter RL, Duncan J, Dwivedi R, Kumbier K, Li X, et al. Curating a COVID-19 data

646 repository and forecasting county-level death counts in the United States. medRxiv. 2020.

Available: https://arxiv.org/abs/2005.07882

648 27. Woody S, Garcia Tec M, Dahan M, Gaither K, Lachmann M, Fox SJ, et al. Projections for first-

wave COVID-19 deaths across the US using social-distancing measures derived from mobile

650   phones. Infectious Diseases (except HIV/AIDS); 2020 Apr. doi:10.1101/2020.04.16.20068163

28.   Kissler SM, Tedijanto C, Goldstein E, Grad YH, Lipsitch M. Projecting the transmission

652   dynamics of SARS-CoV-2 through the postpandemic period. Science. 2020; eabb5793.

        doi:10.1126/science.abb5793

654 29. Chinazzi M, Davis JT, Ajelli M, Gioannini C, Litvinova M, Merler S, et al. The effect of travel

        restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. Science. 2020;

656   eaba9757. doi:10.1126/science.aba9757

30.   Prem K, Liu Y, Russell TW, Kucharski AJ, Eggo RM, Davies N, et al. The effect of control

658   strategies to reduce social mixing on outcomes of the COVID-19 epidemic in Wuhan, China: a

        modelling study. Lancet Public Health. 2020;5: e261–e270. doi:10.1016/S2468-2667(20)30073-6

660 31. Tian L, Li X, Qi F, Tang Q-Y, Tang V, Liu J, et al. Calibrated intervention and containment of

        the COVID-19 pandemic. preprint. 2020; 31. https://covid-

662   19.conacyt.mx/jspui/bitstream/1000/4645/1/1106972.pdf

32.   MN Department of Health. Situation Update for Coronavirus Disease 2019 (COVID-19). 2020.

664   Available: https://www.health.state.mn.us/diseases/coronavirus/situation.html#cases1

33.   Wei WE, Li Z, Chiew CJ, Yong SE, Toh MP, Lee VJ. Presymptomatic transmission of SARS-

666   CoV-2 — Singapore, January 23–March 16, 2020. MMWR Morb Mortal Wkly Rep. 2020;69:

        411–415. doi:10.15585/mmwr.mm6914e1

668 34. Servedio MR, Brandvain Y, Dhole S, Fitzpatrick CL, Goldberg EE, Stern CA, et al. Not just a

        theory—The utility of mathematical models in evolutionary biology. PLoS Biol. 2014;12:

670   e1002017.

35.   Johns S, Shaw AK. Theoretical insight into three disease-related benefits of migration. Popul

672   Ecol. 2016;58: 213–221. doi:10.1007/s10144-015-0518-x

36.   Begon M, Bennett M, Bowers RG, French NP, Hazel SM, Turner J. A clarification of

674   transmission terms in host-microparasite models: numbers, densities and areas. Epidemiol Infect.

2002;129: 147–153. doi:10.1017/S0950268802007148

676   37.   Sanche S, Lin YT, Xu C, Romero-Severson E, Hengartner N, Ke R. High Contagiousness and

Rapid Spread of Severe Acute Respiratory Syndrome Coronavirus 2. Emerg Infect Dis. 2020;26.

678   doi:10.3201/eid2607.200282

  38.   Ling Y, Xu S-B, Lin Y-X, Tian D, Zhu Z-Q, Dai F-H, et al. Persistence and clearance of viral

680   RNA in 2019 novel coronavirus disease rehabilitation patients: Chin Med J (Engl). 2020;133:

1039–1043. doi:10.1097/CM9.0000000000000774

682   39.   Zhang S, Diao M, Yu W, Pei L, Lin Z, Chen D. Estimation of the reproductive number of novel

coronavirus (COVID-19) and the probable outbreak size on the Diamond Princess cruise ship: A

684   data-driven analysis. Int J Infect Dis. 2020;93: 201–204. doi:10.1016/j.ijid.2020.02.033

  40.   Enns EA, Kirkeide M, Mehta A, MacLehose R, Knowlton GS, Smith MK, et al. Modeling the

686   impact of social distancing measures on the spread of SARS-CoV-2 in Minnesota. MN

Department of Health; 2020 Apr p. 16. Available: Enns EA, Kirkeide M, Mehta A, MacLehose

688   R, Knowlton GS, Smith https://mn.gov/covid19/assets/MNmodel_tech_doc_tcm1148-427724.pdf

  41.   Marino S, Hogue IB, Ray CJ, Kirschner DE. A methodology for performing global uncertainty

690   and sensitivity analysis in systems biology. J Theor Biol. 2008;254: 178–196.

doi:10.1016/j.jtbi.2008.04.011

692   42.   Mckay MD, Beckman RJ, Conover WJ. A comparison of three methods for selecting values of

input variables in the analysis of output from a computer code. Technometrics. 1979;42: 55–61.

694   doi:10.1080/00401706.2000.10485979

  43.   Blower SM, Dowlatabadi H. Sensitivity and Uncertainty Analysis of Complex Models of Disease

696   Transmission: An HIV Model, as an Example. Int Stat Rev Rev Int Stat. 1994;62: 229.

doi:10.2307/1403510

698   44.   Csárdi G, Nepusz T. The igraph software package for complex network research. InterJournal

Complex Syst. 2006;1695: 9.

700 45. Pedersen TL. tidygraph: A Tidy API for Graph Manipulation. 2019. Available: https://CRAN.R-
project.org/package=tidygraph

702 46. Pedersen TL. ggraph: An Implementation of Grammar of Graphics for Graphs and Networks.
2020. Available: https://CRAN.R-project.org/package=ggraph

704 47. Pedersen TL, Robinson D. gganimate: A Grammar of Animated Graphics. 2020. Available:
https://CRAN.R-project.org/package=gganimate

706 48. Karin O, Bar-On YM, Milo T, Katzir I, Mayo A, Korem Y, et al. Cyclic exit strategies to
suppress COVID-19 and allow economic activity. medRxiv. 2020 [cited 13 May 2020].
708 doi:10.1101/2020.04.04.20053579

49. Lu J, Gu J, Li K, Xu C, Su W, Lai Z, et al. COVID-19 Outbreak Associated with Air
710 Conditioning in Restaurant, Guangzhou, China, 2020. Emerg Infect Dis. 2020;26.
doi:10.3201/eid2607.200764

712 50. Naven Narayanan, Binning SA, Shaw AK. Infection state can affect host migratory decisions.
Oikos. in press. https://onlinelibrary.wiley.com/doi/abs/10.1111/oik.07188

714 51. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time.
Lancet Infect Dis. 2020;20: 533–534. doi:10.1016/S1473-3099(20)30120-1

716 52. Zhang J, Litvinova M, Liang Y, Zheng W, Shi H, Vespignani A, et al. The impact of relaxing
interventions on human contact patterns and SARS-CoV-2 transmission in China. medRxiv. 2020
718 [cited 31 Aug 2020]. doi:10.1101/2020.08.03.20167056

53. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al.
720 Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. N Engl J Med.
2020;382: 1564–1567. doi:10.1056/NEJMc2004973

722 54. Ghinai I, Woods S, Ritger KA, McPherson TD, Black SR, Sparrow L, et al. Community
Transmission of SARS-CoV-2 at Two Family Gatherings — Chicago, Illinois, February–March
724 2020. MMWR Morb Mortal Wkly Rep. 2020;69: 446–450. doi:10.15585/mmwr.mm6915e1

55. Li W, Zhang B, Lu J, Liu S, Chang Z, Cao P, et al. The characteristics of household transmission of COVID-19. Clin Infect Dis. 2020; ciaa450. doi:10.1093/cid/ciaa450

56. Bi Q, Wu Y, Mei S, Ye C, Zou X, Zhang Z, et al. Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study. Lancet Infect Dis. 2020; S1473309920302875. doi:10.1016/S1473-3099(20)30287-5

57. Morawska L, Cao J. Airborne transmission of SARS-CoV-2: The world should face the reality. Environ Int. 2020;139: 105730. doi:10.1016/j.envint.2020.105730

58. Liu Y, Ning Z, Chen Y, Guo M, Liu Y, Gali NK, et al. Aerodynamic analysis of SARS-CoV-2 in two Wuhan hospitals. Nature. 2020;582: 557–560. doi:10.1038/s41586-020-2271-3

59. Qian H, Miao T, Liu L, Zheng X, Luo D, Li Y. Indoor transmission of SARS-CoV-2. medRxiv. 2020 [cited 11 May 2020]. doi:10.1101/2020.04.04.20053058

60. Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schünemann HJ, et al. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. The Lancet. 2020;395: 1973–1987. doi:10.1016/S0140-6736(20)31142-9

61. Prather KA, Wang CC, Schooley RT. Reducing transmission of SARS-CoV-2. Science. 2020;368: 1422–1424. doi:10.1126/science.abc6197

62. Dowd JB, Andriano L, Brazel DM, Rotondi V, Block P, Ding X, et al. Demographic science aids in understanding the spread and fatality rates of COVID-19. Proc Natl Acad Sci. 2020;117: 9696–9698. doi:10.1073/pnas.2004911117

63. Heape W. Emigration, Migration and Nomadism. Cambridge, UK: Heffer and Sons; 1931.

64. Shaw AK, Craft ME, Zuk M, Binning SA. Host migration strategy is shaped by forms of parasite transmission and infection cost. J Anim Ecol. 2019;88: 1601–1612. doi:10.1111/1365-2656.13050

65. Block P, Hoffman M, Raabe IJ, Dowd JB, Rahal C, Kashyap R, et al. Social network-based

750   distancing strategies to flatten the COVID-19 curve in a post-lockdown world. Nat Hum Behav.

       2020;4: 588–596. doi:10.1038/s41562-020-0898-6

752   66.   Moyer MW. Pods, Microschools and Tutors: Can Parents Solve the Education Crisis on Their

       Own? N Y Times. 2020; 5. https://covid-19.conacyt.mx/jspui/bitstream/1000/4645/1/1106972.pdf

754   67.   Britton T, Ball F, Trapman P. A mathematical model reveals the influence of population

       heterogeneity on herd immunity to SARS-CoV-2. Science. 2020;369: 846–849.

756   doi:10.1126/science.abc6810

    68.   Enright J, Kao RR. Epidemics on dynamic networks. Epidemics. 2018;24: 88–97.

758   doi:10.1016/j.epidem.2018.04.003

    69.   Shaw AK, White LA, Michalska-Smith M, Borer ET, Craft ME, Seabloom EW, et al. Data from:

760   Lessons from movement ecology for the return to work: modeling contacts and the spread of

       COVID-19. 2020. Available: https://doi.org/10.5061/dryad.pg4f4qrmj

762   Table 1. Movement model state variables and their meaning.

Variable	Meaning
S	Total number of susceptible individuals
E	Total number of exposed individuals
I	Total number of infected individuals
R	Total number of removed individuals
S_c	Number of susceptible individuals commuting to campus (during the commute phases)
E_c	Number of exposed individuals commuting to campus (during the commute phases)
I_c	Number of infected individuals commuting to campus (during the commute phases)
R_c	Number of removed individuals commuting to campus (during the commute phases)
S_w	Number of susceptible individuals working from work (during the work phase)
E_w	Number of exposed individuals working from work (during the work phase)
I_w	Number of infected individuals working from work (during the work phase)
R_w	Number of removed individuals working from work (during the work phase)
S_h	Number of susceptible individuals working from home (during the commute and work phases)

E_h	Number of exposed individuals working from home (during the commute and work phases)
I_h	Number of infected individuals working from home (during the commute and work phases)
R_h	Number of recovered individuals working from home (during the commute and work phases)

Table 2. Movement model parameters, meaning, and default value (with units).

Parameter	Meaning	Default values [Units]	Sensitivity analysis range
N	Population size	3000 [people]	(1500 to 6000)
R_0	Basic reproductive number (number of new infections that each infection generates)	2.5 [unitless] [39]	fixed
$R_{e\text{-}c}$	Effective reproductive number while commuting between work and campus	R_0 [unitless]	(1 to 4 R_0)
$R_{e\text{-}h}$	Effective reproductive number while at home	$0.5R_0$ [unitless] [40]	(0.25 R_0 to R_0)
$R_{e\text{-}w}$	Effective reproductive number while at work at campus	R_0 [unitless]	(0.5 R_0 to 2 R_0)
T_c	Fraction of a 24-hour day spent commuting each way for those that commute to campus	1/24 [unitless]	(0.5/24 to 2/24)
T_h	Fraction of a 24-hour day spent not working (everyone is off campus)	$= 1 - 2 T_c - T_w$ [unitless]	$= 1 - 2 T_c - T_w$
T_w	Fraction of a 24-hour day spent at work on campus for those commuting (some individual are on campus)	8/24 [unitless]	(2/24 to 12/24)
β_c	Transmission rate while commuting	$= \gamma\, R_{e\text{-}c}$ [day^{-1}]	$= \gamma\, R_{e\text{-}c}$

β_h	Transmission rate while at home	$= \gamma \, R_{e\text{-}h}$ [day^{-1}]	$= \gamma \, R_{e\text{-}h}$
β_w	Transmission rate while at work	$= \gamma \, R_{e\text{-}w}$ [day^{-1}]	$= \gamma \, R_{e\text{-}w}$
σ	Rate of moving from exposed to infected (inverse of the latent period)	1/4.2 [day^{-1}] [37]	(1/5.1 to 1/3.5)
γ	Recovery rate	1/9.5 [day^{-1}] [38]	(1/11 to 1/6)
θ	Fraction of the campus population commuting to work on campus (instead of continuing to work at home)	1 [unitless]	(0.0001 to 1)

S1 Fig. Movement model monotonicity plots.

772   The relationship between each of the nine model parameters (x-axis) and the model output, final epidemic

size (y-axis) for (a) population size (N); (b) fraction of a 24-hour day spent commuting each way for those

774   that commute to campus (T_c); (c) fraction of a 24-hour day spent on campus for those commuting (T_w);

(d) fraction of the campus population commuting to work on campus (θ); (e) recovery rate (γ); (f)

776   effective reproductive number while at home ($R_{e\text{-}h}$); (g) effective reproductive number while commuting

between work and campus ($R_{e\text{-}c}$); and (h) effective reproductive number while at work on campus ($R_{e\text{-}w}$).

S2 Fig. Movement model monotonicity plots.

780   The relationship between each of the nine model parameters (x-axis) and the model output, epidemic peak

size (y-axis) for (a) population size (N); (b) fraction of a 24-hour day spent commuting each way for those

782   that commute to campus (T_c); (c) fraction of a 24-hour day spent on campus for those commuting (T_w);

(d) fraction of the campus population commuting to work on campus ($\square$); (e) recovery rate (γ); (F)

784   effective reproductive number while at home ($R_{e\text{-}h}$); (g) effective reproductive number while commuting

between work and campus ($R_{e\text{-}c}$); and (h) effective reproductive number while at work on campus ($R_{e\text{-}w}$).

S3 Fig. Movement model sample numbers.

788   Absolute value of PRCC for the final epidemic size model output and each of the nine model parameters

(N, T_c, T_w, $\square$, σ, γ, $R_{e\text{-}h}$, $R_{e\text{-}c}$, $R_{e\text{-}w}$) as a function of different numbers of LHS samples generated. The

790   results seem to stabilize after about 1000 samples.

792   **S4 Fig. Movement model: limiting people, time and contact on campus.**

The epidemic peak size (maximum fraction of the population infected) as a function of (a) the fraction of

794   an 8-hour workday spent on campus (x-axis) and the fraction of the population working on campus (y-

axis) with no physical distancing, (b) the fraction increase in transmission while at work compared to at

796   home (x-axis) and the fraction of the population working on campus (y-axis) with an 8-hour work day, (c)

the fraction increase in transmission while at work compared to at home (x-axis) and the fraction of an 8-

798   hour workday spent on campus (y-axis) with 100% of people on campus.

800   **S5 Fig. Network model simulations**.

Simulations of pathogen spread across networks based on use of both shared office and lab space.

S6 Fig. Network model simulations.

804   Simulations of pathogen spread across networks based on use of only shared lab space.

806   **S7 Fig. Component-wise network structural metrics.**

Measures of the size (number of individuals), diameter (longest shortest path between two individuals),

808   and mean path length (average shortest path length between individuals) for each distinct component of

networks presented in Fig 5a,b. The combined lab and office network (blue points) has 8 distinct

810   components (8 points for each metric), while the shared lab space network contains 31 distinct

components (31 points for each metric).

S8 Supporting Information.

814   Sensitivity analysis of the network model and results.

a)

b)

c)

A) Combined Lab and Office
8 distinct components

B) Just Shared Labspace
31 distinct components

C) Distribution of shortest paths

Susceptible   Exposed   Infectious   Removed

Click here to access/download
Supporting Information
figS1.tiff

Click here to access/download

Supporting Information
figS2.tiff

Click here to access/download
Supporting Information
figS3.tiff

Click here to access/download

Supporting Information
figS4.tiff

Click here to access/download
Supporting Information
figS5.gif

Click here to access/download
Supporting Information
figS6.gif

Supporting Information Fig S7

Click here to access/download
Supporting Information
figS7.tiff

Click here to access/download

Supporting Information

S8 supporting_information_simulations.pdf