A SYSTEMATIC ANALYSIS OF THE TIME COURSE TO DEVELOP TREATMENTS FOR COVID-19

Binh T. Ngo, M.D.[1*], Marc S. Rendell, M.D.[2]

1. Keck USC School of Medicine
 2020 Zonal Avenue
Los Angeles, CA 90033

2. The Rose Salter Medical Research Foundation
34 Versailles
Newport Coast, CA 92657

ABSTRACT

Background. The spread of COVID-19 from Wuhan in China throughout the world has been frighteningly rapid. Epidemiologic techniques succeeded in containing the disease in China, but efforts were not as successful in the rest of the world, particularly the United States where there have been 1,592, 599 cases with 95,863 deaths as of May 25, 2020. Projections are for continued new infections and deaths if no effective treatments can be activated over the next six months. We performed a systematic review to determine the potential time course for development of treatments and vaccines focusing on availability in the last half of 2020.

Methods. Our search was performed during the week of May 25, 2020. Publications: We reviewed up to date information from several sources to identify potential treatments for COVID-19: We used the Reagan-Udall Expanded Access Navigator COVID-19 Treatment Hub to track the efforts of companies to develop treatments. We then used the results to search for publications on pubmed.gov and on MedRxiv, the preprint server. We further used a targeted Google search to find announcements of trial results.

Clinical Trials: We searched for all investigational trials begun in the first quarter of 2020 using several different sources: (A) covid-trials.org, then validated results on (B) clinicaltrials.gov and the (C) World Health Organization's International Clinical Trials Registry Platform (WHO ICTRP). We focused on trials which were completed or currently recruiting for patients, reasoning that the timeline to arrive at treatments by the end of the year would require completion within the next 6 months. We excluded studies which were clearly observational, with no randomization, control or comparison group. We fur-

ther set a cutoff of 100 for numbers of subjects since smaller trial size could lack statistical power to establish superiority of the intervention over the control condition.

Results. Published Data: We found 22 publications reporting findings on 6 classes of agents. There were 11 publications related to hydroxychloroquine (HCQ), three on lopanovir/ritonavir (LPV/R) (one of which was combined with interferon beta-1),four on remdesivir, two on favipiravir, one on meplazumab, and one on tocilizumab. Eleven publications reported randomized controlled trials, the rest were retrospective observational. Only one publication touched on outpatient management, the rest on hospitalized patients.

Clinical Trials: We found 409 trials meeting our minimum requirement of 100 subjects which were recruiting or completed. The WHO has launched the Solidarity megatrial performed in over 100 countries actively comparing HCQ, lopanovir/ritonavir (LPV/R) alone and in combination with interferon beta-1, and remdesivir. That trial is scheduled to complete enrollment in the first quarter of 2021. In addition, we found 46 trials of HCQ, 11 trials of LPV/R and 8 trials of interferons. There were 18 ongoing trials of antiviral agents, 24 immune modulator trials, 9 vaccine trials, and 62 trials of other agents. We excluded a large number of trials of Chinese traditional medications, reasoning that there was insufficient clinical experience with these agents outside China to offer these treatments to the rest of the world. Forty four trials were hoping to complete enrollment by the end of the second quarter of 2020. Of these, only 9 were conducted on outpatients. A few vaccine trials are hoping to complete Phase 3 enrollment by the end of the third quarter, but a prolonged follow-up of patients will likely be required.

Conclusion. Several treatments for severe disease in hospitalized patients have now been granted emergency authorization. However, the disease is propagated primarily by infected ambulatory individuals. There are only a few randomized controlled studies in outpatients which can be expected to yield results in time to impact on the continuing spread of the epidemic in 2020. It will be necessary for public health authorities to make hard decisions with limited data. The choices will be hardest in dealing with potential early release of vaccines.

Keywords: Coronavirus, COVID-19, SARS-Cov-2, remdesivir, hydroxychloroquine

*email: Binh.ngo@med.usc.edu

INTRODUCTION

The coronavirus SARS-Cov-2 pandemic spread through the world with extraordinary speed from its origin in December 2019 in Wuhan City in the Hubei Province of China. As of May 25, 2020, there were 5,307,298 confirmed cases of COVID-19, and 342,070 deaths worldwide affecting 220 countries and territories (1). The illness produces mild or no symptoms in about 80% of infected individuals but, in about 20% manifests as a significant pneumonia which can progress to acute respiratory distress requiring mechanical ventilation in 5% of patients (2). Baud et al calculated a crude mortality ratio of 5.7% using confirmed cases, but that calculation is affected by the high percentage of asymptomatic and mild undiagnosed cases (3). A calculation based on the outbreak among the passengers and crew of the cruise ship Diamond Princess, essentially a captive population, suggested a case fatality ratio of 2.6% and infection fatality ratio of 1.3% (4). However, the mean age of the passengers was about 60, and mortality is definitely higher in the elderly and in those with chronic disease.

COVID-19 is a disease of dual pathogenicity with a characteristic sequence of events related to the primary viral attack and then, subsequently, with an overlapping hyperinflammatory reaction (5). The initial viremia manifests as an influenza like illness with cough, fever, myalgia and fatigue. In patients with severe disease, within seven days of onset of symptoms, bilateral multiple lobular and subsegmental areas of consolidation are typically seen on chest CT (6). The pneumonia causes dyspnea by 8 days which may progress to acute respiratory distress syndrome (ARDS) by 9 days. Cytokine levels in the patients with ARDS are more elevated than in those patients who do not progress, pointing to a severe immune reaction as the probable cause (7,8). There is additionally a coagulopathy which affects the lungs and other organs and can trigger cerebrovascular accidents (9,10).

Recently a new syndrome of pediatric multisystem inflammatory disorder has surfaced in children, previously thought minimally susceptible to severe Covid-19 disease (11,12). This appears to be a delayed autoimmune condition which can occur a month or more after the SARS-Cov-2 infection and has been fatal in a number of children. As of May14th, 2020, New York state health officials were investigating 102 cases, and over 18 states have reported cases in the United States as well (13).

In mainland China, vigorous efforts to contain the spread of COVID19 by home isolation, closure of business activity, travel bans, and tracking and control of contacts succeeded. As of May 25, China had reported 82,985 confirmed cases and 4634 deaths, but the number of new cases has declined to very low levels (1). Similar measures contained the epidemic in Korea, Taiwan and in New Zealand (14). That has not been the case in the rest of the world, particularly the United States , where, despite only a fifth of the population of China, the number of cases was 1,592,599 with 95,863 deaths as of May 25, 2020 (1). In the U.S.A., as in China, the planned response to the pandemic was through reduction of person to person contact. The process of social distancing began March 11, 2020. Schools and businesses were closed. Social gatherings and meetings ended. Recreation areas, parks, and beaches were closed. These efforts have blunted the spread of the disease, but not with the results achieved in China.

Epidemiologic models have varied on prediction of the continued spread of COVID-19 in the United States. The Insttute for Health Metrics and Evaluation IHME projected that the peak of deaths occurred in midApril with lessening by May 1, predicated however on maintenance of strict social distancing measures (15). Kissler et al performed simulations of the persistence of COVID-19 in the United States (16). Their model considered known data on the transmissibility of other less threatening betacoronaviruses, crossimmunity with those viruses, seasonal variations, and the impact of social distancing. They predicted that the peak of infections would occur in May with significant reduction by July. However, a significant resurgence of COVID-19 was then expected to follow the end of social distancing with secondary peaks occurring into 2021. The optimal scenario they simulated was for indefinite social distancing measures, which even early on has triggered massive unemployment and economic collapse in the United States. They stated that medications or vaccines for COVID-19 would obviously reduce the need for reliance on intensive social distancing.

Faced with a growing crisis in early January, Chinese health authorities began treating patients empirically with agents with demonstrated in vitro antiviral activity against coronaviruses and those used during the SARS 2002 and MERS outbreaks in China (17,18). The medications used included the antimalarial drugs hydroxychloroquine (HCQ) and chloroquine, the HIV protease inhibitor lopanovir-ritanovir (LPV/R), the Russian antiviral umifenovir, and traditional Chinese medical approaches. Subsequently, the antimalarial drugs hydroxychloroquine (HCQ) and chloroquine have been widely used to treat COVID-19 patients throughout the world, despite cautions from numerous regulatory bodies. The scientific basis for their use is in vitro evidence of an effect on blocking viral endosomal penetration as well as their known suppressive benefit of undesirable autoimmune effects (19). Thus far, the clinical evidence supporting the use of antimalarials for COVID-19 is limited (20).

In view of the projections for resurgence of infections if current social distancing measures are to be relaxed, our goal was to assess the projected timeline for development and release of pharmacologic agents to treat and to prevent COVID-19 over the next 6 months.

METHODS

Our search was carried on the week of May 25, 2020.

PUBLICATIONS: We reviewed up to date information from several sources to identify potential treatments for COVID-19: We used the Reagan-Udall Expanded Access Navigator which is a joint effort of the Reagan Udall Foundation and the FDA to track the efforts of companies engaging in research to enable access to investigational treatments for individuals who are unable to find or participate in an existing clinical trial. To address the global pandemic of COVID-19, the Foundation enhanced the Expanded Access Navigator to include a comprehensive COVID-19 Treatment Hub. We then identified all publications on proposed treatments using pubmed.gov to find peer reviewed articles but in addition we reviewed medRxiv to find preprint reports. We further carried out Google searches on each poten-

tial treatment to find preliminary reports, typically presented as press releases. We have continued to add any new information forthcoming based on daily searches since May 25.

CLINICAL TRIALS: We searched for all investigational trials involving potential treatments using several different sources:

(A) covid-trials.org This is a registry to collate all trials in real time (21). Data are pulled from the International Clinical Trials Registry Platform, including those from the Chinese Clinical Trial Registry, ClinicalTrials.gov, Clinical Research Information Service - Republic of Korea, EU Clinical Trials Register, ISRCTN, Iranian Registry of Clinical Trials, Japan Primary Registries Network, and German Clinical Trials Register. Both automated and manual searches are done to ensure minimal duplicated entries and appropriateness to the research questions. Identified studies are then manually reviewed by two separate reviewers before being entered into the registry. Concurrently, the registry has developed artificial intelligence (AI)-based methods for data searches to identify potential clinical studies not captured in trial registries.

We cross validated information from covid-trials.org on (B) clinicaltrials.gov This is a registry of clinical trials information maintained by the United States National Library of Medicine for both federally and privately funded trials conducted under investigational new drug applications to test the effectiveness of experimental drugs. Data includes trials in the United States and 210 countries. We further cross referenced the trials on (C) the World Health Organization's International Clinical Trials Registry Platform (WHO ICTRP).

We determined which agents were currently available for clinical use, those with published trial results, those which were in ongoing active randomized clinical trial investigation, those which were in observational trials, those in trials projected to begin but not yet recruiting, agents in preclinical phase, and agents in planning phase. We than reviewed each trial to determine if the trials were completed, recruiting for patients, not recruiting or suspended. We only selected those which had completed or were actively recruiting by May 1, 2020, reasoning that the timeline to implement treatments by the end of the year would require completion within the next 3 months. We excluded studies which were clearly observational for treatment purposes, with no randomization, control or comparison group. We further set a cutoff of 100 for numbers of subjects since smaller trial size could lack statistical power to establish superiority of the intervention over the control condition. For each trial selected, we documented the setting of patient contact, either hospital or outpatient, the intended severity of disease in the trial population, the type of control procedure, the date the trial was initially registered, and the proposed date of completion of enrollment. We further reviewed all inclusion and exclusion requirements to verify intended severity of disease. We focused on any inclusion requirements based on duration of symptoms. We categorized the studies as either prevention (PREV) of SARS-Cov-2 infection in healthy subjects or treatment of established COVID-19 disease (TREAT).

RESULTS

PUBLISHED RESULTS ON COVID 19 TREATMENT. We first searched for all publications reporting on completed investigations of pharmaceutical treatment of COVID-19. We searched on pubmed.gov as well as MedRxiv, the preprint server. As of May 29, we found 22 publications reporting findings on 6 classes of agents (TABLE 1). There were 11 publications related to hydroxychloroquine, three on lopanovir/ritanovir (LPV/R), four on remdesivir, two on favipiravir, one on meplazumab, and one on tocilizumab. Ten of the publications reported randomized controlled or active comparator trials, the rest were observational.

The largest published randomized controlled trials were with remdesivir, an inhibitor of viral RNA polymerase. Two remdesivir trials were randomized controlled placebo trials. The first multicenter trial at ten hospitals in Hubei, China trial was halted in April due to failure to further enroll since no new cases were occurring (22). Eligible patients were adults (aged ≥18 years) admitted to the hospital with laboratory-confirmed SARS-CoV-2 infection, with an interval from symptom onset to enrollment of 12 days or less, radiologically confirmed pneumonia, and hypoxia. Patients were randomly assigned in a 2:1 ratio to intravenous remdesivir (200 mg on day 1 followed by 100 mg on days 2–10 in single daily infusions) or the same volume of placebo infusions for 10 days. Patients were permitted concomitant use of LPV/R, interferons, and corticosteroids. The trial enrolled 237 patients when it was suspended due to lack of recruitment. Remdesivir use was not associated with a difference in time to clinical improvement (hazard ratio 1·23 [95% CI 0·87–1·75]).

A report of compassionate use of remdesivir found an overall mortality of 5% in patients not requiring mechanical ventilation but 18% in those who did (23). The largest trial of remdesivir in 1063 moderate to severely ill patients was a double blind randomized placebo controlled study carried out by the National Institute of Arthritis and Infectious Disease (NIAID). On April 29, Dr. Anthony Fauci, Director of the NIAID, announced that the study had met its endpoint in that a faster time to clinical improvement and discharge was found with remdesivir treatment. On May 22, 2020 the actual trial results were published (24). The patients who received remdesivir had a median recovery time of 11 days (95% confidence interval [CI], 9 to 12), as compared with 15 days (95% CI, 13 to 19) for those who received placebo (rate ratio for recovery, 1.32; 95% CI, 1.12 to 1.55; P<0.001). The Kaplan-Meier estimates of mortality by 14 days after randomization were 7.1% with remdesivir and 11.9% with placebo (hazard ratio for death, 0.70; 95% CI, 0.47 to 1.04). There was no data documenting the rate of viral clearance. On the basis of this trial, the FDA issued an emergency authorization to use remdesivir for severely ill COVID hospitalized patients. Subsequently, a report was published showing that in 397 hospitalized patients with moderate to severe disease, there was no difference in outcome comparing patients who received a 5 day course of remdesivir versus a 10 day course. The analysis was complicated by an imbalance in clinical severity; the patients randomized to 10 days of treatment had significantly more severe disease than those in the 5 day group. The 14 day mortality in the 5 day group was 8% versus 11% in the 10 day group. There was no data on viral clearance reported.

There were two trials of favipiravir (FVP, which like remdesivir is an inhibitor of viral RNA polymerase. Favipiravir is given as an oral tablet whereas remdesivir only has an intravenous route of administration. Both FVP trials were active control studies carried out in China (TABLE 1). In a study of 35 patients receiving favipiravir compared to 45 patients receiving LPV/R plus inhaled interferon alpha, a shorter viral clearance time was found for the FVP arm (median (interquartile range, IQR), 4

(2.5–9) days versus 11 (8–13) days, P < 0.001) (26). The FVP arm also showed significant improve-ment in chest imaging compared with the control arm, with an improvement rate of 91.4% versus 62.2% (P = 0.004). In a larger Chinese trial comparing 116 patients on FPV to 120 on umifenovir, the 7 day clinical recovery rate was 55.9% for the umifenovir group and 71% in the FVP group (27).The Chinese Health Ministry, after reviewing the studies, concluded that favipiravir was efficacious against Covid-19. In Russia, a state sponsored agency confirmed the response to favipiravir in a still unpub-lished study and has approved the drug for clinical treatment beginning on June 12.

Published studies on hydroxychloroquine (HCQ) have been less conclusive. In a small study in Mar-seille, France, COVID-19 patients were included in a single arm protocol to receive 600mg of HCQ daily, with daily viral load testing in nasopharyngeal swabs (28). Depending on their clinical presenta-tion, azithromycin (AZT) was added to the treatment. Untreated patients from another center and cases refusing the protocol were included as negative controls. Presence and absence of virus at Day6-post inclusion was considered the end point. Twenty cases were treated in this study and showed a signifi-cant reduction of the viral carriage at D6-post inclusion compared to controls, and much lower average carrying duration than reported for untreated patients. AZT added to HCQ was significantly more effi-cient for virus elimination.

The same group has subsequently reported their experience with 1061 patients screened positive for SARS-Cov-2 and then treated with HCQ and AZT (29). Patients and asymptomatic contacts with posi-tive viral tests were included. There was no control population. After evaluating electrocardiograms to exclude patients at risk of QT prolongation, they used a combination of HCQ and AZT. Viral clearance was measured at days 2,6, and 10 of treatment and further for patients who were deemed poor clinical responders. The patients were young (mean age 43.6). Good clinical outcome with successful viral clearance by ten days was obtained in 91.7% of the patients. Failure to clear the virus by ten days oc-curred in 47 patients (4.4%). Forty-six patients (4.3%) were classified as poor clinical response due to 10 patients transferred into the ICU and 30 who were hospitalized for ten days or more. There were 10 patients who died (mean age 79) due to respiratory failure giving a case fatality ratio of 0.9%.There were no sudden cardiac deaths.

In a small trial at Renmin Hospital of Wuhan University in a parallel-group randomized trial, 31 pa-tients were assigned to receive 5-days of HCQ (400 mg/d) treatment (30). The body temperature recov-ery time and the cough remission time were significantly shortened in the HCQ treatment group, and 81% (25 of 31) had improvement in pneumonia compared with the control group (54.8%, 17 of 31). All 4 patients who progressed to severe illness were in the control group. However, two other Chinese studies showed no difference in viral clearance in HCQ treated subjects (31,32).

There have been many retrospective large observational trials which suggest that hospitalized patients on HCQ have worse outcomes (33-37) (Table 1).   The largest retrospective observational study was a multinational registry analysis using Surgical Outcomes Collaborative, a data acquisition and ware-housing cloud based system which collected electronic data from 671 hospitals in six continents, in-cluding patients hospitalized between Dec 20, 2019, and April 14, 2020 (37). The analysis com-

pared 14,888 patients who were treated with hydroxychloroquine, chloroquine, or their combination with a macrolide within 48 hours of diagnosis of COVID-19. The control group consisted of 81,144 diagnosed patients who did not receive any of these agents. Using propensity scoring to achieve comparability of baseline characteristics, they found higher mortality in all of the groups receiving antimalarial agents compared to control. They also found a higher incidence of ventricular arrhythmias, a known side effect of HCQ and chloroquine. The difficulty in analyzing retrospective observational data is in the possibility of a selection bias choosing patients who were sicker for treatment. Propensity scoring can help adjust for bias, and they claimed comparability of groups after adjustment. However, in a different study in New York(33), even after adjustment, the (Pao2:Fio2 (223) of HCQ patients was lower than that for the propensity scored non-HCQ patients (273).

In a retrospective comparison study at NYU Langone Health for all patients admitted between March 2 and April 5, 2020 (38), there were a large number of patients treated with HCQ plus AZT. During the one month period, there was a move to add zinc 100 mg daily to the HCQ plus AZT regimen. Zinc supplementation has shown benefit in viral diseases both in inhibiting viral replication and in improvement of systemic immunity (39). Zinc administered within 24 hours of onset of symptoms has been claimed to reduce the duration of common cold symptoms in healthy people (40). There were 411 patients treated with HCQ+AZT+ zinc at NYU. There were only minor differences in the clinical characteristics of these patients compared to the 521 patients on HCQ and AZT who did not receive zinc. After adjustment for the time period when zinc first came to frequent use, there was a significantly lower mortality (13.1%) among the zinc treated patients compared to those who did not receive zinc (22.8%).

There is one reported prevention study with HCQ given for 5 days to individuals with a significant exposure to SARS-Cov-2 (41). The incidence of new illness compatible with Covid-19 did not differ significantly between participants receiving hydroxychloroquine (49 of 414 [11.8%]) and those receiving placebo (58 of 407 [14.3%]); absolute difference −2.4 percentage points (95% confidence interval, −7.0 to 2.2; P=0.35).

Published results with Lopanovir/ritonavir (LPV/R) have been disappointing. A small Chinese trial of 44 patients found no difference in viral clearance compared to standard of care (42). In a randomized trial with 99 patients on LPV and 100 receiving standard care, there was no difference in time to clinical improvement, mortality or viral clearance (43). A potentially more favorable result was obtained in Hong Kong in a study of 86 patients assigned to triple combination therapy with LPV/R plus ribavirin plus interferon beta-1 (44). Compared to a control group of 41 patients receiving LPV/R alone, the combination group had a significantly shorter median time from start of study treatment to negative SARS-Cov-2 nasopharyngeal swab (7 days [IQR 5–11]) than the control group (12 days [8–15];hazard ratio 4·37 [95% CI 1·86–10·24], p=0·0010). The time to complete alleviation of symptoms was 4 days [IQR 3–8] in the combination group vs 8 days [7–9] in the control group; HR 3·92 [95% CI 1·66–9·23. The timing of interferon administration was staggered based on time of symptom onset, so there were 34 patients in the combination group who did not receive interferon at all. Those patients did not show clinical or virological superiority to the LPV/R control group. This study was interpreted as supporting further trials focusing on interferon beta-1 as the primary treatment modality.

A very small pilot study of meplazumab, a monoclonal antibody to CD147, in 17 patients showed a marked improvement in viral clearance time from 13 days (6.5-19.5) in a control group of 11 patients to 3 days (1.5-4.5). This was accompanied by clinical and radiographic improvement (45).

Very hopeful data has emerged for the IL-6 inhibitor tocilizumab in the severe and critically ill phase of the disease. An observational study in China suggested dramatic improvement in 21 patients with abatement of fever, decrease in oxygen requirement and reduction in lung lesions (46). There was no control group in that study, but it was recently announced that tocilizumab treatment had succeeded in the randomized, controlled CORIMUNO-19 – TOCI trial for severely ill COVID-19 patients in France. On the basis of the French still unpublished findings, the CDC issued an emergency use authorization for tocilizumab.

REVIEW OF CURRENT TRIALS: As of May 25, we identified 781 trials worldwide meeting our review criteria. Approximately half the trials were registered in mainland China, and 103 trials were registered in the United States (TABLE 2). There were 553 trials in the recruitment process or completed (SUPPLEMENTARY TABLE 1). We found 409 trials which were recruiting or completed meeting the minimum requirement of 100 subjects. The World Health Organization has organized a megatrial appropriately named Solidarity to assess four separate treatment options: Remdesivir; LPV/R; LPV/R with Interferon beta-1a; and HCQ or chloroquine (47). The Solidarity Trial provides simplified procedures to randomize patients to one of the four agents or alternatively standard care. Brief identifying details and any other conditions are digitally recorded for the patient, who is then randomly allocated to one of the study options. Critical study information is to be collected only at randomization and when the patient is discharged or dies: which study drugs were given (and for how many days); whether ventilation or intensive care was received (and, if so, when it began), date of discharge, or date and cause of death while still in hospital. Data is transmitted centrally to the trial database. There are over 100 countries now participating. Completion of enrollment of 3,100 patients is estimated in the first quarter of 2021. Very recently, the antimalarial drug arm of the study was paused after review of the observational studies previously mentioned pending an interim safety analysis. The Solidarity safety monitoring board then reviewed their own safety data and within a few days found no reason to interrupt randomization to HCQ, so that arm was just restarted

OTHER TRIALS WITH THE SOLIDARITY AGENTS: In addition to and independent from the Solidarity trial, we found 46 trials of HCQ, 11 trials of LPV/R and 8 trials of interferons (TABLE 3). Of these, ten of the HCQ trials focused on prevention of the disease in exposed patients. Only two of these 10 prevention trials proposed completion of enrollment in the second quarter of 2020. There were seven HCQ trials conducted on ambulatory patients with diagnosed mild disease. Only two of these outpatient trials had proposed enrollment completion in the second quarter. There were 21 HCQ trials focused solely on hospitalized patients. Four of these trials had planned enrollment completion in the second quarter of 2020.

Two prevention trials with LPV/R are currently enrolling, with earliest enrollment completion in the second quarter of 2020 for one and the third quarter for the other. There is one outpatient active comparison trial with HCQ with proposed enrollment end in the second quarter of 2020. There are two large active comparison trials with earliest enrollment end date in the first quarter of 2021. There is one large interferon prevention trial scheduled to complete enrollment in the second quarter of 2020.

ANTIVIRAL AGENTS: SARs-CoV-2 attacks cells by binding of the spike protein to the host cell ACE2 receptor. A specific serine protease TMPRSS2 then activates membrane fusion allowing viral RNA to enter the cell to be replicated with release of multiple virions. Numerous antiviral agents are now in active trials (TABLE 4). In addition to remdesivir and favipiravir, trials are ongoing with umifenovir, a membrane lipid fusion inhibitor with broad antiviral properties (48) and with camostat which is a potent serine protease inhibitor (49). A very large multicenter prevention trial has begun for hospital workers exposed to COVID-19 in Spain. This involves 4000 subjects who will be randomized to emicitrabine/tenofovir, an HIV agent, or to HCQ or to placebo. The trial has planned to meet its enrollment goal by the beginning of the third quarter of 2020.

IMMUNE MODULATORS: It is clear that the SARS-Cov-2 infection triggers an immune reaction that leads to the most serious consequences of the disease. Many trials are focusing on several interleukin inhibitors currently approved for treatment for rheumatoid arthritis and other autoimmune conditions to suppress the cytokine storm of COVID-19 (TABLE 5). Tocilizumab has just received emergency authorization. Selinexor has planned completion of enrollment by the end of the second quarter as well. A trial of sarilumab which will run until 2021 recently completed its first phase and announced success in reducing CRP. These trials are all directed at severely ill COVID-19 inpatients. There is a large multicenter placebo-controlled trial of colchicine in non-severely ill outpatients scheduled to enroll 6000 patients by end of the third quarter of 2020.

OTHER STUDIES (SUPPLEMENTARY TABLE 2):

CONVALESCENT PLASMA: Several studies of intravenous globulin as well as intravenous hyperimmune globulin plan completion of enrollment in the third quarter of 2020.

STEM CELLS: Several trials of mesenchymal stem cells are currently recruiting with one study planning completion of enrollment by the end of the second quarter of 2020.

RENIN ANGIOTENSIN SYSTEM: Recognition of the role of the ACE2 receptor in SARS-Cov-2 cell entry has speculated interest that ACE inhibitors or angiotensin receptor blockers might block viral access. There are a number of trials either adding the agents or withdrawing them. One small trial of losartan has been fully enrolled but no results have been posted as yet.

PULMONARY AND RESPIRATORY INTERVENTIONS: These studies are of great interest in managing the ARDS associated with COVID-19. There are many ongoing studies of mechanical and extra-corporeal oxygenation. Most of the studies did not meet the criterion of 100 subjects or more (SUPPLEMENTARY TABLE 1). There was one ongoing study of hyperbaric oxygenation but not due to complete enrollment until the second quarter of 2021. Pirfenidone is an agent approved to treat pulmonary fibrosis, and one study is planning completion of enrollment by the end of the second quarter of 2020.

SUPPLEMENTS: The suggestion that zinc may be a beneficial agent together with HCQ in COVID has generated interest, but there is only one study underway, not including HCQ. A study of HCQ with and without zinc for prevention of COVID infection is soon to begin in Tunisia.

GLUCOCORTICOIDS: A number of studies are trying to determine if steroid use can prevent serious autoimmune consequences without impairing the response to the virus.

ANTICOAGULANTS: Several studies are focusing on prophylaxis for the COVID-19 hypercoagulable state. There are no current studies of the new direct oral anticoagulants.

TRADITIONAL CHINESE MEDICATIONS: There were 48 such studies. We did not include them in our search for agents which could be made available in the next 6 months for two reasons. First, these medications are not familiar to clinicians in the rest of the world. Second, the success of epidemiologic containment in China has made it unlikely that many projected trials would achieve enrollment.

VACCINES: The World Health Organization has identified 110 programs to develop a SARS-Cov-2 vaccine. Of these, seven have reached initial human trials (Table 5). There are four programs active in China, one in the United Kingdom and two in the United States. The conventional wisdom is that it will take a year to 18 months to implement a vaccination program. Clinical vaccine development is typically a multi-year three-phase process. During Phase I, small groups of people receive the trial vaccine. In Phase II, the clinical study is expanded and vaccine is given to people who have characteristics (such as age and physical health) similar to those for whom the new vaccine is intended. In Phase III, the vaccine is given to thousands of people and tested for efficacy and safety. SARS-Cov-2 vaccine developers have approached the regulatory bodies to request acceleration of the process. Optimistic observers have suggested that a vaccine could be available by the beginning of 2021. A bold suggestion has been made to allow emergency authorization of favorable vaccine candidates by the fall. An additional approach has been suggested to attempt to stimulate non-specific immunity prior to availability of a vaccine using oral polio vaccine or BCG vaccination (50). There are two large prevention trials underway in the Netherlands and Australia using BCG vaccine in health care workers. These trials plan conclusion of enrollment, but not until the fourth quarter of 2020.

DISCUSSION

SARS-Cov-2 overwhelmed medical preparedness for the pandemic. Although epidemiologic containment was achieved in China, Korea,Taiwan, and New Zealand, efforts to stop the spread failed in much of the rest of the world. The United States, Brazil, and Europe have been particularly hard hit. In the United States, there were 10,111 new cases and 241 new deaths on May 24, at a time when numerous states and the federal government have relaxed social distancing restrictions. Kissler et al modeled several scenarios of transmission with several periods of variably effective social distancing (16). We reproduce their graphic from Science below (FIGURE 1). Panel B appears to represent the current situation with a relaxation of social distancing, which has begun in midMay. Their projection is for a continuation of new cases and deaths occurring through July, with a possible resurgence in October.

The current pharmaceutical options available in the United States have been limited. In China, it is not the practice to run placebo controlled trials with investigational agents with no alternative choices for the patient. Consequently, their "standard of care" often includes many potentially effective agents, possibly confounding results. Furthermore, their guidance to physicians includes many options which are under active study in certain sites. In contrast to China, in the United States and Europe, there is an emphasis on randomized placebo controlled clinical trials to decide which agents should be widely used for treatment. Those trials are ongoing, but we found that only 40 of 164 trials are proposed to complete enrollment in the second quarter of 2020. It typically may take two months or more to follow patients to completion, verify and analyze data, and our goal was to find treatments to be used during the next six months. In the United States, HCQ, remdesivir, and tocilizumab are currently available on an emergency use basis only for hospitalized severely ill patients. In Russia, it is planned to begin administering favipiravir to patients beginning June 12. There are other large programs now focusing on immune modulators to prevent ARDS due to cytokine storm in hospital. Most of the agents involved are marketed drugs used to treat autoimmune diseases. Therefore, these treatments have known safety profiles and only require proof of benefit in severe and critically ill COVID-19 patients. Tocilizumab has been successful in a French trial. It is hoped that the availability of remdesivir and success of the immune modulators will improve survival in hospitalized severely ill patients over the next few months.

Studies to date have understandably focused on hospitalized patients to try to prevent death. However, rescue from mortality is not the only goal in fighting COVID-19; it is essential to reduce the spread of the virus. Once patients are hospitalized, they only contribute minimally to the development of new cases, primarily in hospital workers. Community not hospital transmission is the major mode of propagation of SARS-Cov-2. We face a dilemma in having no currently approved prevention or treatment for outpatients with mild to moderate disease, the chief spreaders of SARS-Cov-2. Shortages of testing supplies in the United States have thus far limited the ability to positively diagnose the disease, resulting in the recommendation of home quarantine for the majority of patients with suspicious symptoms. Those shortages are being rapidly remedied. Now, there is a need for therapy which can be used to treat outpatients with positive COVID-19 tests. The Marseille group has treated everyone with a positive test with a combination of HCQ and AZT and provided substantial clinical and virologic data but without a

control group. The NYU group treated all hospitalized patients with HCQ and AZT and showed an impressive effect of addition of zinc sulfate, but that was not in mild to moderate disease. They also had no control group for HCQ plus AZT. Remdesivir will be used primarily in severe disease and is administered intravenously, so not a practical outpatient treatment. Favipiravir is an oral tablet, like remdesivir an inhibitor of RNA polymerase, and is now about to be released for treatment in Russia, but not in the United States. It has known embryogenic risks, so its use requires restrictions on women of childbearing potential, as is the case with tretinoin for acne and thalidomide for multiple myeloma. Trials with camostat, the serine protease inhibitor, capable of blocking cell entry of SARS-Cov-2, are not due to end till the third or fourth quarter of the year.

The vaccines have been the principal hope held out by public health authorities in the United States. Progress on vaccine development has been extremely rapid. Moderna prepared the sequence of their proposed mRNA-1273 vaccine on January 13 only two days after China shared the genetic sequence of SARS-Cov-2. On February 7, 2020, the first clinical batch of vaccine was completed and shipped on February 24 to the NIH for use in their Phase 1 clinical study. On March 16, 2020, the NIH announced that the first participant in its Phase 1 study was dosed. On May 7, a successful antibody response was documented. The Company reported that while a commercially-available vaccine is not likely to be available for at least 12-18 months, it is possible that under emergency use, a vaccine could be available to some people, possibly including healthcare professionals, in the fall of 2020.

In the United Kingdom, the Oxford vaccine ChAdOx1 nCoV-19, an attenuated adenovirus expressing the SARS-Cov-2 spike protein, reduced the viral load in a monkey model. A Phase II study involving 1090 healthy volunteers is underway. The INO-4800 DNA vaccine is in Phase I testing. In China, there are two inactivated SARS-Cov-2 vaccines in Phase I testing. One of the vaccines PiCoVacc has protected monkeys from infection (51). Cansino is advancing its recombinant adenovirus vector vaccine Ad5-nCoV to Phase II testing. The rapid progress so far does not reduce the challenges of developing a vaccine to be administered to hundreds of millions of people (52). Although neutralizing antibodies can be produced by the vaccine candidate, the goal is to prove protection from the infection. The demonstration of protection requires Phase III observation and long term follow-up of an exposed vaccinated population. The length of follow-up depends on the degree of active transmission of SARS-Cov-2 during the post vaccination period. No matter how rapidly a vaccine advances to Phase III testing, the duration of follow-up cannot be shortened. In the case of SARS-Cov-2, that follow-up will have to be prolonged due to two factors. First, there is a risk of immune enhancement of infection which occurs when induced antibodies increase entry and internalization of virus into myeloid cells (53). This major complication struck the newly developed dengue vaccine in 2017 (54,55). Second, the sinister autoimmune pathogenicity of SARS-Cov-2 as evidenced by multisystem inflammatory disorder in children raises the risk of delayed long term harm if the virus is not fully and immediately eradicated by the initial vaccine associated immune response. Paradoxically, the large Chinese population is unlikely to serve to test any vaccine due to the success of COVID-19 containment in China

For these reasons of needed long term safety observation, the deployment of an effective vaccine could be delayed for more than the 12-18 months currently contemplated. By that time, the primary pandemic

may well be over due to infection of most of the world's population and ensuing herd immunity. The vaccine would then serve primarily to prevent a new resurgence. An alternative approach to immunization now is to increase the non-specific immune reaction to the initial infection. This suggestion was first publicized in an opinion publication by Drs. Chumakov and Gallo in USA Today. They advocated vaccination with oral polio vaccine (50). Oral polio vaccine is no longer as freely available as their alternative suggestion of BCG vaccine. BCG vaccine has demonstrated heterologous enhancement of immunity (56,57). There are very large prevention trials now being conducted in the Netherlands and Australia using BCG vaccination to increase innate immunity to SARS-Cov-2 in health care workers, but those are not due to complete until the fourth quarter of 2020.

COVID-19 has shaken societies throughout the world. The disease has imposed harsh opposing choices. The Chinese have shown that strict epidemiologic control is possible, but their approach has not been fully successful in Western countries where two months of social distancing and quarantine has devastated the economies. Yet, the alternative is continued spread of the virus by person to person contact. Similarly, there has been a contrast in the medical response comparing China to the Western countries. In China, empirical treatments have been offered and officially endorsed by the China National Health Commisson without proof of efficacy (58). The treatment options they recommend include alpha interferon, LPV/R, chloroquine and umifenovir. They included tociluzimab therapy for patients with cytokine storm prior to the announcement of the CORIMMUNO-TOCI results. They also include many Chinese traditional medications. In comparison, the most recent CDC guidelines developed together with an NIH panel for treatment of COVID-19 now include only remdesivir as well as tocilizumab for severe disease. The U.S. panel recommends against the use of remdesivir for mild to moderate disease outside of a clinical trial. They find no evidence for or against the use of antimalarial drugs and strongly recommend against hydroxychloroquine combined with azithromycin. They strongly recommend against the use of LPV/R and other protease inhibitors. The NIH expert panel has further recommended against the use of interferon. They have no recommended treatments for outpatients.

Ironically, the Western world has much stricter requirements for acceptance of clinical treatments than does the highly regimented Chinese society. That is only one of many incongruities which are affecting the time course of delivering treatments to arrest the spread of SARS-Cov-2. Paradoxically, the success of the Chinese in strict epidemiologic control has jeopardized the completion of the randomized controlled trials begun in China. The remdesivir trials were suspended in China due to failure to accrue new infected patients. Since so many trials were ongoing in China the lack of patients may impair the ability to achieve statistical power to complete the studies. Similarly, the Chinese efforts to develop effective vaccines are affected by the need to demonstrate effectiveness in a population with ongoing exposure to COVID-19 which they no longer have.

On May 25, as this manuscript is being written, there were 100,733 new cases of COVID-19 and 4290 deaths worldwide, and in the United States, there were 24,151 new cases with 1852 deaths (1). Many clinical trials are currently underway. However, the transmission of SARS-Cov-2 has outraced the pace of development of new therapies. There is a need to offer treatment to infected individuals now. HCQ

remdesivir, and tocilizumab have received emergency use authorization for severely ill patients. Hopefully, the mortality of COVID-19 will now be significantly reduced, but that alone will not suppress new infections and propagation of the disease. Hospitalized patients must be rescued, but the goal must be to prevent deterioration requiring hospitalization.

Efforts must now focus on treatments to interfere with the community transmission of the virus. The Marseille study treating patients early at the time of positive diagnosis of SARS-Cov-2 suggests a new emphasis on outpatient management. Favipiravir offers an oral counterpart to remdesivir to block viral propagation prior to hospitalization. The Hong Kong study suggests that interferon beta-1 which can be administered to outpatients has significant benefits in reducing viral replication. The success of oral zinc supplementation in the NYU study offers a simple treatment with minimal potential harm. Of course, none of these available treatments has been proven successful in randomized placebo controlled studies There are strong arguments to avoid emergency use of agents until trials are completed and analyzed, but the agents being tested are not new (59). Most of them are drugs approved for other conditions with well-known safety profiles, like camostat, a Japanese serine protease inhibitor, used for pancreatitis. BCG vaccination is still the norm for children in several countries. The conflict between clinical needs and the demand for rigorous proof of efficacy has resulted in curious events such as the President of the United States announcing that he is taking HCQ and zinc to prevent infection while the CDC and NIH have recommended against such treatment.

The projections suggest continued infections and deaths with a possible serious resurgence as social distancing restrictions are lifted. The Penn-Wharton Budget Model in the United States predicts that removing social distancing restrictions could result in 200,000 more deaths by June 30, but, conversely, save 18,000,000 lost jobs (60). We face these dismal choices if we lack the use of medications to prevent and to treat SARS-Cov-2 infections early before hospitalization. Epidemiologic containment of COVID-19 has not succeeded in the United States and most other countries. With social distancing measures now being relaxed, a large number of new cases and deaths will occur if treatments are not forthcoming in the near future. Several treatments of severe disease in hospitalized patients have now been granted emergency authorization. However, the disease is propagated primarily by infected ambulatory individuals. We have documented that there are only a few randomized controlled studies in outpatients which can be expected to yield results in time to impact on the continuing epidemic in 2020.

There is a precedent to resolve the conflict between immediate clinical needs and the requirements for rigorous controlled trials to prove efficacy. We should remember that the greatest success in fighting a pandemic occurred over the past two decades in the battle against the HIV virus which causes AIDS. AIDS was first recognized in 1981 in the gay community (61). The disease was a death sentence. By 1985, over 20,000 diagnoses and almost 10,000 deaths were recorded. There was widespread fear because there was no treatment, and projections of infection escalated into the millions. The first AIDS remedy was azidothymidine synthesized in 1964 in the hope that it would combat cancer. It was not successful. Twenty years later Dr. Samuel Broder, head of the National Cancer Institute, showed that the drug had activity against the HIV virus in vitro (62). Burroughs Wellcome launched a rapidly conceived trial with just 300 patients. They stopped the trial in 16 weeks claiming that more patients sur-

vived on azidothymidine. The FDA came under enormous pressure from AIDS activists to make the drug available. Azidothymidine was approved by the FDA on March 19, 1987 with only that one trial. It had taken 20 months for the FDA to give approval to release the drug. To this day, the design and results of the trial remain controversial.

The gay community continued to battle for early release of other medications to combat the AIDS pandemic. On October 11, 1988, a massive protest occurred at the FDA. It was back then Dr. Anthony Fauci who publicly advanced the idea of a parallel track to make drugs widely available even while studies are progressing: " Clearly, the standard approach to the design of clinical trials—that is, rigid eligibility criteria as well as the strict regulatory aspects that attend clinical trial investigations and drug approval—was not well-suited to a novel, largely fatal disease such as this with no effective treatments, and we had many intense discussions about how to make that approach more flexible and ethically sound. One example, which I and others worked closely with the AIDS activists to develop, was called a parallel track for clinical trials. The parallel track concept, which the United States Food and Drug Administration ultimately came to support, meant that there would be the standard type of highly controlled admission criteria and data collection for the clinical trial of a particular drug. In parallel, however, the drug also could be made available to those who did not meet the trial's strict admission criteria but were still in dire need of any potentially effective intervention, however unproven, for this deadly disease" (63).

The parallel track advocated by Dr. Fauci was adopted. Today there are 41 drugs or combinations approved by the FDA to treat and to prevent HIV infection. There is still no vaccine. There are now an estimated 1.1 million patients with HIV in the United States. Most have enjoyed near normal life expectancy thanks to the antiviral agents. The CDC has contributed greatly to limit the spread of HIV by advocating safe sex practices, but social distancing is not the norm for HIV. Rather pre-exposure prophylaxis endorsed by the CDC with a daily antiviral combination pill is currently adopted in wide segments of the at risk population (64).

Dr. Fauci has recently launched an NIAID randomized controlled study of the HCQ plus AZT combination. Interferon beta is one of the randomized controlled Solidarity Trial options. Favipiravir is in trials at Massachusetts General Hospital and Japan now. A randomized prevention trial with zinc and HCQ is about to begin in Tunisia, and zinc could be added to the NIAID HCQ plus AZT trial. It makes sense to now invoke the parallel track for outpatient use of interferon beta, favipiravir, and zinc while we await the eventual trial completions. In Russia, the parallel track has been implemented, with favipiravir now offered as treatment, even as they proceed with further confirmatory studies.

Nowhere is the decision more difficult than in the area of vaccination. The developers of several vaccines have made it clear that they can produce large quantities in a short time. Yet, it remains to be tested whether the immunity generated will be protective and, conversely, given the hyperinflammatory nature of SARS-Cov-2 infection, whether certain vaccines could actually be harmful. Acquiring con-

clusive answers will take a long experience. COVID-19 fueled by modern air travel, has engulfed the world, overwhelming containment, outspeeding well designed randomized placebo controlled clinical trials. We call on the public health authorities to make the parallel track immediately available. Cross international boundaries and accept data submitted in China, Russia and other countries, whether officially published or posted on line, and make promising agents available to treating physicians. If vaccines can be brought forward now, it is reasonable to authorize their use in very large parallel track studies in at risk populations, such as healthcare workers and the elderly. It is now necessary for public health authorities to make hard decisions despite limited current data.

REFERENCES

1) WHO Situation Report dashboard May 25, 2020

2) Wu Z, McGoogan JM. Characteristics of and important lessons from the

coronavirus disease 2019 (COVID-19) outbreak in China. Summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. JAMA 2020; 323: 1239-1242

3) Baud D, Qi X, Nielsen-Saines K, Musso D, et al. Real estimates of mortality following COVID-19 infection. Lancet Infect Dis posted on line 2020 Mar 12

4) Russell TW, Hellewell J, Jarvis CI, et al. Estimating the infection and case fatality ratio for coronavirus disease (COVID-19) using age-adjusted data from the outbreak on the Diamond Princess cruise ship, February 2020. Euro Surveill. 2020 Mar 26; 25(12): 2000256.

5) Wang D, Yin Y, Hu C, et al. Clinical course and outcome of 107 patients infected with the novel coronavirus, SARS-CoV-2, discharged from two hospitals in Wuhan, China

Critical Care. 2020; 24: 188

6) Huang C, Wang Y, Li X et al Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China Lancet 2020; 395: 497–506

7) Chen T, Wu D, Chen H, et al.Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. BMJ. 2020;368:m1091.

8) Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with

COVID-19 in Wuhan, China: a retrospective cohort study. Lancet 2020; 395: 1054–62

(9) Connors JM, Levy JH. Thromboinflammation and the hypercoagulability of COVID-19.
J Thromb Haemost. 2020 Apr 17.

(10) Ricardo J Jose, Ari Manuel. COVID-19 cytokine storm: the interplay between inflammation
and coagulation storm: Lancet Respir Med 2020 Published Online April 27, 2020

(11) Verdoni L, Mazza A, Gervasoni A, et al. An outbreak of severe Kawasaki-like disease at the Italian epicentre of the SARS-CoV-2 epidemic: an observational cohort study. Lancet. Published online May 13, 2020

12) Riphagen S, Gomez X, Gonzales-Martinez C, Wilkinson N, Theocharis P. Hyperinflammatory shock in children during COVID-19 pandemic. Lancet. 2020. Advance online publication, doi: 10.1016/S0140-6736(20)31094 https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)31094-1/fulltextexternal icon

13) CDC. Multisystem inflammatory syndrome in children (MIS-C) associated with coronavirus disease 2019 (COVID-19) Distributed via the CDC Health Alert Network May 14, 2020

14) Cousins S New Zealand eliminates COVID-19. Lancet Published:May 09, 2020

(15) https://covid19.healthdata.org/projections

(16) Kissler SM, Tedijanto C, Goldstein E et al. Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period Science 2020;10.1126/science.abb5793 (2020

17) Park M, Ryan S, Thwaites RS, et al. COVID-19: Lessons from SARS and MERS
First published:27 February 2020 Eur J Immun https://doi.org/10.1002/eji.202070035

18) Groneberg DA, Poutanen SM, Low DE, et al.Treatment and vaccines for severe acute respiratory syndrome. Lancet Infect Dis. 2005;5:147-55

(19) Liu J, Cao R, Xu M,et al. Cell Discovery. 2020; 6:16 Cell Discovery https://doi.org/10.1038/s41421-020-0156-0www.nature.com/celldisc

(20) Pastick KA, Okafor EC, Wang F, et al. Review: Hydroxychloroquine and chloroquine for Treatment of SARS-CoV-2 (COVID-19) .Open Forum Infectious Diseases, Volume 7, Issue 4, April 2020

(21) Thorlund KT, Dron L, Park J et al. A real-time dashboard of clinical trials for COVID-19. Lancet Digital Health published on line 24 Apr 20 https://doi.org/10.1016/ S2589-7500(20)30086-8

(22) Wang Y, Zhang D, Guanhua Du G, et al. Remdesivir in adults with severe COVID-19: a randomised,double-blind, placebo-controlled, multicentre trial Lancet Published Online April 29, 2020 https://doi.org/10.1016/ S0140-6736(20)31022-9

23) Grein J, Ohmagari N, Shin D. Compassionate use of remdesivir for patients with severe Covid-19

N Engl J Med April 10, 2020

24) Beigel JH, Tomashek KM, Dodd LE. Remdesivir for the Treatment of Covid-19 — Preliminary Report. May 22, 2020. New Engl J Med. DOI: 10.1056/NEJMoa2007764

25)Goldman JD, Lye DCB, Hui DS, et al. Remdesivir for 5 or 10 days in patients with severe Covid-19. New Engl J Med May 27, 2020 DOI: 10.1056/NEJMoa2015301

26) Cai Q, Yang M, Liu D, et al. Experimental treatment with favipiravir for COVID-19: An open-label control study Engineering (In Press)

27) Chen C, Huang J, Zhenshun Cheng Z, et al. Favipiravir versus Arbidol for COVID-19: A Randomized Clinical Trial medRxiv preprint doi: https://doi.org/10.1101/2020.03.17.20037432.this version posted March 20, 2020.

28) Gautret P ,Lagier J-C , Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. International Journal of Antimicrobial Agents (In Press)

(29) Million M, Lagierab J-C, Gautret P, et al. Early treatment of COVID-19 patients with hydroxychloroquine and azithromycin: A retrospective analysis of 1061 cases in Marseille, France Travel Medicine and Infectious Disease (In Press) Available online 5 May 2020, 101738

(30) Chen Z, Hu J,   Zhang Z, et al. Efficacy of hydroxychloroquine in patients with COVID-19: results of a randomized clinical trial. medRxiv preprint doi:https://doi.org/10.1101/2020.03.22.20040758.this version posted March 30, 2020.

(31) Chen J, Liu D, Liu L, et al. A pilot study of hydroxychloroquine in treatment of patients with common coronavirus disease-19 (COVID-19) J Zhejiang Univ (Med Sci) 2020, Vol. 49 Issue (1): 0-0 DOI: 10.3785/j.issn.1008-9292.2020.03.03

32) Tang W, Cao Z, Han M, et al. Hydroxychloroquine in patients mainly with mild to moderate COVID-19: an open-label, randomized, controlled trial. BMJ 2020;369:m1849

33) Magagnoli J, Narendran S, Pereira F, et al. Outcomes of hydroxychloroquine usage in United States veterans hospitalized with COVID-19. medRxiv 2020; published online April 23. DOI:10.1101/2020.04.16.20065920 (preprint).

(34) Geleris J, Sun Y, Platt J, et al. Observational study of hydroxychloroquine in hospitalized patients with Covid-19. New England Journal of Medicine Downloaded from nejm.org on May 13, 2020.

(35) Rosenberg ES, Dufort EM, Udo T, et al. Association of treatment with hydroxychloroquine or azithromycin with in-hospital mortality in patients with COVID-19 in New York State. JAMA. doi:10.1001/jama.2020.8630 Published online May 11, 2020.

(36) Mahévas M, Tran VT, Roumier M, et al. Clinical efficacy of hydroxychloroquine in patients with covid-19 pneumonia who require oxygen: observational comparative study using routine care data. BMJ 2020;369:m1844

37) Mehra MR, Desai SS, Ruschitzka F, et al. Hydroxychloroquine or chloroquine with or without a macrolide for treatment of COVID-19: a multinational registry analysis. Lancet published on line May 22, 2020 https://doi.org/10.1016/S0140-6736(20)31180-6

38) Carlucci P, Tania Ahuja T, Petrilli CM, et al. Hydroxychloroquine and azithromycin plus zinc vs hydroxychloroquine and azithromycin alone: outcomes in hospitalized COVID-19 patients medRxiv doi: https://doi.org/10.1101/2020.05.02.20080036

39) Read SA, Obeid S,Ahlenstiel C, et al. The role of zinc in antiviral immunity. Adv Nutr. 2019 Jul; 10(4): 696–710.Published online 2019 Apr 22

40) Singh M, Das RR. Zinc for the common cold. Cochrane Database Syst Rev. 2015 Apr; 2015(4): CD001364. Published online 2015 Apr 30.

41) Boulware DR, Pullen MF, Bangdiwala AS, et al. A randomized trial of hydroxychloroquine as postexposure prophylaxis for Covid-19. N Engl J Med June 3, 2020 DOI: 10.1056/NEJMoa2016638

42) Li Y, Xie Z, Lin W, et al. An exploratory randomized, controlled study on the efficacy and safety of lopinavir/ritonavir or arbidol treating adult patients hospitalized with mild/moderate COVID-19 (ELACOI. medRxiv preprint doi: https://doi.org/10.1101/2020.03.19.20038984.this version posted

43) Cao B, Wang Y, Wen D, et al. A Trial of Lopinavir–Ritonavir in adults hospitalized with severe Covid-19. N Engl J Med. 2020 May 7;382(19):1787-1799. doi: 10.1056/NEJMoa2001282.

44) Hung IFN , Lung KC, Tso EYK,Triple combination of interferon beta-1b, lopinavir–ritonavir,

and ribavirin in the treatment of patients admitted to hospital with COVID-19: an open-label, ran-domised, phase 2 trial Lancet, published on line 8 may 2020https://doi.org/10.1016/

45) Bian H, Zhao-Hui Zheng ZH, Ding Wei D. Meplazumab treats COVID-19 pneumonia: an open-la-belled, concurrent controlled add-on clinical trial. medRxiv preprint doi:https://doi.org/10.1101/2020.03.21.20040691

46) Xua X, Hanb M, Lia T, et al. Effective treatment of severe COVID-19 patients with tocilizumab 10970–10975 | PNAS | May 19, 2020 | vol. 117 | no. 20
www.pnas.org/cgi/doi/10.1073/pnas.2005615117

47) "'Solidarity' clinical trial for COVID-19 treatment". www.who.int. World Health Organization. Re-trieved 22 April 2020

(48) Haviernik J,1, Štefánik M, Fojtíková M et al. Arbidol (Umifenovir): A broad-spectrum antiviral drug that inhibits medically important arthropod-borne flaviviruses. Viruses. 2018 Apr; 10(4): 184.

(49) Uno Y. Camostat mesilate therapy for COVID-19. Intern Emerg Med. 2020 Apr 29

50) Chumakov K, Gallo R. Could an old vaccine be a godsend for new coronavirus? Using the oral po-lio vaccine could prevent or reduce the spread of COVID-19 to immunized individuals. USA Today 21 Apr 2020

51) Gao Q,*, Bao L, Mao H, et al. Development of an inactivated vaccine candidate for SARS-CoV-2. Science 06 May 2020

52) Diamond MS, Pierson TC. The challenges of vaccine development against a new virus during a pandemic. Cell Host Microbe. 2020;27(5):699-703.

53) Wan Y, Shang J, Sun S, et al. Molecular mechanism for antibody-dependent enhancement of coronavirus entry. J. Virol. 2020;94:e02015-19.

54) Fatima K, Syed NI. Dengvaxia controversy: impact on vaccine hesitancy. J Glob Health. 2018; 8(2): 020312.

55) Rosenbaum L.Trolleyology and the dengue vaccine dilemma. N Engl J Med. 2018; 26;379:305-307

56) Tanner R, Villarreal-Ramos B, Vordermeier HM, et al. The humoral immune response to BCG vaccination. Front Immunol. 2019;10:1317.

(57) Moorlag SJCFM, Arts RJW, van Crevel R, et al. Non-specific effects of BCG vaccine on viral infections. Clin Microbiol Infect. (2019)

58) China National Health Commission. Chinese Clinical Guidance for COVID-19 Pneumonia Diagnosis and Treatment (7th edition 2020-03-16 12:00:32

59) Goodman JL, Borio L. Finding Effective Treatments for COVID-19. Scientific Integrity and Public Confidence in a Time of Crisis. JAMA. 2020;323(19):1899-1900. doi:10.1001/jama.2020.64

60) https://budgetmodel.wharton.upenn.edu accessed 21 May 2020

61) Sepkowitz KA. AIDS-The first 20 years. N Engl J Med 2001; 344:1764-1772

62) Broder S.The development of antiretroviral therapy and its impact on the HIV-1/AIDS pandemic. Antiviral Res. 2010 Jan; 85(1): 1

63) Fauci AS. Preface: Evolving ethical issues over the course of the AIDS pandemic. Public Health Rev 34, 2 (2012)

64) Harris NS, Johnson AS, Huang YA, et al. Vital Signs: Status of human immunodeficiency virus testing, viral suppression, and HIV preexposure prophylaxis - United States, 2013-2018. MMWR Morb Mortal Wkly Rep. 2019 ;68:1117-1123

TABLE 1. PUBLISHED STUDIES AS OF MAY 25, 2020. Each agent is listed along with the country originating the publication. The type of control procedure: RCT: Randomized controlled study PLAC: Placebo , soc: standard of care which is variable depending on each location. AC: Active control: the control options are listed in parenthesis. OBS: retrospective observation study

TREATMENT	LOCATION	TYPE OF CONTROL	NUMBER SUBJECTS	SETTING	SEVERITY	RESULTS
Remdesivir[22]	China	RCTPLAC vs soc	273	HOSP	SEV	mortality in remdesivir group 14%, control 13% 11% in remdesivir group with <10 days symptoms Vs 14% in placebo group, no difference in viral clearance
Remdesivir[23]	U.S.A.	RCTPLAC	53	HOSP	SEV,CRIT	Mortality 18% (6 0of 34 patients on mechanical ventilation 5% 1 of 19 of those not receiving mechanical ventilation
Remdesivir[24]	U.S.A.	Compassionate Use	1063	HOSP	SEV,CRIT	Mortality 18% (6 of 34 patients on mechanical ventilation 5% 1 of 19 of those not receiving mechanical ventilation
Remdesivir[25]	U.S.A.	AC(5 day vs 10 day, no control)	397	HOSP	MOD,SEV	10 day group had more severe disease, mortality 8% in 5 day group 11% in 10 day group
Favipiravir (FVP)[26]	China	AC(lpv/r vs favipiravir)	70	HOSP	MOD,SEV	4 day viral clearance time FPV vs 11 days lpv/r CT scan improvement 92% fvp vs 62% lpv/r
Favipiravir (FVP)[27]	China	AC (umifenovir vs favipiravir)	240	HOSP	MOD,SEV	Clinical recovery rate 56%umifenovir vs 71% fvp time to fever reduction and cough shorter fvp
HCQ, azithromycin[28]	France	Non randomized control	20	MIXED	MIXED	Treated patients showed more rapid viral clearance than historical controls
HCQ[29]	FRANCE	OBS(hcq+azt, no control group)	1061	MIXED	MILD,MOD	0.9% overall mortality, no sudden death, no cardiac arrythmias
HCQ[30]	China	RCT(soc)	62	HOSP	MOD	shorter time to temperature recovery Pneumonia improvement 81% (HCQ) vs 55% soc no progression to severe status any HCQ patient versus 4 of 17 (soc)

HCQ[31]	China	RCT(soc)	30	HOSP	MOD	no differences in viral conversion
HCQ[32]	China	RCT(hcq vs soc)	150	HOSP	MILD,MOD	No difference in viral clearance at 28 days
HCQ[33]	U.S.A.	OBS(hcq vs no hcq)	338	HOSP	MOD,SEV	Mortality twice as high in patients chosen for HCQ treatment
HCQ[34]	U.S.A.	OBS (hcq vs no hcq)	1376	HOSP	MOD,SEV	no significant association between HCQ use and intubation or death (hazard ratio, 1.04, 95% confidence interval, 0.82 to 1.32). HCQ treated patients were sicker with ppO2 223 vs 360 for controls still lower than controls after propensity score matching 223 vs 273
HCQ[35]	U.S.A.	OBS (hcq+azt vs hcq vs azt)	1438	HOSP	MOD,SEV	Overall in-hospital mortality was 20.3%,25.7% hcq+azt, 19.9% hcq alone neither drug 17.7%, cardiac arrest more likely with hcq+azt
HCQ[36]	France	OBS (hcq vs no hcq)	181	HOSP	MOD,SEV	no difference in mortality or need to transfer to ICU
HCQ, Chloroquine[37]	Multinational	OBS(hcq, chloroquine with or without macrolide vs none of these)	96,032	HOSP	MOD,SEV	Mortality more than doubled vs control group, with increased risk of ventricular arrythmia
HCQ[38]	U.S.A.	OBS (hcq+azt +zinc vs hzq +azt)	932	HOSP	MOD,SEV	Reduction in mortality with addition of zinc OR 0.449, 95% CI 0.271-0.744).
HCQ[41]	U.S.A.	RCTPLAC	821	OUTPAT	HEALTHY	No signficant difference in incidence of subsequent COVID-19 infection (11.8% HCQ, 14.3% PLAC, p=0.35)
LPV/R[42]	China	AC(lpv/r vs umifenovir vs no antiviral)	44	HOSP	MOD,SEV	No differences in viral load, clinical status, or CT scan
LPV/R[43]	China	RCT(soc)	199	HOSP	SEV	No differences in mortality, viral load; time to clinical improvement 1 day less in lpv/r group
LPV/R[44]	Hong Kong	AC(lpv/r+ribavirin+beta interferon vs lpv/r+ribavirin()	127	HOSP	MOD.SEV	Triple combination of interferon beta-1b, LPV/R, and ribavirin yielded more rapid viral clearance but attributable primarily to interferon

Meplazumab[45]	China	RCT(soc)	38	HOSP	MOD,SEV	Viral clearance 3 days (1.5-4.5) compared to 13 days(6.5-19.5) in control group
Tocilizumab[46]	China	OBS(no control)	20	HOSP	SEV,CRIT	no patient died, rapid defervescence, improvement in oxygenation

NUMBER OF
REGISTERED TRIALS

>100	China (323)	United States (103)
>30	Spain (36)	France (35)
>20	Iran (29)	Italy (21)

>10 Germany (19) Canada(18) Netherlands(14) Brazil(13)
Denmark(13) Japan (12) United Kingdom(12)

>5 Israel (9), Pakistan (9),Egypt (8) Mexico (8)
South Korea (8)Belgium (6), Greece (6)

5 Colombia , Norway,Switzerland

4 Australia , Austria, India , Russia , Singapore , Thailand

3 Nigeria, South Africa, Sweden,Tunisia

2 Argentina, Hungary, Ireland, Poland, Qatar,Taiwan

1 Bahrain, Cuba, Czechia, Indonesia, Iraq, Jordan, Macedonia,
Malaysia, Monaco, New Zealand, Peru, Puerto Rico

TABLE 2. TRIALS IN EACH COUNTRY. Number of trials registered in various countries per WHO

TABLE 3. STUDIES WITH ANTIMALARIALS,LOPANOVIR/ RITONAVIR, OR INTERFERONS. The setting of each study, whether hospital or outpatient, the type of control (AC: Active control, RCT: randomized control; soc: standard of care; PLAC: placebo control, whether treatment (TREAT) or prevention (PREV), the severity of illness, number of subjects, quarter study initiated, quarter study projects completion of enrollment, and duration of symptoms for patient inclusion are listed.   Studies projecting completion of enrollment by the end of the second quarter of 2020 are bolded. Outpatient studies with planned completion by the end of the second quarter of 2020 are italicized.

OUTPATIENT OR HOSPITAL	TYPE OF CONTROL	PREVENTION OR TREATMENT	SEVERITY OF ILLNESS	NO OF SUBJECTS	BEGIN DATE	PROPOSED END OF ENROLLMENT	DURATION SYMPTOMS
			HCQ OR CHLOROQUINE				
OUT	RCT,PLAC	PREV (6)	HEALTHY, EXPOSED	360,3500,440,400,350,1700	Q1(20),Q2(20)	3 Q3(20), 2 Q4(20), 1 Q2(21)	N/A
OUT	AC azithromycin, AC LPV/r	PREV (2)	HEALTHY, EXPOSED	900,1200	Q1(20),Q2(20)	1 Q3(20, 1Q4(20)	N/A
OUT	RCT PLAC	PREV,TREAT	HEALTHY EXPOSED, MILD	3000	Q1(20)	*Q2(20)*	4
OUT	AC ascorbic acid	PREV	HEALTHY EXPOSED	2000	Q2(20)	Q3(20)	4
OUT	AC ascorbic acid	TREAT	MILD,MOD	1250	Q2(20)	Q3(21)	2
OUT	AC azithromycin	TREAT	MILD,MOD	1550	Q2(20)	Q1(21)	7
OUT	RCT PLAC	TREAT, PREV	MILD, HEALTHY EXPOSED	3040	Q1(20)	*Q2(20)*	5
OUT	RCT PLAC	TREAT	MILD	210	Q1(20)	NO DATE	7
OUT	MULTIDOSE PLAC	TREAT	MILD	400	Q2(20)	Q2(21)	ANY DUR
OUT	ac (azithromycin+hcq,hcq)	TREAT	MILD	160	Q2(20)	Q2(21)	ANY DUR
OUT	RCT PLAC	TREAT	MILD	1660	Q1(20)	Q3 (20)	4
OUT	AC (lpv/r)	TREAT	MILD	150	Q1(20)	*Q2(20)*	ANY DUR
MIXED	AC azithromycin	TREAT	MOD,SEV	600	Q2(20)	Q2(21)	7
MIXED	RCT PLAC	TREAT	MIXED	210	Q2(20)	Q3 (20)	ANY DUR
HOSP	AC azithromycin	TREAT (4)	MOD, SEV	630;500;226;300	Q2(20)	Q3,Q4(20)	ANY DUR
HOSP	AC, azithromycin	TREAT	SEV,CRIT	440	Q1(20)	Q3 (20)	ANY DUR
HOSP	RCT PLAC	TREAT	MILD,MOD	300	Q2(20)	Q4(20)	3
HOSP	RCT PLAC	TREAT	MIXED	250	Q1(20)	*Q2(20)*	2
HOSP	AC (resdemevir)	TREAT	MOD, SEV	700	Q1(20)	NO DATE	ANY DUR
HOSP	AC (lpv/r,losartan,PLAC)	TREAT	MOD, SEV	4000	Q1(20)	NO DATE	3
HOSP	AC (lpv/r,baricitinib,sarilumab,PLAC)	TREAT	MOD, SEV	1000	Q1(20)	Q1(21)	ANY DUR
HOSP	AC (faviparavir)	TREAT	MOD, SEV	150	Q1(20)	Q1(21)	14
HOSP	RCT PLAC	TREAT	MOD, SEV	202	Q1(20)	*Q2(20)*	ANY DUR
HOSP	RCT PLAC	TREAT	MOD, SEV	350	Q2(20)	*Q2(20)*	5
HOSP	AC (faviparavir)	TREAT	SEV	150	Q1(20)	*Q2(20)*	14
HOSP	AC(chq,hcq)	TREAT	SEV	100	Q1(20)	Q2(21)	14
HOSP	RCT PLAC	TREAT	SEV	510	Q2(20)	Q3 (20)	10
HOSP	AC (azithromycin)	TREAT	SEV,CRIT	276	Q2(20)	Q3 (20)	3
HOSP	RCT PLAC	TREAT	SEV,CRIT	1000	Q2(20)	NO DATE	ANY DUR
HOSP	RCT PLAC	TREAT	CRIT	500	Q1(20)	Q4(20)	ANY DUR
HOSP	AC(high vs low dose HCQ)	TREAT	CRIT	440	Q1(20)	Q3 (20)	ANY DUR
HOSP	AC (hcq +/- famotidine)	TREAT	SEV	1200	Q2(20)	Q3 (20)	1
HOSP	RCT (PLAC)	TREAT	SEV	220	Q2(20)	Q1(21)	ANY DUR
HOSP	RCT (PLAC)	TREAT	SEV	1300	Q2(20)	Q3(20)	2
HOSP	RCT (PLAC)	TREAT	MIXED	626	Q2(20)	Q3 (20)	3
OUT	RCT(soc)	TREAT	MILD	400	Q2(20)	NO DATE	ANY DUR
			LOPANOVIR/RITANOVIR				
OUT	AC HCQ	PREV (2)	HEALTHY	1200,1220	Q2(20)	*Q2(20)*,Q3(21)	N/A
OUT	AC HCQ	TREAT	MILD, MOD	150	03/20/20	*Q2(20)*	ANY DUR
Mixed	AC(Mulitple)	TREAT	MIXED	500	04/20/20	Q4(20)	ANY DUR
Mixed	AC (Ebastine)	TREAT	MIXED	100	02/20/20	Q1(20)	ANY DUR
Mixed	AC (Carrimycin)	TREAT	MIXED	520	02/20/20	Q1(21)	ANY DUR
Mixed	AC HCQ	TREAT	MIXED	112	02/20/20	Q4(20)	ANY DUR
Hosp	AC HCQ,losartan	TREAT	MOD,SEV	4000	04/20/20	Q1(21)	3
Hosp	AC HCQ IFN, STEROID	TREAT	SEV	5000	02/20/20	*Q2(20)*	ANY DUR
HOSP	RCT(soc)	TREAT	SEV	160	Q1(20)	Q1(21)	ANY DUR
HOSP	RCT (soc)	TREAT	MILD	328	Q1(20)	Q1(21)	7
			INTERFERON				
OUT	AC (thymosin)	PREV	HEALTHY	2944	Q1(20)	*Q2(20)*	N/A
OUT	AC(chimed)	TREAT	MILD	238	Q1(20)	Q1(22)	Q2(22)
MIXED	RCT (soc)	TREAT	MIXED	328	Q1(20)	*Q2(20)*	7
MIXED	AC(ebastine)	TREAT	MIXED	100	Q1(20)	Q1(20)	ANY DUR
MIXED	AC (super compunded ifn)	TREAT	MIXED	100	Q1(20)	Q3 (20)	ANY DUR
MIXED	AC(ifn+umifenovir,umifenovir)	TREAT	MIXED	100	Q1(20)	*Q2(20)*	ANY DUR
HOSP	AC(lpv/r, hcq, steroids))	TREAT	MIXED	5000	Q1(20)	Q2(21)	ANY DUR
HOSP	RCT (PLAC)	TREAT	MILD,MOD	400	Q1(20)	NO DATE	1

AGENT	HOSPITAL OR OUTPATIENT	TYPE OF CONTROL	TREAT OR PREVENT	SEVERITY	NUMBER SUBJECTS	BEGIN DATE	PROPOSED ENROLLMENT END	DISEASE DURATION
FAVIPIRAVIR	HOSP	RCT PLAC	TREAT	MIXED	210	Q2(20)	Q3 (20)	ANY DUR
	HOSP	ac (umifenovir)	TREAT	MIXED	240	Q1(20)	**Q2(20)**	12
	HOSP	AC(fvp, fvp combined with tociluzimab,soc)	TREAT	MIXED	150	Q1(20)	**Q2 (20)**	ANY DUR
	HOSP	AC (fvp, fvp combined with chloroquine,soc)	TREAT	MIXED	150	Q1(20)	Q3 (20)	ANY DUR
	HOSP	AC (fvp+hcq, lpv+hcq)	TREAT	MOD,SEV	324	Q2(20)	Q3 (20)	ANY DUR
UMIFENOVIR	OUT	RCT(PLAC)	TREAT	MILD	380	Q1(20)	Q4(20)	ANY DUR
	MIXED	AC(lpv/r)	TREAT	MIXED	125	COMPLETED		ANY DUR
	MIXED	AC (azithromycin)	TREAT	MIXED	100	Q1(20)	**Q2(20)**	ANY DUR
	HOSP	AC(lpv/r, oseltamavir)	TREAT	MOD, SEV	400	Q1(20)	**Q2(20)**	ANY DUR
	HOSP	AC (umifenovir+hcq , hcq+lpv)	TREAT	MIXED	200	Q2(20)	Q3 (20)	ANY DUR
	HOSP	AC (umifenovir+hcq , hcq+lpv)	TREAT	MIXED	100	Q2(20)	Q3 (20)	ANY DUR
CAMOSTAT	MIXED	RCT (PLAC)	TREAT	MIXED	580	Q1(20)	Q4(20)	2
	HOSP	AC hcq+azithromycin, camostat+hcq,soc)	TREAT	MILD, MOD	250	Q2(20)	Q3 (20)	ANY DUR
EMITRICITABINE/ TENOFOVIR	OUT	AC (hcq,PLAC)	PREV	HEALTHY	4000	Q2(20)	Q3 (20)	N/A
RIBAVIRIN	HOSP	AC (lpv/r+ifn b, ribavirin +lpv/r+ifn b)	TREAT	MILD, MOD	108	Q1(20)	Q1(21)	ANY DUR
REMDESIVIR	HOSP	AC(hcq+remdesivir, hcq)	TREAT	MIXED	700	Q1(20)	Q3 (20)	ANY DUR
TRIAZAVIRIN	HOSP	RCT (PLAC)	TREAT	MOD, SEV	240	Q1(20)	**Q2 (20)**	ANY DUR
IVERMECTIN	MIXED	RCT (soc)	TREAT	MIXED	100	Q2(20)	Q3 (20)	ANY DUR

TABLE 4: ANTIVIRAL AGENTS. The setting of each study, whether hospital or outpatient, the type of control (AC: Active control, RCT: randomized control; soc: standard of care; PLAC: placebo control, whether treatment (TREAT) or prevention (PREV), the severity of illness, number of subjects, quarter study initiated, quarter study projects completion of enrollment, and duration of symptoms for patient inclusion are listed. Studies projecting completion of enrollment by the end of the second quarter of 2020 are bolded. Outpatient studies with planned completion by the end of the second quarter of 2020 are italicized.

TABLE 5: IMMUNE MODULATORS. The setting of each study, whether hospital or outpatient, the type of control (AC: Active control, RCT: randomized control; soc: standard of care; PLAC: placebo control, whether treatment (TREAT) or prevention (PREV), the severity of illness, number of subjects, quarter study initiated, quarter study projects completion of enrollment, and duration of symptoms for patient inclusion are listed.   Studies projecting completion of enrollment by the end of the second quarter of 2020 are bolded.

AGENT	HOSPITAL OR OUTPATIENT	TYPE OF CONTROL	TREAT OR PREVENT	SEVERITY	NUMBER SUBJECTS	BEGIN DATE	PROPOSED ENROLLMENT END	DISEASE DURATION
SARILUMAB	HOSP	RCT (PLAC)	TREAT	SEV,CRIT	300	Q1(20)	Q3 (20)	ANY DUR
	HOSP	RCT (PLAC)	TREAT	SEV,CRIT	400	Q1(20)	Q1(21)	ANY DUR
	HOSP	RCT (soc)	TREAT	SEV,CRIT	460	Q2(20)	NO DATE	ANY DUR
CD24F (blocks disinhibition of immune checkpoint)	HOSP	RCT (PLAC)	TREAT	SEV	230	Q1(20)	Q2(21)	ANY DUR
COLCHICINE	OUT	RCT (PLAC)	TREAT	MILD	6000	Q1(20)	Q3 (20)	1
	HOSP	RCT (soc)	TREAT	SEV	2500	Q2(20)	Q3 (20)	ANY DUR
IFX1 (monoclonal ab to complement)	HOSP	RCT (PLAC)	TREAT	SEV	130	Q1(20)	NO DATE	14
SELINEXOR	HOSP	RCT (PLAC)	TREAT	SEV	230	Q2(20)	Q3 (20)	ANY DUR
SILTUXIMAB	HOSP	AC (methlyprednisolone)	TREAT	SEV,CRIT	200	Q2(20)	**Q2 (20)**	ANY DUR
TOCILIZUMAB	MIXED	AC (azithromycin + hcq)	TREAT	SEV	276	Q2(20)	Q3 (20)	ANY DUR
	HOSP	AC sarilumab)	TREAT	SEV	200	Q1(20)	Q2(21)	ANY DUR
	HOSP	RCT (soc)	TREAT	SEV,CRIT	188	Q1(20)	**Q2(20)**	ANY DUR
	HOSP	RCT(PLAC)	TREAT	CRIT	330	Q1(20)	**Q2(20)**	ANY DUR
	HOSP	RCT(PLAC)	TREAT	CRIT	100	Q2(20)	Q3(20)	ANY DUR

	HOSP	AC (anakinra, siltuximab, PLAC)	TREAT	CRIT	342	Q2(20)	Q3(20)	ANY DUR
	HOSP	RCT (PLAC)	TREAT	MIXED	180	Q2(20)	Q2(22)	ANY DUR
LENZILUMAB	HOSP	RCT (PLAC)	TREAT	MOD,SEV	238	Q2(20)	Q3 (20)	ANY DUR
TJ003234 anti-GM-CSF monoclonal ab	HOSP	RCT (PLAC)	TREAT	SEV,CRIT	144	Q2(20)	Q3 (20)	ANY DUR
THYMOSIN	HOSP	AC(PD-1)	TREAT	SEV	120	Q1(20)	Q1(21)	ANY DUR
CM4620 CRAC channel inhibitor	HOSP	RCT (soc)	TREAT	SEV	144	Q2(20)	Q3 (20)	ANY DUR
LERONLIMAB	HOSP	RCT (PLAC)	TREAT	SEV,CRIT	390	Q2(20)	Q4(20)	ANY DUR
ECULIZIMAB	HOSP	RCT soc)	TREAT	SEV,CRIT	120	Q2(20)	Q3 (20)	ANY DUR
CYTOKINE PROTEIN	HOSP	RCT (soc)	TREAT	MIXED	480	Q1(20)	**Q2(20)**	8
AVDORALIMAB	HOSP	RCT (PLAC)	TREAT	SEV,CRIT	108	Q2(20)	Q3 (20)	ANY DUR

COUNTRY	NAME	PROGRESS	NUMBER SUBJECTS	TYPE OF VACCINE	DATE BEGUN	COMPLETION DATE
United Kingdom		PHASE I/11	510,1090	attenuated adenovirus expressing spike protein	Q1(20)	ongoing
China	PicoVacc	PHASE I/11	744	inactivated coronavirus	Q2(20)	Q4(21)
China		PHASE I/11	100	lentiviral vector	Q1(20)	Q3(23)
United States	mRNA-1273	PHASE I/11	105	Lipid nanoparticle (LNP) dispersion containing an mRNA that encodes for the prefusion stabilized spike protein 2019-nCoV.	Q1(20)	Q3(21)
China	Ad5-nCoV,	PHASE II	500	Adenovirus vector recombinant vaccine	Q1(20)	Q1(21)
United States	INO-4800	PHASE 1	40	DNA vaccine with electroporation	Q2(20)	Q2(21)
Netherlands		RCT(PLAC)	1500	BCG Vaccination	Q1(20)	Q4(20)
Australia		RCT(PLAC)	4170	BCG Vaccination	Q1(20)	Q4(20)
China		PHASE I		Inactivated alum vaccine	Q2(20)	ongoing

TABLE 6: VACCINES. The country of origin of each vaccine is listed, with name if established, the number of subjects currently being tested, the type of vaccine, the quarter study initiated, and the quarter and year study projects completion

FIGURE 1 (Reproduced from Kissler et al our ref 16) : A to E) Simulated prevalence of COVID-19 infections (solid) and critical COVID-19 cases (dashed) following establishment on 11 March 2020 with a period of social distancing (shaded blue region) instated two weeks later, with the duration of social distancing lasting (A) four weeks, (B) eight weeks, (C) twelve weeks, (D) twenty weeks, and (E) indefinitely. There is no seasonal forcing; R0 was held constant at 2.2 (see fig. S12 for R0 = 2.6). The effectiveness of social distancing varied from none to a 60% reduction in R0. Cumulative infection sizes are depicted beside each prevalence plot (F to J) with the herd immunity threshold (horizontal black bar). Of the temporary distancing scenarios, long-term (20-week), moderately effective (20%-40%) social distancing yields the smallest overall peak and total outbreak size.

SUPPLEMENTARY TABLE 1: ALL TRIALS REVIEWED. SUPPLEMENTARY TABLE 1: TRIALS REVIEWED. All interventions reviewed are listed with the registration number in the listing database, the country of origin, the xeverity of illness, the clinical setting whether hospital or outpatient, they type of intervention, the number of patients projected, and the category of the intervention. CGRP: Calcitonin gene related peptide CHIMED: Traditional Chinese medication; IFN: Interferons ;LPV: Lopanovir/ritanovir; RAS: Renin angiotensin system

CLINICAL TRIAL REGISTRATION	COUNTRY	SEVERITY	SETTING	INTERVENTION	NO SUBJECTS	CLASS OF AGENT
NCT04355962	Switzerland	NA	ICU	Sevoflurane	64	ANESTHESIA
NCT04348370	United States	Non-severe	Healthcare Workers	BCG Vaccine	1800	ANGIOGENESIS INHIBITOR
NCT04305106	China	Severe	Hospital	Bevacizumab	140	ANGIOGENESIS INHIBITOR
NCT04275414	China	Severe	Hospital	Bevacizumab	20	ANGIOGENESIS INHIBITOR
IR/46977	Iran	NA	Outpatient	Minocycline	40	ANTIBIOTIC
NCT04369365	Austria	NA	Outpatient	AZT	200	ANTIBIOTIC
EU/2020-001640-26/GB/	United Kingdom	NA	ICU	Actilyse	24	ANTICOAG
NCT04365309	China	NA	Unclear	Aspirin	128	ANTICOAG
NCT04348383	Spain	Severe	Hospital	Defibrotide	120	ANTICOAG
NCT04359277	United States	NA	Hospital	Enoxaparin	1000	ANTICOAG
NCT04345848	Switzerland	NA	Hospital, ICU	Enoxaparin	200	ANTICOAG
CHI51265	China	NA	Hospital	LMW Heparin	120	ANTICOAG
NCT04368377	Italy	NA	ICU	Tirofiban, Clopidogrel, Acetylsalicylic acid, Fondaparinux	5	ANTICOAG
CHI50474	China	Unspecified	Unclear	Celecoxib	60	ANTIINFLAMMATORY
NCT04308317	China	Mixed	Unclear	Tetrandrine	60	ANTIINFLAMMATORY
CHI49738	China	Mixed	Outpatient, Hospital	Tranilast	60	ANTIINFLAMMATORY
NCT04339426	United States	NA	Unclear	Atovaquorine, AZT	25	ANTIMALARIAL
IR/47099	Iran	NA	Hospital	Chloroquine + Azithromycin + Melatonin	82	ANTIMALARIAL
NCT04333914	France	NA	Hospital	Chloroquine analog, nivolumab, TCZ	273	ANTIMALARIAL
NCT04321278	Brazil	Severe	Hospital	Azithromycin , HCQ	440	ANTIMALARIAL
NCT04324463	Canada	NA	Outpatient, Hospital	Azithromycin , HCQ	1500	ANTIMALARIAL
NCT04322396	Denmark	NA	Hospital	Azithromycin , HCQ	226	ANTIMALARIAL
EU/2020-001469-35/PL/	Poland	NA	Outpatient	Chlorine phosphate (Arechin)	400	ANTIMALARIAL
NCT04347980	France	Severe	Hospital	Dexamethasone, HCQ	122	ANTIMALARIAL
NCT04351191	Pakistan	Non-severe	Outpatient	HCQ	400	ANTIMALARIAL
NCT04351516	Germany	Mixed	Outpatient	HCQ	350	ANTIMALARIAL
NCT04346667	Pakistan	NA	Outpatient	HCQ	400	ANTIMALARIAL
NCT04362332	Netherlands	NA	Hospital	HCQ	950	ANTIMALARIAL
NCT04353037	United States	NA	Healthcare Workers,	HCQ	850	ANTIMALARIAL

NCT/ID	Country	Severity	Population	Drug	Sample	Class
NCT04354870	United States	NA	Outpatient Healthcare Workers	HCQ	350	ANTIMALARIAL
NCT04349371	United States	Non-severe	Healthcare Workers	HCQ	350	ANTIMALARIAL
IR/47090	Iran	NA	Healthy Exposed	HCQ	1000	ANTIMALARIAL
CHI49420	China	NA	Unclear	HCQ	300	ANTIMALARIAL
NCT04341441	United States	NA	Healthcare Workers	HCQ	3000	ANTIMALARIAL
NCT04308668	United States	Mixed	Healthy Exposed, Healthcare Workers, Outpatient	HCQ	3000	ANTIMALARIAL
NCT04329923	United States	Mixed	Healthcare Workers, Outpatient, Hospital	HCQ	400	ANTIMALARIAL
NCT04333654	United States	Non-severe	Outpatient	HCQ	210	ANTIMALARIAL
NCT04329611	Canada	NA	Outpatient	HCQ	1660	ANTIMALARIAL
NCT04328493	Vietnam	NA	Hospital	HCQ	250	ANTIMALARIAL
NCT04328961	United States	None	Healthy Exposed	HCQ	2000	ANTIMALARIAL
NCT04342169	United States	Non-severe	Outpatient	HCQ	400	ANTIMALARIAL
NCT04323527	Brazil	NA	Hospital	HCQ	440	ANTIMALARIAL
NCT04345692	United States	NA	Hospital	HCQ	350	ANTIMALARIAL
NCT04342650	Brazil	NA	Unclear	HCQ	210	ANTIMALARIAL
NCT04333225	United States	None	Healthcare Workers	HCQ	360	ANTIMALARIAL
NCT04342221	Germany	NA	Hospital	HCQ	220	ANTIMALARIAL
NCT04328467	United States	NA	Healthcare Workers	HCQ	3500	ANTIMALARIAL
NCT04334382	United States	Non-severe	Outpatient	HCQ	1550	ANTIMALARIAL
NCT04331834	Spain	None	Healthcare Workers	HCQ	440	ANTIMALARIAL
CHI49524	China	Mixed	Unclear	HCQ	360	ANTIMALARIAL
CHI49482	China	Severe	Hospital	HCQ	100	ANTIMALARIAL
CHI49536	China	Non-severe	Outpatient	HCQ	100	ANTIMALARIAL
CHI49607	China	NA	Unclear	HCQ	100	ANTIMALARIAL
CHI48880	China	NA	Unclear	HCQ	300	ANTIMALARIAL
EU/2020-001270-29/DK/	Denmark	Mixed	Hospital	HCQ	350	ANTIMALARIAL
NCT04318444	United States	None	Healthy Exposed	HCQ	1600	ANTIMALARIAL
NCT04315896	Mexico	Severe	Hospital	HCQ	500	ANTIMALARIAL
NCT04318015	Mexico	None	Healthcare Workers	HCQ	400	ANTIMALARIAL
NCT04316377	Norway	Mixed	Hospital	HCQ	202	ANTIMALARIAL
ISRCTNISRCTN86534580	United Kingdom	NA	Outpatient	HCQ	3000	ANTIMALARIAL
IR/46932	Iran	NA	Healthy Exposed, Healthcare Workers	HCQ	500	ANTIMALARIAL
NCT04369742	United States	NA	Hospital	HCQ	626	ANTIMALARIAL
NCT04363450	United States	NA	Healthcare Workers	HCQ	1700	ANTIMALARIAL
NCT04332991	United States	Severe	Hospital	HCQ	510	ANTIMALARIAL

Trial ID	Country	Severity	Setting	Intervention	N	Class
NCT04334967	United States	Non-severe	Outpatient	HCQ	1250	ANTIMALARIAL
NCT04351620	United States	Non-severe	Outpatient	HCQ	20	ANTIMALARIAL
NCT04345653	United States	Non-severe	Healthcare Workers	HCQ	45	ANTIMALARIAL
CHI50843	China	NA	Hospital	HCQ	80	ANTIMALARIAL
CHI49317	China	NA	Unclear	HCQ	78	ANTIMALARIAL
CHI49218	China	Severe	Hospital	HCQ	80	ANTIMALARIAL
CHI49612	China	NA	Unclear	HCQ	100	ANTIMALARIAL
CHI48968	China	NA	Hospital	HCQ	20	ANTIMALARIAL
NCT04261517	China	NA	Unclear	HCQ	30	ANTIMALARIAL
IR/46946	Iran	NA	Healthy Unexposed	HCQ	60	ANTIMALARIAL
NCT04322123	Brazil	NA	Hospital, ICU	HCQ , Azithromycin	630	ANTIMALARIAL
IR/46536	Iran	NA	Hospital	HCQ , LPV/R , Oseltamivir	50	ANTIMALARIAL
JPJRCTS031190227	Japan	NA	Hospital	HCQ , LPV/R , Oseltamivir	50	ANTIMALARIAL
IR/47200	Iran	Moderate	Hospital	HCQ, Arbidol	40	ANTIMALARIAL
NCT04336332	United States	Non-severe	Outpatient, Hospital	HCQ, azithromycin	160	ANTIMALARIAL
NCT04335552	United States	Mixed	Hospital	HCQ, azithromycin	500	ANTIMALARIAL
NCT04344379	France	Non-severe	Healthcare Workers	HCQ, azithromycin	900	ANTIMALARIAL
NCT04329832	United States	NA	Hospital	HCQ, Azithromycin	300	ANTIMALARIAL
NCT04345861	France	NA	Hospital	HCQ, AZT	150	ANTIMALARIAL
NCT04354428	United States	NA	Outpatient	HCQ, AZT	630	ANTIMALARIAL
NCT04344444	United States	NA	Hospital	HCQ, Azt	600	ANTIMALARIAL
NCT04341727	United States	NA	Hospital	HCQ, AZT	500	ANTIMALARIAL
EU/2020-001527-14/NL/	Netherlands	NA	Hospital	HCQ, AZT	60	ANTIMALARIAL
IR/46968	Iran	NA	Hospital	HCQ, AZT, Naproxen, LPV/r, Corticosteroids	906	ANTIMALARIAL
NCT04341870	France	NA	Hospital	HCQ, AZT, Sarilumab	60	ANTIMALARIAL
NCT04370262	United States	NA	Hospital	HCQ, Famotidine	1170	ANTIMALARIAL
NCT04344457	United States	Non-severe	Outpatient	HCQ, Indomethacin, azithromycin	80	ANTIMALARIAL
NCT04344457	United States	Non-severe	Outpatient	HCQ, Indomethacin, azithromycin	80	ANTIMALARIAL
NCT04343092	Iraq	Mixed	Hospital	HCQ, Ivermectin	50	ANTIMALARIAL
IR/46926	Iran	NA	Outpatient	HCQ, LPV/r	60	ANTIMALARIAL
IR/47022	Iran	NA	Hospital	HCQ, LPV/r	100	ANTIMALARIAL
NCT04328285	France	NA	Healthcare Workers	HCQ, LPV/r,	1200	ANTIMALARIAL
IR/46561	Iran	Severe	Hospital	HCQ, LPV/r,	20	ANTIMALARIAL
IR/46814	Iran	NA	Hospital	HCQ, LPv/r, Atazanavir/r, Teicoplanin	40	ANTIMALARIAL
NCT04343768	Iran	Mixed	Hospital	HCQ, LPV/r, IFN	60	ANTIMALARIAL
IR/46790	Iran	NA	Hospital	HCQ, LPV/r, Sofosbuvir-Velpatasvir	80	ANTIMALARIAL

IR/47119	Iran	NA	Hospital	HCQ, Thalidomide	60	ANTIMALARIAL
NCT04325893	France	NA	Unclear	HCQ/placebo	1300	ANTIMALARIAL
ISRCTNISRCTN14326006	Canada	NA	Healthcare Workers	Hydroxychloroquine	988	ANTIMALARIAL
IR/46940	Iran	NA	Unclear	LPV/r, HCQ, Herbal medicine	40	ANTIMALARIAL
NCT04328012	United States	NA	Hospital	LPV/r, HCQ, losartan	4000	ANTIMALARIAL
ISRCTNISRCTN50189673	United Kingdom	Mixed	Hospital	LPV/r, IFN, HCQ, steroids	5000	ANTIMALARIAL
NCT04332094	Spain	Non-severe	Hospital	HCQ , TCZ, azithromycin	276	ANTIMALARIAL PLUS IMMUNE MODULATOR
IR/46887	Iran	Mixed	Unclear	Levamisole	50	ANTIPARASITIC
NCT04348409	Brazil	Mixed	Hospital	Nitazoxanide	50	ANTIPARASITIC
NCT04341493	Mexico	NA	Unclear	Nitazoxanide	86	ANTIPARASITIC
IR/46576	Iran	NA	Outpatient, Hospital	Noscapine	125	ANTITUSSIVE
NCT04330638	Belgium	Severe	Hospital, ICU	Anakinra, siltuximab, TCZ	342	ANTIVIRAL
NCT04339712	Greece	Severe	Unclear	Anakinra, TCZ	40	ANTIVIRAL
NCT04324489	China	Severe	Hospital	Antiviral	4	ANTIVIRAL
IR/47087	Iran	NA	Hospital	Arbidol	200	ANTIVIRAL
CHI49165	China	Non-severe	Outpatient	Arbidol	380	ANTIVIRAL
CHI49532	China	NA	Hospital	Azvudine	20	ANTIVIRAL
CHI50507	China	Mixed	Unclear	Azvudine	10	ANTIVIRAL
NCT04358614	Italy	NA	Hospital	Baricitinib	12	ANTIVIRAL
NCT04321096	Denmark	NA	Hospital	Camostat	580	ANTIVIRAL
IR/46573	Iran	Mixed	Hospital	Camostat	40	ANTIVIRAL
NCT04276688	China	NA	Hospital	Combination Antiviral	127	ANTIVIRAL
NCT04291729	China	NA	Hospital	Combination Antiviral	11	ANTIVIRAL
NCT04255017	China	NA	Unclear	Combination Antiviral	400	ANTIVIRAL
CHI50136	China	NA	Unclear	Combination Antiviral	30	ANTIVIRAL
NCT04261270	China	Mixed	Hospital	Combination Antiviral	60	ANTIVIRAL
CHI49918	China	Mixed	Hospital	Danoprevir , Ritonavir	60	ANTIVIRAL
NCT04345276	China	NA	Hospital	Danoprevir, Ritonavir	40	ANTIVIRAL
NCT04252274	China	NA	Unclear	Darunavir And Cobicistat	30	ANTIVIRAL
NCT04304053	Spain	Non-severe	Outpatient	Darunavir/Cobicistat , HCQ	3040	ANTIVIRAL
CHI49790	China	Mixed	Unclear	Ebastine	100	ANTIVIRAL
NCT04334928	Spain	Non-severe	Healthcare Workers	Emtricitabine/tenofovir, HCQ	4000	ANTIVIRAL
NCT04333589	China	NA	Unclear	Favipiravir	210	ANTIVIRAL
NCT04358549	United States	NA	Hospital	Favipiravir	50	ANTIVIRAL
CHI49988	China	NA	Unclear	Favipiravir	30	ANTIVIRAL
CHI49510	China	NA	Hospital	Favipiravir	60	ANTIVIRAL

ID	Country	Severity	Setting	Treatment	N	Category
JPJRCTS031190226	Japan	NA	Hospital	Favipiravir	50	ANTIVIRAL
JPjRCTs041190120	Japan	NA	Outpatient	Favipiravir	86	ANTIVIRAL
NCT04310228	China	NA	Unclear	Favipiravir , TCZ	150	ANTIVIRAL
CHI50137	China	NA	Hospital	Favipiravir Vs. Arbidol	240	ANTIVIRAL
IR/46942	Iran	NA	Hospital	Favipiravir, LPV/r	84	ANTIVIRAL
CHI51126	China	NA	Hospital	Favipiravir, TCZ	150	ANTIVIRAL
CHI49042	China	NA	Outpatient	Favirapir, LPV/r	90	ANTIVIRAL
JapicCTI-205238	Japan	Mixed	Hospital	FVP	96	ANTIVIRAL
NCT04319900	China	NA	Unclear	FVP, HCQ	150	ANTIVIRAL
CHI51329	China	NA	Resolved Discharged	HCQ , /- Favipiravir	150	ANTIVIRAL
IR/46897	Iran	NA	Hospital	HCQ, Arbidol, LPV/r	100	ANTIVIRAL
IR/47078	Iran	NA	Hospital	HCQ, Atazanavir/r, Vitamin C	110	ANTIVIRAL
NCT04355052	Israel	NA	Hospital	HCQ, AZT, Camostat mesilate	250	ANTIVIRAL
IR/46783	Iran	NA	Hospital	HCQ, Favipiravir, LPV/r	324	ANTIVIRAL
CHI51141	China	Non-severe	Outpatient	Interferon, ribavirin, arbidol, ribavirin	30	ANTIVIRAL
IR/47012	Iran	Unclear	Unclear	Ivermecin	100	ANTIVIRAL
CHI50251	China	Mixed	Unclear	Ixekizumab , Antiviral	40	ANTIVIRAL
CHI51112	China	NA	Unclear	Lopinavir/ritonvair	90	ANTIVIRAL
CHI48782	China	Mixed	Outpatient	LPV/r , Interferon	108	ANTIVIRAL
NCT04252885	China	NA	Unclear	LPV/r, Arbidol	125	ANTIVIRAL
NCT04275245	China	NA	Unclear	Mepaluzumab	20	ANTIVIRAL
NCT04280705	United States	Severe	Hospital	Remdesivir	800	ANTIVIRAL
NCT04292899	United States	Severe	Hospital	Remdesivir	6000	ANTIVIRAL
NCT04292730	United States	Mixed	Hospital	Remdesivir	1600	ANTIVIRAL
EU/2020-001052-18/ES/	Denmark	NA	Hospital	Remdesivir	440	ANTIVIRAL
NCT04321616	Norway	Severe	Hospital, ICU	Remdesivir, HCQ	700	ANTIVIRAL
IR/46463	Iran	Mixed	Hospital	Sofosbuvir	70	ANTIVIRAL
IR/46784	Iran	NA	Hospital	Sofosbuvir	60	ANTIVIRAL
IR/46885	Iran	NA	Hospital	Sofosbuvir + ribavirin	48	ANTIVIRAL
IR/46567	Iran	NA	Hospital	Sofosbuvir/Lepdiasvir, HCQ , Oseltamivir, LPV/r	50	ANTIVIRAL
EU/2020-001244-26/AT/	Austria	NA	ICU	Solnatide	40	ANTIVIRAL
CHI49723	China	NA	Hospital	Triazavirin	240	ANTIVIRAL
NCT04321993	Canada	Mixed	Hospital	LPV/r, HCQ, Baricitinib, Sarilumab	1000	ANTIVIRAL + IMMUNE MODULATOR
NCT04347031	Russia	NA	Hospital	Melfoquine, HCQ, AZT, TCZ	320	ANTIVIRAL + IMMUNE MODULATOR
NCT04351763	Poland	Mixed	Hospital	Amiodarone, Verapamil	804	CARDIOVASCULAR
NCT04304313	China	Mixed	Unclear	Sildenafil	10	CARDIOVASCULAR

NCT04319731	United States	NA	Hospital	Amniotic Fluid	10	CELL THERAPY
CHI51404	China	NA	Unclear	Cell Product	20	CELL THERAPY
NCT04324996	China	NA	Unclear	Engineered Cells	90	CELL THERAPY
IR/47264	Iran	NA	Hospital	Natural Killer Cells	70	CELL THERAPY
NCT04280224	China	NA	Unclear	Natural-Killer Cells	30	CELL THERAPY
NCT04361942	Spain	NA	ICU	Stem Cell	24	CELL THERAPY
NCT04339660	China	NA	Unclear	Stem Cell	30	CELL THERAPY
IR/47156	Iran	Mild/moderate	Unclear	Stem Cell Therapy	20	CELL THERAPY
NCT04349631	United States	Non-severe	Healthcare Workers	Stem Cells	56	CELL THERAPY
NCT04355728	United States	NA	Hospital, ICU	Stem Cells	24	CELL THERAPY
IR/47110	Iran	NA	ICU	Stem Cells	10	CELL THERAPY
NCT04333368	France	Severe	ICU	Stem Cells	60	CELL THERAPY
NCT04336254	China	Severe	Hospital	Stem Cells	20	CELL THERAPY
NCT04331613	China	Severe	ICU	Stem Cells	9	CELL THERAPY
CHI51904	China	Severe	Unclear	Stem Cells	36	CELL THERAPY
CHI51050	China	Severe	Hospital	Stem Cells	20	CELL THERAPY
CHI49812	China	NA	Unclear	Stem Cells	20	CELL THERAPY
CHI49901	China	Severe	Hospital	Stem Cells	16	CELL THERAPY
CHI41760	China	Severe	Hospital	Stem Cells	30	CELL THERAPY
CHI49146	China	Severe	Hospital	Stem Cells	63	CELL THERAPY
NCT04252118	China	NA	Unclear	Stem Cells	20	CELL THERAPY
NCT04288102	China	Severe	Hospital	Stem Cells	90	CELL THERAPY
NCT04313322	Jordan	NA	Unclear	Stem Cells	5	CELL THERAPY
NCT04269525	China	Severe	ICU	Stem Cells	10	CELL THERAPY
IR/46789	Iran	NA	Hospital	Stem Cells	5	CELL THERAPY
IR/47030	Iran	NA	Hospital, ICU	Stem Cells	6	CELL THERAPY
CHI49674	China	Mixed	Hospital	Stem Cells	120	CELL THERAPY
NCT04348435	United States	Non-severe	Healthcare Workers	Stem Cells	100	CELL THERAPY
CHI49254	China	Mixed	Hospital	Stem Cells	200	CELL THERAPY
NCT04367077	United States	NA	Hospital	Stem Cells	400	CELL THERAPY
NCT04346615	United States	Mixed	Hospital	Vazegepant	120	CGRP
NCT04333550	Iran	Mixed	Hospital, Unclear	Deferoxamine	50	CHELATING AGENT
CHI49631	China	Non-severe	Unclear	TCM Exercise	160	CHIMED
CHI51030	China	NA	Resolved Discharged	TCM Exercise	28	CHIMED
CHI49762	China	Non-severe	Unclear	TCM Injectable	348	CHIMED
CHI49222	China	NA	Unclear	TCM Injectable	238	CHIMED

CHI49220	China	Non-severe	Outpatient	TCM Injectable	160	CHIMED
CHI50115	China	Non-severe	Unclear	TCM Injectable	80	CHIMED
CHI50306	China	Severe	Hospital	TCM Injectable	60	CHIMED
CHI49748	China	NA	Hospital	TCM Injectable	50	CHIMED
CHI49104	China	NA	Outpatient	TCM Injectable	60	CHIMED
CHI50586	China	Mixed	Unclear	TCM Injectable	60	CHIMED
CHI49297	China	Mixed	Outpatient, Hospital	TCM Injectable	90	CHIMED
CHI50025	China	Severe	Unclear	Traditional Chinese Medicine	120	CHIMED
CHI52699	China	NA	Unclear	Traditional Chinese Medicine	200	CHIMED
CHI52477	China	NA	Unclear	Traditional Chinese Medicine	300	CHIMED
CHI52414	China	NA	Healthy Exposed	Traditional Chinese Medicine	2000	CHIMED
NCT04334265	China	Unspecified	Unclear	Traditional Chinese Medicine	750	CHIMED
CHI52106	China	NA	Unclear	Traditional Chinese Medicine	300	CHIMED
CHI51100	China	NA	Outpatient, Hospital	Traditional Chinese Medicine	2840	CHIMED
CHI51240	China	NA	Resolved Discharged	Traditional Chinese Medicine	144	CHIMED
CHI50202	China	NA	Unclear	Traditional Chinese Medicine	204	CHIMED
CHI49647	China	NA	Unclear	Traditional Chinese Medicine	132	CHIMED
CHI49543	China	Non-severe	Outpatient	Traditional Chinese Medicine	180	CHIMED
CHI49348	China	Mixed	Resolved Discharged	Traditional Chinese Medicine	100	CHIMED
CHI49408	China	NA	Hospital	Traditional Chinese Medicine	408	CHIMED
CHI49453	China	NA	Unclear	Traditional Chinese Medicine	120	CHIMED
CHI49287	China	NA	Unclear	Traditional Chinese Medicine	200	CHIMED
CHI49380	China	Severe	Hospital	Traditional Chinese Medicine	160	CHIMED
CHI48792	China	NA	Unclear	Traditional Chinese Medicine	200	CHIMED
CHI48985	China	NA	Healthy Exposed	Traditional Chinese Medicine	600	CHIMED
CHI48988	China	NA	Outpatient	Traditional Chinese Medicine	400	CHIMED
CHI49080	China	NA	Unclear	Traditional Chinese Medicine	1000	CHIMED
CHI51139	China	NA	Unclear	Traditional Chinese Medicine	400	CHIMED
CHI49051	China	NA	Unclear	Traditional Chinese Medicine	400	CHIMED
CHI48860	China	Mixed	Outpatient, Hospital	Traditional Chinese Medicine	140	CHIMED
CHI48824	China	NA	Unclear	Traditional Chinese Medicine	60	CHIMED
CHI50082	China	Mixed	Hospital	Traditional Chinese Medicine	96	CHIMED
CHI50462	China	NA	Resolved Discharged	Traditional Chinese Medicine	20	CHIMED
CHI51028	China	Non-severe	Unclear	Traditional Chinese Medicine	24	CHIMED
CHI50793	China	NA	Resolved Discharged	Traditional Chinese Medicine	120	CHIMED
CHI49014	China	NA	Unclear	Traditional Chinese Medicine	400	CHIMED

CHI49852	China	Severe	Unclear	Traditional Chinese Medicine	60	CHIMED
CHI49387	China	NA	Unclear	Traditional Chinese Medicine	30	CHIMED
CHI49490	China	Severe	Unclear	Traditional Chinese Medicine	80	CHIMED
CHI48886	China	Severe	Hospital	Traditional Chinese Medicine	42	CHIMED
IR/46775	Iran	NA	Hospital	Traditional medicine	80	CHIMED
IR/46782	Iran	NA	Hospital	Traditional medicine	68	CHIMED
NCT04251871	China	NA	Unclear	Traditional Medicine Oral	150	CHIMED
CHI48898	China	NA	Unclear	Traditional Medicine Oral	240	CHIMED
CHI48889	China	Non-severe	Unclear	Traditional Medicine Oral	240	CHIMED
CHI50476	China	NA	Resolved Discharged	Traditional Medicine Oral	100	CHIMED
CHI50986	China	Non-severe	Unclear	Traditional Medicine Oral	100	CHIMED
CHI51018	China	Non-severe	Outpatient	Traditional Medicine Oral	128	CHIMED
CHI48773	China	None	Healthy Unexposed	Traditional Medicine Oral	20000	CHIMED
CHI44213	China	Mixed	Hospital	Traditional Medicine Oral	100	CHIMED
CHI49301	China	Mixed	Hospital	Traditional Medicine Oral	120	CHIMED
CHI49138	China	Mixed	Hospital	Traditional Medicine Oral	160	CHIMED
CHI49402	China	Mixed	Outpatient, Hospital	Traditional Medicine Oral	300	CHIMED
CHI49502	China	NA	Hospital	Traditional Medicine Oral	110	CHIMED
CHI49422	China	NA	Unclear	Traditional Medicine Oral	600	CHIMED
CHI48861	China	NA	Unclear	Traditional Medicine Oral	100	CHIMED
IR/46678	Iran	Unspecified	Hospital	Traditional Medicine Oral	374	CHIMED
IR/46721	Iran	Unspecified	Outpatient	Traditional Medicine Oral	100	CHIMED
CHI50126	China	NA	Hospital	Traditional Medicine Oral	300	CHIMED
CHI48775	China	NA	Unclear	Traditional Medicine Oral	300	CHIMED
NCT04279197	China	NA	Unclear	Traditional Medicine Oral	136	CHIMED
CHI52958	China	Unclear	Unclear	Traditional medicine oral	120	CHIMED
CHI52942	China	Unclear	Unclear	Traditional medicine oral	200	CHIMED
CHI49435	China	Mixed	Unclear	Traditional Medicine Oral	40	CHIMED
NCT04347382	Pakistan	NA	Hospital	Traditional Medicine Oral	30	CHIMED
CHI49666	China	Mixed	Unclear	Traditional Medicine Oral	72	CHIMED
CHI50662	China	NA	Unclear	Traditional Medicine Oral	50	CHIMED
CHI49753	China	Mixed	Hospital	Traditional Medicine Oral	20	CHIMED
CHI49425	China	Mixed	Hospital	Traditional Medicine Oral	72	CHIMED
CHI49131	China	NA	Hospital	Traditional Medicine Oral	60	CHIMED
IR/46637	Iran	Unspecified	Hospital	Traditional Medicine Oral	60	CHIMED
IR/46715	Iran	Unspecified	Hospital	Traditional Medicine Oral	60	CHIMED

IR/47019	Iran	Unclear	Unclear	Traditional medicine oral	80	CHIMED
CHI50059	China	Severe	Hospital	Plasma Product	100	CONV PLASMA
NCT04355897	United States	NA	Hospital	Convalescent plasma	100	CONV PLASMA
NCT04362176	United States	NA	Hospital	Convalescent plasma	500	CONV PLASMA
NCT04359810	United States	NA	Hospital	Convalescent plasma	105	CONV PLASMA
NCT04357106	Mexico	NA	ICU	Convalescent plasma	10	CONV PLASMA
NCT04347681	Saudi Arabia	NA	Hospital, ICU	Convalescent plasma	40	CONV PLASMA
IR/46973	Iran	NA	Hospital	Convalescent plasma	60	CONV PLASMA
IR/47115	Iran	NA	Hospital	Convalescent plasma	30	CONV PLASMA
IR/47266	Iran	NA	ICU	Convalescent plasma	10	CONV PLASMA
CHI50537	China	Mixed	Hospital	Convalescent plasma	50	CONV PLASMA
IR/46931	Iran	NA	Hospital	Convalescent plasma	12	CONV PLASMA
IR/47212	Iran	NA	Hospital	Immunoglobulin	15	CONV PLASMA
IR/46811	Iran	NA	Hospital	Immunoglobulin	50	CONV PLASMA
IR/47014	Iran	NA	Hospital	Immunoglobulin, HCQ, LPV/r, Ribavirin	100	CONV PLASMA
NCT04342182	Netherlands	Severe	Hospital	Plasma	426	CONV PLASMA
NCT04344535	United States	Mixed	Hospital	Plasma	500	CONV PLASMA
NCT04345523	Spain	NA	Hospital	Plasma	278	CONV PLASMA
IR/46759	Iran	Severe	Hospital	Plasma	200	CONV PLASMA
NCT04364737	United States	NA	Hospital	Plasma	300	CONV PLASMA
NCT04346446	India	Severe	Hospital	Plasma	20	CONV PLASMA
NCT04343755	United States	Severe	Hospital	Plasma	55	CONV PLASMA
NCT04327349	Iran	NA	Unclear	Plasma	30	CONV PLASMA
IR/47058	Iran	NA	ICU	Plasma	32	CONV PLASMA
IR/47163	Iran	Severe	Hospital, ICU	Plasma	357	CONV PLASMA
CHI49081	China	Severe	Hospital	Plasma Product	200	CONV PLASMA
NCT04343261	United States	NA	Hospital	Plasma Product	15	CONV PLASMA
NCT04340050	United States	NA	Hospital	Plasma Product	10	CONV PLASMA
CHI51072	China	Severe	ICU	Plasma Product	10	CONV PLASMA
CHI50727	China	Severe	Hospital	Plasma Product	30	CONV PLASMA
CHI49544	China	Mixed	Outpatient, Hospital	Plasma Product	90	CONV PLASMA
CHI49861	China	Mixed	Hospital	Plasma Product	10	CONV PLASMA
CHI49533	China	NA	ICU	Plasma Product	20	CONV PLASMA
NCT04321421	Italy	Severe	ICU	Plasma Product	49	CONV PLASMA
IR/47169	Iran	NA	Unclear	Purified protein derivative	50	DIAGNOSTIC
CHI49664	China	NA	Unclear	Electrotherapy	40	ELECTROTHERAPY

IR/47015	Iran	NA	ICU	Hemodialysis	10	HEMODIALYSIS
IR/47234	Iran	NA	Outpatient	Echinacea + Ginger	100	HERBAL
NCT04322344	Italy	NA	Hospital	Escin	120	HERBAL
IR/47018	Iran	NA	Hospital	Foot bath	60	HYGIENE
NCT04344236	United States	NA	Unclear	Nasal rinses	48	HYGIENE
NCT04254874	China	Mixed	Unclear	Combination Antiviral	100	IFN
NCT04350281	China	Severe	Hospital, ICU	IFN, HCQ	80	IFN
NCT04350671	Iran	Severe	Hospital	IFN, HCQ, LPV/r,	40	IFN
CHI50470	China	NA	Unclear	Interferon Therapy	328	IFN
EU/2020-001023-14/GB/	United Kingdom	NA	Hospital	Interferon Therapy	400	IFN
NCT04331899	United States	Non-severe	Outpatient	Interferon Therapy	120	IFN
NCT04320238	China	None	Healthcare Workers	Interferon Therapy	2944	IFN
CHI49224	China	Mixed	Unclear	Interferon Therapy	100	IFN
IR/46665	Iran	Unspecified	Hospital	Interferon Therapy	40	IFN
NCT04365153	United States	NA	Hospital	Canakinumab	45	IMMUNE MODULALTOR
CHI50934	China	NA	ICU	Blood purification	20	IMMUNE MODULATOR
NCT04347226	United States	NA	Hospital	BMS-986253	138	IMMUNE MODULATOR
NCT04371367	France	NA	ICU	Avdoralimab	108	IMMUNE MODULATOR
EU/2020-001736-95/GB/	United Kingdom	Mixed	Unclear	Bemcentinib	825	IMMUNE MODULATOR
NCT04317040	United States	Severe	Hospital	Cd24f	230	IMMUNE MODULATOR
NCT04343989	United States	NA	ICU	Clazakizumab	90	IMMUNE MODULATOR
NCT04345614	United States	Severe	Hospital	CM4620	120	IMMUNE MODULATOR
NCT04328480	Argentina	Severe	Hospital	Colchicine	2500	IMMUNE MODULATOR
NCT04322682	United States	NA	Outpatient	Colchicine	6000	IMMUNE MODULATOR
NCT04360980	Iran	NA	Hospital	Colchicine	80	IMMUNE MODULATOR
NCT04363437	United States	NA	Hospital	Colchicine	70	IMMUNE MODULATOR
CHI48809	China	NA	Hospital	Cyotkine protein injection	90	IMMUNE MODULATOR
NCT04361526	Spain	NA	ICU	Cytokine Adsorption	40	IMMUNE MODULATOR
CHI49065	China	NA	Outpatient, Hospital	Cytokine Protein Injection	480	IMMUNE MODULATOR
NCT04344080	Germany	NA	ICU	Cytosorb	24	IMMUNE MODULATOR
NCT04346797	France	NA	Hospital, ICU	Eculizumab	120	IMMUNE MODULATOR
NCT04324021	Italy	Mixed	Healthcare Workers, Unclear	Emapalumab, Anakinra	54	IMMUNE MODULATOR
IR/46837	Iran	Severe	Hospital, ICU	Extracorporeal Blood Purification	10	IMMUNE MODULATOR
CHI49632	China	NA	ICU	Extracorporeal Blood Purification	60	IMMUNE MODULATOR
IR/46786	Iran	Unspecified	Hospital	Fingolimod	40	IMMUNE MODULATOR
NCT04280588	China	Mixed	Hospital	Fingolimod	30	IMMUNE MODULATOR

NCT04333420	Netherlands	Severe	Hospital	IFX-1	130	IMMUNE MODULATOR
CHI50017	China	Severe	Hospital	Jakotinib Hydrochloride	16	IMMUNE MODULATOR
NCT04351152	United States	Mixed	Hospital	Lenziulumab	238	IMMUNE MODULATOR
NCT04343651	United States	NA	Unclear	Leronimab	75	IMMUNE MODULATOR
NCT04343651	United States	NA	Unclear	Leronimab	75	IMMUNE MODULATOR
NCT04343651	United States	NA	Unclear	Leronimab	75	IMMUNE MODULATOR
NCT04347239	United States	NA	Hospital	Leronlimab	390	IMMUNE MODULATOR
NCT04342897	United States	NA	Hospital	LY3127804	200	IMMUNE MODULATOR
CHI48907	China	NA	Unclear	M1 Macrophage Suppressor	45	IMMUNE MODULATOR
NCT04334044	Mexico	NA	Unclear	Ruxolitinib	20	IMMUNE MODULATOR
NCT04338958	Germany	Severe	Hospital	Ruxolitinib	200	IMMUNE MODULATOR
NCT04326920	Belgium	Severe	Hospital	Sargramostim	80	IMMUNE MODULATOR
EU/2020-001162-12/FR/	Canada	Severe	Hospital	Sarilumab	460	IMMUNE MODULATOR
NCT04327388	Canada	Severe	Hospital	Sarilumab	300	IMMUNE MODULATOR
NCT04315298	United States	Severe	Hospital	Sarilumab	400	IMMUNE MODULATOR
NCT04359901	United States	NA	Hospital	Sarilumab	120	IMMUNE MODULATOR
NCT04357808	Spain	NA	Hospital	Sarilumab	30	IMMUNE MODULATOR
NCT04349098	United States	Severe	Hospital	Selinexor	230	IMMUNE MODULATOR
NCT04329650	Spain	Mixed	Hospital	Siltuximab	200	IMMUNE MODULATOR
NCT04348695	Spain	Mixed	Hospital	Simvastatin, ruxolitinib	94	IMMUNE MODULATOR
NCT04341675	United States	Mixed	Hospital	Sirolimus	30	IMMUNE MODULATOR
NCT04371640	United States	NA	Hospital	Sirolimus	40	IMMUNE MODULATOR
NCT04322773	Denmark	Severe	Hospital	TCZ	200	IMMUNE MODULATOR
GERDRKS00021238	Germany	NA	ICU	TCZ	200	IMMUNE MODULATOR
NCT04335071	Switzerland	Severe	Hospital	TCZ	100	IMMUNE MODULATOR
CHI49409	China	Mixed	Hospital	TCZ	188	IMMUNE MODULATOR
NCT04320615	United States	Severe	Hospital	TCZ	330	IMMUNE MODULATOR
NCT04317092	Italy	Severe	Hospital	TCZ	400	IMMUNE MODULATOR
NCT04361552	United States	NA	Hospital	TCZ	180	IMMUNE MODULATOR
NCT04363853	Mexico	Severe	Hospital, ICU	TCZ	200	IMMUNE MODULATOR
NCT04331795	United States	NA	Hospital	TCZ	50	IMMUNE MODULATOR
NCT04346355	Italy	Mixed	Hospital	TCZ	398	IMMUNE MODULATOR
IR/47021	Iran	NA	ICU	TCZ	10	IMMUNE MODULATOR
NL8504	Netherlands	NA	Hospital	TCZ	354	IMMUNE MODULATOR
EU/2020-001770-30/BE/	Belgium	Severe	Hospital	TCZ	60	IMMUNE MODULATOR
CHI50693	China	Severe	Hospital	TCZ , Adalimumab	60	IMMUNE MODULATOR

NCT04335305	Spain	Severe	Hospital	TCZ, pembrolizumab	24	IMMUNE MODULATOR
CHI49161	China	Severe	Hospital	Thymosin	120	IMMUNE MODULATOR
NCT04341116	United States	Severe	Hospital	TJ003234	144	IMMUNE MODULATOR
CHI49215	China	NA	Hospital	vMIP	40	IMMUNE MODULATOR
NCT04350580	France	Severe	ICU	Immunoglobulin	138	IVIG
IR/47219	Iran	Severe	Hospital, ICU	IVIG, LPV/r	20	IVIG
CHI49075	China	Mixed	Hospital	Asc09/Ritonavir	160	LPV
CHI49514	China	Mixed	Outpatient, Hospital, ICU	Carriomycin	520	LPV
CHI49263	China	NA	Outpatient	HCQ, LPV/r	112	LPV
NCT04364022	Switzerland	NA	Healthy Exposed	HCQ, LPV/r	420	LPV
CHI48684	China	NA	Hospital	LPV/r	160	LPV
CHI48991	China	NA	Outpatient	LPV/r	328	LPV
NCT04330690	Canada	Mixed	Hospital	LPV/r	440	LPV
NCT04321174	Canada	None	Healthy Exposed	LPV/r	1220	LPV
NCT04307693	South Korea	Non-severe	Outpatient	LPV/r	150	LPV
CHI50057	China	NA	Unclear	LPV/r	60	LPV
NCT04365127	United States	NA	Hospital	Progesterone	40	MATABOLIC
NCT04365699	United States	NA	Hospital	AT-001	500	METABOLIC
IR/46639	Iran	NA	Hospital	Atorvastatin	100	METABOLIC
NCT04350593	United States	Mixed	Hospital	Dapagliflozin	900	METABOLIC
NCT04359329	United States	NA	Outpatient	Estradiol	110	METABOLIC
IR/46762	Iran	NA	Outpatient	Medium-chain triglyceride	120	METABOLIC
IR/46690	Iran	Mixed	Hospital	Metformin	200	METABOLIC
IR/47334	Iran	NA	Hospital	Sitagliptin	84	METABOLIC
IR/47141	Iran	Unclear	Outpatient	Lithium	40	MINERAL
IR/47244	Iran	Unclear	Hospital	Azithromycin + Oseltamivir + LPV/r + Atazanavir + Hydroxychloroquine	110	MULTITHERAPY
IR/46538	Iran	NA	Hospital	HCQ , Oseltamivir , LPV/r , IFN	30	MULTITHERAPY
NCT04366089	Italy	NA	Hospital	HCQ, Azithromycin, Oxygen, Probiotic, Dietary supplement	152	MULTITHERAPY
IR/46810	Iran	NA	Hospital	HCQ, Corticosteroids, Interferon, Immunoglobulin	105	MULTITHERAPY
IR/46853	Iran	NA	Hospital	HCQ, Sof+Dac, Lithium, Trifluoprazine + trihexyphenidyl	80	MULTITHERAPY
NCT04331470	Iran	NA	Unclear	Levamisole, LPV/r, HCQ	30	MULTITHERAPY
NCT04350684	Iran	Mixed	Hospital	IFN, HCQ, LPV/R, Umifenovir	40	MULTITHERAPY
IR/46681	Iran	Mixed	Hospital	TCZ, LPV/r, HCQ, oseltamivir	40	MULTITHERAPY
IR/47052	Iran	Unclear	Unclear	LPV/r, Naproxen, HCQ, Oseltamivir	101	MULTITHERAPY
IR/46533	Iran	NA	Unclear	Pirfenidone, Chloroquine, LPV/r, Oseltamivir	40	MULTITHERAPY
NCT04366245	Spain	NA	Hospital	Plasma, HCQ, LPV/r, Azithromycin	72	MULTITHERAPY

NCT04315948	France	Severe	Hospital	Remdesivir, LPV/r, Interferon, HCQ	3100	MULTITHERAPY
CHI49088	China	Severe	Hospital	Stem Cell, Ruxolitinib	70	MULTITHERAPY
NCT04326426	United States	Severe	Hospital	Tradipitant	300	NEURO
NCT04343963	Mexico	NA	Hospital	Pyridostigmine Bromide	436	NEUROGENIC
NCT04368156	Spain	NA	Hospital	Vagus nerve stimulation	90	NEUROGENIC
NCT04346693	Russia	NA	Hospital	Dalargin	320	OPIOID
CHI49342	China	Non-severe	Unclear	Polyinosinic- Polycytidylic Acid	40	POLYNUCLEOTID
NL8491	Netherlands	Severe	Hospital	Imatinib	386	PROTEIN KINASE INHIBITOR
EU/2020-001236-10/NL/	Netherlands	Severe	Hospital	Imatinib mesilate	304	PROTEIN KINASE INHIBITOR
CHI49952	China	NA	Outpatient	Psychological	180	PSYCHOLOGIC
NCT04342663	United States	NA	Outpatient	Fluvoxamine	152	PSYCHOLOGIC
IR/47006	Iran	NA	Hospital	Trifluoperazine, HCQ	70	PSYCHOLOGIC
NCT04326452	United States	Severe	Hospital	Bidirectional oxygen	15	PULMONARY
IR/46969	Iran	NA	Hospital	Bromhexine	60	PULMONARY
NCT04355364	France	NA	ICU	Dornase Alfa Inhalation Solution	100	PULMONARY
CHI50910	China	Severe	ICU	ECMO	30	PULMONARY
GERDRKS00021300	Germany	Severe	Hospital	Ecmo	30	PULMONARY
IR/47173	Iran	Unclear	Unclear	Formoterol	50	PULMONARY
NCT04325906	United States	Severe	ICU	High nasal flow cannula	346	PULMONARY
EU/2020-001391-15/IE/	Ireland	NA	Hospital	HUMAN ALPHA1-PROTEINASE INHIBITOR	36	PULMONARY
NCT04332081	United States	NA	Hospital	Hyperbaric oxygen	40	PULMONARY
CHI49799	China	NA	Unclear	Inhaled solution	60	PULMONARY
NCT04340414	China	NA	Hospital	Low flow ECMO	10	PULMONARY
NCT04359251	China	NA	Hospital	Mechanical Support	20	PULMONARY
NCT04344730	France	NA	ICU	O2 support, Corticosteroids	550	PULMONARY
NCT04359862	Spain	NA	Hospital, Unclear	Sedation	50	PULMONARY
NCT04359862	Spain	NA	Hospital, Unclear	Sedation	50	PULMONARY
NCT04355026	Slovenia	NA	Hospital	HCQ, Bromhexine	90	PULMONARY
NCT04336462	China	Severe	Hospital	Hydrogen , oxygen	100	PULMONARY
NCT04344431	France	NA	Hospital	Hyperbaric oxygen	100	PULMONARY
EU/2020-001329-30/AT/	China	NA	ICU	Nitric oxide	200	PULMONARY
NCT04305457	United States	Mixed	Hospital	Nitric Oxide	240	PULMONARY
NCT04306393	United States	Severe	ICU	Nitric Oxide	200	PULMONARY
CHI49283	China	Mixed	Hospital	Oxygen	440	PULMONARY
NCT04320056	Canada	Severe	Hospital	Oxygen	216	PULMONARY
CHI49747	China	Mixed	Outpatient, Hospital	Ozone	180	PULMONARY

ID	Country	Severity	Setting	Intervention	N	Category
CHI49947	China	Mixed	Outpatient, Hospital	Ozone	60	PULMONARY
CHI49737	China	NA	Hospital	Ozone	60	PULMONARY
IR/47149	Iran	NA	Unclear	Ozone Major Autohemotherapy	80	PULMONARY
NCT04359797	United States	NA	Hospital	Prone positioning	300	PULMONARY
NCT04358939	France	NA	Hospital	Prone positioning	248	PULMONARY
NCT04344106	France	NA	Hospital, ICU	Prone positioning	25	PULMONARY
NCT04337541	Denmark	Non-severe	Healthcare Workers, Outpatient	Surgical facial mask	6000	PULMONARY
NCT04331366	United States	Mixed	Hospital	Bidirectional oxygen	5	PULMONARY
CHI48801	China	Severe	Hospital, ICU	Pirfenidone	292	PULMONARY FIBROSIS RX
NCT04282902	China	NA	Hospital	Pirfenidone	294	PULMONARY FIBROSIS RX
CHI51416	China	NA	Unclear	Pirfenidone	40	PULMONARY FIBROSIS RX
CHI51118	China	NA	Unclear	Pirfenidone	40	PULMONARY FIBROSIS RX
NCT04366791	United States	NA	Hospital	Radiation	10	RADIATION
NCT04330300	Ireland	NA	Healthy Unexposed	Antihypertensives	2414	RAS
NCT04364893	Brazil	NA	Hospital	ARB and ACEi	500	RAS
NCT04329195	France	NA	Hospital	Discontinuation of RAS	554	RAS
NCT04340557	United States	NA	Hospital	Losartan	200	RAS
IR/46824	Iran	Unspecified	Hospital	Losartan	100	RAS
NCT04312009	United States	Severe	Hospital	Losartan	200	RAS
NCT04311177	United States	Non-severe	Outpatient	Losartan	580	RAS
NCT04335123	United States	Severe	Hospital, ICU	Losartan	50	RAS
NCT04355936	Argentina	NA	Unclear	Telmisartan	400	RAS
NCT04335786	Netherlands	Mixed	Hospital	Valsartan	651	RAS
CHI50341	China	NA	Unclear	Physical rehabiliation	80	REHABILITATION
NCT04365738	Turkey	NA	Outpatient	Respiratory coaching	270	REHABILITATION
CHI50973	China	Severe	Hospital	Statin	100	SERINE PROTEASE INHIBITOR
ISRCTNISRCTN83971151	Argentina	NA	Hospital	LPV/r, IFN, HCQ, remdesivir	10000	SOLIDARITY
JPjRCTs031190269	Japan	Unspecified	Outpatient	Ciclesonide	100	STEROID
NCT04348305	Denmark	Severe	Hospital, ICU	Corticosteroids	1000	STEROID
NCT04355637	Spain	NA	Hospital	Corticosteroids	300	STEROID
NCT04343729	Brazil	NA	Unclear	Corticosteroids	420	STEROID
NCT04331054	France	NA	Hospital	Corticosteroids	436	STEROID
NCT04327401	Brazil	Mixed	Unclear	Corticosteroids	290	STEROID
CHI50453	China	NA	Outpatient, Hospital	Corticosteroids	200	STEROID
NCT04263402	China	Severe	Unclear	Corticosteroids	100	STEROID
NCT04344288	France	NA	Hospital	Corticosteroids	304	STEROID

ID	Country	Severity	Setting	Intervention	N	Category
IR/46904	Iran	NA	Hospital	Methylprednisolone	40	STEROID
IR/47088	Iran	NA	ICU	Methylprednisolone	40	STEROID
NCT04325061	Spain	Severe	Hospital	Corticosteroids	200	STEROIDS
IR/46776	Iran	NA	Hospital	Corticosteroids	48	STEROIDS
NCT04355247	Puerto Rico	NA	Hospital	Corticosteroids	20	STEROIDS
CHI48777	China	Severe	Hospital	Corticosteroids	48	STEROIDS
NCT04244591	China	Severe	ICU	Corticosteroids	80	STEROIDS
NCT04323592	Italy	Severe	Hospital	Corticosteroids	104	STEROIDS
NCT04341038	Spain	Severe	ICU	Corticosteroids, tacrolimus	84	STEROIDS
CHI49534	China	Severe	Hospital	Alpha Lipoic Acid	68	SUPPLEMENT
IR/47139	Iran	NA	Outpatient	Asafoetdida	60	SUPPLEMENT
IR/46868	Iran	NA	Outpatient	Berberine	40	SUPPLEMENT
IR/47193	Iran	NA	Hospital	Black myrobalan + Mastic + Sugarcane	10	SUPPLEMENT
IR/46875	Iran	NA	Hospital	Calcifediol	260	SUPPLEMENT
IR/47010	Iran	NA	Healthy Exposed, Healthcare Workers	Calcifediol	540	SUPPLEMENT
IR/46978	Iran	NA	Outpatient	Crocetin	40	SUPPLEMENT
IR/46554	Iran	NA	Hospital	Crostop/Croguard	150	SUPPLEMENT
IR/46879	Iran	NA	Hospital	Herbal	110	SUPPLEMENT
IR/46913	Iran	NA	Unclear	Herbal	130	SUPPLEMENT
IR/46828	Iran	NA	Hospital	Herbal	60	SUPPLEMENT
IR/46850	Iran	NA	Unclear	Herbal	60	SUPPLEMENT
IR/46907	Iran	NA	Unclear	Herbal	60	SUPPLEMENT
IR/46958	Iran	NA	Outpatient	Herbal	40	SUPPLEMENT
IR/46961	Iran	NA	Hospital	Herbal	15	SUPPLEMENT
IR/47016	Iran	NA	Hospital	Herbal	60	SUPPLEMENT
IR/46733	Iran	NA	Hospital	Herbal medicine	64	SUPPLEMENT
IR/46893	Iran	NA	Outpatient	Licorice	40	SUPPLEMENT
CHI50421	China	Mixed	Outpatient, Hospital	Lipoic Acid	394	SUPPLEMENT
IR/46903	Iran	NA	Hospital	Marshmallow + Licorice	60	SUPPLEMENT
IR/46963	Iran	NA	ICU	Melatonin, Vitamin C	20	SUPPLEMENT
NCT04323345	Egypt	NA	Unclear	Nutritional Support	1000	SUPPLEMENT
NCT04323514	Italy	Severe	Hospital	Nutritional Support	500	SUPPLEMENT
IR/47251	Iran	NA	Hospital	Pomegranate syrup	66	SUPPLEMENT
NCT04366180	Spain	NA	Healthcare Workers	Probiotic	314	SUPPLEMENT
CHI49321	China	Mixed	Outpatient, Hospital	Probiotics	300	SUPPLEMENT
CHI49530	China	Severe	Hospital	Regulating Intestinal Flora Agent	60	SUPPLEMENT

IR/47318	Iran	NA	Hospital	Syrup zofa	100	SUPPLEMENT
IR/46974	Iran	NA	Unclear	Vitamin A	140	SUPPLEMENT
NCT04264533	China	Severe	ICU	Vitamin C	140	SUPPLEMENT
NCT04357782	United States	NA	Hospital	Vitamin C	20	SUPPLEMENT
IR/46804	Iran	NA	Hospital	Vitamin C	60	SUPPLEMENT
NCT04342728	United States	Non-severe	Outpatient	Vitamin C, Zinc	520	SUPPLEMENT
NCT04344041	France	NA	Unclear	Vitamin D	260	SUPPLEMENT
IR/46751	Iran	Unspecified	Hospital	Vitaminc C	40	SUPPLEMENT
IR/46838	Iran	NA	ICU	Vitamins	60	SUPPLEMENT
NCT04328441	Netherlands	NA	Healthcare Workers	BCG Vaccine	1500	VACCINE
NCT04327206	Australia	NA	Healthcare Workers	BCG Vaccine	4170	VACCINE
NCT04352608	China	NA	Healthy Unexposed	Vaccine	744	VACCINE
EU/2020-001072-15/GB/	United Kingdom	NA	Healthy Unexposed	Vaccine	510	VACCINE
NCT04283461	United States	None	Healthy Unexposed	Vaccine	105	VACCINE
NCT04324606	United Kingdom	None	Healthy Unexposed	Vaccine	1112	VACCINE
NCT04276896	China	NA	Hospital	Vaccine	100	VACCINE
NCT04299724	China	NA	Healthy Unexposed, Healthy Exposed, Outpatient	Vaccine	100	VACCINE
CURPCEC00000306-En	Cuba	NA	Healthy Exposed	Vaccine	80	VACCINE
NCT04336410	United States	Non-severe	Healthy Unexposed	Vaccine	40	VACCINE
NCT04336410	United States	Non-severe	Healthy Unexposed	Vaccine	40	VACCINE

SUPPLEMENTARY TABLE 2: OTHER AGENTS. The interventions are listed by class. The setting of each study, whether hospital or outpatient, the type of control (AC: Active control, RCT: randomized control; soc: standard of care; PLAC: placebo control, whether treatment (TREAT) or prevention (PREV), the severity of illness, number of subjects, quarter study initiated, quarter study projects completion of enrollment, and duration of symptoms for patient inclusion are listed. Studies projecting completion of enrollment by the end of the second quarter of 2020 are bolded. Outpatient studies with planned completion by the end of the second quarter of 2020 are italicized.

TREATMENT	TYPE OF CONTROL	TREAT OR PREVENT	SEVERITY	NUMBER SUBJECTS	BEGIN DATE	PROPOSED END ENROLLMENT	DURATION SYMPTOMS
IMMUNE GLOBULIN							
CONVALESCENT PLASMA	RCT PLAC	TREAT	MIXED	500	Q2(20)	Q2(21)	14
	RCT (soc)	TREAT	MIXED	426	Q2(20)	Q3 (20)	4
	RCT (soc)	TREAT	MIXED	278	Q2(20)	Q3 (20)	12
	RCT (soc)	TREAT	SEV	200	Q1(20)	Q1(21)	3
	RCT (soc)	TREAT	SEV,CRIT	100	Q1(20)	**Q2(20)**	ANY DUR
IVIG	RCT (PLAC)	TREAT	SEV,CRIT	138	Q2(20)	Q3 (20)	ANY DUR
CELL THERAPY							
STEM CELL	RCT(PLAC)	PREV	HEALTHY	100	Q2(20)	Q4(20)	N/A
	Case control	TREAT	MIXED	200	Q1(20)	Q4(21)	ANY DUR
	RCT(PLAC)	TREAT	SEV	120	Q1(20)	Q2(20)	ANY DUR
CARDIOVASCULAR							
DISCONTINUATION OF RAS	AC (continue or discontinue RAS)	TREAT(2) PREV(1)	MIXED, HEALTHY	554;500;2414	Q2(20),Q1(20)	Q3 (20);Q4(20);Q1(21)	ANY DUR, N/A
LOSARTAN	RCT (soc)	TREAT	MIXED	100	COMPLETED		
	RCT (soc)	TREAT	MILD,MOD	200	Q2(20)	Q3 (20)	ANY DUR
	RCT (soc)	TREAT	MILD,MOD	580	Q2(20)	Q2(21)	3
	RCT (soc)	TREAT	SEV,CRIT	200	Q2(20)	Q2(21)	ANY DUR
TELMISARTAN	RCT(soc)	TREAT	MIXED	400	Q2(20)	Q3 (20)	ANY DUR
VALSARTAN	RCT (PLAC)	TREAT	MOD	651	Q2(20)	Q3 (20)	1

AMIODARONE	AC(verapamil)	TREAT	MIXED	804	Q2(20)	Q1(21)	ANY DUR

PULMONARY TREATMENT

PIRFENIDONE	RCT (soc)	TREAT	MIXED	294	Q1(20)	**Q2(20)**	7
	RCT (soc)	TREAT	SEV,CRIT	292	Q1(20)	Q3(20)	14
HYPERBARIC OXYGEN	RCT (soc)	TREAT	MIXED	100	Q2(20)	Q2(21)	ANY DUR
INHALED NITRIC OXIDE	RCT(soc)	TREAT	MIXED	240	Q1(20)	Q2(21)	ANY DUR
	RCT(soc)	TREAT	SEV,CRIT	200	Q1(20)	Q2(21)	ANY DUR
DORNASE ALFA (cystic fibrosis agent)	RCT (soc)	TREAT	CRIT	100	Q2(20)	Q4(20)	ANY DUR
PRONE POSITION	AC(supine)	TREAT(2)	CRIT; MIXED	248;300	Q2(20)	Q4(20); Q1(21)	ANY DUR
OZONE	RCT (soc)	TREAT	MIXED	180	Q1(20)	**Q2 (20)**	7
OXYGEN/HYDROGEN	AC (oxygen)	TREAT(2)	MOD,SEV	100;440	Q1(20)	Q3 (20)	ANY DUR
SURGICAL FACE MASK	AC(face mask or none)	PREV	HEALTHY	6000	Q2(20)	Q3 (20)	N/A
CLOSED LOOP OXYGEN	AC (normal O2 delivery vs closed loop)	TREAT	MOD,SEV	216	Q2(20)	Q4(20)	ANY DUR
RESPIRATORY COACHING	RCT (soc)	TREAT	MILD,MOD	270	Q1(20)	Q3 (20)	ANY DUR

ONCOLOGIC AGENTS

IMATINIB	RCT (PLAC)	TREAT(2)	SEV	304;386	Q1(20)	**Q2 (20)**	ANY DUR
LY3127804	RCT (PLAC)	TREAT	MIXED	200	Q2(20)	Q3 (20)	ANY DUR
BEVACIZUMAB	RCT (soc)	TREAT	SEV,CRIT	140	Q1(20)	**Q2(20)**	ANY DUR

METABOLIC AGENTS

METFORMIN	RCT (soc)	TREAT	MIXED	200	Q1(20)	**Q2 (20)**	ANY DUR
DAPLIGLIFLOZIN	RCT (PLAC)	TREAT	MILD,MOD	900	Q2(20)	Q4(20)	3
ESTRADIOL	RCT (soc)	TREAT	MILD,MOD	110	Q2(20)	Q4(20)	7
ATORVASTATIN	RCT (soc)	TREAT	MIXED	100	Q2(20)	NO DATE	ANY DUR
MEDIUM CHAIN TG	RCT (soc)	TREAT	MILD	120	Q2(20)	*Q2 (20)*	ANY DUR

SUPPLEMENTS

ECHINACEA AND GINGER	RCT (soc)	TREAT	MILD	100	Q2(20)	***Q2 (20)***	ANY DUR
VITAMIN D	AC (200,000 IU vs 50,000 IU), RCT(soc)	TREAT(3)	MILD,MOD	260;260;540	Q2(20)	Q3 (20), Q4(20)	3, ANY DUR
ASCORBIC ACID	RCT (PLAC)	TREAT	CRIT	140	Q1(20)	Q3 (20)	ANY DUR
ASCORBIC ACID, ZINC	AC(ascorbic acid, zinc, soc)	TREAT	MILD	520	Q2(20)	Q3 (20)	ANY DUR
LIPOIC ACID	RCT (soc)	TREAT	MIXED	394	Q1(20)	***Q2 (20)***	ANY DUR
PROBIOTICS	RCT (soc);RCT(PLAC)	TREAT;PREV	MIXED;HEALTHY	300;314	Q1(20);Q2(20)	Q3 (20)	ANY DUR
NATURAL HONEY	RCT (soc)	TREAT	MIXED	1000	Q1(20)	Q4(20)	ANY DUR
ESCIN	RCT (soc)	TREAT	MIXED	120	Q1(20)	**Q2 (20)**	ANY DUR
Crostop/Croguard	RCT (soc)	TREAT	MIXED	150	Q1(20)	**Q2 (20)**	ANY DUR
VITAMIN A	RCT (soc)	TREAT	MIXED	140	Q2(20)	NO DATE	ANY DUR

NEUROGENIC

TRADIPITANT	RCT (PLAC)	TREAT	SEV,CRIT	300	Q2(20)	Q3 (20)	ANY DUR

PYRIDOSTIGMINE	RCT (PLAC)	TREAT	SEV	436	Q2(20)	Q2(21)	ANY DUR
VAZEGEPANT	RCT (PLAC)	TREAT	SEV	120	Q2(20)	Q3 (20)	ANY DUR

PSYCHOLOGICAL

FLUVOXAMINE	RCT (PLAC)	TREAT	MILD	152	Q2(20)	Q3 (20)	ANY DUR
PSYCHOLOGICAL THERAPY	RCT (soc)	TREAT	MIXED	180	Q1(20)	Q1(22)	ANY DUR

GLUCOCORTICOIDS

METHYLPREDNISOLONE	AC (high vs low dose)	TREAT	SEV	100	Q1(20)	Q2 (20)	ANY DUR
CORTICOSTEROID	AC (early vs late vs no steroid)	TREAT	MIXED	200	Q1(20)	Q2 (20)	ANY DUR
DEXAMETHASONE	RCT (soc)	TREAT(2)	CRIT	290;200	Q2(20)	Q3 (20),Q4(20)	1,ANY DUR
CICLESONIDE	RCT (PLAC)	TREAT	SEV	120	Q2(20)	Q3 (20)	ANY DUR
SYMBICORT	RCT (soc)	TREAT	MIXED	436	Q2(20)	Q3 (20)	2

ANTICOAGULATION

ASPIRIN	RCT (soc)	TREAT	MIXED	128	Q1(20)	Q2 (20)	ANY DUR
LOW MOLECULAR WEIGHT HEPARIN	AC(lmw vs mechanical VTE prev)	PREV	MIXED	120	Q1(20)	Q2 (20)	ANY DUR
DEFIBROTIDE	RCT (PLAC)	TREAT	SEV,CRIT	120	Q2(20)	Q3 (20)	ANY DUR
ULINASTATIN	RCT (soc)	TREAT	MIXED	100	Q1(20)	Q2 (20)	ANY DUR