# Comprehensive mycotoxin exposure biomonitoring in breastfed and non-exclusively breastfed Nigerian children

Chibundu N. Ezekiel[1,2*], Wilfred A. Abia[2,3,4], Dominik Braun[5], Bojan Šarkanj[2,6], Kolawole I. Ayeni[1], Oluwawapelumi A. Oyedele[1], Emmanuel C. Michael-Chikezie[7,8], Victoria C. Ezekiel[9], Beatrice Mark[10], Chinonso P. Ahuchaogu[11], Rudolf Krska[2,3], Michael Sulyok[2], Paul C. Turner[12] and Benedikt Warth[5,*]

[1]*Department of Microbiology, Babcock University, Ilishan Remo, Ogun State, Nigeria. [2]Institute of Bioanalytics and Agro–Metabolomics, Department of Agrobiotechnology (IFA–Tulln), University of Natural Resources and Life Sciences Vienna (BOKU), Konrad Lorenzstr. 20, A–3430 Tulln, Austria. [3]Institute for Global Food Security, School of Biological Sciences, Queen's University Belfast, University Road, Belfast, BT7 1NN, Northern Ireland, United Kingdom. [4]Laboratory of Pharmacology and Toxicology, Department of Biochemistry, Faculty of Science, University of Yaounde I, P.O. Box 812, Yaounde, Cameroon. [5]University of Vienna, Faculty of Chemistry, Department of Food Chemistry and Toxicology, Währinger Str. 38, A-1090 Vienna, Austria. [6]Department of Food Technology, University North, Center Koprivnica, Trg dr. Zarka Dolinara 1, HR, 48000, Koprivnica, Croatia. [7]Clifford University, Owerrinta (Ihie Campus), Abia State, Nigeria. [8]Benjamin Carson (Snr.) School of Medicine, Babcock University, Ilishan Remo, Ogun State, Nigeria. [9]Independent Researcher, Ilishan Remo, Ogun State, Nigeria. [10]Department of Community Health, Babcock University Teaching Hospital, Ilishan Remo, Ogun State, Nigeria. [11]Department of Clinical Sciences, Babcock University Teaching Hospital, Ilishan Remo, Ogun State, Nigeria. [12]MIAEH, School of Public Health, University of Maryland, College Park, MD 20742, USA.

Corresponding Authors

*Chibundu N. Ezekiel, Department of Microbiology, Babcock University, Ilishan Remo, Ogun State, Nigeria. Email: chaugez@gmail.com

*Benedikt Warth, University of Vienna, Faculty of Chemistry, Department of Food Chemistry and Toxicology, Währinger Straße 38, 1090 Vienna, Austria. Phone: +43 1 4277 70806. E-mail: benedikt.warth@univie.ac.at

# Abstract

**BACKGROUND**: Infants are a vulnerable population whose nutrition changes as complementary foods are introduced; a process which may modify patterns of exposure to dietary mycotoxins. However, exposure monitoring of dietary mycotoxin mixtures in several biological samples obtained from breastfed and non-exclusively breastfed children is scarce.

**OBJECTIVES**: To examine mycotoxin co-exposure patterns in infants using a multi-specimen, multi-mycotoxin approach.

**METHODS**: Breast milk, complementary food and urine obtained from 65 infants, aged 1–18 months, in Ogun state, Nigeria, were analyzed for mycotoxins using ultra-sensitive LC-MS/MS approaches.

**RESULTS**: Complementary food was contaminated with seven distinct classes of mycotoxins including aflatoxins (9/42 samples; range: 1.0–16.2µg/kg) and fumonisins (14/42; range: 8–167µg/kg). Aflatoxin $M_1$ was detected in breast milk (4/22), while six other classes of mycotoxins were quantified; including dihydrocitrinone (6/22; range: 14.0–59.7ng/L) and sterigmatocystin (1/22; 1.2ng/L) detected for the first time. Mycotoxins were detected in 64/65 of the urine samples, with seven distinct classes of mycotoxins observed demonstrating ubiquitous exposure. Two aflatoxin metabolites ($AFM_1$ and $AFQ_1$) and $FB_1$ were detected in 6/65, 44/65 and 17/65 samples, respectively. Mixtures of mycotoxin classes were common, including 14/42, 22/22 and 56/65 samples having 2–4, 2–6 or 2–6 mycotoxins present, for complementary food, breast milk and urine, respectively. Aflatoxin and/or fumonisin was detected in 12/14, 4/22 and 46/56 for complimentary foods, breast milk and urine, respectively. Furthermore, the detection frequency, mean concentrations and occurrence of mixtures were typically greater in urine of non-exclusively breastfed compared to breastfed children.

50  **CONCLUSIONS**: The study provides novel insights into mycotoxin co-exposures in children in a mycotoxin high-risk country without proper food safety measures. Albeit a small sample set, it highlights significant transition to higher levels of infant mycotoxin exposure as complementary foods are introduced, providing impetus to mitigate during this critical early-life period and encourage breastfeeding.

## Introduction

Nutrition within the first 1000 days of life is crucial to the growth, development and performance of children in their later years and into adulthood as postulated in the 'developmental origins of health and disease' (DOHaD) concept (Barker 2000; Bateson et al. 2004; Mandy and Nyirenda 2018). However, children's diets within this period may be diverse and complex, and depend on the age of the children. The diet categories can include exclusively consumed breast milk, breast milk combined with complementary foods, and exclusively consumed complementary foods. Human breast milk composition varies widely depending on diet, age, lactation stage, number of pregnancies, and other physiological parameters (Innis 2014; Butts et al. 2018; Wu et al. 2018). Complementary foods are mainly made from single or mixed cereals as a porridge, sometimes in combination with nuts, animal milk and other mashed items. These foods also become more diverse as children get older (Kimanya et al. 2009, 2010, 2014; Alvito et al. 2010; Kamala et al. 2016; Chuisseu Njamen et al. 2018; Ojuri et al. 2018, 2019). Unfortunately, in some parts of the world, such foods may contain a number of fungal metabolites known as mycotoxins (Kimanya et al. 2009; 2010, 2014; Alvito et al. 2010; Juan et al. 2014; Cherkani-Hassani et al. 2016; Kamala et al. 2016; Warth et al. 2016; Chuisseu Njamen et al. 2018; Braun et al. 2018; Ojuri et al. 2018, 2019; Preindl et al. 2019).

Mycotoxins such as aflatoxin (AF), fumonisin (FUM), ochratoxin A (OTA), deoxynivalenol (DON), and zearalenone (ZEN) are considered priority mycotoxins and are regulated in food. AFs are categorized as class 1 human liver carcinogen (IARC 2002, 2012a). AF is additionally implicated in immune system suppression, interference with micronutrient absorption, impairment of gut integrity and reduction in normal child growth (Gong et al. 2002, 2003, 2004, 2012; Turner et al. 2003, 2007; Turner 2013; IARC 2015; Shirima et al. 2015; Watson et al. 2016, 2018). FUM intake has been associated with birth defects (Missmer et al. 2006), poor

child growth (Shirima et al. 2015) and the incidence of esophageal cancer (Chu et al. 1994;

Yoshizawa et al. 1994; Sun et al. 2007, 2011); OTA is linked to renal toxicity (Heussner and

Bingle, 2015), while immune system modulation was reported for trichothecenes such as DON

(Pestka 2010). Furthermore, *in vitro* data suggest interactions of regulated mycotoxins with

emerging ones such as alternariol (AOH) and its methylated ether form (alternariol monomethyl

ether; AME), beauvericin (BEA), enniatins and moniliformin (MON) (Vejdovszky et al. 2017a,

2017b; Aichinger et al 2019; Woelflingseder 2019).

Complementary foods in Africa can be contaminated with mycotoxins (Chuisseu Djamen

et al. 2018; Kimanya et al. 2009, 2010, 2014; Juan et al. 2014; Kolakowski et al. 2016;

Cappozzo et al. 2017; Oueslati et al. 2018; Ojuri et al. 2018, 2019), and ingested mycotoxins or

their metabolites may be found in various bio-fluids including blood, urine and breast milk

(Cherkani-Hassani et al. 2016; Warth et al. 2016; Braun et al. 2018; Ferrufino-Guardia et al.

2019; Polychronaki et al. 2008; Ediage et al. 2013; Ezekiel et al. 2014; Shirima et al. 2015;

Gerding et al. 2015; Heyndrickx et al. 2015; Schwartzbord et al. 2016; Ayelign et al. 2017; Chen

et al. 2017; Papageorgiou et al. 2018; Sanchez and Diaz 2019, Turner et al. 2012a; Vidal et al.

2018). Only a limited number of studies have assessed multiple mycotoxin exposure in Nigerian

children using biomarkers of exposure (Adejumo et al. 2013; Ezekiel et al. 2014, 2018b; Braun

et al. 2018; McMillan et al. 2018; Sarkanj et al. 2018), while none in any country has assessed

infant's exposures to multi-mycotoxins using multiple specimens (complementary foods and

breast milk) and additionally via urine.

This study comprehensively elucidated mycotoxin co-exposure patterns in infants using

three measures (food, breast-milk and urine) all with quantitative multiple mycotoxin analysis to

compare exposure levels in exclusively breastfed (EB) with non-exclusively breastfed (NEB)

infants.

## Methods

### *Study area and population*

The study population was 65 infants aged 1–18 months from Ilishan and Ikenne in Remo land of

Ogun state (south-western Nigeria). Both are small sister communities within 1 km distance of

each other, with rich diversity of foods available in households. Both communities are semi-

urban due to the presence of campuses of tertiary institutions and they share similar

characteristics as detailed below. Indigent families in these communities are farmers and

traders, and both communities consist of approximately 15,000 residents. The food sources for

families are local markets, small stores and own farms. Major foods consumed are cereal-based

products (rice, maize, wheat), tubers (cassava and yam), and groundnuts, melon seeds and

cowpea. The predominant breastfeeding practice in both communities involves the introduction

of complementary foods from the first month after birth; though timing depends on family

income, mother's health and work status, and knowledge on breastfeeding practices. A majority

of the families associated with the tertiary institutions practice exclusive breastfeeding from birth

till end of the sixth month.

### *Sampling design and ethical considerations*

This pilot, cross-sectional survey was conducted between January and February 2016. Families

that participated were identified through local health centres and post-natal clinics in the two

communities where mother/infant pairs were registered. The study consisted of two cohorts:

exclusively breastfed (EB) children (age: 1–6 months) and non-exclusively breastfed (NEB)

children (age: 3–18 months) to which 23 and 42 children (female/male ratio = 1:1) were

recruited, respectively. A total of 50 and 15 infants were recruited into the study from Ilishan and

Ikenne communities, respectively. The age of infants recruited into the study were ascertained

from their health records. Infants with known medical conditions (e.g. jaundice, HIV positive or

visibly malnourished) were excluded during the recruitment stage as it was the purpose of this

study to investigate typical background exposure levels and compare EB with NEB rather than

targeting the impact of exposure on health status. Prior to recruitment or inclusion into the study,

the mother of each child was informed in her preferred language (English or Yoruba) on the

purpose of the study. Only mothers who signed the informed consent document (and on behalf

of their infant) to participate in the research were included into the study.

Complementary foods were obtained as "plate-ready" samples ($n$=42; 50 g each), by a

trained team member, for the NEB children ($n$=42) on the day prior to infant urine sample

collection. The food samples collected included industrially-processed infant cereal ($n$=6),

fermented maize gruel *ogi* ($n$=26) and *tombran* (home-made pudding from mixed cereal and

nut; $n$=10). One food sample was collected per infant. Breast milk samples ($n$=22; 3–5 mL each)

from mothers of EB infants were collected on the day preceding the sampling of infant urine.

One mother did not provide breast milk sample due to intake of medications. Other details of

sampling and handling of the breast milk samples were described by Braun et al. (2018). In this

paper, the breast milk samples have been previously analysed for mycotoxins utilizing a less

sensitive method. First morning urine samples (10–40 mL each) were collected in 50 falcon

tubes from all 65 infants recruited into the study by the mothers, using their experience and

knowledge of the time at which their infant pass urine in the morning. All samples (plate-ready

complementary food, breast milk and urine) were immediately frozen at –20 $^{\circ}$C and couriered on

dry ice to Austria for LC-MS/MS analyses.

Ethical permission for this study was granted by the Babcock University Health

Research Ethics Committee under two approval numbers: BUHREC294/16 for the EB children

and BUHREC156/15 for the NEB children. In adherence to ethical standards, all samples were

blinded and coded to exclude participant information prior to analytical measurements.

***Food consumption questionnaires***

The mothers received a copy of a well-structured questionnaire for completion prior to food,

breast milk and urine sample collection. The questionnaires were administered by trained

interviewers and translation to the local language (Yoruba) was available on request of some

mothers. The administered questionnaire was designed to elicit information on participant

demography, family socio-economic status, individual dietary preferences and food

consumption patterns and associated health implications among the infants.

***Mycotoxin determination in breast milk samples***

Mycotoxins were quantified in breast milk samples as described in Braun et al. (2020). Briefly,

each homogenized breast milk sample (1 mL) was extracted using 1 mL acidified ACN (1%

formic acid) on a shaker for 3 min, followed by addition of anhydrous magnesium sulfate (0.4 g)

and sodium chloride (0.1 g). Sequel to two rounds of cold centrifugation (4,750 x g at 10 °C for

10 min; 14,000 x g, 4 °C for 2 min), a SPE clean up step using SPE column (Oasis PRiME

HLB®, Waters, Milford, MA, USA) followed. Mycotoxins were then concentrated on a vacuum

concentrator (Labconco, Missouri, USA) prior to reconstitution using 81 µL MeOH/ACN and

9 µL of the internal standard mixture. Mycotoxins in breast milk samples were measured on a

Sciex QTrap 6500$^+$ LC-MS/MS system (Foster City, CA, USA) equipped with a Turbo-V™

electrospray ionization (ESI) source coupled to an Agilent 1290 series UHPLC system

(Waldbronn, Germany). Analytes were separated on an Acquity HSS T3 column (2.1×100mm;

Waters, Vienna, Austria) with 1.8 µm particle size at 40 °C. ESI-MS/MS was performed in

scheduled multiple reaction monitoring (MRM) mode using fast polarity switching. For each

analyte two individual mass transitions were acquired.

## LC-MS/MS-based mycotoxin analysis of food samples

Plate-ready complementary food samples were analysed for the presence of multiple mycotoxins according to the dilute and shoot LC-MS/MS-based method described by Sulyok et al. (2020). This method covers the detection of more than 500 metabolites, inclusive of mycotoxins, in food samples. Briefly, 5 g of each food sample was extracted with 20 mL of acetonitrile/water/acetic acid 79:20:1, (v/v/v) in a 50 mL polypropylene tube (Sarstedt, Nümbrecht, Germany) for 90 min on a GFL 3017 rotary shaker (GFL, Burgwedel, Germany). Afterwards, the extract was diluted 1:1 (v/v) in dilution solvent (acetonitrile/water/acetic acid 20:79:1, v/v/v) prior to measurement on a Sciex QTrap 5500 LC-MS/MS System (Applied Biosystem, Foster City, CA, USA) equipped with TurboIonSpray electrospray ionisation (ESI) source and a 1290 Series HPLC System (Agilent, Waldbronn, Germany). Chromatographic separation was performed at 25°C on a Gemini® C18-column, 150×4.6 mm i.d., 5 µm particle size, equipped with a C18 4×3 mm i.d. security guard cartridge (Phenomenex, Torrance, CA, USA). ESI-MS/MS was performed in the scheduled MRM mode both in positive and negative polarities in two separate chromatographic runs per sample by scanning two fragmentation reactions per analyte. The MRM detection window of each analyte was set to its expected retention time ±20 and ±26 s in the positive and the negative modes, respectively. The identification of each positive analyte was confirmed when two MRMs per analyte was obtained (EC 2002). Further details on the method performance including data obtained through inter-laboratory comparison are described in Sulyok et al. (2020).

## Urinary mycotoxin measurements by UPLC-MS-MS

Mycotoxins in the urine samples were determined following a stable-isotope dilution assay-based LC-MS/MS method described in detail by Sarkanj et al. (2018). Briefly, each urine sample

was centrifuged for 3 min at 5600 x *g*, then treated    β-glucuronidase from *E. coli* Type IX-A

(Sigma-Aldrich, G7396-2MU) of prior to a SPE cleanup on Oasis PRiMEHLB® SPE columns

(Waters, Milford, MA, USA). Extracts were then evaporated under nitrogen, reconstituted with

470 µL dilution solvent (10% acetonitrile, 0.1% glacial acetic acid) and fortified with 30 µL of an

IS mixture. Measurement of mycotoxins in urine samples was performed on a Sciex QTrap

6500$^+$ LC-MS/MS system (Foster City, CA) equipped with a Turbo-V™ ESI source coupled to

an Agilent 1290 series UHPLC system (Waldbronn, Germany). Analytes were separated on an

AtlantisT3 HSS column (2.1×100mm; Waters, Wexford, Ireland) with 1.8 µm particle size at 35

°C. ESI-MS/MS was performed in scheduled MRM mode and with a 180 sec detection window.

At least two individual transitions were monitored for each analyte.

***Data analysis***

The IBM SPSS Statistics v21.0 (SPSS, Inc., Chicago, IL, USA) was applied in the analyses of

all data. Boxplots were created for the distribution of urinary mycotoxins in the two cohorts of

children. Urinary mycotoxin concentrations were logarithm transformed by the equation y =

$Log_{10}$ (1 + ng/L) to create a normal distribution for the boxplots. For comparison of urinary

mycotoxin levels in the population, means were separated by the Duncan's Multiple Range test

(DMRT) and tested for significance by analysis of variance (ANOVA) at α = 0.05. Whereas for

the comparison of all other data between the two cohorts of children in the study, the

independent sample *t*-test using the Levene's test for equality of variances were applied to test

for significance at α = 0.05.

## Results

### *Demography and food preference of children*

The demographic characteristics of the infants as well as their food consumption data are given in Table 1. The NEB infants were significantly ($p<0.01$) older (9.0±3.6 months) and heavier (8.0±1.6 Kg; $p<0.05$) than the EB children (3.7±1.6 months and 6.3±1.0 Kg, respectively). All infants received breast milk at least once daily, with higher consumption frequency of 8−10 times per day in the EB cohort compared to 1−3 times per day in the NEB cohort who received complementary food at least thrice per day. NEB infants received a variety of foods, in most cases (>90%) including infant cereal (mix of animal milk with maize, rice, or wheat), *ogi* (fermented maize gruel) and *tombran* (mixed cereal and nut pudding).

### *Occurrence of mycotoxins in breast milk*

Breast milk samples were examined for the presence of 34 mycotoxins (or their metabolites), of which nine were observed. All 22 breast milk samples contained mycotoxins although mostly at very low concentrations (Table 2). The most frequently detected mycotoxins were BEA (incidence: 100%; range: 0.99−12.0 ng/L; mean: 3.0 ng/L) and AME (incidence: 96%; range: 0.50−11.7 ng/L; mean: 3.03 ng/L), whereas DHC (incidence: 27%; range: 14.0−59.7 ng/L; mean: 25.2 ng/L) and OTA (incidence: 64%; range: 2.0−67.6 ng/L; mean: 9.58 ng/L) had higher mean levels than all other mycotoxins found in the samples. $AFM_1$ was the only aflatoxin metabolite detected in breast milk, occurring in 18% of the samples at levels below the LOQ; and thus, the data points were assigned LOQ/2 values of 2.0 ng/L (Table 2). Enniatins (EnnB and $EnnB_1$) occurred in 77% of the samples, with EnnB dominating in 73% of samples at mean concentration of 5.0 ng/L (range: 0.70−10.1 ng/L). Sterigmatocystin (STER) was detected in one

sample at 1.20 ng/L. The chromatograms of DHC and STER detected in breast milk are shown

in Figure 1 (C–D).

***Distribution of mycotoxins in plate-ready complementary food***

Sixty-five fungal metabolites, of which 14 were mycotoxins (Table 3), and 25 other secondary

metabolites (Table S1) were quantified in the samples of plate-ready complementary meals.

Aflatoxin $B_1$, $B_2$ and $G_1$ were detected, but not $AFG_2$. Total AFs contaminated 21% of food

samples of which the mean was 6.0 µg/kg (range: 1.0–16.0 µg/kg). The mean level of $AFG_1$ (12

µg/kg) deriving from two *ogi* samples was three-fold higher than that of $AFB_1$ (3 µg/kg). The

incidence of FUM (sum of fumonisins $B_1$ and $B_2$) was 29%, of which the mean was: 49 µg/kg

(range 7.9–194 µg/kg). BEA (incidence: 79%; mean: 3.8 µg/kg) was the most frequently

detected mycotoxin in the food samples. Enniatins, ZEN, MON and DON also occurred in 19,

10, 7 and 5% of the foods, respectively. With respect to occurrence of the toxins in food types,

the highest mean concentrations of BEA (19 µg/kg), enniatins (10 µg/kg) and FUM (63 µg/kg),

were found in infant cereal. Of the three food types, *ogi* contained the highest levels of AF (6.5

µg/kg) and DON (62 µg/kg). AF and MON were not found in infant cereal. *Ogi* and *tombran*

contained more diverse mycotoxins than infant cereal (Table 3).

***Mycotoxins in urines of infants***

*Overview of the distribution of mycotoxins in infant urines*

At least one mycotoxin was detected in 64/65 (98.5%) of the urine samples, with 10 distinct

mycotoxins observed, including aflatoxins ($AFM_1$ and $AFQ_1$), fumonisin $B_1$, DON, DHC, NIV,

OTA, zearalenones (ZEN, $\alpha$-ZEL, and $\beta$-ZEL), representing seven classes of mycotoxin, see

Table 4. The detected aflatoxins were two metabolites: $AFM_1$ (6/65, 9% of samples; mean of

detected (MoD): 140 ng/L; range: 20–510 ng/L) and $AFQ_1$ (44/65, 68% of samples; MoD: 210

ng/L; range: 3–900ng/L). Two infants had only $AFM_1$ detected, with no $AFQ_1$, thus in total 46/65

(71%) of the infants had been exposed to dietary aflatoxin. Only one of the fumonisin class of

mycotoxins was detected, $FB_1$ (17/65, 26% of samples; MoD: 810 ng/L; range: 40–1590 ng/L).

OTA, ZEN, DON and DHC were also frequently detected (Table 4). The chromatograms of

$AFQ_1$ and DHC detected in urine are given in Figure 1(A–B). Mycotoxins in urine tended to be

higher in females versus males, though no statistically significant differences in frequency or

concentration for individual toxins was observed (data not shown).

*Mycotoxins in exclusively breastfed (EB) and non-exclusively breastfed (NEB) infants' urine*

Eight of the 10 mycotoxins were observed frequently in NEB infants compared to EB infants:

$AFM_1$ (12 vs 4%), DON (55 vs 30%), DHC (74 vs 26%), $FB_1$ (31 vs 17%), OTA (38 vs 26%),

ZEN (83 vs 57%), respectively, and NIV and α-ZEL which were only detected in NEB infants'

urine (Table 4). The recorded detection frequencies of the mycotoxins resulted in an 46–300%

increase from EB to NEB cohorts (Figure 2). There were however reductions in the frequencies

of detection for $AFQ_1$ (53 vs 87%) and β-ZEL (14 vs 30%). The mean concentrations in urine

with detectable mycotoxins were greater for seven of the 10 mycotoxins, in NEB versus EB

infants: $AFM_1$ (166 vs 23 ng/L), $AFQ_1$ (225 vs 182 ng/L), DON (5,282 vs 3,194 ng/L), DHC (202

vs 135 ng/L), $FB_1$ (454 vs 136 ng/L), NIV (161 ng/L vs non-detect) and α-ZEL (20 ng/L vs non

detect), respectively, see Table 4; resulting in 124–721% increase (Figure 2). However,

significant higher mean levels were only recorded for $AFQ_1$, DON, DHC and $FB_1$ in favour of

NEB infants, and for OTA in favour of EB infants (Table 4 and Figure S1).

**Mycotoxin co-occurrence and co-exposures**

The numbers of different classes of mycotoxins detected in each of breast milk, complementary

food and urine were determined (Table 5). All breast milk samples contained more than one and

up to five mycotoxin classes; 41% of samples had four or more different mycotoxins detected,

though overall mycotoxin concentrations were low. The primary combinations in the mycotoxin

mixtures in breast milk included AME, BEA enniatins and OTA. Aflatoxins were observed as

part of 4/22 mixtures of toxins, and fumonisins were not measured in this study.

Mycotoxin mixtures, of 2–5 toxins, were observed in just over half (22/42) of the plate-

ready complementary food samples. Mixtures predominantly involved three classes of

mycotoxins: BEA in 12 mixtures, FUM in nine mixtures and AF in eight mixtures. Of these, five

samples with three or more toxins contained both AF and FUM.

Mixtures of 2–6 mycotoxins were observed in about 86% of the urines, with NEB

typically having more urine samples with mixtures (93%, with 2–6 toxins) compared to EB

infants (74% with 2–5 toxins). No urine from EB infants had six mycotoxins detected. The most

frequently occurring classes of toxins in mixtures were ZEN (45/65), AF (42/65), DHC (41/65)

and DON (30/65). AF and FB in the same urine occurred in 13/65 (20%). One infant from the

NEB cohort was co-exposed to seven mycotoxins and metabolites belonging to six of the

assessed mycotoxin classes except NIV (Figure 3).

**Discussion**

Infants transition from breastfeeding to complementary food is essential for nutritional

requirements, however, it is also important to understand if this transition creates opportunities

for differential exposure to diverse classes of fungal toxins and other environmental

contaminants (i.e. the exposome). The diversity of mycotoxins in breast milk is still in the

exploratory stage, though several mycotoxins have been observed, most notably aflatoxins;

however, food measurements are by far more established to date despite several limitations

(IARC 2012a, 2012b, 2015). This study provides a measure of multiple mycotoxins in both

breast milk and complementary plate ready foods, and then compares patterns of these

mycotoxins in urine from infants that are either EB or partially receiving breast milk and

complementary foods, NEB.

In this study the analytical sensitivity for measuring mycotoxins in breast milk was

extremely low, and thus the frequency of toxin detection was high providing unique insights into

chronic background level exposure patterns. All 22 samples had two or more toxins, though

overall the concentrations would be regarded as low. The lipophilic aflatoxins such as $AFB_1$ or

$AFG_1$ are commonly detected in grains in sub-Saharan Africa (Ezekiel et al. 2018a), but were

not observed in breastmilk. The $AFB_1$ metabolite, $AFM_1$, was observed in four samples and at a

level below the LOQ (4 ng/L). This data suggests that breast milk is not providing significant

exposure to the most carcinogenic of the mycotoxins for participants of this study. $AFM_1$ in

breast milk occurred at lower mean concentrations than those previously reported from Sierra

Leone (800 ng/L, Jonsyn et al. 1995), Egypt (60 ng/L, Polychronaki et al. 2007), Nigeria (35

ng/L, Adejumo et al. 2013) and Ecuador (45 ng/L, Ortiz et al. 2018), but similar to levels

reported in two samples from Brazil (0.3 and 0.8 ng/L, Iha et al. 2014).

Prior to the present study, there are no reports on the occurrence of CIT and its

metabolites as well as STER in human breast milk (Cherkani-Hassani et al. 2016; Warth et al.

2016; Ali and Degen 2019). Thus, to the best of our knowledge, we present the first report of

DHC and STER occurrence in human breast milk. DHC, mostly considered a detoxification

product of CIT but also co-detected with CIT in food samples (Ojuri et al. 2018, 2019; Ezekiel et

al. 2020), is indicative of CIT exposure (Follmann et al. 2014; Ali and Degen, 2019). The

concentrations of DHC in breast milk samples analyzed in the present study were low, and as

much as 10 times lower than the concentrations found in urine. However, the presence of DHC

in both biological fluids compel us to hypothesize that CIT intake is common in Nigeria (Ojuri et

al. 2018, 2019; Akinfala et al. 2020; Ezekiel et al. 2020). Furthermore, it agrees with the

suggestions of Ali and Degen (2019) who retrospectively estimated high CIT exposure

assessment/intake above the "level of no concern for nephrotoxicity" of 0.2 µg/kg bwt/day set by

EFSA (EFSA 2012; EC 2014) in participants in a previous urinary biomarker study in Nigeria

(Sarkanj et al. 2018). Due to sparse data on its occurrence and toxicity, there is no regulation on

maximum levels (ML) of STER in food, including infant diets, in most countries, Nigeria inclusive

(EC 2006; Oplatowska-Stachowiak et al. 2018). Nonetheless, this mycotoxin may be

carcinogenic in humans (IARC 1987; EFSA 2013), as such, its detection in breast milk samples

provides a basis for the consideration of more investigations into the occurrence and regulation

of this mycotoxin in foods intended for infants and young children.

The mean level of the nephrotoxic OTA in the analyzed breast milk was 50 times lower

than the maximum tolerable limit of 500 ng/L set for infant food in Europe (EC 2006).

Additionally, the recorded level was four to five times lower than those reported in Bolivia

(mean: 53 ng/L, Ferrufino-Guardia et al. 2019) and Chile (mean: 44 ng/L, Munoz et al. 2014) but

nearly two times higher than the mean level reported in Italy (mean: 6.01 ng/L, Turconi et al.

2004). The identification of BEA and EnnB in human breast milk agrees with previous reports

from Nigeria (Braun et al. 2018) and Spain (Rubert et al. 2014). The concentration level of AME

in the present study is, however, lower than those reported in a previous study from Nigeria

(Braun et al. 2020). Due to poor analytical recoveries at the outset, fumonisins were not

included in the breast milk analysis.

Mycotoxins have been previously reported in complementary foods (Juan et al. 2014;

Kolakowski et al. 2016; Kamala et al. 2016; Cappozzo et al. 2017; Chuisseu Njamen et al. 2018;

Oueslati et al. 2018; Ojuri et al. 2018, 2019). The detection of more diverse spectra of

mycotoxins (AF, BEA, EnnB, FUM, MON and ZEN) in the traditionally processed

complementary foods (*ogi* and *tombran*) compared to the industrially processed infant cereal

agrees with an earlier report from our group on mycotoxins in complementary foods from

Nigeria (Ojuri et al. 2018, 2019). AFs were detected in about 20% of the food samples, but

notably not in the infant cereal. The mean of these samples with detectable AFs in traditionally

processed complementary foods exceeded the 4 µg/kg maximum level for total AF in Nigerian

baby food. The presence of AF in locally processed complementary foods but not in commercial

infant cereal is in keeping with the report from Cameroon (Chuisseu Njamen et al. 2018).

However, the levels found in this present study were lower than those previously reported in

locally processed baby foods from Nigeria (mean: 104 µg/kg, Ojuri et al. 2018), Cameroon

(mean: 177 ug/kg, Chuisseu Njamen et al. 2018) but similar to Ghana (mean: 8.3 µg/kg,

Blankson et al. 2019). Unlike most of the aforementioned reports, herein we analyzed plate-

ready meal for a more precise food contamination data because it considers food ingested after

final preparatory steps such as dilution and cooking (Chuisseu Njamen et al. 2018; Ezekiel et al.

2019). Clearly, the methods employed at household levels, often characterized by poor food

processing practices, influenced the mycotoxin contents in the traditionally processed

complementary foods. The fact that we could not detect AF in the infant cereal could be

attributed to tailoring of production by industrial processors to meet regulatory standards for AF.

FUM was detected in all the complementary food types. However, the mean

concentrations we found were lower than the European Union ML of 200 µg/kg for baby food

(EC 2006) and the mean level (2808 µg/kg) reported in Tanzania (Kamala et al. 2016). FUM is

not regulated in baby foods in Nigeria. Nevertheless, the observed co-occurrence of FUM with

AF in about 20% of the complementary foods is of concern, because these two mycotoxins are

suggested to potentially interact to negatively influence linear growth in children (IARC 2015;

Shirima et al. 2015). Meanwhile, a potential role of additional toxins in mixtures is unknown.

Several mycotoxins were detected in the children's urine similar to previous reports from

Cameroon (Ediage et al. 2013), Nigeria (Ezekiel et al. 2014), Spain (Rodriguez-Carrasco et al.

2014) and Belgium (Heyndrickx et al. 2015) where more than two mycotoxins were detected.

We compared the incidences of commonly found mycotoxins in children urine with data from

previous studies. Overall, ZEN (incidence: 74%), a potent estrogenic mycotoxin able to cross

the human placenta (Warth et al. 2019), was the most frequently detected mycotoxin in the

urine samples. Two ZEN metabolites, α-ZEL and β-ZEL, were detected in the urine samples

albeit at lower frequencies (11% and 20%, respectively) compared to the parent compound.

This is in contrast to the reports of Heyndrickx et al. (2015) wherein ZEN and its metabolites

were not detected in children urine from Belgium. The incidence of ZEN in the present study

was, however, higher than the incidence (4%, 9/220) previously reported in Cameroonian

children employing a method of lower sensitivity (Ediage et al. 2013). Similarly, DHC was found

in more than one half (57%) of the 65 children in the present study. This finding is much higher

than the 6% (7/124) and 14% (20/142) incidences reported for DHC in urine of Belgian children

(Heyndrickx et al. 2015) and those of adults and children in Haiti (Gerding et al. 2015),

respectively. We did not detect the parent toxin CIT in any urine sample, which agrees with the

reports from Haiti but negates the Belgian study that found both parent compound (CIT) and its

metabolite (DHC). The higher incidence recorded in the present study further suggests that CIT

exposure is widespread in Nigerian children. DON (46%) and OTA (34%) occurrences in urine

in the present study were lower than those previously reported in adults from France (99%,

Turner et al. 2010), The United Kingdom (95–100%, Hepworth et al. 2012; Papageorgiou et al.

2018) and Portugal (93%, Silva et al. 2019), but more similar to that observed in Egypt and Iran

(Piekkola et al. 2012; Turner et al. 2012b). Infant data on DON in urine are mostly lacking.

Similarly, urinary $FB_1$ incidence of 26%, which we recorded, was lower than the 96% incidence

reported in a study comprising 147 Tanzanian children's urine (Shirima et al. 2013) but was

higher than the report involving Nepalese children (incidence: 2% (1/50), Mitchell et al. 2016).

To date, $AFM_1$ has been the most reported urinary AF metabolite and biomarker in

children (Polychronaki et al. 2008; Ediage et al. 2013; Ezekiel et al. 2014; Gerding et al. 2015;

Schwartzbord et al. 2016; Ayelign et al. 2017; Chen et al. 2017; Sanchez and Diaz 2019).

However, in the present study, we found $AFQ_1$ as the major aflatoxin metabolite (both in

frequency and concentration), occurring in 44/65 (68%) of the children's urine samples, whereas

$AFM_1$ was detected in 6/65 (9%) of urine samples, four of which had both $AFM_1$ and $AFQ_1$.

Previous reports on $AFQ_1$ occurrence in human urine were on adolescents and adults 15–64

years of age in China and The Gambia (Groopman et al. 1992a, 1992b; Wang et al. 2001;

Mykkanen et al. 2005). Consequently, to the best of our knowledge, $AFQ_1$ is reported for the

first time in the urine of children in any setting. Urinary $AFM_1$ and AF-N7-guanine are

established biomarkers for aflatoxin exposure, however these were established in adults

(Groopman et al. 1992a). Given the role of CYP3Aa4 in $AFQ_1$ formation (Wild and Turner 2002)

and the known significant ontogeny in xenobiotic metabolizing enzymes (Blake et al. 2005),

involved in $AFB_1$ biotransformation, it would be useful to better understand the relationship

between dose and the various metabolites of aflatoxins in urine of the very young. Nevertheless,

the recorded higher incidence of $AFQ_1$ in the urine samples compared with $AFM_1$ informs us that

aflatoxin exposure in young children may be underreported.

Comparing the two cohorts of infants, higher mean levels of seven out of the 10 detected

mycotoxins were found in the NEB urine compared to the EB urine, with statistically significant

levels recorded for four of these mycotoxins including $AFQ_1$, DON, DHC and $FB_1$. Notably,

increases in the mean mycotoxin levels in urine moving from EB regime to the NEB regime

reached 721 and 334% for $AFM_1$ and $FB_1$, respectively. Additionally, the NEB cohort generally

recorded higher diversities of mycotoxin mixtures in urine compared to the EB cohort. Here, we

show that transition from EB to NEB results in a large variation of multiple mycotoxin exposures

in infants. This primarily owes to the higher intake of complementary foods, which are usually

made from mycotoxin-prone cereals, nuts and oil seeds (Kimanya et al. 2009; 2010, 2014; Juan

et al. 2014; Okeke et al. 2015; Kamala et al. 2016; Chuisseu Njamen et al. 2018; Ojuri et al.

2018, 2019). These observations are strongly suggestive that exclusive breastfeeding provides

a relatively, but not completely protected period from mycotoxin exposure.

The spectra and combinations of mycotoxins detected in the three specimens examined

in this study further suggest frequent exposure in Nigerian children, especially in the NEB

category. Obviously, urine contained higher number of mycotoxins and their mixtures compared

to complementary foods and breast milk. Further, some mycotoxins occurred only in the

biological fluids reiterating the importance of preferring biomonitoring approaches over food

contamination data for exposure assessment in humans (Turner et al. 2012; Warth et al. 2013;

Vidal et al. 2018; Ali and Degen 2019). A typical example is DHC and OTA that were detected

only in the biological fluids (breast milk for the EB group and urine from both cohorts). Although

CIT/DHC was not detected in the examined complementary foods in the present study it was

found in 74% of the urines of EB children. CIT levels in grains (especially maize) and

complementary food in Nigeria have recently been shown to be high reaching 16,000 µg/kg

(Okeke et al. 2015, 2018; Ojuri et al. 2018, 2019; Ezekiel et al. 2020). Fermented cocoa beans

in Nigeria, which could also be processed and used in the preparation of foods for infants and

young children, have also recently been reported to contain CIT (Akinfala et al. 2020).

An additional important finding was the detection of BEA and EnnB in complementary

food and breast milk but not in urine because the urinary analytical method did not include both

mycotoxins. EnnB was recently detected in adult human urine from Italy (Rodriguez-Carrasco et

al. 2018); however, there is limited data on urinary BEA and EnnB in children. Considering the

widespread detection of BEA and EnnB in foods in Nigeria, albeit at low concentrations

(Adetunji et al. 2014; Abdus-Salaam et al. 2015; Ezekiel et al. 2016; Oyedele et al. 2017; Ojuri

et al. 2018), and the documented *in vitro* data suggesting their roles in decreasing immune

response to infections (Ficheux et al. 2013) as well as their interactions to cause additive

myelotoxicity (Ficheux et al. 2012), we recommend the inclusion of both mycotoxins in future

urinary biomonitoring studies, especially for children.

The overall health implication for children in the study region and generally where

mycotoxin exposure may be high are enormous suggesting the need for urgent interventions

targeted at subsistence farmers and household food processing. In addition to the documented

effects of the widely studied aflatoxins on child growth, *in vitro* studies using human intestinal

cell lines revealed that DON selectively modulated intestinal transporters (Maresca et al. 2002)

while $FB_1$ decreased intestinal cell viability and proliferation (Minervini et al. 2014).

Consequently, these mycotoxins may play interactive roles to affect nutrient absorption (Liew

and Mohd-Redzwan 2018) which may cumulatively contribute to growth faltering. Thus, future

studies may consider exploring the type and extent of interactive effects the various recorded

mixtures may have on this highly vulnerable population and their growth into adulthood.

Conclusively, this study provides a comprehensive account on and novel insights into

multiple mycotoxin exposures in EB and NEB infants in a high-risk country, Nigeria, based on

examination of three specimens (breast milk, complementary food and urine). First, we show

the presence of several uncommon mycotoxins in biological fluids (DHC and STER in breast

milk; and $AFQ_1$, being the dominant urinary aflatoxin metabolite). The higher incidence of $AFQ_1$

in infant urine compared with $AFM_1$ suggests that aflatoxin exposure in young children may be

underreported. Second, the presence of up to six distinct mycotoxin classes are reported for the

first time in breast milk and infant urine. Third, urines of NEB infants contained higher levels of

mycotoxins and several mixtures of these toxins more frequently than in urines of EB infants.

Thus, exposure to mycotoxins for infants less than 18 months of age is common in the region,

with a significant transition in higher frequency and levels of exposure with the introduction of

complementary foods. Consequently, it is recommended that mothers adhere strictly to

measures with potential to reduce mycotoxin contamination in household diets; including

maintaining compliance to WHO recommendations for exclusive breastfeeding for 6 months.

Once foods are being introduced, adequate drying and proper storage of grains in airtight

containers at household levels, sorting out of moldy, damaged and discolored grains prior to

meal preparation, and diet diversification to include less mycotoxin prone food crops such as

tubers are advised. In addition, prioritizing partnerships focused on driving interventions to

mitigate exposures in this vulnerable population and investigating mycotoxin exposure and

toxicity holistically in the context of the exposome paradigm is urgent.

**Conflict of interest**

Authors do not have any conflict to declare.

**Acknowledgements**

Authors are thankful to Chidinma Akwarandu and Linda Ogu for their assistance during sample collection, and Xiaomin Han, during urine sample preparation. Families who donated samples are deeply appreciated. The authors want to gratefully acknowledge the Mass Spectrometry Centre (MSC) of the Faculty of Chemistry at the University of Vienna and Sciex for providing mass spectrometric instrumentation to BW. This work was performed with the financial support of the University of Vienna, the City of Vienna Jubilee Funds (BOKU Research Funding, project MycoMarker) and the Austrian Science Fund (FWF): P 33188-B.

**References**

Abdus-Salaam RS, Fanelli F, Atanda O, Sulyok M, Cozzi G, Bavaro S, et al. 2015. Fungal and bacterial metabolites associated with natural contamination of locally processed rice (*Oryza sativa* L.) in Nigeria. Food Addit Contam Part A 32(6):950–959, PMID: 25767914, https://doi.org/10.1080/19440049.2015.1027880.

Adejumo O, Atanda O, Raiola A, Somorin Y, Bandyopadhyay R, Ritieni A. 2013. Correlation between aflatoxin $M_1$ content of breast milk, dietary exposure to aflatoxin $B_1$ and socioeconomic status of lactating mothers in Ogun state, Nigeria. Food Chem Toxicol 56:171–177, PMID: 23462105, https://doi.org/10.1016/j.fct.2013.02.027.

Adetunji M, Atanda O, Ezekiel C, Sulyok M, Warth B, Beltran E, et al. 2014. Fungal and bacterial metabolites of stored maize (*Zea mays*, L.) from five agro-ecological zones of Nigeria. Mycotoxin Res 30:89–102, PMID: 24643458, https://doi.org/10.1007/s12550-014-0194-2.

Aichinger G, Kruger F, Puntscher H, Preindl K, Warth B, Marko D. 2019. Naturally occurring mixtures of *Alternaria* toxins: anti-estrogenic and genotoxic effects *in vitro*. Arch Toxicol 93 (10):3021–3031, PMID: 31559443 https://doi.org/10.1007/s00204-019-02545-z.

Akinfala TO, Houbraken J, Sulyok M, Adedeji AR, Odebode AC, Krska R, et al. 2020. Moulds and their secondary metabolites associated with the fermentation and storage of two

cocoa bean hybrids in Nigeria. Int J Food Microbiol 316:108490, PMID: 31874327,
https:// doi.org/10.1016/j.ijfoodmicro.2019.108490

Ali N, Degen GH. 2019. Citrinin biomarkers: a review of recent data and application to human
exposure assessment. Arch Toxicol 93(11):3057–3066, PMID: 31501918,
https://doi.org/10.1007/s00204-019-02570-y.

Alvito PC, Sizoo EA, Almeida CMM, Egmond HPV. 2010. Occurrence of aflatoxins and
ochratoxin A in baby foods in Portugal. Food Anal Methods 3, 22–30,
https://doi.org/10.1007/s12161-008-9064-x.

Ayelign A, Woldegiorgis AZ, Adish A, De Boevre M, Heyndrickx E, De Saeger S.
2017. Assessment of aflatoxin exposure among young children in Ethiopia using urinary
biomarkers. Food Addit Contam Part A 34(9):1606–1616, PMID: 28670952,
https://doi.org/10.1080/19440049.2017.1350290.

Barker, D. 2000. Fetal programming: influences on development and disease in later life. In:
Series NM. ed Dekker M. New York, NY, USA.

Bateson P, Barker D, Clutton-Brock T, Deb D, D'Udine B, Foley R.A., et al. 2004.
Developmental plasticity and human health. Nature 430:419–421, PMID: 15269759,
https://doi.org/10.1038/nature02725.

Blake MJ. Castro L. Leeder JS Kerans GL. 2005. Ontogeny of drug metabolizing enzymes in
the neonate. Semin Fetal Neonatal Med 10(2):123–138, PMID: 15701578,
https://doi.org/10.1016/j.siny.2004.11.001.

Blankson GK, Mills-Robertson FC, Ofosu IW. 2019. Survey of occurrence levels of aflatoxins in
selected locally processed cereal-based foods for human consumption from Ghana.
Food Control 95:170–175, https://doi.org/10.1016/j.foodcont.2018.08.005.

Braun D, Ezekiel CN, Abia WA, Wisgrill L, Degen GH, Turner PC, et al. 2018. Monitoring early-
life mycotoxin exposures via LC-MS/MS breast milk analysis. Anal Chem 90:14569–
14577, PMID: 30449087, https://doi.org/10.1021/acs.analchem.8b04576.

Braun D, Ezekiel CN, Marko D, Warth B 2020. Exposure to mycotoxin-mixtures via breast milk:
an ultra-sensitive LC-MS/MS biomonitoring approach. Front Chem. 8:423,
https://doi.org/10.3389/fchem.2020.00423.

Butts CA, Hedderley DI, Herath TD, Paturi G, Glyn-Jones S, Wiens F, et al. 2018. Human milk
composition and dietary intakes of breastfeeding women of different ethnicity from the
Manawatu Wanganui region of New Zealand. Nutrients 10:1231, PMID: 30181524,
https://doi.org/10.3390/nu10091231.

Cappozzo J, Jackson L, Lee HJ, Zhou, W, Al-Taher F, Zweigenbaum J, et al. 2017. Occurrence
of ochratoxin A in infant foods in the United States. J Food Prot 80(2):251–256, PMID:
28218865, https://doi.org/10.4315/0362-028X.JFP-16-339.

Chen G, Gong YY, Kimanya ME, Shirima CP, Routledge MN. 2017. Comparison of urinary
aflatoxin $M_1$ and aflatoxin albumin adducts as biomarkers for assessing aflatoxin
exposure in Tanzanian children. Biomarkers 23(2):131–136, PMID: 28114823,
https://doi.org/10.1080/1354750X.2017.1285960.

Cherkani-Hassani A, Mojemmi B, Mouane N. 2016. Occurrence and levels of mycotoxins and
their metabolites in human breast milk associated to dietary habits and other factors: A
systematic literature review, 1984–2015. Trends Food Sci Technol 50:56–69,
https://doi.org/10.1016/j.tifs.2016.01.024.

Chuisseu Djamen DP, Abia WA, Zibi SB, Kuo NS, Ngantchouko Ngalemo CB, Tambo E, et al.
2018. Safety of breast milk vis-à-vis common infant formula and complementary foods
from mycotoxin perspective. RAdvFoodSci 1(1):23–31.

Chu FS, Li GY. 1994. Simultaneous occurrence of fumonisin $B_1$ and other mycotoxins in moldy
corn collected from the People's Republic of China in regions with high incidences of
esophageal cancer. Appl Environ Microbiol 60:847–852, PMID: 8161178.

EC (European Commission). 2002. Commission Decision 2002/657 of 12 August 2002
implementing Council Directive 96/23/EC concerning the performance of analytical
methods and the interpretation of results. Off J Euro Union L221:8–36.

EC (European Commission). 2006. Commission regulation (EC) No 1881/2006 of 19 December
2006 setting maximum levels for certain contaminants in foodstuffs. 364. Off J Euro
Union, 5–24. Available at https://eur-lex.europa.eu/. Accessed 01/ 02/2020.

EC (European Commission). 2014. Commission regulation (EU) No 212/2014 of 6 March 2014
amending Regulation (EC) No 1881/2006 as regards maximum levels of the
contaminant citrinin in food supplements based on rice fermented with red yeast

*Monascus purpureus*. Available at http://faolex.fao.org/docs/pdf/eur131703.pdf. Accessed on 01/02/2020.

Ediage EN, Diana Di Mavungu J, Song S, Sioen I, De Saeger S. 2013. Multimycotoxin analysis in urines to assess infant exposure: a case study in Cameroon. Environ Int 57–58:50–59, PMID: 23669720, https://doi.org/10.1016/j.envint.2013.04.002.

EFSA (European Food Safety Authority). 2012. Scientific opinion on the risks for public and animal health related to the presence of citrinin in food and feed. EFSA J 10:2605. https://doi.org/10.2903/j.efsa.2012.2605.

EFSA (European Food Safety Authority). 2013. Scientific opinion on the risk for public and animal health related to the presence of sterigmatocystin in food and feed. EFSA J 11(6):3254. https://doi.org/10.2903/j.efsa.2013.3254.

Ezekiel CN, Ayeni KI, Misihairabgwi JM, Somorin YM, Chibuzor-Onyema IE, Oyedele OA, et al. 2018a. Traditionally processed beverages in Africa: A review of the mycotoxin occurrence patterns and exposure assessment. Compr Rev Food Sci Food Saf 17(2):334–351, https://doi.org/10.1111/1541-4337.12329.

Ezekiel CN, Oyedele OA, Kraak B, Ayeni KI, Sulyok M, Houbraken J, et al. 2020. Fungal diversity and mycotoxins in low moisture content ready-to-eat foods in Nigeria. Front Microbiol, 11:615, https://doi.org/10.3389/fmicb.2020.00615.

Ezekiel CN, Oyeyemi OT, Oyedele OA, Ayeni KI, Oyeyemi IT, Nabofa W, et al. 2018b. Urinary aflatoxin exposure monitoring in rural and semi-urban populations in Ogun state, Nigeria. Food Addit Contam Part A. 35(8):1565–1572, PMID: 29843566 https://doi.org/10.1080/19440049.2018.1475752.

Ezekiel CN, Sulyok M, Ogara IM, Abia WA, Warth B, Šarkanj B, et al. 2019. Mycotoxins in uncooked and plate-ready household food from rural northern Nigeria. Food Chem Toxicol 128:171–179, PMID: 30965105, https://doi.org/10.1016/j.fct.2019.04.002.

Ezekiel CN, Sulyok M, Somorin Y, Odutayo FI, Nwabekee SU, Balogun AF, et al. 2016. Mould and mycotoxin exposure assessment of melon and bush mango seeds, two common soup thickeners consumed in Nigeria. Int J Food Microbiol 237, 83–91, PMID: 27543818, https://doi.org/10.1016/j.ijfoodmicro.2016.08.019.

Ezekiel CN, Warth B, Ogara IM, Abia WA, Ezekiel VC, Atehnkeng J, et al. 2014. Mycotoxin
exposure in rural residents in northern Nigeria: A pilot study using multi-urinary
biomarkers. Environ Int 66:138–145, PMID: 24583186,
https://doi.org/10.1016/j.envint.2014.02.003.

Ferrufino-Guardia E, Chavez-Rico V, Larondelle Y. 2019. Ochratoxin A in human breast milk,
maternal and placental blood from Cochabamba-Bolivia. Rev Toxicol 36:116–125.

Ficheux AS, Sibiril Y, Parent-Massin D. 2012. Co-exposure of *Fusarium* mycotoxins: *in vitro*
myelotoxicity assessment on human hematopoietic progenitors. Toxicon 60(6):1171–
1179, PMID: 22921581, https://doi.org/10.1016/j.toxicon.2012.08.001.

Ficheux AS, Sibiril Y, Parent-Massin D. 2013. Effects of beauvericin, enniatin B and
moniliformin on human dendritic cells and macrophages: An *invitro* study. Toxicon 71:1–
10, PMID: 23685117, https://doi.org/10.1016/j.toxicon.2013.04.024.

Gerding J, Ali N, Schwartzbord J, Cramer B, Brown DL, Degen GH, Humpf H-U. 2015. A
comparative study of the human urinary mycotoxin excretion patterns in Bangladesh,
Germany, and Haiti using a rapid sensitive LC-MS/MS approach. Mycotoxin Res
31(3):127–136, PMID: 25957672, https://doi.org/10.1007/s12550-015-0223-9.

Gong YY, Cardwell K, Hounsa A, Egal S, Turner PC, Hall AJ, et al. 2002. Dietary aflatoxin
exposure and impaired growth in young children from Benin and Togo: cross sectional
study. Br Med J 325:20–21, PMID: 12098724, https://doi.org/10.1136/bmj.325.7354.20.

Gong YY, Egal S, Hounsa A, Turner PC, Hall AJ, Cardwell K, et al. 2003. Determinants of
aflatoxin exposure in young children from Benin and Togo, West Africa: the critical role
of weaning. Int J Epidemiol 32:556–562, PMID: 12913029,
https://doi.org/10.1093/ije/dyg109.

Gong Y, Hounsa A, Egal S, Turner PC, Sutcliffe AE, Hall AJ, et al. 2004. Postweaning exposure
to aflatoxin results in impaired child growth: a longitudinal study in Benin, West Africa.
Environ Health Perspect 112:1334–1338, PMID: 15345349,
https://doi.org/10.1289/ehp.6954.

Gong YY, Wilson S, Mwatha JK, Routledge MN, Castelino JM, Zhao B, et al. 2012. Aflatoxin
exposure may contribute to chronic hepatomegaly in Kenyan school children. Environ
Health Perspect 120:893–896, PMID: 22370114, https://doi.org/10.1289/ehp.1104357.

Groopman JD, Hall AJ, Whittle H, Hudson GJ, Wogan GN, Montesano R, et al. 1992a.
658            Molecular dosimetry of aflatoxin-N7-guanine in human urine obtained in the Gambia,
659            west Africa. Cancer Epidemiol Biomarkers Prev 1:221–227, PMID: 1339082.

660    Groopman JD, Jiaqi Z, Donahue PR, Pikul A, Lisheng Z, Jun-shi C, et al. 1992b. Molecular
661            dosimetry of urinary aflatoxin-DNA adducts in People living in Guangxi autonomous
662            region, People's Republic of China. Cancer Res 52:45–52. PMID: 1727385.

663    Hepworth SJ, Hardie LJ, Fraser LK, Burley VJ, Mijal RS, Wild CP, et al. 2012. Deoxynivalenol
664            exposure assessment in a cohort of pregnant women from Bradford, UK. Food Addit
665            Contam Part A: Chem Anal Control Expo Risk Assess 29(2):269–276. PMID: 21462017,
666            https://doi.org/1080/19440049.2010.551301.

667    Heussner AH and Bingle LEH 2015. Comparative ochratoxin toxicity: a review of the available
668            data. Toxins 7(10):4253-4282, PMID: 26506387, https://doi.org/10.3390/toxins7104253.

669    Heyndrickx E, Sioen I, Huybrechts B, Callebaut A, De Henauw S, De Saeger S. 2015. Human
670            biomonitoring of multiple mycotoxins in the Belgian population: results of the BIOMYCO
671            study. Environ Int 84:82–89, PMID: 26233555,
672            https://doi.org/10.1016/j.envint.2015.06.011.

673    IARC (International Agency for Research on Cancer). 1987. Some naturally occurring
674            substances. In: IARC monographs on the evaluation of carcinogenic risks to humans.
675            Summaries and evaluations. 10:72.

676    IARC (International Agency for Research on Cancer). 2002. Traditional herbal medicines, some
677            mycotoxins, napthalene, and styrene. IARC monographs on the evaluation carcinogenic
678            risk to humans, 82: 1–556, PMID: 12687954

679    IARC (International Agency for Research on Cancer) 2012a Monographs on the evaluation of
680            carcinogenic risks to humans: chemical agents and related occupations. A review of
681            human carcinogens. Lyon, France: 100F:224–248

682    IARC (International Agency for Research on Cancer). 2012b. IARC Scientific publication No.
683            158. In: Pitt, J.I., Wild, C.P., Baan, R.A., Gelderblom, W.C.A., Miller, J.D., Riley, R.T.,
684            Wu, F. (Eds.), Improving public health through mycotoxin control. ISBN-13:978-92-832-
685            2158-6.

IARC (International Agency for Research on Cancer). 2015. In: Wild, Christopher P., David
Miller, J., Groopman, John D. (Eds.), Mycotoxin control in low- and middle-income
countries (IARC Working Group Reports; 9), (NLM Classification: W1). ISBN-13: 978-
9283225102.

Iha MH, Barbosa CB, Heck AR, Trucksess MW. 2014. Aflatoxin $M_1$ and ochratoxin A in human
milk  in  Ribeirão  Preto-SP,  Brazil.  Food  Control  40:310–313,
https://doi.org/10.1016/j.foodcont.2013.12.014.

Innis SM. 2014. Impact of maternal diet on human milk composition and neurological
development of infants. Am J Clin Nutr 99(3):734S-741S, PMID: 24500153,
https://doi.org/10.3945/ajcn.113.072595.

Jonsyn FE, Maxwell SM, Hendrickse RG. 1995. Ochratoxin A and aflatoxins in breast milk
samples  from  Sierra  Leone.  Mycopathologia,  131:121–126,  PMID:  8532055,
https://doi.org/10.1007/BF01102890.

Juan C, Raiola A, Mãnes J, Ritieni A, 2014. Presence of mycotoxin in commercial infant
formulas  and  baby  foods  from  Italian  market.  Food  Control  39:227–236,
https://doi.org/10.1016/j.foodcont.2013.10.036.

Kamala A, Kimanya M, Lachat C, Jacxsens L, Haesaert G, Kolsteren P, et al. 2016. Risk of
exposure to multiple mycotoxins from maize-based complementary foods in Tanzania. J
Agr  Food  Chem  65(33):7106–7114,  PMID:  28830150,
https://doi.org/10.1021/acs.jafc.6b03429.

Kimanya ME, Meulenaer BD, Baert K, Tiisekwa B, Camp JV, Samapundo S, et al. 2009.
Exposure of infants to fumonisins in maize-based complementary foods in rural
Tanzania.  Mol  Nutr  Food  Res  53:667–674,  PMID:  18837467,
https://doi.org/10.1002/mnfr.200700488.

Kimanya ME, Meulenaer BD, Baert K, Tiisekwa B, Camp JV, Samapundo S, et al. 2010.
Fumonisin exposure through maize in complementary foods is inversely associated with
linear growth of infants in Tanzania. Mol Nutr Food Res 54:1659–1667, PMID:
20521269, https://doi.org/10.1002/mnfr.200900483.

Kimanya ME, Shirima CP, Magoha H, Shewiyo DH, De Meulenaer B, Kolsteren P, et al. 2014.
Co-exposures of aflatoxins with deoxynivalenol and fumonisins from maize-based

complementary foods in Rombo, northern Tanzania. Food Control, 41:76–81,
https://doi.org/10.1016/j.foodcont.2013.12.034.

Kolakowski B, O'rourke SM, Bietlot HP, Kurz K, Aweryn B. 2016. Ochratoxin A concentrations in
a variety of grain-based and non–grain-based foods on the Canadian retail market from
2009 to 2014. J Food Prot 79(12):2143–2159, PMID: 28221957,
https://doi.org/10.4315/0362-028X.JFP-16-051.

Liew WPP, Mohd-Redzwan S. 2018. Mycotoxin: Its Impact on gut health and microbiota. Front
Cell Infect Microbiol 8:60, PMID: 29535978, https://doi.org/10.3389/fcimb.2018.00060.

Mandy M, Nyirenda M. 2018. Developmental origins of health and disease: The relevance to
developing nations. Int Health 10:66-70, PMID: 29528398,
https://.doi.org/10.1093/inthealth/ihy006

Maresca M, Mahfoud R, Garmy N, Fantini J. 2002. The mycotoxin deoxynivalenol affects
nutrient absorption in human intestinal epithelial cells. J Nutr 132(9):2723–2731, PMID:
12221236, https://doi.org/10.1093/jn/132.9.2723.

McMillian A, Renaud JB, Burgess KMN, Orimadegun AE, Akinyinka OO, Allen SJ, et al. 2018.
Aflatoxin exposure in Nigerian children with severe acute malnutrition. Food Chem
Toxicol 111:356–362. PMID: 29175577, https://doi.org/10.1016/j.fct.2017.11.030.

Minervini F, Garbetta A, D'Antuono I, Cardinali A, Martino NA, Debellis L, et al. 2014. Toxic
mechanisms induced by fumonisin $B_1$ mycotoxin on human intestinal cell line. Arch
Environ Con Tox 67(1):115–123, PMID: 24549592, https://doi.org/10.1007/s00244-014-
0004-z.

Missmer SA, Suarez L, Felkner M, Wang E, Merrill Jr AH, Rothman KJ, et al. 2006. Exposure to
fumonisins and the occurrence of neural tube defects along the Texas– Mexico border.
Environ Health Perspect 114:237–241, PMID: 16451860,
https://doi.org/10.1289/ehp.8221.

Mitchell NJ, Riley RT, Egner PA, Groopman JD, Wu F. 2016. Chronic aflatoxin exposure in
children living in Bhaktapur, Nepal: extension of the MAL-ED study. J Exp Sci Env Epid
27(1):106–111, PMID: 26732375, https://doi.org/10.1038/jes.2015.87.

Muñoz K, Blaszkewicz M, Campos V, Vega M, Degen GH. 2014. Exposure of infants to
ochratoxin A with breast milk. Arch Toxicol 88(3):837–846, PMID: 24270973,
https://doi.org/10.1007/s00204-013-1168-4.

Mykkänen H, Zhu H, Salminen E, Juvonen RO, Ling W, Ma J, et al. 2005. Fecal and urinary
excretion of aflatoxin $B_1$ metabolites ($AFQ_1$, $AFM_1$ and AFB-N7-guanine) in young
Chinese males. Int J Cancer 115(6):879–884, PMID: 15723309,
https://doi.org/10.1002/ijc.20951.

Ojuri OT, Ezekiel CN, Eskola MK, Šarkanj B, Babalola AD, Sulyok M, et al. 2019. Mycotoxin co-
exposures in infants and young children consuming household and industrially-
processed complementary foods in Nigeria and risk management advice. Food Control
98: 312–322, https://doi.org/10.1016/j.foodcont.2018.11.049.

Ojuri OT, Ezekiel CN, Sulyok M, Ezeokoli OT, Oyedele OA, Ayeni KI, et al. 2018. Assessing the
mycotoxicological risk from consumption of complementary foods by infants and young
children in Nigeria. Food Chem Toxicol 121:37–50, PMID: 30118820,
https://doi.org/10.1016/j.fct.2018.08.025.

Okeke CA, Ezekiel CN, Nwangburuka CC, Sulyok M, Ezeamagu CO, Adeleke RA, et al. 2015.
Bacterial diversity and mycotoxin reduction during maize fermentation (steeping) for *ogi*
production. Front Microbiol 6:1402, PMID: 26697001,
https://doi.org/10.3389/fmicb.2015.01402.

Okeke CA, Ezekiel CN, Sulyok M, Ogunremi OR, Ezeamagu CO, Šarkanj B, et al.
2018. Traditional processing impacts mycotoxin levels and nutritional value of ogi – A
maize-based complementary food. Food Control 86:224–233,
https://doi.org/10.1016/j.foodcont.2017.11.021.

Oplatowska-Stachowiak M, Reiring C, Sajic N, Haasnoot W, Brabet C, Campbell K, et al. 2018.
Development and in-house validation of a rapid and simple to use ELISA for the
detection and measurement of the mycotoxin sterigmatocystin. Anal Bioanal Chem
410(12):3017–3023, PMID: 29523941, https://doi.org/10.1007/s00216-018-0988-8.

Ortiz J, Jacxsens L, Astudillo G, Ballesteros A, Donoso S, Huybregts L, et al. 2018. Multiple
mycotoxin exposure of infants and young children via breastfeeding and

complementary/weaning foods consumption in Ecuadorian highlands. Food Chem
Toxicol 118: 541–548, PMID: 29883782, https://doi.org/10.1016/j.fct.2018.06.008.

Oueslati S, Berrada H, Mañes J, Juan C. 2018. Presence of mycotoxins in Tunisian infant foods
samples and subsequent risk assessment. Food Control 84:362–369,
https://doi.org/10.1016/j.foodcont.2017.08.021.

Oyedele OA, Ezekiel CN, Sulyok M, Adetunji MC, Warth B, Atanda, OO, et al. 2017. Mycotoxin
risk assessment for consumers of groundnut in domestic markets in Nigeria. Int J Food
Microbiol, 251:24–32, PMID: 28380344,
https://doi.org/10.1016/j.ijfoodmicro.2017.03.020.

Papageorgiou M, Wells L, Williams C, White K, De Santis B, Liu Y, et al. 2018. Assessment of
urinary deoxynivalenol biomarkers in UK children and adolescents. Toxins 10(2):50,
PMID: 29360781, https://doi.org/10.3390/toxins10020050.

Pestka J.J. 2010. Deoxynivalenol: Mechanisms of action, human exposure, and toxicological
relevance. Arch. Toxicol 84:663–679. https://doi.org/10.1007/s00204-010-0579-8.

Piekkola S, Turner PC, Abdel-Hamid M, Ezzat S, El-Daly M, El-Kafrawy S, et al. 2012.
Characterisation of aflatoxin and deoxynivalenol exposure among pregnant Egyptian
women. Food Addit Contam Part A: Chem Anal Control Expo Risk Assess 29(6):962–
971, PMID: 22376138, https://doi.org/10.1080/19440049.2012.658442.

Polychronaki N, West RM, Turner PC, Amra H, Abdel-Wahhab M, Mykkänen H, El-Nezami H.
2007. A longitudinal assessment of aflatoxin $M_1$ excretion in breast milk of selected
Egyptian mothers. Food Chem Toxicol 45 (7):1210–1215, PMID: 17306915
https://doi.org/10.1016/j.fct.2007.01.001.

Polychronaki N, Wild CP, Mykkänen H, Amra H, Abdel-Wahhab M, Sylla A, et al. 2008. Urinary
biomarkers of aflatoxin exposure in young children from Egypt and Guinea. Food Chem
Toxicol 46(2):519–526, PMID: 17920747, https://doi.org/10.1016/j.fct.2007.08.034.

Preindl K, Braun D, Aichinger G, Sieri S, Fang M, Marko D, et al. 2019. A generic LC-MS/MS
exposome method for the determination of xenoestrogens in biological matrices. Anal
Chem, 91(17):11334–11342, PMID: 31398002,
https://doi.org/10.1021/acs.analchem.9b02446.

802 Rodriguez-Carrasco Y, Izzo L, Gaspari A, Graziani G, Mañes J, Ritieni A. 2018. Urinary levels
803    of enniatin B and its phase I metabolites: first human pilot biomonitoring study. Food
804    Chem Toxicol 118:454–459, PMID: 29802944, https://doi.org/10.1016/j.fct.2018.05.052.

805 Rodriguez-Carrasco Y, Molto JC, Manes J, Berrada H. 2014. Exposure assessment approach
806    through mycotoxin/creatinine ration evaluation in urine by GC-MS/MS. Food Chem
807    Toxicol 72:69–75, PMID: 25042512, https://doi.org/10.1016/j.fct.2014.07.014.

808 Rubert J, León N, Sáez C, Martins CPB, Godula M, Yusà V, et al. 2014. Evaluation of
809    mycotoxins and their metabolites in human breast milk using liquid chromatography
810    coupled to high resolution mass spectrometry. Anal Chim Acta 820:39–46,
811    PMID: 24745736, https://doi.org/10.1016/j.aca.2014.02.009.

812 Sanchez EM, Diaz GJ. 2019. Frequency and levels of aflatoxin $M_1$ in urine of children in Bogota,
813    Colombia. Mycotoxin Res 35(3):271–278, PMID: 30915618,
814    https://doi.org/10.1007/s12550-019-00355-x.

815 Sarkanj B, Ezekiel CN, Turner PC, Abia WA, Rychlik M, Krska R, et al. (2018). Ultra-sensitive,
816    stable isotope assisted quantification of multiple urinary mycotoxin exposure biomarkers.
817    Anal Chim Acta 1019:84–92, PMID: 29625687,
818    https://doi.org/10.1016/j.aca.2018.02.036.

819 Schwartzbord JR, Leroy JL, Severe L, Brown DL. 2016. Urinary aflatoxin $M_1$ in Port-au-Prince
820    and a rural community in north-east Haiti. Food Addit Contam Part A 33(6):1036–1042,
821    PMID: 27146636, https://.doi.org/10.1080/19440049.2016.1185899.

822 Shirima CP, Kimanya ME, Kinabo JL, Routledge MN, Srey C, Wild CP, et al. 2013. Dietary
823    exposure to aflatoxin and fumonisin among Tanzanian children as determined using
824    biomarkers of exposure. Mol Nutr Food Res 57:1874–1881, PMID: 23776058,
825    https://doi.org/10.1002/mnfr.201300116.

826 Shirima CP, Kimanya ME, Routledge MN, Srey C, Kinabo JL, Humpf H-U, et al. 2015. A
827    prospective study of growth and biomarkers of exposure to aflatoxin and fumonisin
828    during early childhood in Tanzania. Environ Health Perspect 123(2):173–178,
829    PMID: 25325363, https://doi.org/10.1289/ehp.1408097.

Silva LJG, Macedo L, Pereira AMPT, Duarte S, Lino CM, Pena A. 2019. Ochratoxin A and
Portuguese children: urine biomonitoring, intake estimation and risk assessment. Food
Chem Toxicol 135:110883, PMID: 31610259, https://doi.org/10.1016/j.fct.2019.110883.

Sulyok M, Stadler D, Steiner D, Krska R. 2020. Validation of an LC-MS/MS-based dilute-and-
shoot approach for the quantification of >500 mycotoxins and other secondary
metabolites in food crops: challenges and solutions. Anal Bioanal Chem 412(11): 2607–
2620, PMID: 32078002, https://doi.org/10.1007/s00216-020-02489-9.

Sun G, Wang S, Hu X, Su J, Huang T, Yu J, et al. 2007. Fumonisin $B_1$ contamination of home-
grown corn in high-risk areas for esophageal and liver cancer in China. Food Addit
Contam 24:181–185, PMID: 17364919, https://doi.org/10.1080/02652030601013471.

Sun G, Wang S, Hu X, Su J, Zhang Y, Xie Y, et al. 2011. Co-contamination of aflatoxin $B_1$ and
fumonisin $B_1$ in food and human dietary exposure in three areas of China. Food Addit
Contam      Part      A      28:461–470,      PMID: 21259142,
https://doi.org/10.1080/19440049.2010.544678.

Turconi G, Guarcello M, Livieri C, Comizzoli S, Maccarini L, Castellazzi AM, et al.
2004. Evaluation of xenobiotics in human milk and ingestion by the newborn. Eur J Nutr
43(4):191–197, PMID: 15309437, https://doi.org/10.1007/s00394-004-0458-2.

Turner PC, Collinson AC, Cheung YB, Gong YY, Hall AJ, Prentice AM, et al. 2007. Aflatoxin
exposure in utero causes growth faltering in Gambian infants. Int J Epidemiol 36:1119–
1125, PMID: 17576701, https://doi.org/10.1093/ije/dym122.

Turner PC, Flannery B, Isitt C, Ali M, Pestka J. 2012a. The role of biomarkers in evaluating
human health concerns from fungal contaminants in food. Nutr Res Rev 25:162–179,
PMID: 22651937, http://doi.org/10.1017/S095442241200008X.

Turner PC, Gong YY, Pourshams A, Jafaric E, Routledge MN, Malekzadeh R, et al. 2012b. A
pilot survey for *Fusarium* mycotoxin biomarkers in women from Golestan, northern Iran.
World Mycotoxin J 5:195–199, https://doi.org/10.3920/WMJ2011.1337.

Turner PC, Hopton R, Lecluse Y, White KLM, Fisher J, Lebailly P (2010) Determinants of
urinary deoxynivalenol in male farmers from Normandy, France J. Agri. Food Chem. 58:
5206–5212, PMID: 20349912, https://doi.org/10.1021/jf100892v.

Turner PC, Moore SE, Hall AJ, Prentice AM, Wild CP. 2003. Modification of immune function
through exposure to dietary aflatoxin in Gambian children. Environ Health Perspect
111:217–220, PMID: 12573908, https://doi.org/10.1289/ehp.5753.

Turner PC. 2013. The molecular epidemiology of chronic aflatoxin driven impaired child growth.
Scientifica 2013, 152879, PMID: 24455429, http://doi.org/10.1155/2013/152879.

Vejdovszky K, Hahn K, Braun D, Warth B, Marko D. 2017a. Synergistic estrogenic effects of
*Fusarium* and *Alternaria* mycotoxins *in vitro*. Arch Toxicol 91(3):1447–1460, PMID:
27401186, https://doi.org/10.1007/s00204-016-1795-7.

Vejdovszky K, Schmidt V, Warth B, Marko D. 2017b. Combinatory estrogenic effects between
the isoflavone genistein and the mycotoxins zearalenone and alternariol *in vitro*. Mol
Nutr     Food     Res     61(3):1600526,     PMID:     27739238,
https://doi.org/10.1002/mnfr.201600526.

Vidal A, Mengelers M, Yang S, De Saeger S, De Boevre M. 2018. Mycotoxin biomarkers of
exposure: a comprehensive review. Compr Rev Food Sci Food Saf 17(5):1127–1155,
https://doi.org/10.1111/1541-4337.12367.

Wang J-S, Huang T, Su J, Liang F, Wei Z, Liang Y, et al. 2001. Hepatocellular carcinoma and
aflatoxin exposure in Zhuqing village Fusui county, People's Republic of China. Cancer
Epidemiol Biomarkers Prev 10:143–146, PMID: 11219772.

Warth B, Braun D, Ezekiel CN, Turner PC, Degen GH, Marko D. 2016. Biomonitoring of
mycotoxins in human breast milk: current state and future perspectives. Chem Res
Toxicol                 29(7):1087–1097,                 PMID: 27300310,
https://doi.org/10.1021/acs.chemrestox.6b00125.

Warth B, Preindl K, Manser P, Wick P, Marko D, Buerki-Thurnherr T. 2019. Transfer and
metabolism of the xenoestrogen zearalenone in human perfused placenta. Environ
Health Perspect 127(10):107004, PMID: 31596610, https://doi.org/10.1289/EHP4860.

Warth B, Sulyok M, Krska R. 2013. LC-MS/MS-based multibiomarker approaches for the
assessment of human exposure to mycotoxins. Anal Bioanal Chem 405(17):5687–5695,
PMID: 23774829, https://doi.org/10.1007/s00216-013-7011-1.

Watson S, Chen G, Sylla A, Routledge MN, Gong YY. 2016. Dietary exposure to aflatoxin and
micronutrient status among young children from Guinea. Mol Nutr Food Res 60:511–
518, PMID: 26603511, https://doi.org/10.1002/mnfr.201500382.

Watson S, Moore S, Darboe MK, Chen G, Tu Y-K, Huang Y-T, et al. 2018. Impaired growth in
rural Gambian infants exposed to aflatoxin: a prospective cohort study. BMC Public
Health 18:1247, PMID: 30413157, https://doi.org/10.1186/s12889-018-6164-4.

Wild CP, Turner PC. 2002. The toxicology of aflatoxins as a basis for public health decisions.
Mutagenesis 17:471–481, PMID: 12435844, https://doi.org/10.1093/mutage/17.6.471.

Woelflingseder L, Warth B, Vierheilig I, Schwartz-Zimmermann H, Hametner C, Nagl V, et al.
2019. The *Fusarium* metabolite culmorin suppresses the *in vitro* glucuronidation of
deoxynivalenol. Arch Toxicol 93(6):1729–1743, PMID: 31049613,
https://doi.org/10.1007/s00204-019-02459-w.

Wu X, Jackson RT, Khan SA, Ahuja J, Pehrsson PR. 2018. Human milk nutrient composition in
the United States: current knowledge, challenges, and research needs. Curr Develop
Nutr 2(7):nzy025, PMID: 30087951, https://doi.org/10.1093/cdn/nzy025.

Yoshizawa T, Yamashita A, Luo Y. 1994. Fumonisin occurrence in corn from high- and low-risk
areas for human esophageal cancer in China. Appl Environ Microbiol 60:1626–1629,
PMID: 8017941.

**Table 1.** Demographic characteristics and food preference of study participants.

| Characteristics | All | Exclusively breastfed | Non-exclusively breastfed |
|---|---|---|---|
| Subjects ($n$ (%)) | 65 | 23 (35.4) | 42 (64.6) |
| Gender ($n$ (%)) | | | |
|   Male | 31 (47.7) | 11 (47.8) | 20 (47.6) |
|   Female | 34 (52.3) | 12 (52.2) | 22 (52.4) |
| Age (months) | | | |
|   Mean ± SD[a] | 7.1 ± 3.9 | 3.7 ± 1.6 | 9.0 ± 3.6* |
|   Range | 1–18 | 1–6 | 3–18 |
| Body weight (kg) | | | |
|   Mean ± SD[a] | 7.4±1.7 | 6.3±1.0 | 8.0±1.6** |
|   Range | 4.5–11.2 | 4.5–8.0 | 5.0–11.2 |
| | | | |
| Frequency of food consumption/day | | | |
|   Breast milk | 1–>10 | 8–>10[b] | 1–3 |
|   Complementary food | 3–5 | 0 | 3–5 |
| Complementary food (n (%)) preference | | | |
|   Single food (e.g. *ogi*) | 3 (4.6) | - | 3 (7.1) |
|   Mixed foods[c] | 39 (60.0) | - | 39 (92.9) |

[a]Standard deviation.

[b]As reported by Braun et al. (2018).

[c]Combination of complementary foods (e.g. infant cereal, *ogi* and *Tombran*) consumed in a day.

*Significant at $p<0.01$, ** $p<0.05$

**Table 2.** Mycotoxins in breast milk ($n$ = 22) consumed by exclusively breastfed infants in Ogun

state, Nigeria.

| Mycotoxins | LOD (ng/L) | LOQ (ng/L) | $N$ (%)[a] | Concentration (ng/L) | | |
|---|---|---|---|---|---|---|
| | | | | Range | Median | Mean |
| Aflatoxin $M_1$ | 2.0 | 4.0 | 4 (18.2) | 2.0[b] | 2.0 | 2.0 |
| Alternariol monomethyl ether | 0.5 | 1.0 | 21 (95.5) | 0.5[b]–11.7 | 1.8 | 3.0 |
| Beauvericin | 0.1 | 0.3 | 22 (100) | 1.0–12.0 | 2.2 | 3.0 |
| Dihydrocitrinone | 14.0 | 28.0 | 6 (27.3) | 14.0[b]–59.7 | 14.0 | 25.2 |
| Enniatin B | 0.7 | 1.4 | 16 (72.7) | 0.7[b]–10.1 | 4.6 | 5.0 |
| Enniatin $B_1$ | 0.5 | 1.0 | 5 (22.7) | 0.5[b]–1.15 | 0.5 | 0.6 |
| Ochratoxin A | 2.0 | 4.0 | 14 (63.6) | 2.0[b]–67.6 | 2.0 | 9.6 |
| Ochratoxin B | 2.5 | 5.0 | 2 (9.1) | 5.3–6.7 | 6.0 | 6.0 |
| Sterigmatocystin | 0.5 | 1.0 | 1 (4.6) | 1.2 | 1.2 | 1.2 |

912    [a]Number (and percentage) of samples containing the mycotoxin.
913    [b]<LOQ values were considered as positive data and substituted with LOQ/2.

914 **Table 3.** Mycotoxins in plate-ready complementary food fed to non-exclusively breastfed infants in Ogun state, Nigeria.

| Mycotoxins | LOD (µg/kg) | LOQ (µg/kg) | All samples ($n$=42) | | | Infant cereal ($n$=6) | | Ogi ($n$=26) | | Tombran ($n$=10) | |
| | | | $N$ (%)[a] | Range[b] | Mean[b]±SD | $N$ | Mean[b] | $N$ | Mean[b] | $N$ | Mean[b] |
|---|---|---|---|---|---|---|---|---|---|---|---|
| Aflatoxin $B_1$ | 0.24 | 0.72 | 9 (21.4) | 1.0–7.0 | 3.0 ± 2.0 | 0 | <LOD | 6 | 2.5 | 3 | 4.1 |
| Aflatoxin $B_2$ | 0.4 | 1.2 | 1 (2.4) | 1.2 | 1.2 | 0 | <LOD | 0 | <LOD | 1 | 1.2 |
| Aflatoxin $G_1$ | 0.32 | 0.96 | 2 (4.8) | 12.0–12.1 | 12.1 ± 0.0 | 0 | <LOD | 2 | 12.1 | 0 | <LOD |
| *Total aflatoxin[c]* | - | - | *9 (21.4)* | *1.0–16.2* | *5.8 ± 6.3* | *0* | *<LOD* | *6* | *6.5* | *3* | *4.5* |
| Beauvericin | 0.008 | 0.024 | 33 (78.6) | 0.1–116 | 3.8 ± 20.1 | 6 | 19.4 | 19 | 0.3 | 8 | 0.3 |
| Deoxynivalenol | 1.2 | 3.6 | 2 (4.8) | 52.9–61.5 | 57.2 ± 6.1 | 1 | 52.9 | 1 | 61.5 | 0 | <LOD |
| Enniatin $A_1$ | 0.032 | 0.096 | 7 (16.7) | 0.1–5.7 | 1.3 ± 2.0 | 2 | 3.3 | 1 | 0.9 | 4 | 0.4 |
| Enniatin B | 0.024 | 0.072 | 8 (19.0) | 0.03–6.9 | 2.1 ± 3.0 | 2 | 6.9 | 1 | 1.1 | 5 | 0.4 |
| Enniatin $B_1$ | 0.04 | 0.12 | 6 (14.3) | 0.4–14.3 | 4.1 ± 5.3 | 2 | 9.6 | 1 | 2.2 | 3 | 1.1 |
| Enniatin $B_2$ | 0.04 | 0.12 | 3 (7.1) | 0.09–0.34 | 0.22 ± 0.13 | 2 | 0.29 | 0 | <LOD | 1 | 0.09 |
| Fumonisin $A_1$ | 2 | 6.0 | 1 (2.4) | 2.0 | 2.0 | 0 | <LOD | 1 | 2.0 | 0 | <LOD |
| Fumonisin $B_1$ | 3.2 | 9.6 | 11 (26.2) | 13.8–114 | 35.6 ± 32.2 | 2 | 54.2 | 7 | 35.3 | 2 | 18.0 |
| Fumonisin $B_2$ | 2.4 | 7.2 | 8 (19.0) | 7.9–80.3 | 24.3 ± 24.0 | 1 | 17.8 | 6 | 26.7 | 1 | 15.8 |
| *Total fumonisin[d]* | - | - | *12 (28.6)* | *7.9–194* | *48.8 ± 52.9* | *2* | *63.1* | *8* | *50.9* | *2* | *25.9* |
| Moniliformin | 1.6 | 4.8 | 3 (7.1) | 7.2–10.6 | 8.6 ± 1.8 | 0 | <LOD | 0 | <LOD | 3 | 8.6 |
| Zearalenone | 0.12 | 0.36 | 4 (9.5) | 0.5-6.8 | 2.4 ± 3.0 | 1 | 0.6 | 2 | 1.1 | 1 | 6.8 |

915 [a]Number (and percentage) of samples containing the mycotoxin. [b]Mean (and standard deviation) of mycotoxin concentrations expressed in µg/kg.

916 [c]Sum of aflatoxins $B_1$, $B_2$ and $G_1$.

917 [d]Sum of fumonisins $B_1$ and $B_2$.

918 **Table 4.** Mycotoxins in urines of exclusively breastfed (EB) and non-exclusively breastfed (NEB) infants in Ogun state, Nigeria.

| Mycotoxins | LOD (ng/L) | LOQ (ng/L) | Quantified levels (ng/L) | | | | | |
| | | | Exclusively breastfed (n=23) | | | Non-exclusively breastfed (n=42) | | |
| | | | N (%)[a] | Range | Mean[b] | N (%)[a] | Range | Mean[b] |
|---|---|---|---|---|---|---|---|---|
| Aflatoxin $M_1$ | 0.3 | 1.0 | 1 (4) | 23 | 23 | 5 (12) | 32–504 | 166 |
| Aflatoxin $Q_1$ | 0.3 | 1.0 | 20 (87) | 3–791 | 182 | 24 (57) | 14–898 | 225* |
| Deoxynivalenol | 50 | 150 | 7 (30) | 225–19,781 | 3194 | 23 (55) | 226–21,344 | 5282* |
| Dihydrocitrinone | 3.0 | 10 | 6 (26) | 5–944 | 135 | 31 (74) | 5–1,377 | 202* |
| Fumonisin $B_1$ | 1.0 | 10 | 4 (17) | 35–188 | 136 | 13 (31) | 47–1,587 | 454* |
| Nivalenol | 50 | 100 | nd | nd | nd | 1 (2) | 161 | 161 |
| Ochratoxin A | 0.3 | 1.0 | 6 (26) | 0.5–76 | 19* | 16 (38) | 0.5–34 | 9 |
| Zearalenone | 1.0 | 3.0 | 13 (57) | 17–784 | 148 | 35 (83) | 13–678 | 140 |
| α-Zearalenol | 3.0 | 10 | nd | nd | nd | 7 (17) | 5–55 | 20 |
| β-Zearalenol | 1.0 | 3.0 | 7 (30) | 8–648 | 122 | 6 (14) | 14–68 | 32 |

919 [a]Number (percentage) of children with quantifiable mycotoxin levels in urine.

920 [b]Mean values were calculated to include <LOQ data points substituted with half LOQ.

921 *Significant at $p<0.05$.

923 **Table 5.** Co-occurrence of mycotoxin classes[a] in breast milk, complementary food and urine.

| Numbers of mycotoxins | Incidence [$N$ (%)][b] in infant's food | | Incidence [$N$ (%)][b] in infant's urine | | |
| --- | --- | --- | --- | --- | --- |
| | Breast milk ($n$=22) | Complementary food ($n$=42) | Exclusively breastfed ($n$=23) | Non-exclusively breastfed ($n$=42) | All ($n$=65) |
| 0 | 0 (0) | 0 (0) | 1 (4.3) | 0 (0) | 1 (1.5) |
| 1 | 0 (0) | 20 (47.6) | 5 (21.7) | 3 (7) | 8 (12.3) |
| 2 | 3 (13.6) | 13 (31.0) | 5 (21.7) | 5 (11.9) | 10 (15.4) |
| 3 | 5 (22.8) | 4 (9.5) | 3 (13.0) | 12 (28.6) | 15 (23.1) |
| 4 | 9 (40.9) | 4 (9.5) | 7 (30.4) | 12 (28.6) | 19 (29.2) |
| 5 | 2 (9.1) | 1 (2.4) | 2 (8.7) | 9 (21.4) | 11 (16.9) |
| 6 | 3 (13.6) | 0 (0.0) | 0 (0.0) | 1 (2.4) | 1 (1.5) |

[a]Mycotoxin classes: aflatoxins (breast milk: $AFM_1$; food: $AFB_1$, $AFB_2$ and $AFG_1$; urine: $AFM_1$ and $AFQ_1$); alternariolmethylether; beauvericin; dihydrocitrinone;
deoxynivalenol; dihydrocitrinone (breast milk and urine); enniatins (breastmilk: ENNB and $ENNB_1$; food: $ENNA_1$, ENNB, $ENNB_1$ and $ENNB_2$); fumonisins (food: $FB_1$
and $FB_2$; urine: $FB_1$); moniliformin; nivalenol; ochratoxins (breast milk: OTA and OTB; urine: OTA); sterigmatocystin (breast milk); zearalenone (food: zearalenone;
urine: zearalenone, α-zearalenol and β-zearalenol).
[b]Number (and percentage) of samples containing co-occurring mycotoxin classes. <LOQ values were considered as positive data.

**Table S1.** Fungal metabolites in plate-ready complementary food fed to non-exclusively breastfed infants in Ogun state, Nigeria.

| Metabolites | LOD (µg/kg) | All samples (n=42) | | | Infant cereal (n=6) | | Ogi (n=26) | | Tombran (n=10) | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | N (%)[a] | Range[b] | Mean[b]±SD | N | Mean[b] | N | Mean[b] | N | Mean[b] |
| 3-Nitropropionic acid | 0.8 | 2 (4.76) | 11.7-20.6 | 16.2±6.30 | 0 | <LOD | 2 | 16.2 | 0 | <LOD |
| Agroclavine | 0.12 | 4 (9.52) | 12.6-532 | 243±216 | 0 | <LOD | 1 | 12.6 | 3 | 320 |
| Asperglaucide | 0.08 | 31 (73.8) | 0.08-90.3 | 9.87±22.1 | 6 | 2.15 | 16 | 0.90 | 9 | 31.0 |
| Asperphenamate | 0.04 | 24 (57.1) | 0.06-32.2 | 4.11±8.90 | 2 | 3.45 | 14 | 0.42 | 8 | 10.7 |
| Averantin | 0.04 | 2 (4.76) | 0.38-0.88 | 0.63±0.35 | 0 | <LOD | 2 | 0.63 | 0 | <LOD |
| Averufin | 0.04 | 7 (16.7) | 0.04-2.54 | 0.88±1.10 | 1 | 0.04 | 4 | 1.34 | 2 | 0.38 |
| Brevianamid F | 0.16 | 12 (28.6) | 0.35-24.1 | 9.06±8.10 | 2 | 9.39 | 5 | 3.18 | 5 | 14.8 |
| cyclo(L-Pro-L-Tyr) | 0.8 | 26 (61.9) | 0.81-25.9 | 7.00±8.13 | 7 | 5.07 | 11 | 2.57 | 8 | 14.8 |
| cyclo(L-Pro-L-Val) | 0.64 | 39 (92.9) | 0.65-71.0 | 16.8±18.7 | 7 | 22.6 | 22 | 8.35 | 10 | 31.2 |
| Elymoclavine | 0.4 | 7 (16.7) | 0.99-45.9 | 16.6±16.9 | 0 | <LOD | 2 | 1.10 | 5 | 22.8 |
| Emodin | 0.056 | 2 (4.76) | 0.98-1.01 | 1.00±0.02 | 0 | <LOD | 2 | 1.00 | 0 | <LOD |
| Fellutanine A | 0.2 | 4 (9.52) | 6.00-10.0 | 8.22±1.99 | 0 | <LOD | 2 | 7.89 | 2 | 8.55 |
| Festuclavine | 0.08 | 3 (7.14) | 2.65-4.74 | 3.53±1.08 | 0 | <LOD | 0 | <LOD | 3 | 3.53 |
| Flavoglaucin | 0.24 | 7 (16.7) | 0.54-57.5 | 17.0±24.4 | 1 | 0.67 | 2 | 52.3 | 4 | 3.43 |
| Kojic acid | 16 | 9 (21.4) | 17.0-956 | 142±306 | 0 | <LOD | 4 | 260 | 5 | 46.7 |
| Macrosporin | 0.04 | 3 (7.14) | 0.07-1.10 | 0.42±0.59 | 0 | <LOD | 0 | <LOD | 3 | 0.42 |
| N-Benzoyl-Phenylalanine | 0.064 | 22 (52.4) | 0.33-11.9 | 2.03±3.13 | 2 | 0.92 | 12 | 0.90 | 8 | 4.00 |
| Neoechinulin A | 0.8 | 3 (7.14) | 4.33-17.4 | 11.7±6.71 | 0 | <LOD | 2 | 15.5 | 1 | 4.33 |
| O-MethylSterigmatocystin | 0.12 | 2 (4.76) | 0.35-0.41 | 0.38±0.04 | 0 | <LOD | 2 | 0.38 | 0 | <LOD |
| Questiomycin A | 2 | 1 (2.38) | 5.58 | 5.58 | 0 | <LOD | 1 | 5.58 | 0 | <LOD |
| Quinolactacin A | 0.08 | 5 (11.9) | 0.10-0.66 | 0.35±0.25 | 1 | 0.23 | 3 | 0.48 | 1 | 0.10 |
| Rugulosovin | 0.64 | 9 (21.4) | 0.71-8.33 | 3.72±2.79 | 1 | 3.85 | 5 | 4.48 | 3 | 2.41 |
| Skyrin | 0.4 | 1 (2.38) | 0.91 | 0.91 | 0 | <LOD | 1 | 0.91 | 0 | <LOD |
| Tryptophol | 8 | 27 (64.3) | 8.88-792 | 87.2±180 | 4 | 49.0 | 14 | 88.7 | 9 | 102 |
| Versicolorin C | 0.24 | 2 (4.76) | 0.58-0.62 | 0.60±0.02 | 0 | <LOD | 2 | 0.60 | 0 | <LOD |

930 [a]Number (percentage) of samples containing the metabolite.
931 [a]Concentrations expressed in µg/kg.

**Figure 1.** MRM-chromatograms of a matrix-matched standard in urine (A), and an infant urine sample (B) of aflatoxin $Q_1$ (AFQ$_1$) and dihydrocitrinone (DHC), respectively. In addition, MRM-chromatograms of a matrix-matched standard in breast milk (C) and a breast milk sample (D) of DHC and sterigmatocystin (STER), respectively, are shown. Quantifier, qualifier and retention time are given for each mycotoxin: AFQ$_1$ (*m/z* 329.0–282.9; *m/z* 329.0–175.0; RT: 15.1 min) and DHC (*m/z* 265.0–221.1; *m/z* 265.0–246.9; RT: 16.1 min) in urine; DHC (*m/z* 265.0–177.0; *m/z* 265.0–203.0; RT: 4.5 min) and STER (*m/z* 325.1–281.1; *m/z* 325.1–310.2; RT: 8.1 min) in breast milk.

942 **Figure 2.** Scatter plot for percentage increases in detection frequencies and means of detects

943 of mycotoxins in urine of non-exclusively breastfed (NEB) vs. exclusively breastfed (EB) infants

944 in Ogun state, Nigeria. Numerals on x-axis denote mycotoxins: 1 = aflatoxin $M_1$; 2 = aflatoxin $Q_1$;

945 3 = deoxynivalenol; 4 = dihydrocitrinone; 5 = fumonisin $B_1$; 6 = ochratoxin A; 7 = zearalenone; 8

946 = β-zearalenol.

**Figure 3.** MRM-chromatogram of one non-exclusively breastfed infant urine sample from Ogun state, Nigeria, containing a mixture of seven mycotoxins belonging to six mycotoxin classes. Quantifier, qualifier, internal standard (IS) and retention time are given for each mycotoxin: DON (*m/z* 355.1–246.9; *m/z* 355.1–137.9; *m/z* 370.1–278.8; RT: 9.0 min), AFQ$_1$ (*m/z* 329.0–282.9; *m/z* 329.0–175.0; no IS; RT: 15.0 min), AFM$_1$ (*m/z* 329.1–273.2; *m/z* 329.1–229.1; *m/z* 346.0–288.2; RT: 15.1 min), FB$_1$ (*m/z* 722.5–334.4; *m/z* 722.5–352.3; *m/z* 756.3–356.3; RT: 15.1 min), DHC (*m/z* 265.0–221.1; *m/z* 265.0–246.9; no IS; RT: 15.9 min), OTA (*m/z* 404.0–239.0; *m/z* 404.0–358.0; *m/z* 424.2–250.1; RT: 20.6 min), and ZEN (*m/z* 317.1–175.0; *m/z* 317.1–131.0; *m/z* 335.2–185.1; RT: 20.8 min).