

1 **Prognostic value of visual quantification of lesion severity at initial chest CT in**
2 **confirmed Covid-19 infection: a retrospective analysis on 216 patients**

3
4 Elias TAIEB^{1*} MD, Aïssam LABANI¹ MD, Yvon RUCH² MD, François DANION² MD, Pascal BILBAULT³ MD
5 PhD, Mathieu OBERLIN³ MD, Pierre LEYENDECKER¹ MD, Catherine ROY¹ MD, Mickaël OHANA¹ MD
6 PhD

- 7 1. *Hôpitaux Universitaires de Strasbourg, Service de Radiologie, Nouvel Hôpital Civil, Strasbourg,*
8 *France*
9 2. *Hôpitaux Universitaires de Strasbourg, Service de Maladies infectieuses et tropicales et*
10 *médecine interne, Nouvel Hôpital Civil, Strasbourg, France*
11 3. *Hôpitaux Universitaires de Strasbourg, Service d'Accueil des Urgences, Nouvel Hôpital Civil,*
12 *Strasbourg, France*

13
14
15
16
17
18 ****Corresponding author***

19 *Élias TAIEB*

20 *Nouvel Hôpital Civil - Service de Radiologie*

21 *1 place de l'Hôpital, 67000 Strasbourg, FRANCE*

22 elias.taieb@chru-strasbourg.fr

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39 **ABSTRACT**

40

41

42 **Rationale and Objectives**

43 Studies suggest an association between chest CT findings assessed with semi-quantitative CT
44 score and gravity of Covid-19. The objective of this work is to analyze potential correlation
45 between visual quantification of lesion severity at initial chest CT and clinical outcome in
46 confirmed Covid-19 patients.

47

48

49 **Materials and Methods**

50 From March 5th to March 21st, 2020, all consecutive patients that underwent chest CT for
51 clinical suspicion of Covid-19 at a single tertiary center were retrospectively evaluated for
52 inclusion. Patients with lung parenchyma lesions compatible with Covid-19 and positive RT-
53 PCR were included.

54 Global extensiveness of abnormal lung parenchyma was visually estimated and classified
55 independently by 2 readers, following the European Society of Thoracic Imaging Guidelines.

56 Death and/or mechanical ventilation within 30 days following the initial chest CT was chosen
57 as the primary hard endpoint.

58

59

60 **Results**

61 216 patients (124 men, 62 years-old \pm 16.5, range 22 – 94 yo) corresponding to 216 chest CT
62 were included.

63 Correlation between lesion severity and percentage of patients that met the primary
64 endpoint was high, with a coefficient ρ of 0.87 ($p = 0.05$).

65 A greater than 25% involvement was significantly associated with a higher risk of mechanical
66 ventilation or death at 30 days, with a Risk Ratio of 5.00 (95% CI [3.59-6.99]).

67

68

69 **Conclusion**

70 This retrospective cohort confirms a correlation between visual evaluation of lesions severity
71 at initial chest CT and the 30 days clinical outcome of Covid-19 patients and suggests using a
72 threshold of greater than 25% involvement to identify patients at risk.

73

74

75 **INTRODUCTION**

76

77

78 Covid-19 is due to the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), first
79 described in China in December 2019 and that has spread as a worldwide pandemic in
80 March 2020. Even though most cases are benign, 17 to 29% of hospitalized patients may
81 develop a severe pneumonia with acute respiratory distress syndrome (ARDS) necessitating
82 aggressive resuscitative measures such as mechanical ventilation with prone position (1,2).

83 Reverse-transcription polymerase chain reaction (RT-PCR) assay is the reference standard to
84 confirm Covid-19 (3). Nonetheless, results are delayed and initial false negatives are
85 frequent (4), resulting in the need for alternative triage methods in Emergency Department
86 (ED) patients during high prevalence of the disease. Chest computed tomography (CT) is an
87 established alternative for the early screening and triage of suspected Covid-19 patients with
88 moderate to severe clinical symptoms. Indeed, chest-CT typical findings of Covid-19
89 pneumonia are now well described with ground-glass opacities (GGO), consolidation and
90 crazy-paving pattern that are bilateral, multifocal with peripheral predominant distribution
91 (4)(5).

92

93 Initial studies suggest an association between chest CT findings and gravity of Covid-19. Yuan
94 et al. described on 27 consecutive patients a higher semi-quantitative CT score in the 10 that
95 died compared to the 17 that recovered (6,7). Zhao et al. described on 101 cases a
96 correlation between lesion severity on CT and clinical severity of the disease (8). Yang et al.
97 also found on 102 Covid-19 patients a higher CT score in those with severe clinical disease
98 (9). Correlation between initial chest-CT findings and a precise clinical outcome such as

99 admission in intensive care unit (ICU) or mortality has been assessed in 236 patients by
100 Colombi et al. They showed that software quantification of well-aerated lung parenchyma
101 on initial chest-CT was predictive of ICU admission or death, with a suggested cut-off at 73%
102 (10). This method would however require the use of dedicated quantification software,
103 which is time-consuming and difficult to integrate in routine workflow. On the contrary,
104 visual evaluation of lesion severity as recommended by the European Society of Thoracic
105 Imaging Guidelines (11) is simple and easy to integrate in the everyday workflow. Yet, and to
106 the best of our knowledge, no study has reported the potential predictive value of this
107 approach.

108

109 The objective of this work is to analyze potential correlation between visual quantification of
110 lesion severity at initial chest CT and the clinical outcome in confirmed Covid-19 patients.

111

112

113 **MATERIALS AND METHODS**

114 This retrospective study was approved by the local ethics committee of Strasbourg
115 University Hospital and the need of written informed consent was waived.

116

117

118 **Population**

119 From March 5th to March 21st, 2020, all consecutive patients that underwent chest CT for
120 clinical suspicion of Covid-19 at our ED (Nouvel Hôpital Civil, Strasbourg University Hospital,
121 France) were retrospectively evaluated for inclusion. Patients were addressed for moderate
122 to severe respiratory symptoms and chest CT was systematically used at their ED arrival to

123 expedite triage. Systematic RT-PCR sample obtained by nasopharyngeal swab were done
124 concomitantly to the CT. Some patients could have a second or third sampling using
125 expectorations or bronchoalveolar lavage if the first sample was negative.

126 Inclusion criteria were:

- 127 - availability of chest CT at the time of ED admission;
- 128 - presence of lung parenchyma lesions compatible with Covid-19 on the initial chest
129 CT;
- 130 - at least one positive RT-PCR result.

131 Exclusion criteria were:

- 132 - delay between initial chest CT and the first positive RT-PCR greater than 14 days,
- 133 - lack of follow-up data at 30 days after initial chest CT.

134

135

136 **Clinical Evaluation**

137 Death and/or mechanical ventilation within 30 days following the initial chest CT was chosen
138 as the primary hard endpoint.

139 Age, sex, onset of symptoms at the time of chest CT and hospitalization status were
140 retrieved from the Electronic Health Record.

141

142

143 **CT Acquisition Protocol**

144 All CT examinations were acquired using an 80-row CT scanner (Aquilion PRIME SP, Canon
145 Medical Systems), without injection of contrast media. Acquisition protocol was adapted
146 according to the patient's morphotype, with a tube voltage of 100-135kV and a tube current

147 of 2-350mAs. Patients were instructed to hold their breath and raise their arms above their
148 head to minimize artifacts. Images were reconstructed with a slice-thickness of 1 mm in
149 mediastinal and parenchymal windows and transmitted to post-processing workstations for
150 multiplanar and maximum intensity projection reconstructions.

151

152

153 **CT Image Analysis**

154 Two radiologists (ET with a 3 years' experience in CT and AL with 10 years' experience)
155 blinded to the clinical data, reviewed CT images independently and resolved discrepancies
156 by consensus, after an initial training on a separate set of 20 cases.

157 Global extensiveness of abnormal lung parenchyma – *i.e.* lesions consistent with Covid-19
158 such as GGO, consolidation, crazy paving and organizing pneumonia – was visually estimated
159 and classified following the European Society of Thoracic Imaging Guidelines as <10%, 11-
160 25%, 26-50%, 51-75% or >75% (11). This visual quantification was obtained after visual
161 analysis of 5 zones on axial images: at the level of the supra-aortic trunks, the azygos cross,
162 the carina, the inferior pulmonary veins and the pleural recesses.

163

164

165 **Statistical analysis**

166 Continuous variables were expressed as median and standard deviation (SD). Normality of
167 distribution was assessed using a Shapiro-Wilk test. Variables were compared using Student
168 t-test when normally distributed, or Wilcoxon.

169 Categorical variables were expressed as counts and percentage, with corresponding 95%
170 confidence interval (95% CI) using Wilson method, and compared using Pearson's χ^2 tests or
171 Fisher's exact tests.

172 Inter-reader agreement was assessed with a Cohen's Kappa test.

173 Correlation between the severity group and the percentage of patients that met the primary
174 endpoint was assessed with a Spearman test.

175 Risk ratio for primary endpoint was calculated between patients with less than 25% lesions
176 and those with more than 25% lesions.

177 All the analyses were performed using R software version 3.6.3. (R Core Team (2019). R: A
178 language and environment for statistical computing. R Foundation for Statistical Computing,
179 Vienna, Austria. URL <https://www.R-project.org/>).

180

181

182 **RESULTS**

183

184

185 **Population**

186 598 patients were admitted in the ED for a clinical suspicion of Covid-19 over the inclusion
187 period; 216 patients (124 men, 62 years-old \pm 16.5, range 22 – 94yo) corresponding to 216
188 chest CT were finally retrospectively included in this work (**Fig 1**).

189

190

191 **Clinical Evaluation**

192 76/216 patients (35%, 95% CI: 29.1-41.8) met the primary endpoint of death (N = 21, with 4
193 after an ICU stay) or mechanical ventilation (N = 55) in the 30 days following the initial chest
194 CT.

195 Patients in mechanical ventilation or death group were older than patients in group without
196 need of mechanical ventilation or death (66 years-old vs 59 years-old; $p < 0.001$) (**Table 1**).

197

198

199

200 **CT Image Analysis**

201 The most common subgroup of severity was 11-25% (89/216; 41%) (**Table 2**).

202

203

204 Inter-reader agreement was high, with a discrepancy rate of 3% (6/216) and a weighted
205 Cohen's Kappa of 0.97 [94.8-99.4].

206 Correlation between lesion severity and percentage of patients that met the primary
207 endpoint was high, with a Spearman coefficient ρ of 0.87 ($p = 0.05$).

208 Occurrence of the primary outcome with regards to the lesion severity subgroups at the
209 initial chest CT presented a major gap at a 25% threshold (**Table 3**). A greater than 25%
210 involvement was significantly associated with a higher risk of mechanical ventilation or
211 death at 30 days, with a Risk Ratio of 5.00 (95% CI [3.59-6.99]).

212

213

214 Occurrence of ICU and/or death was 90% (95%CI [79-96]) in patients with more than 25%
215 extensiveness versus 18% (95%CI [13-25]) in patients with less than 25% extensiveness (**Fig**
216 **2**).

217

218 Demonstrative examples are shown in **Figs 3 and 4**.

219

220

221 **DISCUSSION**

222 The results of the present study showed that the rate of death or mechanical ventilation in
223 Covid-19 patients increased significantly with the degree of lesion severity at initial chest CT.
224 Patients with more than 25% of lung parenchyma involved had a risk ratio of 5.00 for this
225 primary endpoint. This study is, to the best of our knowledge, the first to validate the use of
226 the ESR/ESTI CT visual assessment scale for Covid-19 lesions. Compared to semi-quantitative
227 scores or dedicated quantification software, visual assessment has the advantage of
228 simplicity, availability and rapidity.

229

230 Our results are in accordance with those of Colombi et al. in which a significant association
231 between quantification of well-aerated lung parenchyma on admission chest-CT less than
232 73% and ICU admission or death was described (OR 5,4; $p < 0,001$) (10), which is consistent
233 with our proposed threshold of 25%.

234 Covid-19 is a novel disease, so far not fully understood and can present multiple forms and
235 courses with a quite unpredictable outcome. Identifying prognostic factors is of major
236 interest. In our ED workflow, chest CT is used as a triage tool for patients with moderate to
237 severe respiratory clinical manifestations. As is, lesions severity could be used to further
238 guide patient's management right at the ED.

239 One could fear that the limiting factor of visual evaluation would be the poor reproducibility,
240 yet we achieved a high inter-reader agreement in our work. This could be related to the fact
241 that readers did an initial consensus training to limit this variability.

242

243 This study still presents some important limitations. First, it was a retrospective and
244 monocentric study, with a limited number of cases. Second, other clinical data were
245 unavailable and an independent analysis with other possible confounding factors (12) taken
246 into account (such as age, sex, body mass index, comorbidities, smoking status...) could not
247 be done. Therefore, we cannot assume that CT visual scoring is an independent predictor
248 factor of Covid-19 evolution. Third, this study was done on patients from the initial part of
249 the epidemic wave, when pressure on ICU space was the most important and when patient's
250 management was not completely defined and optimized. This could partly explain the high
251 fatality rate in our cohort. Fourth, we decided to include only patients with CT lesions, which
252 could constitute an inclusion bias. Last, we did not perform software-based quantitative
253 measurement to validate our visual evaluation. However, the high inter-reader agreement is
254 in favor of the validity of this visual approach.

255

256 To conclude, this retrospective cohort confirms a correlation between visual evaluation of
257 lesion severity at initial chest CT and the 30 days clinical outcome of Covid-19 patients, thus
258 reinforcing the role of chest CT in ED triage. Future studies will help to refine knowledge
259 about CT prognostic factors in Covid-19.

260

261

262 **REFERENCES**

263

264

- 265 1. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected
266 with 2019 novel coronavirus in Wuhan, China. *The Lancet*. 2020 Feb 15;395(10223):497–
267 506.
- 268 2. Chavez S, Long B, Koyfman A, Liang SY. Coronavirus Disease (COVID-19): A primer for
269 emergency physicians. *Am J Emerg Med* [Internet]. 2020 Mar 24 [cited 2020 Apr 15];
270 Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7102516/>
- 271 3. Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DKW, et al. Detection of
272 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill Bull Eur Sur Mal*
273 *Transm Eur Commun Dis Bull*. 2020;25(3).
- 274 4. Xie X, Zhong Z, Zhao W, Zheng C, Wang F, Liu J. Chest CT for Typical 2019-nCoV
275 Pneumonia: Relationship to Negative RT-PCR Testing. *Radiology*. 2020 Feb 12;200343.
- 276 5. Chung M, Bernheim A, Mei X, Zhang N, Huang M, Zeng X, et al. CT Imaging Features of
277 2019 Novel Coronavirus (2019-nCoV). *Radiology*. 2020 Feb 4;295(1):202–7.
- 278 6. Yuan M, Yin W, Tao Z, Tan W, Hu Y. Association of radiologic findings with mortality of
279 patients infected with 2019 novel coronavirus in Wuhan, China. *PLOS ONE*. 2020 Mar
280 19;15(3):e0230548.
- 281 7. Li K, Fang Y, Li W, Pan C, Qin P, Zhong Y, et al. CT image visual quantitative evaluation
282 and clinical classification of coronavirus disease (COVID-19). *Eur Radiol* [Internet]. 2020
283 Mar 25 [cited 2020 Apr 15]; Available from: [https://doi.org/10.1007/s00330-020-06817-](https://doi.org/10.1007/s00330-020-06817-6)
284 6
- 285 8. Zhao W, Zhong Z, Xie X, Yu Q, Liu J. Relation Between Chest CT Findings and Clinical
286 Conditions of Coronavirus Disease (COVID-19) Pneumonia: A Multicenter Study. *AJR Am J*
287 *Roentgenol*. 2020 Mar 3;1–6.
- 288 9. Yang R, Li X, Liu H, Zhen Y, Zhang X, Xiong Q, et al. Chest CT Severity Score: An Imaging
289 Tool for Assessing Severe COVID-19. *Radiol Cardiothorac Imaging*. 2020 Mar
290 30;2(2):e200047.
- 291 10. Colombi D, Bodini FC, Petrini M, Maffi G, Morelli N, Milanese G, et al. Well-aerated Lung
292 on Admitting Chest CT to Predict Adverse Outcome in COVID-19 Pneumonia. *Radiology*.
293 2020 Apr 17;201433.
- 294 11. Revel M-P, Parkar AP, Prosch H, Silva M, Sverzellati N, Gleeson F, et al. COVID-19 patients
295 and the radiology department – advice from the European Society of Radiology (ESR)

296 and the European Society of Thoracic Imaging (ESTI). Eur Radiol [Internet]. 2020 Apr 20
297 [cited 2020 May 8]; Available from: <https://doi.org/10.1007/s00330-020-06865-y>

298 12. Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, et al.
299 Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients
300 Hospitalized With COVID-19 in the New York City Area. JAMA [Internet]. 2020 Apr 22
301 [cited 2020 May 14]; Available from:
302 <https://jamanetwork.com/journals/jama/fullarticle/2765184>

303

304 Tables

305 **Table 1. Patient demographic and clinical data.**

	All patients (n=216)	Patients with mechanical ventilation or death (n=76)	Patients without mechanical ventilation or death (n=140)	P value
Female	92 (43%)	26 (34%)	66 (47%)	p = 0.0664
Male	124 (57%)	50 (66%)	74 (53%)	
Median age in years (SD)	62 (16.5)	66 (14)	59 (17)	p = 0.00079
Median onset of symptoms in days (SD)	6 (3.25)	5 (3.5)	6 (3.13)	p = 0.461

306

307

308

309

310

311

312

313 **Table 2. Patient demographic and clinical data in severity subgroups.**

	1-10% <i>(n=76)</i>	11-25% <i>(n=89)</i>	26-50% <i>(n =37)</i>	51-75% <i>(n=11)</i>	76-100% <i>(n=3)</i>
Female (n=92)	32 <i>(42%)</i>	42 <i>(47%)</i>	13 <i>(35%)</i>	3 <i>(27%)</i>	2 <i>(67%)</i>
Male (n=124)	44 <i>(58%)</i>	47 <i>(53%)</i>	24 <i>(65%)</i>	8 <i>(73%)</i>	1 <i>(33%)</i>
Median age in years (SD)	58 (17.3)	65 (16.2)	62 (15.4)	62 (14.1)	57 (3.4)
Median onset of symptoms in days (SD)	4 (2.4)	7 (3.1)	7 (3.9)	4 (2.8)	7 (5.9)

314

315 **Table 3. Occurrence of primary endpoint in severity subgroups.**

All patients (n=216)	Patients with mechanical ventilation or death (n=76)	Patients without mechanical ventilation or death (n=140)	Rates of mechanical ventilation or death with 95%CI
1-10% (n=76)	9	67	11% [6-23]
11-25% (n=89)	21	68	24% [16-33]
26-50% (n=37)	32	5	87% [72-94]
51-75% (n=11)	11	0	100% [74-100]
76-100% (n=3)	3	0	100% [44-100]

316

317

318 **Figure legends**

319 **Fig 1.** Flowchart of patients.

320

321

322 **Fig 2.** Graphical abstract.

323

324

325 **Fig 3.** 45-year-old male at day 3 of fever with acute dyspnea.

326 Chest-CT showing limited peripheral ground-glass opacity in the left lower lobe, with a visual

327 quantification of lesions severity of <10%. Patient was discharged home after ED admission.

328

329

330 **Fig 4.** 58-year-old woman at day 4 of dyspnea.

331 Chest-CT showing extensive bilateral ground-glass opacities, with a visual quantification of

332 lesions severity >75%. The patient was intubated and admitted in intensive care unit few

333 hours after CT.

334

335


336

337


338

339

340



ESR/ESTI **visual** scale



In 216 confirmed Covid-19 patients (124 men, 62 years-old \pm 16.5, range 22 – 94)

