

1 **Computerized physical and cognitive training improves the**
2 **functional architecture of the brain in adults with Down**
3 **Syndrome: a longitudinal network science EEG study**

4
5 Alexandra Anagnostopoulou^{1*}, Charis Styliadis^{1*}, Panagiotis Kartsidis¹, Evangelia
6 Romanopoulou¹, Vasiliki Zilidou¹, Chrysi Karali², Maria Karagianni¹, Manousos Klados³,
7 Evangelos Paraskevopoulos^{1**}, Panagiotis D. Bamidis^{1**}

8
9 ¹ Medical Physics Laboratory, School of Medicine, Faculty of Health Sciences, Aristotle
10 University of Thessaloniki, Greece.

11 ² School of Biology, Faculty of Science, Aristotle University of Thessaloniki, Greece.

12 ³ Department of Psychology, The University of Sheffield International Faculty, City College,
13 Thessaloniki, Greece.

14
15 * Joined first authorship

16 ** Joined senior authorship

17
18 **Corresponding Author:** Dr. Charis Styliadis, School of Medicine, Faculty of Health Sciences,
19 Aristotle University of Thessaloniki, Thessaloniki, Greece. Aristotle University of Thessaloniki;
20 University Campus; P.C. 54124; Thessaloniki; Greece. Email: cstyliadis@auth.gr

21
22 **Keywords:** Down Syndrome, Physical Training, Cognitive Training, Electroencephalography,
23 Network Science Indices, Adaptive Neuroplasticity

27 **Abstract**

28 Understanding the neuroplastic capacity of people with Down Syndrome (PwDS) can potentially
29 reveal the causal relationship between aberrant brain organization and phenotypic characteristics.
30 We used resting-state EEG recordings to identify how a neuroplasticity-triggering training
31 protocol relates to changes in the functional connectivity of the brain's intrinsic cortical networks.
32 Brain activity of 12 PwDS before and after a ten-week protocol of combined physical and
33 cognitive training was statistically compared to quantify changes in directed functional in
34 conjunction with psycho-somatometric assessments. PwDS showed increased connectivity within
35 the left hemisphere and from left to right hemisphere, as well as increased physical and cognitive
36 performance. Our results reveal a strong adaptive neuroplastic reorganization, as a result of the
37 training that leads to a more complex and less-random network, with a more pronounced
38 hierarchical organization. Our results go beyond previous findings by indicating a transition to a
39 healthier, more efficient, and flexible network architecture, with improved integration and
40 segregation abilities in the brain of PwDS. Resting-state electrophysiological brain activity is
41 used here for the first time to display meaningful longitudinal relationships to underlying DS
42 processes and outcomes of importance in a translational inquiry. This trial is registered with
43 ClinicalTrials.gov Identifier [NCT04390321](https://clinicaltrials.gov/ct2/show/study/NCT04390321).

44

45 **Author Summary**

46 The effects of cognitive and physical training on the neuroplasticity attributes of people with
47 and without cognitive impairment have been well documented via neurophysiological evaluations
48 and network science indices. However, there is still insufficient evidence for people with Down
49 Syndrome (PwDS). We investigated the effects of a combinational training protocol on the brain
50 network organization of 12 adult PwDS using EEG and network indices coupled with tests
51 assessing their cognitive and physical capacity. We report evidence of adaptational neuroplastic
52 effects, pointing to a transitional state towards a healthier organization with an increased ability to
53 integrate and segregate information. Our findings underline the ability of the DS brain to respond
54 to the cognitive demands of external stimuli, reflecting the possibility of developing independent-
55 living skills.

56 Introduction

57 Neuroplasticity can emerge in both typical and atypical brains and allows for either
58 development (evolutionary plasticity), reaction (reactive plasticity), recovery (reparation
59 plasticity), or adaptation to internal and external stimuli (adaptational plasticity)¹. The different
60 aspects of plasticity are suggested to have the same molecular basis¹. In contrast to the brain of
61 typically developed (TD) individuals, the Down Syndrome (DS) brain presents atypical levels of
62 inhibition, due to gene over-expression, which lead to prolonged failure of synaptic plasticity and
63 a reduced capacity for remodeling². These characteristics, along with morphogenetic
64 modifications, have been identified as some of the leading causes of brain disability in people
65 with DS (PwDS)³.

66 Children and young adults with DS, in comparison to age-matched individuals of typical
67 development, have smaller brain volumes (i.e., decreased volume in frontal gyrus, amygdala, and
68 cerebellar structures but also increased parahippocampal volume)^{4,5}, a pattern which over-
69 pronounces after their 50th year of age⁶. The current consensus has associated the cognitive
70 capabilities and deficiencies of PwDS with several distinct brain regions emphasizing an
71 abnormal, and less efficient DS brain organization⁷.

72 Electroencephalography (EEG) and magnetoencephalography (MEG) studies have
73 complemented the notion of this atypical organization; PwDS when compared to TD controls
74 exhibit slow brain wave, especially in left posterior areas^{8,9}, with higher delta band and lower
75 alpha and beta band activity, a pattern also evident in patients with Alzheimer's Disease (AD)⁸⁻¹¹.
76 fMRI studies have shown that the DS network has a rather-simplified organization¹², lacking the
77 appropriate efficiency and flexibility¹³. Given the limited capacity of the DS brain to consolidate
78 information due to its disorganized architecture of reduced segregation and impaired integration,
79 diffused connectivity (hyper-synchrony), and decreased long-range connectivity^{12,13}, the DS brain
80 network's potential for plasticity is in question¹³.

81 Despite these DS-related atypical deviations, the DS brain does possess neuroplastic
82 capabilities, at least in the form of compensatory events¹⁴⁻¹⁶. The emergence of AD-related
83 characteristics (i.e., altered theta band activity and power) in PwDS, at least a decade before the
84 manifestation of dementia and much earlier than in TD people^{11,17}, reaffirm the possibility of
85 compensatory mechanisms, before and during the expression of AD^{14,16}. Characterizing
86 neuroplasticity in PwDS is vital in understanding causality between aberrant brain circuitry and
87 the cognitive and behavioral phenotype. This understanding would allow the quantification of the
88 remodeling potential of evidence-based interventions that aim to purposely reawake neural
89 plasticity and permit improved cognition.

90 Longitudinal and interventional studies are required to investigate whether it is possible to
91 overcome, at least partially, the cognitive disability through novel therapies. Such interventional
92 approaches have shifted from a pharmaceutical concept, with, so far, inconclusive results
93 regarding beneficial effects¹⁸⁻²², to non-pharmaceutical interventions of physical and behavioral
94 components, aiming to trigger neuroplasticity and enhance brain health or to protect against
95 neurodegenerative events¹⁴. Such interventions have shown promise in healthy aging, and
96 populations with cognitive impairments, and have provided mounting evidence for lifelong brain
97 plasticity²³⁻³¹.

98 Despite evidence supporting the positive impact of exercise-based^{32,33}, as well as behavioral
99 and cognitive^{34,35} interventions for PwDS, DS literature features a substantial knowledge gap, as
100 there are no reports of neurophysiological investigations of the training-induced effects, and a
101 network theory based assessment of the training-induced plasticity. Deviations in the brain
102 functionality of PwDS have been previously addressed with the investigation of resting-state
103 networks (RSNs)^{8,9,12,36-38}.

104 We hypothesized that our training protocol of combined physical (PT) and cognitive training
105 (CT), will trigger neuroplasticity and reorganize the DS brain network to adapt to the increased
106 cognitive and physical demands. Resting-state (eyes-open) EEG data were acquired to
107 characterize plasticity in the DS brain through network science and to investigate how the
108 training-induced neuroplasticity affects the state of the DS network, in terms of organization and
109 characteristics, since it is known to maintain a random-like architecture³⁸ with impaired
110 integration and segregation capabilities. Could the triggering of neuroplasticity bring about the
111 emergence of a more efficient, complex, and specialized network organization?

112 Source analysis was performed using low resolution electromagnetic tomography
113 (LORETA)³⁹ and directed functional connectivity was estimated via phase transfer entropy (PTE)
114 in the 0.53-35 Hz frequency band. This measure quantifies the direction of the information flow,
115 allowing for a whole-head analysis without requiring an *a priori* head model definition. An FDR
116 statistical model was applied to extract the significant, within-network differences (post- vs. pre-
117 intervention) in connectivity. A graph-theoretical analysis was performed to index the training's
118 neuroplastic effects through the statistical comparison of graph measures between the two time-
119 points.

120

121 Results

122 Psychometric and somatometric results

123 Psychosomatometric score comparisons between the two time-points were performed with the
124 use of non-parametric Wilcoxon tests and paired t-tests. Physical assessments exhibited a
125 significant improvement in score for the arm curl test (Table 1). Significant differences in the
126 time of completion (decrease in duration) were evident in the Time Up and Go assessment test
127 (Table 1). The comparison of psychometric assessments revealed an increase in the Digits
128 Forward score (Digits Span test), the Mazes test, as well as the Ravens AB, and Ravens Total
129 Score (Table 1). The rest of the tests showed no significant post-pre score changes.

130

131 **Table 1.** Somatometric and psychometric tests with significant differences post vs. pre, along
132 with the post-pre mean difference. Results were considered significant for $p < 0.05$.

Somatometric	p-value	mean difference
Arm Curl	0.017	1.75
Time Up and Go	0.007	-1.67
Psychometric		
Digits Span (Digit Forward)	0.032	1.50
Mazes	0.032	0.97
Ravens AB	0.017	2.12
Ravens Total Score	0.015	3.13

133

134 PTE results

135 Statistical comparisons of the PTE matrices between the pre- and post-intervention EEG
136 measurements indicate the reorganization of cortical connections (19 nodes and 19 edges), due to
137 training, in the DS resting-state network ($p < 0.05$, FDR corrected, 10000 permutations) (Figure
138 1). Specifically, the cortical reorganization in the DS brain is characterized by the strengthening
139 of connections within: i) the left parietal lobe (paracentral, postcentral, precuneus, superior and
140 inferior parietal), with most connections originating from superior parietal nodes, and ii) left
141 occipital gyrus, with a modulating node at the lingual gyrus, and between: i) left superior/inferior
142 parietal nodes to precentral and middle temporal nodes in the right hemisphere, and ii) left
143 superior/inferior parietal node to left superior frontal nodes (Figure 1). In respect to the core

144 RSNs, as defined by Yeo et al.⁴⁰, we have noted increased connectivity within the visual network,
 145 dorsal attention network (DAN), and default mode network (DMN), and between the
 146 frontoparietal network (FPN) and DMN, DAN and somatosensory network (SMN), as well as
 147 DMN and DAN (Table 2).

148 [place Figure 1 here]

149 **Table 2.** Significant increase in connectivity of RSNs as defined by Yeo et al.⁴⁰. The black color
 150 signifies the within-network connectivity, the green color the bilateral connectivity between-
 151 networks, and the blue color the connectivity between-networks from the RSN in the first column
 152 to the RSN in the first row.

RSN	VIS	SMN	DAN	VAN	LIM	FPN	DMN
VIS							
SMN							
DAN							
VAN							
LIM							
FPN							
DMN							

153

154 Graph measures

155 The significant changes ($p < 0.05$) in global efficiency (GE), transitivity (TS), characteristic
 156 path length (CPL), clustering coefficient (CC), local efficiency (LE), and betweenness centrality
 157 (BC) between the pre- and post-networks were investigated using Analysis of Covariance, where
 158 graph measures served as the dependent values and density as the covariate (Table 3). Regarding
 159 the global measures, we report increases in GE, and TS, and a decrease in CPL. From the
 160 comparison of local measures, CC and LE show significant differences in all nodes, with
 161 increased values in 739 out of 863 nodes (Figure 2). From these, 374 nodes belong to the left
 162 hemisphere (91 in the frontal lobe (FL), 55 in the central midline (CM), 64 in the parietal lobe
 163 (PL), 69 in the temporal lobe (TL), 40 in the limbic cortex (LC), 55 in the occipital lobe (OL))
 164 and 365 to the right hemisphere (99 in FL, 50 in CM, 63 in PL, 57 in TL, 36 in LC, 58 in OL, 1
 165 in the uvula, 1 in the insula) (Figure 2). From the remaining 124 nodes (significant decrease in
 166 value), 49 nodes are localized in the left hemisphere (37 in FL, 2 in CM, 4 in TL, 2 in LC, 2 in
 167 OL) and 75 in the right hemisphere (27 in FL, 15 in CM, 2 in PL, 19 in TL, 11 in LC, 1 in OL).
 168 BC increased in one node of the left fusiform gyrus and decreased in three nodes of the left
 169 superior frontal and one of the inferior frontal gyrus

170 **Table 3.** Significant post-pre changes in global (Global efficiency, Transitivity, Characteristic
171 Path Length) and local (Local Clustering Coefficient, Local Efficiency, Node Betweenness
172 Centrality) graph measures. Results were considered significant for $p < 0.05$. *For the local graph
173 measures the p-value reported is the mean value of the nodes showing a significant increase and a
174 significant decrease, respectively.

Graph Measure	p-value	Direction of Effect
Global Efficiency	0.00	Increase
Transitivity	0.00	Increase
Characteristic Path Length	0.00	Decrease
Clustering Coefficient	1.63e-07/8.59e-07*	Increase/Decrease
Local Efficiency	1.63e-07/8.59e-07*	Increase/Decrease
Node Betweenness Centrality	0.0221/0.0487*	Increase/Decrease

175

176

[place Figure 2 here]

177

178 **Discussion**

179 We present novel evidence, indicating that a ten-week intervention of combined physical and
180 cognitive training, in adults with DS, can trigger neuroplasticity causing a cortical reorganization,
181 quantified via EEG indices and graph measures. Our results reveal that: i) short-term training can
182 modify the physical and cognitive performance of adults with DS, ii) the DS brain can adapt to
183 novel stimulation and challenges by utilizing neuroplasticity and reorganizing itself, and iii)
184 adaptive neuroplasticity may lead to a more complex, flexible and healthier functional network.

185

186 **Short-term training modifies physical and cognitive performance**

187 The physical training (aerobic, flexibility, strength, and balance exercises) followed here can
188 significantly improve physical capacity for upper body strength and endurance (Arm Curl Test),
189 mobility, static and dynamic balance (Time Up and Go). Our findings for PT are in line with
190 previously reported findings^{33,41} on adult PwDS. However, our protocol targets multiple physical
191 domains, while most studies investigating the effect of PT on the somatic capacity of adult
192 PwDS^{33,41} have mainly focused on a specific domain, i.e. resistance training or aerobic training
193 and, less often, on a combination of two or more domains. The changes reported here can result in

194 improved health status and decreased risks for several DS related complications, such as
195 cardiovascular disease⁴².

196 The cognitive module of the intervention targets memory, attention, cognitive processing
197 speed, and orientation, as well as social skills. Our participants with DS exhibited an improved
198 general cognitive capacity, i.e., in the level of general intelligence (Raven AB and Raven Total),
199 planning and organization skills (WISC-III, Mazes), and in short-term memory, attention, and
200 concentration (WISC-III, Digit Span, Digit Forward). Previous cognitive interventions on adult
201 PwDS have mainly focused on memory³⁴ and, less often, on executive functions (i.e., planning,
202 attention, working memory, problem-solving, and processing speed)³⁵. In line with the findings of
203 McGlinchey and colleagues³⁵, we also report that adult PwDS can complete a computerized CT
204 program that targets multiple domains of cognition, and we similarly report significant
205 improvements.

206

207 **Cortical reorganization in the DS brain and neuroplastic capacity**

208 Our findings on the changes (i.e., increased connectivity) of the eyes-open resting-state
209 network indicate that PwDS present capacity for neuroplasticity. Regarding the type of
210 neuroplasticity that is expressed here, we can rule out the triggering of evolutionary and reactive
211 plasticity, since the first is most active during the development of the brain¹, and the latter is most
212 commonly triggered by a single stimulus¹ (our training utilizes repetitive stimuli). Adaptational
213 and reparation plasticity can both cause cortical reorganization through long-term stimulation.

214 In the present study, combined physical and cognitive training has increased the engagement
215 of the left hemispheric nodes in information processing and has also increased communication
216 between the nodes of the two hemispheres (Figure 1A&C). The DS cortical networks are
217 characterized by a simplified organization¹² with hyper-synchrony between brain regions^{12,13,36}.
218 Therefore, the reported shifts in connectivity indicate the emergence of a more complex network
219 organization which can be the result of either adaptational or reparation plasticity. Our finding
220 regarding the increased left intra-hemispheric information flow can be regarded as the result of
221 adaptational neuroplasticity being triggered in the DS brain given the atypical right-hemispheric
222 functional preference⁴³. However, the left-to-right hemisphere directionality is a pattern that has
223 been previously evidenced in young PwDS⁸, and its enhancement in adult PwDS may indicate
224 that the training has triggered reparation neuroplasticity.

225 The increased connectivity within (visual, DAN, DMN) and between (FPN and DMN, DAN
226 and SMN, DMN and DAN, DMN and SMN) the core RSNs⁴⁰, further highlights the DS network

227 complexity due to training. Given the evidence pointing towards an inverse relationship between
228 the level of cognitive impairment of PwDS and the within DAN^{36,37} connectivity (i.e. the lower
229 the DAN connectivity, the higher the cognitive impairment for DS), we suggest that the increased
230 connectivity within DAN may associate with the increased cognitive capacity due to training in
231 our DS sample. This would rule out the possibility that the type of plasticity represents reparation.
232 In essence, future studies could potentially confirm or rule out this possibility.

233 The cortical reorganization in conjunction with the increases in general intelligence indicates
234 that the DS brain has entered a more flexible state. The DS network has been previously reported
235 to lack the flexibility that characterizes the development of the TD brain¹³ and subsequently
236 higher intelligence^{44,45}. Flexibility can reflect the brain's capacity to adapt to novel stimulations⁴⁴.
237 It is plausible that the transition to a healthier organization emerges from the increased DS
238 network flexibility and further points to the triggering of adaptational neuroplasticity.

239

240 **Graph theory characteristics support the neuroplastic transition towards a** 241 **healthier DS network organization**

242 We suggest that the network's reorganization, due to training and triggering of neuroplasticity,
243 indexes a transitional state from a random-like network towards a healthier functional architecture
244 (please see Figure 3 for an illustration of our theoretical proposal). Considering the DS brain's
245 simplified architecture, and the lack of functional specialization with impaired integration and
246 segregation¹², the significant changes (i.e., increase) in GE and TS hint that an adult DS
247 functional network, similar to the one here before training, retains features of a random network³⁸.
248 This is not surprising given that basic cognitive abilities (low level of general intelligence, which
249 also characterizes PwDS) are exemplified by random networks, while broad cognitive abilities
250 (high level of general intelligence) would require a small-world (SW) network^{44,45}. SW networks
251 maintain an optimal balance of integration and segregation (combining characteristics of random
252 and regular networks), to support not only general but also specific abilities^{44,45}. In contrast, the
253 absence of SW properties in the DS brain, namely reduced GE, TS, and therefore robustness,
254 indicates the increased risk of communication loss between connected regions and their causality
255 for cognitive decline^{38,46}. The simultaneous increment in global efficiency and transitivity denotes
256 the increase of both integration and segregation and the transition to a less-random, healthier
257 network organization. Potentially, the training could further induce cortical reorganization,
258 resulting in a more modular network capable of facilitating higher cognitive abilities and
259 engaging broad and perhaps specific abilities.

260 The decrease of CPL in TD individuals could be interpreted as a contradicting shift towards a
261 more random network that disturbs the SW balance. In our case, the reduction of CPL further
262 supports our hypothesis of a transitional stage, since the adult DS network deviates from its
263 random-like characteristics by displaying a greater CPL. Previous fMRI studies report that the
264 brain network of PwDS presents increased local connectivity^{12,13} indicating increased CPL.
265 Hence, the decrease of CPL indeed indicates the transition to a less-random network and could
266 point to the deceleration or even reversal of degenerative processes in the DS brain.

267 If the optimal function of the brain is to be attributed to SW network architecture, it is
268 plausible that the adaptational neuroplastic transition has characteristics that would eventually
269 lead to such a network. Here, these characteristics are evident not only on a global level (i.e.,
270 changes in CPL and GE) but also from a local network perspective. The increases in CC and LE
271 support our hypothesis, as they indicate an increase in segregation and robustness, less
272 randomness in functional clustering organization, and additionally, an increase of fault tolerance
273 in the network⁴⁷. The post-training network robustness and segregation can potentially serve in
274 neuroprotection for PwDS³⁸. Still, the effectiveness of such a mechanism would rely on the trade-
275 off between integration and segregation, mainly maintaining SW network characteristics⁴⁸. These
276 shifts signal the decrease in wiring cost and subsequent increase in cost-efficacy^{47,49} (GE-wiring
277 cost), rendering the network more efficient as a whole. Finally, the transition towards SW
278 network characteristics is backed up by the changes in BC (increase and decrease), which point to
279 a more hierarchical and less random organization. Previously, it was reported that the DS brain is
280 characterized by decreased centrality¹⁰.

281 [place Figure 3 here]

282

283 **Future directions and limitations**

284 A limitation of the study was the small sample size of the experimental group. Our results
285 could potentially be interpreted differently in reference to a control group (TD or passive DS).
286 Though this study is the first to provide insights into the benefits of combined PT and CT in
287 adults with DS through the use of neuropsychological and neurophysiological assessments, a gap
288 remains in our basic understanding of the sole contributions of each training to DS related
289 cognition and brain function. Further studies are necessary to index the influence of each
290 component on the triggering of neuroplasticity and subsequently network organization and
291 cognition. Similarly, the stability of the reported neuroplastic shifts can only be addressed

292 through additional and more in-depth investigations, including follow-up and further longitudinal
293 studies, to fill the gaps in the existing literature.

294

295 **Conclusion**

296 The rise of DS prevalence sets a significant challenge in developing innovative health care
297 interventions that provide the best quality of care. Neurocognitive treatments can augment the
298 functional and cognitive abilities of PwDS and thus allow for a more independent, productive,
299 and fulfilling daily life. Our results are based on brain imaging, connectivity, and graph-theory
300 analysis, and emphasize the significance of introducing stimulation and adaptable challenges in
301 the environment of PwDS. We provide evidence that triggering adaptational neuroplasticity in the
302 DS brain provokes the emergence of a less-random, more complex, hierarchical network,
303 accompanied by an increased ability to integrate and segregate information and improved
304 efficiency, robustness and flexibility. Designing focalized neurobehavioral interventions can aid
305 in the development of a balanced and stable functional DS phenotype, in terms of SW
306 characteristics.

307

308 **Materials and Methods**

309 **Subjects**

310 The study's pool consisted of 12 subjects with DS (age: 29±11, 6 females). Participants were
311 recruited from a variety of local organizations (please see Acknowledgements). The training
312 procedure took place in the Thessaloniki Active and Healthy Ageing Living Lab (Thess-
313 AHALL)⁵⁰, and the premises of the Greek Association of Down Syndrome. This study is part of
314 the dsLLM clinical trial, registered with ClinicalTrials.gov, with identifier code [NCT04390321](https://clinicaltrials.gov/ct2/show/study/NCT04390321).
315 The study protocol was approved by the Bioethics Committee of the Medical School of the
316 Aristotle University of Thessaloniki and was conducted per the Helsinki Declaration of Human
317 Rights. The participants' legal guardians signed a written informed consent prior to their
318 inclusion in the study.

319

[place Figure 4 here]

320

321 **LLM Care Intervention**

322 The intervention protocol consisted of physical and cognitive training (LLM Care⁵¹,
323 <http://www.llmcare.gr/en>). The study protocol was further developed to improve the quality of
324 life and aid in the development of independent living skills in PwDS^{52,53}. LLM Care also aims to
325 further improve brain functionality⁵¹. All training sessions were computerized, center-based, and
326 conducted under supervision. The sequence of training methods was pseudo-randomized and
327 counterbalanced. The details of each training intervention have been previously described in
328 detail^{54,55} and are summarized in Figure 4.

329

330 **Cognitive training**

331 The cognitive training (CT) component of LLM Care uses the BrainHQ software (Posit
332 Science Corporation, San Francisco, CA, USA), an online interactive environment in Greek
333 language⁵⁶. It consists of six categories (29 exercises in total), with customizable difficulty levels,
334 utilizing audiovisual stimuli. CT targets memory, attention, cognitive-processing speed,
335 navigation, and people skills. This regime was selected since patients with DS exhibit
336 deficiencies in these processes. CT was conducted for half an hour, with a frequency of 2 days
337 per week for 10 weeks. In every CT session, the participants were required to complete at least
338 one task from each category. During the training, participants were urged to complete as many
339 exercises as they could from each category.

340

341 **Physical training**

342 The physical training (PT) component of LLM Care is based on the WebFitForAll
343 protocol^{55,57}, adequately adjusted to the needs and capacity of PwDS. It utilizes motion sensor
344 devices (i.e., Kinect). Games and physical exercise are combined, providing a pleasant experience
345 throughout the training. PT sessions lasted for half an hour and were conducted with the same
346 frequency as CT. The training consists of aerobic (cycling, in-place-hiking), flexibility
347 (stretching), strength (resistance, weightlifting), and balance (static, dynamic) exercises. The
348 warm-up and cool-down routines (5-minutes duration), signify the start and completion of every
349 session, respectively. During aerobic exercises, the participants entered a virtual environment, set
350 up in Google maps, and explored cities and landscapes. Upon correct completion of the flexibility
351 and strength exercises, the trainees were progressively rewarded with an array of pleasing images.
352 The scope of balance exercises was to move their bodies either horizontally or vertically, which
353 was achieved through games.

354

355 **Psychometric and somatometric assessments**

356 The participants' cognitive and physical capacity was assessed before and after the
357 intervention. The psychometric evaluation consisted of a set of neurocognitive tests that measure
358 memory attention, concentration (WISC-III⁵⁸: Digits Span), verbal and non-verbal mental
359 capabilities (Raven⁵⁹), processing speed (WISC-III: Digits Span and Picture Arrangement),
360 problem-solving, visuospatial processing, organization skills (WISC-III: Block Design, Picture
361 Arrangement, Mazes), social intelligence (WISC-III: Picture Arrangement), and identification of
362 emotions (Reading the mind in the eyes^{60,61}, and a variation: Reading the mind in the face
363 (emotion recognition from video)).

364 The somatometric evaluation included the Short Physical Performance Battery (SPPB)⁶², 10
365 Meter Walk⁶³, Back Scratch⁶⁴, Sit and Reach⁶⁵, Arm Curl⁶⁴, Four Square Step (FSST)⁶⁶, Stork
366 Balance (for both legs)⁶⁷, Timed Up and Go⁶⁸ tests, and Body Mass Index (BMI). These tests
367 appraise functioning mobility, flexibility (in specific areas), dynamic stability, strength, and static
368 and dynamic balance.

369

370 **EEG recording**

371 Pre- and post-intervention resting-state EEG activity was recorded for 5 minutes, using a high-
372 density Nihon-Kohden EEG device (128 active scalp electrode) at a sampling rate of 1000Hz.
373 The EEG recordings were performed in an electrically shielded, sound, and light attenuated
374 booth. The electrode impedances were lower than 10 k Ω . The participants were instructed to
375 remain in a resting position while keeping their eyes open. Eyes-closed EEGs were not measured
376 due to the limited capacity of the participants with DS to stay relaxed with their eyes closed.

377

[place Figure 5 here]

378

379 **EEG data analysis**

380 **Pre-processing**

381 The raw EEG data were visually inspected and bad channels were interpolated while any eye-
382 movement related artifacts (blinks and horizontal movement) were corrected, through adaptive
383 artifact correction⁷⁰ using the Brain Electrical Source Analysis software (BESA research, version
384 6, Megis Software, Heidelberg, Germany) (Figure 5, blue section). The artifact-corrected data
385 from each measurement were imported in the Fieldtrip Matlab toolbox⁷¹ for additional processing

386 (Figure 5, blue section). The signals were denoised by filtering the data (0.53 Hz high-pass IIR
387 filter, 48-52 Hz notch IIR filter, 97 Hz low-pass IIR filter). The filtered signals were analyzed
388 into independent components⁷², and the artifactual components were rejected. After signal
389 reconstruction, the data was, once again, visually inspected for any remaining artifacts. Randomly
390 selected fifteen segments (4 seconds each) from each artifact-free EEF recording for additional
391 processing.

392

393 **Source reconstruction**

394 Each subject's segments (15 segments, 4000 samples, 4 seconds duration each) were imported
395 into BESA (Figure 5, orange section). The current density reconstructions (CDR) were estimated
396 for each sample point, solving the inverse problem using LORETA in the 0.5-35 Hz frequency
397 range. LORETA³⁹ was utilized, as it does not require the a priori declaration of the number of
398 sources and is suitable for whole cortex analysis. The CDRs were exported as four-dimensional
399 images (4-D) in the Analyze format (keeping all sampling points), which were, in turn, imported
400 into Matlab. A cortex mask was superimposed on the images. The mask includes only grey
401 matter, and excludes the subcortex, the brainstem, and cerebellum, to limit the source space⁶⁹.
402 The source space consisted of 863 voxels.

403

404 **Functional connectivity**

405 After extracting the time-series of each voxel from the 4-D images, they were used to compute
406 the Phase Transfer Entropy (PTE)⁷³ (Figure 5, green section). The computation resulted in
407 863×863 adjacency matrices. The metric is calculated independently for every pair of voxels in
408 each segment. Each voxel represented a node of the brain network, with the node's coordinates
409 corresponding to the center of each voxel. PTE is the estimation of TE between phase time-series,
410 i.e., it evaluates the influence of one signal's phase on another signal's phase⁷³. PTE was selected
411 because its results are not based on a specific data model since its computation is reliant on non-
412 linear probability distributions. Therefore, it allows the detection of higher-order relations in the
413 phase information flow and renders the measure resistant to source leakage⁷³. The algorithm
414 applies the Hilbert transform to estimate the phases of each signal. It determines the number of
415 bins by utilizing the Scott methodology⁷⁴, resulting in 37 bins on average (13 samples per bin).
416 The adjacency matrices of each subject were averaged, resulting in two sets of 12 adjacency
417 matrices (Figure 5, green section).

418

419 **Graph measures computation**

420 Using the Brain Connectivity Toolbox⁷⁵, the graph measures of global efficiency (GE),
421 transitivity (TS, a variation of global clustering coefficient), characteristic path length (CPL),
422 clustering coefficient (CC), local efficiency (LE) and node betweenness centrality (BC) were
423 computed for each participant. In a later step, the centrality Matlab function, which measures
424 node importance, was used to compute the node degree centrality (DC) of the network mapping
425 the pre- and post-intervention differences. The density of each graph was estimated by
426 summarizing the weights of each graph.

427 These measures were chosen to examine the influence of our intervention on the
428 characteristics of the DS brain and its functional organization. To that aim, GE and CPL were
429 utilized to index the effects on integration⁷⁵, TS, CC, and LE to measure the effects on
430 segregation⁷⁵ and BC and DC to check for hierarchical changes in the network. GE is the average
431 inverse shortest path length of the network⁷⁶ and quantifies the efficacy of information
432 transference and its assimilation in the network⁷⁵. CPL is the average shortest path length of the
433 edges connecting the nodes of the network⁷⁷ and characterizes its robustness⁷⁵. TS is the ratio of
434 closed triplets to the maximum number of triplets (open (three nodes with up to two connections
435 between them) and closed (three nodes with three connections between them))⁷⁸, while CC of a
436 node is the ratio of its connected neighbors to the maximum number of possible connections⁷⁷.
437 These two measures reflect the clustering organization of the DS brain network on a global and
438 local level, respectively, and also quantify its robustness⁷⁵. LE of a node is the computation of GE
439 on a local level, strongly related to CC^{76,79}, and measures the efficiency of information
440 transference to the node's neighbors. BC corresponds to the fraction of shortest paths that pass
441 through a node⁸⁰ and is a measure of centrality, measuring the importance of a node in the
442 information flow between nodes⁷⁵. So, the higher the value of BC is, the more vital that node is
443 for the transfer of information in the network. Lastly, DC calculates the number of links
444 connected to a specific node, i.e., it measures if a node exhibits higher or lower connectivity to
445 other nodes.

446

447 **Statistical analysis**

448 The pre- and post-intervention somatometric and psychometric assessment scores, were
449 compared with the use of non-parametric Wilcoxon tests and paired t-tests. The statistical
450 comparisons were performed using the IBM SPSS 25.0 software. Wilcoxon tests were performed

451 on the psychometric battery tests and somatometric tests with discrete-values score, while paired
452 t-tests were applied on the remaining somatometric assessment tests.

453 The Network Based Statistics (NBS)⁸¹ MATLAB toolbox was employed to estimate the
454 statistically significant differences between the whole-head network of the pre- and post-
455 intervention connectivity networks of our subjects. A paired samples t-test corrected for 10000
456 random comparisons via False Discovery Rate (FDR) correction⁸² was performed. This
457 methodology evaluates the significance of each edge independently, providing an independent p-
458 value for each connection. Significant differences between the two time-points were visualized as
459 weighted graphs through the BrainNet Viewer⁸³ toolbox (Figure 1). DC was estimated from the
460 outcome of this comparison, and the results were depicted in the same graph.

461 For the graph measures, Analysis of Covariance (ANCOVA) was used, where each measure
462 serves as the dependent variable and density as the covariate, because density seriously affects the
463 values of the other measures. For local measures (i.e., CC, LE, and BC), ANCOVA and FDR
464 correction were performed for each of the 863 nodes and p-values for Type I errors, respectively.
465 Results below the 5 percent threshold were considered significant.

466

467 **Acknowledgments**

468 This study is an extension of the European CIP-ICTPSP. 2008.1.4 Long Lasting Memories
469 (LLM) project (Project no. 238904) (<http://www.longlastingmemories.eu/>). The authors would
470 like to thank all the participants, their families, and caregivers, the Down Syndrome Association
471 of Greece, Spring Children Mixed Living Center – Daily Care Activity & Training for Disabled
472 People, The Down Syndrome Association of Serres, Center for the Rehabilitation of Social
473 Support & Creative Employment of People with Disabilities, Child Care Center in Ptolemaida,
474 Centre For Persons With Learning Difficulties - Zoodohos Pigi, Center of Creative Activities -
475 Municipality of Neapoli and Sykies, Special laboratory for vocational education and training in
476 Alexandria, Center for the Rehabilitation of Social Support & Creative Employment of People
477 with disabilities "O Sotir", Center of Creative Activities for People with Disabilities-Municipality
478 of Neapoli and Sykes. Also, the authors would like to thank a list of people assisting in the trial
479 facilitation, as follows: Dimitris Bamidis, Foteini Dolianiti, Ioanna Dratsiou, Sotiria Gilou, Maria
480 Karagianni, Katerina Katsouli, Eleni Manthou, Maria Metaxa, Annita Varella (in alphabetical
481 order), as well as, students from the Dept of Psychology, The University of Sheffield
482 International Faculty, City College (Greece) who were practicing under the supervision of Dr.
483 Anna Vivas and Dr. Elissavet Chrysochoou who were also advising the authoring team on the

484 psychological aspects of the whole endeavor. Last but not least, the authors would like to thank
485 Emeritus Professor of Pediatrics and Genetics Dr. Charikleia Chatzisevastou-Loukidou for
486 advising the team on all medical aspects of the project and for linking the team with many of the
487 pilot sites.

488

489 **Conflict of Interest**

490 There are potential conflicts of interest (other, not financial, outside the scope of the submitted
491 work) for the author P. Bamidis in respect of PositScience and the Aristotle University of
492 Thessaloniki. There is a co-marketing agreement between PositScience and the Aristotle
493 University of Thessaloniki to exploit Brain HQ within the LLM Care self-funded initiative that
494 emerged as the non-for-profit business exploitation of the Long Lasting Memories (LLM Project)
495 (www.longlastingmemories.eu) originally funded by the ICT-CIP-PSP Program of the European
496 Commission. Brain HQ now forms part of LLM Care, a technology transfer/self-funded initiative
497 that emerged as the non-for-profit business exploitation of LLM. Additionally, FitForAll (FFA)
498 has been developed in the Aristotle University of Thessaloniki during the Long Lasting Memories
499 (LLM Project) (www.longlastingmemories.eu) originally funded by the ICT-CIP-PSP Program of
500 the European Commission. It now forms part of LLM Care, a technology transfer/self-funded
501 initiative that emerged as the non-for-profit business exploitation of LLM.

502

503 **References**

- 504 1. Trojan, S. & Pokorny, J. Theoretical aspects of neuroplasticity. *Physiol. Res.* **48**, 87–98
505 (1999).
- 506 2. Baroncelli, L. *et al.* Brain plasticity and disease: A matter of inhibition. *Neural Plast.*
507 **2011**, (2011).
- 508 3. Dierssen, M., Herault, Y. & Estivill, X. Aneuploidy: From a Physiological Mechanism of
509 Variance to Down Syndrome. *Physiol. Rev.* **89**, 887–920 (2009).
- 510 4. White, N. S., Alkire, M. T. & Haier, R. J. A voxel-based morphometric study of
511 nondemented adults with Down Syndrome. *Neuroimage* **20**, 393–403 (2003).
- 512 5. Pinter, J. D., Eliez, S., Schmitt, J. E., Capone, G. T. & Reiss, A. L. Neuroanatomy of
513 Down's Syndrome: A High-Resolution MRI Study. *Am. J. Psychiatry* **158**, 1659–1665
514 (2001).

- 515 6. Dierssen, M. Down syndrome: the brain in trisomic mode. *Nat. Rev. Neurosci.* **13**, 844–
516 858 (2012).
- 517 7. Menghini, D., Costanzo, F. & Vicari, S. Relationship Between Brain and Cognitive
518 Processes in Down Syndrome. *Behav. Genet.* **41**, 381–393 (2011).
- 519 8. Babiloni, C. *et al.* Inter-hemispheric functional coupling of eyes-closed resting EEG
520 rhythms in adolescents with Down syndrome. *Clin. Neurophysiol.* **120**, 1619–1627 (2009).
- 521 9. Babiloni, C. *et al.* Cortical sources of EEG rhythms are abnormal in down syndrome. *Clin.*
522 *Neurophysiol.* **121**, 1205–1212 (2010).
- 523 10. Hemmati, S., Ahmadi, M., Gharib, M., Vameghi, R. & Sajedi, F. Down syndrome’s
524 brain dynamics: analysis of fractality in resting state. *Cogn. Neurodyn.* **7**, 333–40 (2013).
- 525 11. García-Alba, J. *et al.* Neuropsychological and neurophysiological characterization of mild
526 cognitive impairment and Alzheimer’s disease in Down syndrome. *Neurobiol. Aging* **84**,
527 70–79 (2019).
- 528 12. Anderson, J. S. *et al.* Abnormal brain synchrony in Down Syndrome. *NeuroImage Clin.* **2**,
529 703–715 (2013).
- 530 13. Edgin, J. O., Clark, C. A. C., Massand, E. & Karmiloff-Smith, A. Building an adaptive
531 brain across development: Targets for neurorehabilitation must begin in infancy. *Front.*
532 *Behav. Neurosci.* **9**, (2015).
- 533 14. Head, E., Lott, I. T., Patterson, D., Doran, E. & Haier, R. J. Possible Compensatory Events
534 in Adult Down Syndrome Brain Prior to the Development of Alzheimer Disease
535 Neuropathology: Targets for Nonpharmacological Intervention. *J. Alzheimer’s Dis.* **11**,
536 61–76 (2007).
- 537 15. Haier, R. J., Head, K., Head, E. & Lott, I. T. Neuroimaging of individuals with Down’s
538 syndrome at-risk for dementia: Evidence for possible compensatory events. *Neuroimage*
539 **39**, 1324–1332 (2008).
- 540 16. Takashima, M. *et al.* Neuronal Plasticity in the Developing and Aging Cerebral Cortices
541 of Patients with Down Syndrome. *Int. J. Neurorehabilitation* **01**, (2014).
- 542 17. Katada, A. *et al.* On chronological changes in the basic EEG rhythm in persons with
543 Down syndrome - With special reference to slowing of alpha waves. *Brain Dev.* **22**, 224–
544 229 (2000).
- 545 18. Costa, A. C. S. On the promise of pharmacotherapies targeted at cognitive and
546 neurodegenerative components of Down syndrome. *Dev. Neurosci.* **33**, 414–27 (2011).
- 547 19. Lott, I. T. *et al.* Down syndrome and dementia: A randomized, controlled trial of
548 antioxidant supplementation. *Am. J. Med. Genet. Part A* **155**, 1939–1948 (2011).

- 549 20. Mohan, M., Carpenter, P. K. & Bennett, C. Donepezil for dementia in people with Down
550 syndrome. *Cochrane Database Syst. Rev.* (2009).
- 551 21. Kishnani, P. S. *et al.* Donepezil for treatment of cognitive dysfunction in children with
552 Down syndrome aged 10-17. *Am. J. Med. Genet. Part A* **152**, 3028–3035 (2010).
- 553 22. Kálmán, J. *et al.* Serum interleukin-6 levels correlate with the severity of dementia in
554 Down syndrome and in Alzheimer’s disease. *Acta Neurol. Scand.* **96**, 236–240 (2009).
- 555 23. Erickson, K. I. *et al.* Exercise training increases size of hippocampus and improves
556 memory. *Proc. Natl. Acad. Sci.* **108**, 3017–3022 (2011).
- 557 24. Colcombe, S. J. *et al.* Cardiovascular fitness, cortical plasticity, and aging. *Proc. Natl.*
558 *Acad. Sci. U. S. A.* **101**, 3316–21 (2004).
- 559 25. Cotman, C., Berchtold, N. C. & Christie, L. A. Exercise builds brain health: key roles of
560 growth factor cascades and inflammation. *Trends Neurosci.* **30**, 464–472 (2007).
- 561 26. Styliadis, C., Kartsidis, P., Paraskevopoulos, E., Ioannides, A. A. & Bamidis, P. D.
562 Neuroplastic Effects of Combined Computerized Physical and Cognitive Training in
563 Elderly Individuals at Risk for Dementia: An eLORETA Controlled Study on Resting
564 States. *Neural Plast.* **2015**, 1–12 (2015).
- 565 27. Styliadis, C., Kartsidis, P. & Paraskevopoulos, E. Neuroimaging approaches for elderly
566 studies. in *Handbook of Research on Innovations in the Diagnosis and Treatment of*
567 *Dementia* (eds. Bamidis, P. D., Tarnanas, I., Hadjileontiadis, L. & Tsolaki, M.) 47–86 (IGI
568 Global, 2015).
- 569 28. Klados, M. A., Styliadis, C., Frantzidis, C. A., Paraskevopoulos, E. & Bamidis, P. D.
570 Beta-Band Functional Connectivity is Reorganized in Mild Cognitive Impairment after
571 Combined Computerized Physical and Cognitive Training. *Front. Neurosci.* **10**, 55 (2016).
- 572 29. Bamidis, P. D. *et al.* A review of physical and cognitive interventions in aging. *Neurosci*
573 *Biobehav Rev.* **44**, 206–220 (2014).
- 574 30. Kramer, A. F. *et al.* Ageing, fitness and neurocognitive function. *Nature* **400**, 418–419
575 (1999).
- 576 31. Colcombe, S. J. *et al.* Aerobic Exercise Training Increases Brain Volume in Aging
577 Humans. *Journals Gerontol. Ser. A Biol. Sci. Med. Sci.* **61**, 1166–1170 (2006).
- 578 32. Bertapelli, F., Pitetti, K., Agiovlasis, S. & Guerra-Junior, G. Overweight and obesity in
579 children and adolescents with Down syndrome—prevalence, determinants, consequences,
580 and interventions: A literature review. *Res. Dev. Disabil.* **57**, 181–192 (2016).
- 581 33. Hardee, J. P. & Fetters, L. The effect of exercise intervention on daily life activities and
582 social participation in individuals with Down syndrome: A systematic review. *Res. Dev.*

- 583 *Disabil.* **62**, 81–103 (2017).
- 584 34. Fonseca, L. M., Navatta, A. C. R., Bottino, C. M. C. & Miotto, E. C. Cognitive
585 Rehabilitation of Dementia in Adults with Down Syndrome: A Review of Non-
586 Pharmacological Interventions. *Dement. Geriatr. Cogn. Dis. Extra* **5**, 330–40 (2015).
- 587 35. McGlinchey, E., McCarron, M., Holland, A. & McCallion, P. Examining the effects of
588 computerised cognitive training on levels of executive function in adults with Down
589 syndrome. *J. Intellect. Disabil. Res.* **63**, 1137–1150 (2019).
- 590 36. Vega, J. N., Hohman, T. J., Pryweller, J. R., Dykens, E. M. & Thornton-Wells, T. A.
591 Resting-State Functional Connectivity in Individuals with Down Syndrome and Williams
592 Syndrome Compared with Typically Developing Controls. *Brain Connect.* **5**, 461–475
593 (2015).
- 594 37. Pujol, J. *et al.* Anomalous brain functional connectivity contributing to poor adaptive
595 behavior in Down syndrome. *Cortex* **64**, 148–156 (2015).
- 596 38. Ahmadi, M., Gharib, M., Hemmati, S., Vameghi, R. & Sajedi, F. Disrupted small-world
597 brain network in children with Down Syndrome. *Clin. Neurophysiol.* **124**, 1755–1764
598 (2013).
- 599 39. Pascual-Marqui, R. D., Michel, C. M. & Lehmann, D. Low resolution electromagnetic
600 tomography: a new method for localizing electrical activity in the brain. *Int. J.*
601 *Psychophysiol.* **18**, 49–65 (1994).
- 602 40. Yeo, B. T. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic
603 functional connectivity. *J. Neurophysiol.* **106**, 1125–65 (2011).
- 604 41. Ruiz-González, L., Lucena-Antón, D., Salazar, A., Martín-Valero, R. & Moral-Munoz, J.
605 A. Physical therapy in Down syndrome: systematic review and meta-analysis. *J. Intellect.*
606 *Disabil. Res.* **63**, 1041–1067 (2019).
- 607 42. Rimmer, J. H., Heller, T., Wang, E. & Valerio, I. Improvements in Physical Fitness in
608 Adults With Down Syndrome. *Am. J. Ment. Retard.* **109**, 165 (2004).
- 609 43. Elliott, D., Weeks, D. J. & Chua, R. Anomalous Cerebral Lateralization and Down
610 Syndrome. *Brain Cogn.* **26**, 191–195 (1994).
- 611 44. Barbey, A. K. Network Neuroscience Theory of Human Intelligence. *Trends Cogn. Sci.*
612 **22**, 8–20 (2018).
- 613 45. Hilger, K., Fukushima, M., Sporns, O. & Fiebach, C. J. Temporal stability of functional
614 brain modules associated with human intelligence. *Hum. Brain Mapp.* **41**, 362–372
615 (2019).
- 616 46. Sanz, J. M. *et al.* Loss of ‘Small-World’ Networks in Alzheimer’s Disease: Graph

- 617 Analysis of fMRI Resting-State Functional Connectivity. *PLoS One* **5**, (2010).
- 618 47. Achard, S. & Bullmore, E. Efficiency and cost of economical brain functional networks.
619 *PLoS Comput. Biol.* **3**, e17 (2007).
- 620 48. Stam, C., Jones, B., Nolte, G., Breakspear, M. & Scheltens, P. Small-World Networks and
621 Functional Connectivity in Alzheimer's Disease. *Cereb. Cortex* **17**, 92–99 (2006).
- 622 49. Bullmore, E. & Sporns, O. The economy of brain network organization. *Nat. Rev.*
623 *Neurosci.* **13**, 336–49 (2012).
- 624 50. Konstantinidis, E. I., Billis, A. S., Bratsas, C., Siountas, A. & Bamidis, P. D. Thessaloniki
625 Active and Healthy Ageing Living Lab: the roadmap from a specific project to a living lab
626 towards openness. in *9th International Conference PErsasive Technologies Related to*
627 *Assistive Environments (PETRA)* 1–4 (2016).
- 628 51. Bamidis, P. D. *et al.* Gains in cognition through combined cognitive and physical training:
629 the role of training dosage and severity of neurocognitive disorder. *Front. Aging Neurosci.*
630 **7**, 152 (2015).
- 631 52. Bamidis, P. D., Konstantinidis, E. I., Billis, A. S. & Siountas, A. From e-homes to living
632 labs: founding and organising the Greek active and healthy ageing living lab (Thess-
633 AHALL) and its networked services. *Hell. J. Nucl. Med.* **20**, 112–126 (2017).
- 634 53. Romanopoulou, E., Zilidou, V., Savvidis, T., Chatzisevastou-Loukidou, C. & Bamidis, P.
635 D. Unmet Needs of Persons with Down Syndrome: How Assistive Technology and Game-
636 Based Training May Fill the Gap. *Stud. Health Technol. Inform.* **251**, 15–18 (2018).
- 637 54. Smith, G. E. *et al.* A cognitive training program based on principles of brain plasticity:
638 Results from the improvement in memory with plasticity-based adaptive cognitive training
639 (IMPACT) study. *J. Am. Geriatr. Soc.* **57**, 594–603 (2009).
- 640 55. Billis, A. S. *et al.* A game-like interface for training seniors' dynamic balance and
641 coordination. in *IFMBE Proceedings* 691–694 (2010).
- 642 56. Romanopoulou, E. D., Zilidou, V. I. & Bamidis, P. D. Creating and sustaining a social
643 health care ecosystem: the case of LLM care services in Greece. *Hell. J. Nucl. Med.* **20**,
644 40–48 (2017).
- 645 57. Konstantinidis, E. I., Bamparopoulos, G. & Bamidis, P. D. Moving Real Exergaming
646 Engines on the Web: The webFitForAll case study in an active and healthy ageing living
647 lab environment. *IEEE J Biomed Heal. Inform.* **21**, 859–866 (2016).
- 648 58. Woolger, C. Wechsler intelligence scale for children-(WISC-III). in *Understanding*
649 *Psychological Assessment* 219–233 (Springer US, 2001). doi:10.1007/978-1-4615-1185-
650 4_11

- 651 59. John & Raven, J. Raven Progressive Matrices. in *Handbook of Nonverbal Assessment*
652 223–237 (Springer US, 2003).
- 653 60. Baron-Cohen, S., Jolliffe, T., Mortimore, C. & Robertson, M. Another advanced test of
654 theory of mind: evidence from very high functioning adults with autism or asperger
655 syndrome. *J. Child Psychol. Psychiatry.* **38**, 813–22 (1997).
- 656 61. Baron-Cohen, S., Wheelwright, S., Hill, J., Raste, Y. & Plumb, I. The “Reading the
657 Mind in the Eyes” Test revised version: a study with normal adults, and adults with
658 Asperger syndrome or high-functioning autism. *J. Child Psychol. Psychiatry.* **42**, 241–51
659 (2001).
- 660 62. Guralnik, J. M. *et al.* A short physical performance battery assessing lower extremity
661 function: association with self-reported disability and prediction of mortality and nursing
662 home admission. *J. Gerontol.* **49**, M85-94 (1994).
- 663 63. Bohannon, R. W. Comfortable and maximum walking speed of adults aged 20-79 years:
664 reference values and determinants. *Age Ageing* **26**, 15–19 (1997).
- 665 64. Jones, C. J. & Rikli, R. E. Measuring Functional Fitness in Older Adults. *J. Act. Ageing* **1**,
666 24–30 (2002).
- 667 65. Wells, K. F. & Dillon, E. K. The Sit and Reach—A Test of Back and Leg Flexibility. *Res.*
668 *Quarterly. Am. Assoc. Heal. Phys. Educ. Recreat.* **23**, 115–118 (1952).
- 669 66. Whitney, S. L., Marchetti, G. F., Morris, L. O. & Sparto, P. J. The Reliability and Validity
670 of the Four Square Step Test for People With Balance Deficits Secondary to a Vestibular
671 Disorder. *Arch. Phys. Med. Rehabil.* **88**, 99–104 (2007).
- 672 67. Johnson, B. L. & Jack K. Nelson. *Practical Measurements for evaluation in physical*
673 *education.* (1979).
- 674 68. Shumway-Cook, A., Brauer, S. & Woollacott, M. Predicting the probability for falls in
675 community-dwelling older adults using the Timed Up & Go Test. *Phys. Ther.* **80**,
676 896–903 (2000).
- 677 69. Paraskevopoulos, E., Kraneburg, A., Herholz, S. C., Bamidis, P. D. & Pantev, C. Musical
678 expertise is related to altered functional connectivity during audiovisual integration. *Proc.*
679 *Natl. Acad. Sci.* **112**, 201510662 (2015).
- 680 70. Ille, N., Berg, P. & Scherg, M. Artifact correction of the ongoing EEG using spatial filters
681 based on artifact and brain signal topographies. *J. Clin. Neurophysiol.* **19**, 113–24 (2002).
- 682 71. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J. M. FieldTrip: Open source software
683 for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput.*
684 *Intell. Neurosci.* **2011**, 156869 (2011).

- 685 72. Lee, T.-W., Girolami, M. & Sejnowski, T. J. Independent Component Analysis using an
686 Extended Infomax Algorithm for Mixed Sub - Gaussian and Super - Gaussian Sources.
687 *Neural Comput.* **11**, 409–433 (1999).
- 688 73. Lobier, M., Siebenhühner, F., Palva, S. & Palva, J. M. Phase transfer entropy: A novel
689 phase-based measure for directed connectivity in networks coupled by oscillatory
690 interactions. *Neuroimage* **85**, 853–872 (2014).
- 691 74. Scott, D. W. *Multivariate density estimation* □: theory, practice, and visualization. (Wiley,
692 1992).
- 693 75. Rubinov, M. & Sporns, O. Complex network measures of brain connectivity: uses and
694 interpretations. *Neuroimage* **52**, 1059–69 (2010).
- 695 76. Latora, V. & Marchiori, M. Efficient Behavior of Small-World Networks. *Phys. Rev. Lett.*
696 **87**, 198701 (2001).
- 697 77. Watts, D. J. & Strogatz, S. H. Collective dynamics of ‘small-world’ networks. *Nature* **393**,
698 440–442 (1998).
- 699 78. Newman, M. E. J. The structure and function of complex networks. *SIAM Rev.* **45**, 167–
700 256 (2003).
- 701 79. Bullmore, E. & Sporns, O. Complex brain networks: graph theoretical analysis of
702 structural and functional systems. *Nat. Rev. Neurosci.* **10**, 186–198 (2009).
- 703 80. Freeman, L. C. Centrality in social networks conceptual clarification. *Soc. Networks* **1**,
704 215–239 (1978).
- 705 81. Zalesky, A., Fornito, A. & Bullmore, E. T. Network-based statistic: identifying differences
706 in brain networks. *Neuroimage* **53**, 1197–207 (2010).
- 707 82. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
708 Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B* **57**, 289–300 (1995).
- 709 83. Xia, M., Wang, J. & He, Y. BrainNet Viewer: a network visualization tool for human
710 brain connectomics. *PLoS One* **8**, e68910 (2013).

711

712 **Figure Captions**

713 **Figure 1.** Cortical connectivity between post- and pre-intervention networks and for each time
714 point. The color scales represent t-values. **A: Post vs. Pre.** Significant post-pre connectivity
715 differences. Information direction is depicted though line arrows. The visualized networks are
716 significant at a level of $p < 0.05$, FDR corrected. The difference in nodal size depicts the increase
717 in the node degree centrality; the nodes with the most increased connectivity are located at the left






718 parietal lobe. **B**: Depiction of pre- and post-intervention networks in comparison to the null
719 hypothesis. The t-values indicate that the post resting-state network has significantly stronger
720 connections than the pre network. **C**: Circular graph depicting the cortical reorganization in the
721 DS brain (cyan: left hemisphere, green: right hemisphere). The cortical reorganization is
722 characterized by the strengthening of direct connections within: i) left parietal lobe (paracentral,
723 postcentral, precuneus, superior and inferior parietal), and ii) left occipital gyrus, and between: i)
724 left superior/inferior parietal nodes to precentral and middle temporal nodes in the right
725 hemisphere, and ii) left superior/inferior parietal node to left superior frontal nodes.


726 **Figure 2.** Graph depicting nodes with a significant increase in the clustering coefficient (CC).
727 Node size quantifies the increase of the CC value, while the color map indicates the significance
728 of the shift (p-value).

729 **Figure 3.** Illustration of our theoretical proposal. Changes in connectivity and graph-theory
730 characteristics, as an outcome of the adaptational neuroplasticity in the DS brain, characterize the
731 post-intervention DS network (**B** → **C**) as a transitional state from the random-like organization
732 of the pre-intervention DS network³⁸ (**B**) towards a healthier functional structure. Random
733 networks (**A**) are characterized by high global efficiency (over-integration) and low clustering
734 (under-segregation) and exemplify the general abilities of general intelligence (low intelligence
735 level). Small-world networks (**C**) feature the functional organization of TD brain networks, and
736 incorporate characteristics of both random and regular networks, achieving an optimal balance
737 between global and local characteristics. SW networks (**C**), are at large associated with the broad
738 abilities component of general intelligence (higher intelligence level). The pre-intervention DS
739 network (**B**) showed a random-like, simplified architecture, with impaired segregation and
740 integration, as evidenced by the low CC and LE (random network characteristics) and decreased
741 GE and increased CPL, respectively. This is in line with previous literature^{12,36-38}. The DS pre-
742 network's (**B**) integration-related characteristics (lower GE, higher CPL), are not common in
743 random networks. Hence, the DS pre-network (**B**) is classified as random-like, and not entirely
744 random, maintaining an equal distance from random (**A**) and SW (**C**) networks. The DS post-
745 network (**B** → **C**), exhibits an increase in integration (random network characteristic), as well
746 as segregation (regular network characteristic), so it is interpreted as a step towards an SW-like
747 architecture that highlights a healthier brain organization.


748 **Figure 4.** DS-LLM Care design and flow of participants with DS.

749 **Figure 5.** EEG data analysis schematic. **Pre-processing** (blue): EEG data were interpolated and
750 artifact corrected, visually inspected, high-pass, bandpass, and low-pass filtered. Independent
751 component analysis (ICA), as well as visual inspection, were used to reject artifactual data.
752 Following, 15 segments of 4-s were randomly selected. **Source reconstruction** (orange): The
753 data were processed within 0.53-35 Hz frequency range, source reconstructed (4-D LORETA, for
754 all time points), and a previously used 863-node atlas⁶⁹ was applied to extract the time-series of
755 every voxel from every segment per subject. **Functional connectivity** (green): Functional
756 connectivity was computed for every segment of every subject, using the phase transfer entropy
757 metric (PTE), and the 15 matrices of every subject were averaged into one. A network science
758 approach was taken for the computation of graph measures per subject. Group average statistics
759 were calculated to identify the statistically significant differences of post- and pre-intervention
760 networks.

A**POST vs. PRE****B****PRE****POST****C****POST vs. PRE CIRCULAR**



A
RANDOM NETWORK




over-integration
under-segregation
increased wiring cost

small CPL
high GE
low CC
low LE

GENERAL ABILITIES


B
RANDOM-LIKE NETWORK
PRE-INTERVENTION
DS NETWORK



impaired integration
under-segregation
increased wiring cost

greater CPL
smaller GE
low CC
low LE

B → **C**
TRANSITIONAL STATE
POST-INTERVENTION
DS NETWORK



increased integration
increased segregation
reduced wiring cost

decreased CPL
increased GE
increased CC
increased LE

C
SMALL-WORLD NETWORK



balanced integration
balanced segregation
optimal wiring cost

small CPL
high GE
high CC
high LE

BROAD ABILITIES

Subjects

12 DS individuals
age: 29 ± 11 , 6 females
10 week protocol

Pre-measurements

Psychometric and somatometric assessment tests
5-minute resting state EEG measurement

Intervention details

2 CT sessions/week, 30 minutes each
audiovisual stimuli, memory, attention,
cognitive processing speed and orientation
average 20 ± 3 sessions

2 PT sessions/week, 30 minutes each
aerobic, flexibility, strength and balance
average 18 ± 4 sessions


Post-measurements

Psychometric and somatometric assessment tests
5-minute resting state EEG measurement


Pre-processing

5-min RS EEG recording

128 channels



Interpolations & Artefact corrections




Filtering

ICA

Visual Inspection

15 random
4s epochs

Source Reconstruction



4-D LORETA calculation for every segment (0.53-35Hz)



Functional Connectivity

Identification of statistically significant network differences




Group average statistics

PTE adjacency matrices for each segment of every subject

Voxel Time-series for each segment/subject



averaged
PTE per
subject



Graph measures computation