COVID-19 higher morbidity and mortality in Chinese regions with lower air quality

Riccardo Pansini[1,2,3] & Davide Fornacca[1,4]

[1] Institute of Eastern-Himalaya Biodiversity Research, Dali University, Yunnan, China
[2] Behavioral and Experimental Economics Research Center, Statistic and Mathematics College, Yunnan University of Finance and Economics, Kunming, China
[3] Department of Economics and Finance, Global Research Unit, City University of Hong Kong
[4] Institute for Environmental Sciences, University of Geneva, Switzerland

r.pansini@gmail.com   fornacca@eastern-himalaya.cn

Highlights
• By correlating levels of air pollution with COVID-19 spread and mortality controlled for varying population densities, we found a significant correlation in China.
• We infer that living in an area with low air quality seems to be a risk factor for becoming infected and dying from this new form of coronavirus.

Abstract - We investigated the geographical character of the COVID-19 infection in China and correlated it with satellite- and ground-based measurements of air quality. Controlling for population size, we found more viral infections in those areas afflicted by high Carbon Monoxide, formaldehyde, PM 2.5, and Nitrogen Dioxide values. Higher mortality was also correlated with relatively poor air quality. Air pollution appears to be a risk factor for the incidence of this disease, similar to smoking. This suggests the detrimental impact of air pollution in these types of respiratory epidemics.

Keywords: air pollution; SARS-CoV-2; risk factors; virulence; climate change.

COVID-19, initially detected in China and rapidly spread to the rest of the world, has ignited a pandemic at an exorbitant human cost. In a span of just over three months, eastern and western doctors, biologists, and sociologists alike have turned their attention at disentangling the aetiology of this disease. Various risk factors have been implicated with the fast spread of the virus, assuming different characters whether considered within or between countries. On an individual level, older age, male gender and smoking status have all been shown to increase the virulence of SARS-CoV-2[1].

From the standpoint of the natural sciences and evolution, we can take a broader perspective beyond virological and medical mechanisms to appreciate how a coronavirus transmitted once more from an animal species to us. Elements including human and livestock overpopulation, biodiversity loss and climate change played a critical role in making the ground suitable for a new epidemics to flourish[2,3]. Zoonoses should not be treated so that wildlife is blamed, the actual problem is the human exploitation of wildlife[4,5].

Pertaining to climate change, air pollution is notoriously known to cause health problems and, in particular, viral respiratory infections and pneumonia to individuals exposed for several days a year[6,7].

From our perspective as biologists, economists and geographers, we have investigated the expansion of the infection in China (with Hong Kong and Taiwan included – data from the Chinese government health commission[8]) and we have correlated it with the annual tropospheric column measurements of several air pollutants sampled from the Sentinel-5 satellite time series (data from the European Space Agency portal[9]) as well as data derived from traditional air quality stations[10]. The data were updated on 23 May 2020, well after the first, main infection wave and they include the major 17 April update (at the time, an increase of 1,290 casualties, about 50% from the previous figure, and an increase of 325 infections for the city of Wuhan only). See Table 1 for a comprehensive summary of the data.

Controlling for varying population sizes[11], we find more viral infections and fatalities in those Chinese prefectures afflicted by common pollutants of the air: CO, HCHO, PM 2.5, PM 10, and NO_2. All correlation tests were performed using Kendall's Tau with the significance threshold set to 0.05. Aerosol data from the satellite, which include PM 2.5 and PM 10, were not associated with a higher mortality. This is not surprising, they in fact comprise non-pollutant, inert particulates such as dust, sand and sea salt. On the other hand, statistics suggested that higher levels of O_3 and SO_2 were not associated to more COVID-19 deaths, which goes against the trends of the other pollutants, an aspect that requires further investigation. A comprehensive statistical output is reported in Table 2.

Despite the fact the SARS-CoV-2 was first detected in Wuhan and that the first location of the pathogen assumes a key role in the geographical spread of the infection, the detrimental effect that air pollution assumes at a national level remains evident. This trend in fact holds after removing (1) Wuhan city and (2) the whole Hubei province from the dataset in succession (see Table 2S.I. and £S.I.). Although the measures to contain the virus taken by the Chinese government were effective when compared to other countries affected by what later became a pandemic, the strong association between air pollution and higher morbidity and mortality rates can be detected even where the series of outbreaks originated from.

Although free health care was dispensed to everyone in the exceptional case of this epidemics, the Chinese health system, as those ones of many other countries, is not adequate at including risk factors as comorbidity elements to prevent diseases and allow for free health care accordingly[12]. We hypothesise that the correlational significance we have found points towards air pollution as a critical risk cofactor for COVID-19, also because of other studies. (1) Testing the more proximal hypothesis that COVID-19 outbreaks could follow with a temporal delay from days with high NO_2 presence in the air, colleagues in Shanghai are publishing detailed time series suggesting a delay of 12 days before hospitalisations for the Hubei province[13]. This may even suggest the role of air pollutants as airborne vectors for this virus[14], later also suggested by another study[15] run in three cities in Hubei province. At a Chinese national level, PM 2.5 and NO_2 pollution from ground stations was recently found correlated with this pathology incidence, after adjusting for some socio-economic factors and human movements following the Spring festival[16]. (2) In the United States, chronic exposure to high levels of PM 2.5 have been found co-responsible for 20-time higher mortality rate of respiratory diseases, specifically in 2020, that is in presence of COVID-19[17]. This is a rate much higher than other American demographic co-variables. (3) In Italy, a similar positive correlation controlling for further demographic variables was also reported[18], with the additional and novel evidence that fragments of the RNA from this virus were found in the particulate matter of the harshly hit Italian city of Bergamo[14]. (4) Finally, our comprehensive study including eight countries assessed with a similar analysis, confirms this same trend in six out of eight countries[19] (we were in fact the very first to release an analysis comparing chronic exposure to air pollution and a higher morbidity and mortality from SARS-CoV-2 in early April[20]).

As a clear and immediate action to prevent the trajectory of this and future epidemics, curbing climate change[21,22] must be endorsed way more seriously. With no exceptions, it must be endorsed now. Will the smallest of the parasites be able to wake us up this time, so that we start caring about the health of the environment as much as we have been caring about our own and public health?

Data	Measuring Unit	Time period	Format	Source
COVID-19	No. of infections, No. of deaths	Updated on 23 May 2020	Tabular Prefecture level	DXY - DX Doctor: http://ncov.dxy.cn/ncovh5/view/en_pneumonia from Chinese government health commission
Population	No. of residents	Estimates 2017	Tabular Prefecture level	https://www.citypopulation.de/ Data from provincial governments
Air quality, ground stations:				
PM2.5, PM10, O_3,	AQI	2014	Tabular	University of Harvard Dataverse:

NO₂, SO₂, CO			GPS points	https://dataverse.harvard.edu
				Data from http://aqicn.org
Air quality, satellite:				
UV Aerosol Index CO, HCHO, NO₂, O₃, SO₂	Qualitative Index mol/m^2	2019	Continuous grid (0.01 arc deg.)	Sentinel-5 Atmospheric variables https://developers.google.com/earth-engine/datasets/tags/air-quality

Table 1. The datasets information with their sources.

Figure 1 Distribution of COVID-19 infections, fatalities and mortality rates across the prefectures of China (updated on 23 May 2020), and the distribution of the tropospheric column amounts of three representative air pollutants derived from satellite (2019 averages): Nitrogen Dioxide (NO₂), Carbon Monoxide (CO), and Formaldehyde (HCHO). The values in the square brackets show the COVID-19 cases' counts of administrative units.

China	Infections (/100k pop)			Fatalities (/100k pop)			Mortality (fatalities/infections)		
	df (N-2)	tau	*p*-value	df (N-2)	tau	*p*-value	df (N-2)	tau	*p*-value
CO sat	337	.28	**<.001**	337	.19	**<.001**	313	.16	**<.001**
NO2 sat	337	.23	**<.001**	337	.14	**.001**	313	.12	**.006**
O3 sat	337	-.08	**.030**	337	.00	.967	313	.02	.635
SO2 sat	337	-.10	**.005**	337	-.02	.634	313	.00	.964
Aerosol sat	337	-.12	**.001**	337	-.03	.488	313	.00	.950
HCHO sat	337	.34	**<.001**	337	.20	**<.001**	313	.17	**<.001**
PM25 ground	302	.15	**<.001**	302	.18	**<.001**	285	.18	**<.001**
PM10 ground	302	.04	.330	302	.12	**.006**	285	.13	**.005**
CO ground	302	-.01	.840	302	.11	**.012**	285	.12	**.007**
NO2 ground	302	.12	**.002**	302	.12	**.005**	285	.12	**.007**
O3 ground	302	-.03	.477	302	-.02	.585	285	-.03	.482
SO2 ground	302	-.01	.843	302	.04	.409	285	.06	.178
population	337	.23	**<.001**	337	.17	**<.001**			
pop density	337	.32	**<.001**	337	.16	**<.001**			

Table 2. Correlation between satellite- and ground-based air quality variables and cumulated COVID-19 infections per 100,000 inhabitants, fatalities per 100,000 inhabitants, and mortality rate in China, until 23 May 2020.

Acknowledgements

No funding was obtained for the completion of this study. Livia Ottisova improved and revised the manuscript. Chun Chen commented on statistics. RP did this work while in isolation in the Italian Po valley, due to the ongoing pandemic.

References

Vardavas, C. I. & Nikitara, K. 2020. COVID-19 and smoking: A systematic review of the evidence. *Tobacco Induced Diseases*, 18, 10.18332/tid/119324.

Bedford, J., Farrar, J., Ihekweazu, C., Kang, G., Koopmans, M. & Nkengasong, J. 2019. A new twenty-first century science for effective epidemic response. *Nature*, 575, 130-136, 10.1038/s41586-019-1717-y.

Keesing, F., Belden, L. K., Daszak, P., Dobson, A., Harvell, C. D., Holt, R. D., Hudson, P., Jolles, A., Jones, K. E., Mitchell, C. E., Myers, S. S., Bogich, T. & Ostfeld, R. S. 2010. Impacts of biodiversity on the emergence and transmission of infectious diseases. *Nature*, 468, 647-652, 10.1038/nature09575.

Editorial. 2020. Prevent and predict. *Nature Ecology & Evolution*, 4, 283-283, 10.1038/s41559-020-1150-5.

Morand, S., Owers, K. & Bordes, F. 2014. Biodiversity and emerging zoonoses. In: *Confronting Emerging Zoonoses*, pp. 27-41: Springer, 10.1007/978-4-431-55120-1_3.

Krewski, D. 2009. Evaluating the effects of ambient air pollution on life expectancy. *N Engl J Med*, 360, 413-415, 10.1056/NEJMe0809178.

Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D. & Pozzer, A. 2015. The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature*, 525, 367-371.

Doctor, D.-D. 2020. COVID-19 Global pandemic real-time report. Access date: 25/05/2020, http://ncov.dxy.cn/ncovh5/view/en_pneumonia.

ESA. 2018. European Space Agency - Sentinel-5. Access date: 25/03/2020, https://developers.google.com/earth-engine/datasets/tags/air-quality.

World Air Quality Index. Access date: 25/03/2020, https://aqicn.org/.

Brinkhoff, T. 2020. City Population. Access date: 25/04/2020, https://www.citypopulation.de/en/china/admin/.

Zhai, T. & Goss, J. 2020. Health system reform in China: the challenges of multimorbidity. *The Lancet Global Health*, 8, e750-e751, 10.1016/S2214-109X(20)30225-4.

Yao, Y., Pan, J., Liu, Z., Meng, X., Wang, W., Kan, H. & Wang, W. 2020. Ambient nitrogen dioxide pollution and spread ability of COVID-19 in Chinese cities. *medRxiv*, 10.1101/2020.03.31.20048595.

Setti, L., Passarini, F., Gennaro, G. D., Baribieri, P., Perrone, M. G., Borelli, M., Palmisani, J., Gilio, A. D., Torboli, V., Pallavicini, A., Ruscio, M., Piscitelli, P. & Miani, A. 2020. SARS-Cov-2 RNA Found on Particulate Matter of Bergamo in Northern Italy: First Preliminary Evidence. *medRxiv*, 10.1101/2020.04.15.20065995.

Jiang, Y., Wu, X.-J. & Guan, Y.-J. 2020. Effect of ambient air pollutants and meteorological variables on COVID-19 incidence. *Infection Control & Hospital Epidemiology*, 1-11, 10.1017/ice.2020.222.

Tian, H., Liu, Y., Song, H., Wu, C.-H., Li, B., Kraemer, M. U. G., Zheng, P., Yan, X., Jia, G., Zheng, Y., Stenseth, N. C. & Dye, C. 2020. Risk of COVID-19 is associated with long-term exposure to air pollution. *medRxiv*, 10.1101/2020.04.21.20073700.

Wu, X., Nethery, R. C., Sabath, B. M., Braun, D. & Dominici, F. 2020. Exposure to air pollution and COVID-19 mortality in the United States. *medRxiv*, 10.1101/2020.04.05.20054502.

Pluchino, A., Inturri, G., Rapisarda, A., Biondo, A. E., Le Moli, R., Zappala', C., Giuffrida, N., Russo, G. & Latora, V. 2020. A Novel Methodology for Epidemic Risk Assessment: the case of COVID-19 outbreak in Italy. In: *arXiv e-prints*, pp. 2004.02739

Pansini, R. & Fornacca, D. 2020. Higher virulence of COVID-19 in the air-polluted regions of eight severely affected countries. *medRxiv*, 10.1101/2020.04.30.20086496.

Pansini, R. & Fornacca, D. 2020. Initial evidence of higher morbidity and mortality due to SARS-CoV-2 in regions with lower air quality. *medRxiv*, 10.1101/2020.04.04.20053595.

Haines, A. & Ebi, K. 2019. The Imperative for Climate Action to Protect Health. *N Engl J Med*, 380, 263-273, 10.1056/NEJMra1807873.

Haines, A., Scheelbeek, P. & Abbasi, K. 2020. The health case for urgent action on climate change. *British Medical Journal*, 368, m1103, 10.1136/bmj.m1103.

Author Contributions

Concept and design: RP.
Acquisition and statistical analysis and of data: DF.
Interpretation: DF and RP.
Drafting of the manuscript: RP.
Critical revision of the manuscript: RP and DF.

Data Availability

The data are available in Github upon acceptance in a journal.

Competing Interests

The authors declare no competing interests.

Digital supplement

	unit	count	mean	std	min	25%	50%	75%	max	range	iqr
CO sat	µmol/m2	347	40721	8751	15826	36202	42348	47034	60605	44778	10833
NO2 sat	µmol/m2	347	57.58	42.62	5.00	26.00	42.70	83.85	185.39	180.42	57.85
O3 sat	µmol/m2	347	130602	14073	115053	118774	126147	138924	170378	55325	20150
SO2 sat	µmol/m2	347	58.61	65.11	-60.00	10.43	42.10	102.70	316.53	376.43	92.27
Aerosol sat	index	347	-0.97	0.17	-1.30	-1.06	-1.00	-0.92	-0.02	1.28	0.14
HCHO sat	µmol/m2	347	142.04	42.65	50.00	107.63	150.06	178.36	219.85	169.55	70.74
PM25 ground	AQI	308	110.78	27.79	38.20	92.63	111.55	128.51	186.96	148.76	35.88
PM10 ground	AQI	308	63.61	22.98	19.96	46.26	60.90	75.67	170.27	150.31	29.40
CO ground	AQI	308	9.71	3.97	2.39	7.11	8.74	11.85	27.01	24.63	4.74
NO2 ground	AQI	308	13.72	5.40	3.17	9.63	13.40	17.41	28.57	25.40	7.78
O3 ground	AQI	308	25.83	5.87	14.22	21.70	25.17	28.98	50.54	36.31	7.28
SO2 ground	AQI	308	13.24	7.96	1.14	7.88	10.97	16.46	40.58	39.44	8.58

Table 1 S.I. Descriptive statistics for the different satellite- and ground-based air quality measurements.

	Population density		
	df (N-2)	tau	p-value
CO sat	345	.39	<.001
NO2 sat	345	.37	<.001
O3 sat	345	.36	<.001
SO2 sat	345	.36	<.001
Aerosol sat	345	-.07	.039
HCHO sat	345	.00	.890
PM25 ground	345	-.07	.046
PM10 ground	345	.40	<.001
CO ground	306	.24	<.001
NO2 ground	306	.13	.001
O3 ground	306	.04	.327
SO2 ground	306	.30	<.001

Table 2 S.I. Correlation between satellite- and ground-based air quality variables and population density in China.

China without Wuhan	Infections (/100k pop)			Fatalities (/100k pop)			Mortality (fatalities/infections)		
	df (N-2)	tau	p-value	df (N-2)	tau	p-value	df (N-2)	tau	p-value
CO sat	336	.28	<.001	336	.18	<.001	312	.16	<.001
NO2 sat	336	.22	<.001	336	.14	.001	312	.12	.008
O3 sat	336	-.08	.031	336	.00	.943	312	.02	.617
SO2 sat	336	-.10	.006	336	-.02	.690	312	.00	.975
Aerosol sat	336	-.12	.001	336	-.03	.537	312	.00	.992
HCHO sat	336	.34	<.001	336	.19	<.001	312	.16	<.001
PM25 ground	301	.15	<.001	301	.18	<.001	284	.17	<.001
PM10 ground	301	.04	.361	301	.12	.008	284	.13	.006
CO ground	301	-.01	.798	301	.11	.013	284	.12	.008
NO2 ground	301	.11	.003	301	.12	.009	284	.11	.012
O3 ground	301	-.03	.512	301	-.02	.637	284	-.03	.528
SO2 ground	301	-.01	.851	301	.04	.396	284	.06	.171
population	336	.22	<.001	336	.16	<.001			
pop density	336	.32	<.001	336	.15	<.001			

Table 3 S.I. Correlation between satellite- and ground-based air quality variables and cumulated COVID-19 infections per 100,000 inhabitants, fatalities per 100,000 inhabitants, and mortality rate in China (without the administrative unit of Wuhan), until 23 May 2020.

China without Hubei prov.	Infections (/100k pop)			Fatalities (/100k pop)			Mortality (fatalities/infections)		
	df (N-2)	tau	*p*-value	df (N-2)	tau	*p*-value	df (N-2)	tau	*p*-value
CO sat	320	.26	**<.001**	320	.13	**.003**	296	.11	**.013**
NO2 sat	320	.26	**<.001**	320	.17	**<.001**	296	.15	**.001**
O3 sat	320	.26	**<.001**	320	.13	**.002**	296	.11	**.011**
SO2 sat	320	.24	**<.001**	320	.15	**.001**	296	.13	**.005**
Aerosol sat	320	-.07	.054	320	.03	.551	296	.04	.417
HCHO sat	320	-.08	**.044**	320	.04	.386	296	.05	.275
PM25 ground	320	-.10	**.008**	320	.03	.539	296	.05	.299
PM10 ground	320	.32	**<.001**	320	.14	**.002**	296	.11	**.016**
CO ground	286	.10	**.011**	286	.11	**.018**	269	.11	**.022**
NO2 ground	286	.00	.992	286	.07	.108	269	.08	.087
O3 ground	286	-.03	.395	286	.09	.052	269	.10	**.032**
SO2 ground	286	.13	**.001**	286	.15	**.002**	269	.14	**.003**
population	320	.26	**<.001**	320	.21	**<.001**			
pop density	320	.35	**<.001**	320	.19	**<.001**			

Table 4 S.I. Correlation between satellite- and ground-based air quality variables and cumulated COVID-19 infections per 100,000 inhabitants, fatalities per 100,000 inhabitants, and mortality rate in China (without Hubei province), until 23 May 2020.