Determinants of self-reported symptoms and testing for COVID in Canada using a nationally representative survey

- 3 Daphne C Wu MSc (1), Prabhat Jha DPhil (1), Teresa Lam BA (2), Patrick Brown PhD (1, 3),
- 4 Hellen Gelband MHS (1), Nico Nagelkerke PhD (1), H. Chaim Birnboim MD (1, 4), Angus Reid
- 5 PhD (2) on behalf of the Action to Beat Coronavirus in Canada/Action pour Battre le
- 6 Coronavirus (Ab-C) Study Group
- 7 1. Centre for Global Health Research, St. Michael's Hospital, Unity Health Toronto, and
- 8 Dalla Lana School of Public Health, University of Toronto
- 9 2. Angus Reid Institute
- 10 3. Department of Statistical Sciences, University of Toronto
- 11 4. deltaDNA Biosciences, Inc., Toronto
- 12
- 13 Prabhat.jha@utoronto.ca
- 14
- 15
- 16 Abstract: 149 words
- 17 Text: 2160 words, 12 references, 3 tables

18 Abstract

- 19 In April 2020, the first-ever nationally representative survey in Canada polled 4,240 adults age
- 20 18 years and older about their COVID experience in March, early in the epidemic. We examined
- 21 determinants of COVID symptoms, defined as fever plus difficulty breathing/shortness of
- 22 breath, dry cough so severe that it disrupts sleep, and/or loss of sense of smell; and testing for
- 23 SARS-CoV-2 by respondents and/or household members. About 8% of Canadians reported that
- 24 they and/or one or more household members experienced COVID symptoms. Symptoms were
- 25 more common in younger than older adults, and among visible minorities. Overall, only 3% of
- respondents and/or household members reported testing for SARS-CoV-2. Being tested was
- associated with having COVID symptoms, Indigenous identity, and living in Quebec. Periodic
- 28 nationally representative surveys—including high-risk older populations—of symptoms, as well
- as SARS-CoV-2 antibodies, are required in many countries to understand the pandemic and
- 30 prepare for the future.

31 Introduction

- 32 The pandemic of SARS-CoV-2 infection causing coronavirus disease-2019 (COVID-19, hereafter
- 33 "COVID") has affected Canada and many other high-income countries (Eggertson and Wolfville,
- 34 2020). In these settings where reliable data can be gathered, a combination of population-
- 35 based surveys (including surveys and testing), hospitalizations, and mortality data can produce
- 36 an accurate profile of the impact of the epidemic.
- 37 We report on the results of the first nationally-representative poll of self-reported COVID
- 38 symptoms conducted in Canada by the Angus Reid Forum in early April 2020 covering
- 39 symptoms reported mostly in March 2020, prior to the peak month of cases in April. Our goal
- 40 was to understand the distribution and determinants of Canadians reporting possible COVID
- 41 symptoms. We also sought to understand who underwent testing for SARS-CoV-2 using the
- 42 current standard (PCR-based) test. We discuss these findings in the context of the age
- 43 distribution of COVID hospitalizations and deaths, and a planned survey of antibodies to SARS-
- 44 CoV-2 in a random sample of Canadians.

45 Results

- 46 To determine the representativeness of the respondents, we compared their socio-
- 47 demographic characteristics to those of the Canadian population in 2019 (Table 1). Overall, the
- 48 survey respondents were broadly representative of Canadian society in terms of gender, age,
- 49 regional distribution, and numbers of household members. Survey respondents were less
- 50 representative of Canada in terms of ethnicities other than Indigenous. The survey had fewer
- 51 single-member households than in the Canadian census, and had slightly higher education
- 52 levels than did the 2019 Canadian population. Approximately 0.3 million (out a total population
- of 38 million) Canadians are estimated to live in nursing homes/long-term care institutions
- 54 (Canadian Life and Health Insurance Association, 2014). This would represent less than 1% of
- the expected population in the survey (or about 40 participants). However, such populations
- 56 were under-represented in the survey.

Characteristics	Completed survey, sample size		Canadian population, 2019 (%) (Statistics Canada, 2016; Statistics Canada, 2020a)	
	n	%		
Gender				
Male	2017	47.9	49.4	
Female	2205	52.0	50.6	
Other	18	0.4		
Age group				
18-45 years	2007	47.3	46.4	
46-65 years	1511	35.6	33.4	
66+ years	722	17.0	20.2	
Education				
High school and under	1064	25.1	35.2	
Some college/university and higher	3176	74.9	64.8	
Visible minority				
No	3686	86.9	71.4	
Yes	554	13.1	28.7	
Number of household members				
Lived alone	672	15.9	28.2	
Two people	1615	38.1	34.4	
Three people	804	19.0	15.2	
Four or more people	1149	27.1	22.2	
Ethnicity				
Indigenous ⁺	226	5.3	5.0	
English and other European	3567	84.1	73.1	
Others	365	8.6	22.0	
Rather Not Say	82	1.9		
Province/region				
Ontario	1600	37.7	38.9	
British Columbia	554	13.1	13.8	
Quebec	1020	24.1	22.7	
Alberta	475	11.2	11.2	
Manitoba	150	3.5	3.5	
Saskatchewan	133	3.1	3.0	
Atlantic provinces	308	7.3	6.6	

Table 1. Socio-demographic characteristics of respondents (N=4,240) as compared to the Canadian population in 2019.

*All comparisons are from the 2016 Census, except Province/region which is from 2019 Statistics Canada population data (Statistics Canada, 2020a).

⁺Indigenous populations including First Nations, Inuit, or Métis.

- 57 Of the 4,240 respondents, 334 (7.9% after applying survey weights) reported COVID symptoms,
- 58 defined as the respondent reporting himself/herself and/or at least one of the household
- 59 members experiencing a combination of fever (with or without hallucinations) and any of i)
- 60 difficulty breathing/shortness of breath <u>or</u> ii) dry cough so severe that it disrupts sleep <u>or</u> iii) a
- loss of a sense of smell. Of these, 210 (5.0%) reported COVID symptoms only in themselves. The
- 62 adjusted odds ratio (OR) of the respondent having symptoms when at least one of the
- household members reported symptoms was 1.45 (95% CI 1.41-1.49); the OR was similar for at
- 64 least one household member having symptoms if the respondent reported symptoms. In terms
- of testing for SARS-CoV-2, 126 (or 3.0%) reported some household testing or being scheduled
- 66 for testing, and only 68 (or 1.6%) reported that they have been or are scheduled for testing.
- 67 Details of the variation in COVID symptoms and SARS-CoV-2 testing in this sample across
- 68 provinces have already been published (Angus Reid Institute, 2020).
- Table 2 shows the OR of respondents or a member of the household, and respondents only
- 70 having COVID symptoms after adjustment for other variables examined. We excluded 99 (or
- 71 2.3%) respondents who did not report on at least one of the variables. The proportion of
- respondents reporting COVID symptoms within the household decreased with age: 11.2% of
- those aged 18-45 years, 5.6% among those aged 46-65 years, and 3.1% among those aged 66
- 74 years and older. The lower prevalence at higher ages was similar among those reporting COVID
- 75 symptoms only themselves. After controlling for gender, province, age, ethnicity, visible
- 76 minority, and number of household members, older adults were significantly less likely to
- report having COVID symptoms themselves or within the household (age 46-65, OR=0.55, 0.42-
- 0.71; age 66+, OR=0.30, 0.18-0.47). Those who reported themselves as a visible minority were
- real significantly more likely to have COVID symptoms (12.9%) compared to those who did not
- 80 (7.1%; adjusted OR=1.55; 1.08-2.20). Similar results were found when we examined COVID
- 81 symptoms only among respondents. The associations changed substantially between
- 82 unadjusted and adjusted analyses, most notably for ethnicity, and somewhat for being a visible
- 83 minority, suggesting that some residual confounding factors were present. Results using a
- 84 narrower definition of COVID symptoms, namely having fever, difficulty breathing/shortness of
- 85 breath, and severe dry cough were similar (data not shown).
- 86 Table 3 shows the adjusted odds ratios of respondents or a member of the household being
- 87 tested or scheduled for SARS-CoV-2 testing. The strongest determinant of testing was having
- 88 COVID symptoms among members of the household, of whom about 16.5% were tested,
- 89 compared to 2.1% among those without COVID symptoms among household members
- 90 (OR=6.63, 4.46-9.79). Testing rates fell with age, but not significantly so. Testing rates were
- 2.7% in English and other European ethnicity, and higher in indigenous people (7.2%; OR=2.07;
- 1.13-3.64). However, the ORs fell notably (from 2.94 to 2.07) after adjustment for co-variates,
- 93 suggesting residual confounding (data not shown). Testing rates in Quebec were roughly double

- 94 those in Ontario, the reference province (OR=2.41; 1.49-3.96). Findings were similar among
- 95 those who reported having been tested or being scheduled to be tested (data not shown).

Characteristics	Respondents or a member of			Respondents only having symptoms			
	household having symptoms						
	COVID sympt	oms	Adjusted odds	COVID symptoms		Adjusted odds	
	positive/nega	ative	ratios (95% CI)	positive/neg	gative	ratios (95% CI)	
	N=324/3817	%		N=202/393	%		
				9			
Gender							
Male	149/1840	7.5	Ref	85/1904	4.3	Ref	
Female	175/1977	8.1	1.11 (0.89-1.40)	117/2036	5.4	1.30 (0.97-1.74)	
Age group							
18-45 years	218/1724	11.2	Ref	126/1816	6.5	Ref	
46-65 years	84/1403	5.6	0.55 (0.42-0.71)	61/1426	4.1	0.61 (0.44-0.84)	
66+ years	22/690	3.1	0.30 (0.18-0.47)	15/697	2.1	0.29 (0.16-0.50)	
Education							
Some college/university	237/2874	8.4	Ref	150/2961	4.8	Ref	
and higher							
High school and under	87/943	7.6	1.25 (0.96-1.62)	52/978	5.1	1.16 (0.83-1.61)	
Visible minority							
No	256/3359	7.1	Ref	155/3460	4.3	Ref	
Yes	68/458	12.9	1.55 (1.08-2.20)	47/479	8.9	2.08 (1.34-3.15)	
Number of household							
members							
Two people	108/1479	6.8	Ref	79/1507	5.0	Ref	
Lived alone	29/626	4.4	0.69 (0.45-1.02)	29/626	4.5	1.01 (0.64-1.53)	
Three people	77/704	9.9	1.17 (0.85-1.61)	45/736	5.8	0.86 (0.57-1.28)	
Four or more people	110/1008	9.8	1.19 (0.89-1.59)	48/1070	4.3	0.72 (0.49-1.05)	
Ethnicity							
English and other	252/3305	7.1	Ref	158/3399	4.4	Ref	
European							
Indigenous*	27/195	12.2	1.31 (0.83-2.01)	19/203	8.5	1.34 (0.76-2.24)	
Others	45/317	12.4	1.20 (0.79-1.80)	25/337	7.0	0.87 (0.50-1.47)	
Province/region							
Ontario	129/1448	8.2	Ref	83/1493	5.3	Ref	
British Columbia	44/495	8.2	1.08 (0.77-1.51)	23/516	4.2	0.80 (0.51-1.24)	
Quebec	65/922	6.6	0.93 (0.66-1.29)	42/945	4.2	0.85 (0.56-1.27)	
Prairie provinces	63/674	8.5	1.09 (0.79-1.49)	39/697	5.2	0.98 (0.66-1.45)	
Atlantic provinces	23/278	7.6	1.11 (0.68-1.76)	15/287	5.0	0.99 (0.53-1.74)	

Table 2. Logistic regression models for respondents or a member of household, and respondents only having COVID symptoms.

*Indigenous populations including First Nations, Inuit, or Métis.

Bolded values indicate significance at 95% confidence interval. Adjustment was for the other variables in the table. The number reporting COVID symptoms was 341 in unweighted analyses and 334 after survey weights were applied, showing the sampling frame as robust.

Table 3. Logistic regression models for respondent or a member of household having been tested for SARS-CoV-2.

Characteristics	Respondents or a member of			Respondents only having been tested			
	household having been tested		en tested				
	SARS-CoV-2		Adjusted odds	SARS-CoV-2		Adjusted odds	
	tested/not tested		ratios (95% CI)	tested/not tested		ratios (95% CI)	
	N=126/4015	%		N=66/4075	%		
Respondent and/or household member had COVID symptoms							
No	80/3737	2.1	Ref	42/3775	1.1	Ref	
Yes	46/278	16.5	6.63 (4.46-9.79)	24/300	7.5	6.63 (3.82-11.32)	
Gender							
Male	68/1921	3.5	Ref	34/1955	1.7	Ref	
Female	58/2094	2.7	0.72 (0.51-1.03)	32/2120	1.5	0.85 (0.52-1.38)	
Age group							
18-45 years	84/1858	4.5	Ref	41/1900	2.1	Ref	
46-65 years	32/1455	2.2	0.69 (0.45-1.05)	19/1468	1.3	0.75 (0.42-1.31)	
66+ years	10/702	0.7	0.54 (0.25-1.05)	5/707	0.7	0.44 (0.14-1.08)	
Education							
Some college/university	96/3015	3.1	Ref	52/3059	1.7	Ref	
and higher							
High school and under	29/1000	2.8	0.88 (0.56-1.35)	14/1016	1.3	0.74 (0.37-1.35)	
Visible minority							
No	96/3519	2.7	Ref	52/3563	1.4	Ref	
Yes	30/496	5.7	0.83 (0.36-1.05)	14/512	2.7	0.54 (0.27-1.15)	
Number of household							
members							
Two people	38/1548	2.4	Ref	19/1567	1.2	Ref	
Lived alone	13/642	2.0	0.83 (0.43-1.52)	13/642	2.0	1.77 (0.86-3.57)	
Three people	28/754	3.6	1.18 (0.69-1.97)	18/764	2.3	1.28 (0.62-2.56)	
Four or more people	47/1071	4.2	1.32 (0.84-2.09)	16/1102	1.4	0.93 (0.47-1.84)	
Ethnicity							
English and other	95/3462	2.7	Ref	54/3503	1.5	Ref	
European							
Indigenous*	16/206	7.2	2.07 (1.13-3.64)	9/213	3.9	2.04 (0.89-4.28)	
Others	15/347	4.1	0.99 (0.49-1.89)	4/359	1.0	0.47 (0.13-1.32)	
Province/region							
Ontario	40/1537	2.5	Ref	22/1554	1.4	Ref	
British Columbia	15/524	2.8	1.08 (0.59-1.93)	7/532	1.3	0.81 (0.34-1.81)	
Quebec	47/940	4.8	2.41 (1.49-3.96)	26/961	2.6	1.98 (1.05-3.81)	
Prairie provinces	19/717	2.6	1.13 (0.65-1.95)	7/730	1.0	0.74 (0.32-1.63)	
Atlantic provinces	5/297	1.7	0.92 (0.34-2.14)	4/299	1.2	1.03 (0.29-2.87)	

*Indigenous populations including First Nations, Inuit, or Métis.

Bolded values indicate significance at 95% confidence interval. Adjustment was for the other variables in the table.

96 Discussion

A nationally representative survey of Canadians finds that about 8% of adults report that they 97 98 or someone in their household reported symptoms suggestive of COVID in March 2020. Being a visible minority was associated with higher self-reported COVID symptoms. Self-reported 99 100 symptoms were notably less common at older ages than in younger adults. Only 3% of Canadian adults reported that they or someone in the household had been tested for SARS-101 102 CoV-2. The main determinants of being tested were the presence of COVID symptoms, being of Indigenous identity, and living in Quebec. Testing rates were somewhat lower in older adults. 103 104 There are surprisingly few nationally representative studies, and despite some limitations, this 105 study represents the first to document self-reported symptoms in a reasonably representative 106 sample.

107 COVID symptoms overlap with some other infections, notably seasonal influenza, which could 108 have inflated the coronavirus rates in this survey. However, a study comparing the COVID 109 symptom syndrome with other infections in the United States suggests that most are actually due to COVID (Centers for Disease Control and Prevention, 2020). We found possible COVID 110 111 symptoms to be less prevalent and the levels of testing marginally less prevalent in older adults than other age groups, despite the certainty that the vast majority of COVID hospitalizations 112 and deaths occur at older ages. However, a weakness of our sample is the lack of 113 representation from nursing and long-term care residents, in whom more than three-quarters 114 115 of all COVID deaths occur (Eggertson and Wolfville, 2020). There may be additional reasons for the age-specific findings, however. Anecdotal reports suggest that older adults do not 116 117 experience the symptoms used to define COVID infection in the poll, but may report vaguer symptoms such as dizziness and confusion (Graham, 2020). Further surveys focused on 118 syndromes that might occur in older adults are warranted. The testing results are broadly 119 120 consistent with reports of the general levels of access to SARS-CoV-2 testing during the survey 121 time period, including a higher level of testing in Quebec than in Ontario (Angus Reid Institute, 2020). 122

123 This syndromic survey provides some insights into the prevalence of actual SARS-CoV-2 in 124 Canada. The survey period was mostly in March, whereas models and case reports suggest that 125 peak exposure and incidence of COVID occurred in mid- to late April (Eggertson and Wolfville, 126 2020). Hence, we would expect that a repeated syndromic survey should report a higher 127 prevalence in April. On the other hand, Forum Research and Mainstreet Research reported an 128 estimated prevalence of 5-8% in early-to-mid April in Ontario. However, they used a narrow range of symptoms from those we defined in our study (Forum Research and Mainstreet 129 130 Research, 2020).

In the UK, pilot results from the COVID-19 Infection Survey being carried out by the Office for 131 132 National Statistics found that 0.25% of the community population in England above the age of 133 2, tested positive for the SARS-CoV-2 antigen in early-to-mid May 2020. Testing involved home self-tests of nasal swabs, and excluded those in hospitals, care homes, or other institutional 134 settings (Office for National Statistics, 2020). The prevalence was lower than expected in the 135 136 pilot, and larger studies are planned, along with antibody surveys. A nationally representative 137 sero-epidemiological study is needed to establish the population distribution of cumulative 138 SARS-CoV-2 infection, which would capture both symptomatic and asymptomatic cases. We will 139 soon conduct such a study in Canada called Action to Beat Coronavirus/Action pour Battre le Coronavirus (Ab-C). This study will determine the cumulative prevalence of SARS-CoV-2 140 141 infection during March-May 2020 in Canada through antibody testing, paired with household questionnaire data on COVID experience. A second round of questionnaires and antibody 142 testing four to six months later in the same individuals will provide information on ongoing 143 144 transmission, changes in the immune status of the population, and the durability of the 145 immune response. The Angus Reid Forum is partnering with the Centre for Global Health Research at Unity Health Toronto and the University of Toronto on this study. The expected 146 147 sample frame is 12-18,000 individuals surveyed, of which we expect 10-12,000 to agree to

- answer questions and provide a blood sample. 148
- The present analysis informs the design of antibody surveys. For example, since the strongest 149
- 150 determinant of COVID testing was the presence of symptoms, a similar higher prevalence of
- antibodies might be expected among those reporting symptoms (who may be more likely to 151
- 152 enroll in the survey). Thus, ensuring that the sample size is sufficiently large to examine the
- seroprevalence in those without symptoms is key. Similarly, while the current study did not 153
- 154 examine geographic clustering of COVID symptoms, this might well be the case with actual
- 155 SARS-CoV-2 infection. This also necessitates a large and as diversely representative survey as
- 156 possible. Older adults bear the brunt of COVID hospitalization and deaths. However, this
- 157 population reported a lower prevalence of symptoms than younger adults. Thus, the Ab-C study
- 158 will oversample the population age 60 or older, particularly to understand the possible role of
- asymptomatic infections at older ages. As the Ab-C study is not likely to capture the prevalence 159
- 160 of infection among nursing home residents, ancillary studies are needed to quantify hazards
- among this important group. 161

162 Limitations

- 163 Self-reported symptoms are, by their nature, subject to limitations and misclassification.
- However, the trends over time, even with misclassification, are useful for understanding trends 164
- in the actual underlying prevalence of infection. This is because the "noise" of COVID symptoms 165
- 166 should change little from one survey to the next, and provided there is large "signal" due to the

- 167 COVID pandemic—certainly the case with this virus—reliable estimation of the trends of
- 168 infection are possible. Similar methodological insights have arisen from HIV-1 testing in
- pregnant populations in various countries (Kumar et al., 2006). The biases in enrollment are
- also inherent in any polling, but the Angus Reid polling showed reasonably good consistency
- 171 with the Canadian 2016 census. Moreover, the testing results did not appear to be biased
- 172 greatly, and the more common testing reported in Quebec is consistent with public health
- 173 reports.
- 174 In conclusion, COVID surveillance using nationally representative surveys is essential, and
- ideally needs to be accompanied by antibody determination of infection. Particular attention
- 176 must be paid to symptoms and testing levels in older adults.

177 Materials and Methods

- 178 The Angus Reid Institute (ARI) conducted an online survey from April 1-5, 2020, among a
- 179 representative randomized sample of 4,240 Canadian adults who are members of Angus Reid
- 180 Forum, a national online sample of 50,000 Canadians used for political and other social polling
- 181 (Angus Reid Institute, 2020). A probability sample of this size carries a margin of error of +/- 2
- 182 percentage points, 19 times out of 20. The survey was commissioned and paid for by ARI.
- 183 Questions were related to COVID and socio-demographic characteristics among both the
- 184 respondents and members of their households (<u>http://angusreid.org/covid-epidemiology-</u>
- 185 <u>study/</u>).
- 186 The outcomes were self-reported symptoms suggestive of COVID infection and testing for the
- 187 virus. To understand the socio-demographic determinants of COVID symptoms, we conducted a
- 188 logistic regression analysis, comparing COVID symptoms to the explanatory variables of gender,
- 189 education level, province, age, ethnicity, visible minority (defined as persons, other than
- Aboriginal peoples, who are non-Caucasian in race or non-white in colour) (Statistics Canada,
- 191 2020b), and number of household members. We defined Indigenous as whether the person
- 192 reported identifying with the Aboriginal peoples of Canada which includes First Nations, Métis
- or Inuit and/or those who reported Registered or Treaty Indian status (Statistics Canada, 2019).
- 194 We also used logistic regression analysis to identify determinants of testing, which included the
- above explanatory variables and also COVID symptoms in the respondent or a family member.
- 196 Respondents who did not report on any of the explanatory variables were excluded from these
- analyses. Discrepancies in or between totals are due to rounding. We applied to prevalences
- the survey weights, which are described earlier (Angus Reid Institute, 2020). We used RStudio
- 199 Version 1.1.453 for analyses.
- 200 Ethics approval

- 201 The Angus Reid Forum obtains consent from all participants, and the polling data without any
- 202 individual identifiers are made available openly to bona-fide researchers. Ethics approval was
- 203 not required as per Unity Health Toronto Research Ethics Board.

204

- 205 Funding: Canadian Institutes of Health Research, Angus Reid Institute
- 206 We declare no competing interests

207

208 References

- Angus Reid Institute. 2020. The incidence of COVID-19 infection in Canada? New survey points to over 100,000
 households. <u>http://angusreid.org/covid-epidemiology-study/</u> [Accessed May 10, 2020]
- Canadian Life and Health Insurance Association. 2014. *CLHIA report on long-term care policy: Improving the accessibility, quality and sustainability of long-term care in Canada.*
- https://www.clhia.ca/web/clhia_lp4w_Ind_webstation.nsf/page/3C342451F891CF1D85257A240044F961/\$file/L
 TC Policy Paper 1 EN.pdf [Accessed May 19, 2020]
- 215 Centers for Disease Control and Prevention. 2020 May 8. COVIDView: A weekly surveillance summary of U.S. covid-
- 19 activity. <u>https://www.cdc.gov/coronavirus/2019-ncov/covid-data/covidview/index.html</u> [Accessed May 13, 2020]
- Eggertson L, Wolfville NS. 2020. COVID-19: Recent updates on the coronavirus pandemic. CMAJ News, May 12,
 2020. <u>https://cmajnews.com/2020/05/12/coronavirus-1095847/</u> [Accessed May 13, 2020]
- Forum Research, Mainstreet Research. 2020. COVID-19 Study Ontario- Wave 2. April 22, 2020 for immediate
 release. https://ipolitics.ca/wp-content/uploads/2020/04/Wave-2-Release-FINAL-2-April-22-2020.pdf [Accessed]
- 222 May 12, 2020]
- 223 Graham, J. 2020 Apr 23. Seniors with covid-19 show unusual symptoms, doctors say. CNN.
- https://www.cnn.com/2020/04/23/health/seniors-elderly-coronavirus-symptoms-wellness-partner/index.html
 [Accessed May 13, 2020].
- 226 Kumar R, Jha P, Arora P, Mony P, Bhatia P, Millson P, Dhingra N, Bhattacharya M, Remis RS, Nagelkerke N,
- International studies of HIV/AIDS (ISHA) investigators. 2006. Trends in HIV-1 in young adults in south India from
 2000 to 2004: a prevalence study. The Lancet **367**:1164-1172.
- Office for National Statistics. 2020. Coronavirus (COVID-19) infection survey pilot: England, 21 May 2020.
 https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/bulletins/
- 231 <u>coronaviruscovid19infectionsurveypilot/england21may2020</u> [Accessed May 26, 2020].
- 232 Statistics Canada. 2016. Census of population. [Accessed May 12, 2020].
- 233 Statistics Canada. 2019. Aboriginal identity of person.
- 234 https://www23.statcan.gc.ca/imdb/p3Var.pl?Function=DECI&Id=59224 [Accessed May 14, 2020]
- Statistics Canada. 2020a. Table 17-10-0005-01 Population estimates on July 1st, by age and sex. [Accessed May
 12, 2020]
- 237 Statistics Canada. 2020b. Visible minority of person. Retrieved from
- 238 <u>https://www23.statcan.gc.ca/imdb/p3Var.pl?Function=DEC&Id=45152</u> [Accessed May 14, 2020]