Resting-state network plasticity induced by music therapy after traumatic brain injury

Noelia Martínez-Molina[1,✉], **Sini-Tuuli Siponkoski**[1], **Linda Kuusela**[3,4], **Sari Laitinen**[5], **Milla Holma**[6], **Mirja Ahlfors**[7], **Päivi Jordan-Kilkki**[8], **Katja Ala-Kauhaluoma**[9], **Susanna Melkas**[10], **Johanna Pekkola**[4], **Antoni Rodriguez-Fornells**[11-13], **Matti Laine**[14], **Aarne Ylinen**[10,15], **Pekka Rantanen**[16], **Sanna Koskinen**[2], **Benjamin Ultan Cowley**[17,18], and **Teppo Särkamö**[1]

[1]Department of Psychology and Logopedics, Cognitive Brain Research Unit, Music, Ageing and Rehabilitation Team, University of Helsinki, Helsinki, Finland.
[2]Department of Psychology and Logopedics, University of Helsinki, Helsinki, Finland.
[3]Department of Physics, University of Helsinki, Helsinki, Finland.
[4]HUS Medical Imaging Center, Department of Radiology, Helsinki Central University Hospital and University of Helsinki, Helsinki, Finland.
[5]Espoo Hospital, Espoo, Finland.
[6]Musiikkiterapiaosuuskunta InstruMental (Music Therapy Cooperative InstruMental), Helsinki, Finland.
[7]Private music therapy entrepreneur, Finland.
[8]Dialogic Partner Oy, Espoo, Finland.
[9]Ludus Oy Tutkimus- ja kuntoutuspalvelut (Assessment and Intervention Services), Helsinki, Finland.
[10]Department of Neurology and Brain Injury Outpatient Clinic, Helsinki University Central Hospital, Helsinki, Finland.
[11]Cognition and Brain Plasticity Group, Bellvitge Biomedical Research Institute, L'Hospitalet de Llobregat, Barcelona, Spain.
[12]Department of Cognition, Development and Educational Psychology, University of Barcelona, Barcelona, Spain.
[13]Catalan Institution for Research and Advanced Studies, Barcelona, Spain.
[14]Department of Psychology, Åbo Akademi University, Turku, Finland.
[15]Tampere University Hospital, Tampere, Finland.
[16]Kanta-Häme Central Hospital, Hämeenlinna, Finland.
[17]Department of Education, Faculty of Education, University of Helsinki, Finland.
[18]Cognitive Science, Department of Digital Humanities, Faculty of Arts, University of Helsinki, Finland.

Traumatic brain injury (TBI) is characterized by a complex pattern of abnormalities in resting-state functional connectivity (rsFC), and neuropathology focused on network dysfunction. Here we report a fMRI study of brain network changes induced during a randomised controlled trial of neurological music therapy in 23 moderate/severe TBI patients. Our ROI-to-ROI approach used four networks as sources: the frontoparietal (FPN), dorsal attention (DAN), default mode (DMN), and salience (SAL) networks. These networks include high-degree nodes or network hubs, and have all been associated with cognitive impairment after TBI. Furthermore, we investigated the correlation between brain network changes and executive function (EF). Lastly, we implemented a seed-to-voxel analysis to cross-link whole-brain rsFC with brain morphometry results obtained in our previous study of this data (1). The neurological music therapy increased the coupling between the FPN and DAN as well as between these networks and primary sensory networks that were engaged during musical training. By contrast, the DMN was less connected with sensory networks after the intervention. Similarly, there was a shift towards a less connected state within the FPN and SAL networks, which are typically hyperconnected following TBI. Improvements in EF were correlated with rsFC within the FPN and between the DMN and sensorimotor networks. Finally, the increase in grey matter volume in frontal regions was associated with greater rsFC in areas implicated in music processing. This study is the largest of its kind, and suggests that rsFC in response to music-based rehabilitation may provide sensitive biomarkers of cognitive recovery after TBI.

executive function | music therapy | traumatic brain injury | resting-state MRI

Correspondence: *noelia.martínezmolina@helsinki.fi*

Introduction

Each year there are over 50 million diagnosed cases of traumatic brain injury (TBI), and it has been estimated that approximately half of the world's population will sustain at least a minor TBI during their lifetime (2). The consequences of TBI can be fatal; it is the leading cause of mortality in young adults and a major cause of death and disability across all ages worldwide. Despite the wide variety of symptoms that can follow a TBI, the most prominent cognitive impairments affect the domains of attention, memory, communication, and executive functioning (EF) (2–6). In fact, deficits in EF are deemed to be the core symptoms of TBI (7, 8). Although there is no consensus on the exact definition of EF, it is thought to encompass several cognitive processes including the core of set shifting, inhibition, and updating (9). Given the heterogeneous and complex nature of TBI, there is an urgent need to develop novel and motivating rehabilitation strategies that target multiple deficits simultaneously, yet with a primary focus on EF.

It has been shown that musical training enhances EF and increases the engagement of the cognitive control network (10–19). Since brain injury patients are still able to enjoy and participate in musical activities (20), neurological music therapy can potentially contribute to restore the EF deficits observed in TBI patients. Until recently, this question had been addressed by only three studies exploring the cognitive effects of music-based interventions after TBI (21–23). Evidence from these studies indicated that music-based rehabilitation can indeed lead to cognitive recovery after brain injury, especially in the domain of mental flexibility. However, these studies presented important limitations with regard to the sample size, lack of proper randomized controlled designs, and inclusion of patients with brain injury not caused by trauma.

We have conducted the first-ever randomized controlled trial (RCT) of neurological music therapy in a sample of 40 moderate/severe TBI patients, where different domains

of EF, attention, and memory were systematically analysed. We used a single-blind crossover design with 3 time-points (baseline/3-months/6-months) for neuropsychological assessment and s/fMRI acquisition. The neurological music therapy consisted of 20 individual therapy sessions held by a trained music therapist over a 3-month period (see Methods section for more details) and was targeted primarily to the rehabilitation of EF, attention, and working memory. In a previous publication, we reported that the music-based intervention induced a cognitive improvement in general EF performance as well as in set shifting (1).

In addition to demonstrating this improvement in neuropsychological performance, in (1) we conducted a voxel-based morphometry (VBM) analysis to investigate the volumetric changes induced by the music therapy. This analysis was motivated by previous work showing that environmental enrichment, such as that provided by musical activities, can increase cognitive reserve and promote adaptive structural neuroplasticity changes in TBI (24–26) and also in stroke patients (27). Our VBM results indicated that TBI patients showed an increase in grey matter volume (GMV) in different brain regions involved in music processing and cognitive function after the intervention. A therapy-induced increase in GMV was seen especially in the right inferior frontal gyrus (IFG) and was correlated with enhanced set shifting ability.

Resting-state functional connectivity (rsFC) is characterized by task-free spontaneous fluctuations in brain activity that occur synchronously across spatially distant brain regions in humans and non-human primates (28, 29). These fluctuations can be measured with the blood-oxygen-level-dependent response at low frequencies (usually under 0.15Hz) and are spatially organized in resting-state networks (RSNs) (30) that mirror activity evoked during cognitive tasks (31, 32). Examining rsFC after TBI is an active area of investigation motivated by the fact that diffuse axonal injury, which is a common pathology reported in all severities of TBI (33, 34), damages the structural connectivity that partly underlies functional connectivity across RSNs (4). This phenomenon makes the study of rsFC after brain injury especially relevant as the loss of integration of information in large-scale brain networks impairs performance in high-level cognitive functions (30, 35–38).

rsFC studies of TBI patients have revealed both increases and decreases in network connectivity, including the default mode (DMN) and salience (SAL) networks as well as multiple sensory and cognitive networks across the spectrum of injury severity (39–44). For example, decreased rsFC has been observed within five network pairs (DMN-basal ganglia, attention-sensorimotor, frontal-DMN, attention-frontal, and sensorimotor-sensorimotor) (43); also within the motor-striatal network, in contrast to increased connectivity within the right frontoparietal network (FPN) (42). In several cases, these abnormalities correlated with post-concussive symptoms (39, 40) and cognitive impairment (45).

Despite the complex abnormalities of interactions within and

between RSNs following TBI, it is possible to identify some distinctive patterns. Within networks, reduced rsFC within the nodes of the DMN predicts attentional impairment (36). Such association could be driven in turn by damage to the cingulum bundle connecting the nodes of the DMN. The temporal coordination between networks, which is important for efficient high-level cognitive function, has also shown consistent abnormalities after TBI. According to an influential model of cognitive control (46, 47), the switching from automatic to controlled behaviour is mediated by the interaction between the SAL and DMN networks, including deactivation of the DMN to attend unexpected external events. Indeed, TBI patients exhibit a failure to appropriately deactivate the DMN, which is associated with impaired response inhibition to a stop signal (36).

The paradoxical yet well-documented finding that functional connectivity may increase secondary to TBI (40, 41, 44, 48–52) has given rise to the 'hyperconnectivity' hypothesis to explain the evolution of brain network reorganization after neurological disruption (53). In this context, hyperconnectivity is defined as enhanced functional connectivity in the number or strength of connections. It is thought to predominantly affect RSNs with high-degree nodes, also known as network hubs, such as the FPN, DMN and SAL networks. Although this hyperconnected state may be adaptive in the short-term in order to re-establish network communication through network hubs, it has been argued that it may have negative long-term consequences due to the chronic enhancement of brain resource utilization, and increased metabolic stress. In the case of TBI patients, this may lead to late pathological complications including Alzheimer´s disease (54, 55), where amyloid beta deposition has been linked to the neurodegeneration of posterior DMN hubs with high metabolic rate (56–58).

In the present study, we extend our previous findings by analysing rsFC from the same sample of moderate/severe TBI patients (see Methods for more details) as the music therapy RCT (1). We used a seed-to-target approach to analyse the reconfiguration of RSNs induced by the neurological music therapy, selecting seeds from the nodes of four key networks: the DMN, SAL, FPN and dorsal attention (DAN) networks.

The current work is grounded in two main hypotheses: (H1) the intervention leads to enhanced coordination activity between attention and executive function (DAN, FPN)-supporting networks, and sensory networks that are engaged during the music therapy and (H2) the intervention leads to reduced connectivity of nodes from the SAL, DMN and FPN networks. In addition, we examined the relationship between the changes in rsFC within and between networks, and the therapy-induced improvement in EF shown previously with neuropsychological testing (1). We anticipated that the FPN would be less connected in those TBI patients with better EF performance; as the hyperconnectivity hypothesis predicts. Lastly, in a similar vein to the approach adopted by Han et al. (59), we conducted an exploratory seed-to-voxel analysis to elucidate the link between whole-brain rsFC and GMV

enhancements induced by the neurological music therapy.

Methods

We conducted a single-blinded cross-over RCT (trial number: NCT01956136) with a 6-month follow-up phase. Upon recruitment, patients were randomly assigned to one of two groups: AB and BA. During the first 3-month period, the AB group received neurological music therapy in addition to standard care, whereas the BA group received only standard care. During the second 3-month period, the BA group received the music therapy intervention and standard care and the AB group received only standard care. Baseline measurements were administered at time point 1 (TP1), and follow-up measurements were conducted at the 3-month crossover point (TP2) and at the 6-month completion point (TP3).

Subjects and study design. The subjects analysed herein are a subset of our previous study (1). In the RCT, 40 TBI patients were recruited through the Brain Injury Clinic of the Helsinki University Central Hospital (HUCH), Validia Rehabilitation Helsinki, and the Department of Neurology of Lohja Hospital during 2014–2017. The inclusion criteria were: 1) diagnosed [Statistical Classification of Diseases and Related Health Problems, 10th revision (ICD-10)] TBI fulfilling the criteria of at least moderate severity [Glasgow Coma Scale (GCS) score: ≤ 12 and/or post-traumatic amnesia (PTA) ≥ 24h]; 2) time since injury ≤ 24 months at the time of recruitment; 3) cognitive symptoms caused by TBI (attention, executive function, memory); 4) no previous neurological or severe psychiatric illnesses or substance abuse; 5) age 16–60 years; 6) native Finnish speaking or bilingual with sufficient communication skills in Finnish; 7) living locally (in the Helsinki–Uusimaa region of Finland); and 8) understanding the purpose of the study and being able to give an informed consent.

Randomization of patients was performed using an online random number generator (https://www.random.org/) by a person not involved in patient recruitment or assessments. To ensure steady allocation to both groups across the trial, the randomization was done in batches of two consecutive patients. The randomization was stratified for lesion laterality (left/right/bilateral). The trial was conducted according to the Declaration of Helsinki and was consistent with good clinical practice and the applicable regulatory requirements. The trial protocol was approved by the Coordinating Ethics Committee of the Hospital District of Helsinki and Uusimaa (reference number 338/13/03/00/2012) and all subjects signed an informed consent.

Of the 40 TBI patients enrolled to the trial, there were three drop-outs before TP2 and another three drop-outs before TP3. The dropouts were mainly due to lack of energy and motivation. Of the remaining 34 patients, one was excluded from the analyses due to intensive self-implemented piano training, which was not part of the trial protocol, and 10 were excluded from the analyses due to lack of fMRI data owing to MRI contraindications of the patients preventing scanning, technical difficulties, and geometric distortion artifacts in fMRI. This yielded a total sample of 23 patients (AB: n = 15, BA: n = 8) for the present study.

Intervention. The neurological music therapy intervention consisted of 20 individual therapy sessions (2 times/week, 60 min/session) held by a trained music therapist (authors S.L., M.H., and M.A.) at Validia Rehabilitation Helsinki. The intervention required no previous musical experience and was adaptable to different degrees of injury across TBI patients. The intervention focused on active musical production with drums and piano. Each session included three 20 min modules: (i) rhythmical training (playing sequences of musical rhythms and coordinated bimanual movements on a djembe drum and on own body), (ii) structured cognitive-motor training (playing musical exercises on a drum set with varying levels of movement elements and composition of drum pads), and (iii) assisted music playing (learning to play own favorite songs on the piano with the help of the therapist and using a special musical notation system called Figure Note). The difficulty level of the exercises was initially adjusted, and then increased in a step-wise manner within and across the sessions, to meet the skill level and progression of the patient. Musical improvisation was also included in all modules and encouraged throughout the therapy to facilitate creative expression.

Neuropsychological assessment. An extensive battery of neuropsychological tests was used to measure the potential cognitive outcomes of the music therapy intervention. The battery is fully described in our previous article (1). Briefly, it comprised standard neuropsychological tests of general EF [Frontal Assessment Battery (FAB) (60)], reasoning ability [Wechsler Adult Intelligence Scale IV (WAIS-IV) / Similarities and Block design subtests (61)], and verbal memory [Wechsler Memory Scale III (WMS-III) / Word Lists I-II subtests (62) and WAIS-IV / Digit Span subtest (61)] as well as computerized tests measuring different EF subcomponents, including set shifting [Number-Letter Task (NLT) (63)], working memory updating [Auditory N-back Task (64)], and inhibition [Simon Task (65)], and sustained attention [Sustained Attention to Response Task (SART)(66)]. In addition, a self-report questionnaire measuring the severity of executive deficits [Behavioural Rating Inventory of Executive Function – Adult version (BRIEF-A) (67)] was included.

In the present study, we analysed the fMRI data by correlating with the general EF (FAB total score), set shifting (NLT switching cost errors), and self-reported executive deficits (BRIEF-A / Self-Monitor and Inhibition scales), as these instruments showed a significant positive effect of the music therapy intervention in our previous publication (1) (as well as in another publication currently under preparation). The FAB (60) consists of six subtests exploring conceptualization, mental flexibility, motor programming, sensitivity to interference, inhibitory control, and environmental autonomy. The computerized tests of executive function (EF) were chosen to reflect aspects of EF (set shifting, updating, inhibi-

tion) defined by Miyake and colleagues [68], as well as sustained attention. Briefly, set shifting was measured with the Number-Letter Task (NLT) [63]; updating with an Auditory N-back task [64]; inhibition with the Simon Task [65]; and sustained attention with the Sustained Attention to Response Task (SART) [66]. In the NLT, switching costs for both reaction times and error percentages were calculated by comparing performance on switching and nonswitching trials.

MRI Data Acquisition. Patients underwent structural MRI (sMRI) and resting-state functional (rs-fMRI) scanning, using an 8-channel SENSE head coil, in a 3T Philips Achieva MRI scanner (Philips Medical Systems) of the HUS Helsinki Medical Imaging Center at Helsinki University Central Hospital. In sMRI, we acquired 2 T1-weighted structural images, using: (A) a magnetization-prepared rapid acquisition gradient echo (MPRAGE, hereafter referred to as T1) sequence [repetition time (TR) = 9.9 ms; echo time (TE) = 4.60 ms; flip angle (FA) = 8^o; field of view (FOV) = 24.0×24.0 cm, matrix = 272×272, 187 slices, slice thickness 0.88 mm with no gap]; and (B) a fluid-attenuated inversion recovery (FLAIR) sequence [repetition time (TR) = 8 ms; echo time (TE) shortest; angle= 50^o; field of view (FOV) = 9.8×9.7 cm, matrix = 228×226, 300 slices, slice thickness 0.60 mm with no gap]. In rs-fMRI, images were acquired with a T2*-weighted image sequence (TR/TE = 2000/35 ms; FA = 90^o, FOV = 23.0×23.0 cm, matrix = 80×80, 31 slices, slice thickness 4 mm with 0.5 mm gap). During the rs-fMRI acquisition, the patients were instructed to remain still, eyes open and fixated to a cross.

rs-fMRI Preprocessing. Data were preprocessed using a standard pipeline in Statistical Parametric Mapping software (SPM12, Wellcome Department of Cognitive Neurology, University College London) running under Matlab Release 2017a (The MathWorks, Inc., Natick, MA). The preprocessing methods applied included slice timing correction, realignment, segmentation, normalization to the MNI template (final voxel size 1 mm isotropic), and smoothing with a Gaussian kernel of 8 mm full width at half maximum (FWHM). Focal brain lesions were detected in 11 patients. As the presence of lesions may influence the normalization algorithm, cost function masks (CFM) were defined for these 11 lesioned patients, to achieve optimal normalization with no postregistration lesion shrinkage or out-of-brain distortion [69]. Binary masks of the lesioned areas were obtained by manually drawing, on a slice-by-slice basis, the precise boundaries of the lesion directly into the T1 image from the TP1 session with MRIcron May 2, 2016 release (https://www.nitrc.org/projects/mricron). Accuracy of the CFM was validated by an expert neuroradiologist (author J.P.) who assessed multiple modalities of neuroimaging data acquired at TP1 (MPRAGE, FLAIR).

Next, within-subject T1 images from all time-points (TP1, TP2, TP3) were coregistered using the T1 images from TP1 as reference to ensure that they remained in spatial alignment with the T1 images and CFM from this acquisition. All

images and CFM were oriented to the anterior commissure before this coregistration step. Unified segmentation with medium regularization was applied to the T1 images (masked with CFM for those patients with visible lesions on the T1); grey matter (GM), white matter (WM), and cerebrospinal fluid (CSF) probability maps were obtained for each individual [70]. This procedure is common practice for patients with lesions [71].

After preprocessing, outlier identification was performed using the Artifact Detection Tools (ART) toolbox (https://www.nitrc.org/projects/artifact$_$detect). We identified outliers as scans that exceeded 3 standard deviations from the global mean BOLD signal, or with framewise displacement >2 mm.

rs-fMRI Denoising. We used denoising to minimize the variability due to physiological, outlier, and residual subject-motion effects. The preprocessed output files from SPM and ART were imported into the CONN toolbox v(18.b) (https://www.nitrc.org/projects/conn), including the following subject- and session-specific files: T1 and rs-fMRI scans; segmented GM, WM, and CSF images; 6 realignment parameters after rigid body motion correction; and the regressor files containing the outliers. We used the default denoising pipeline in the CONN toolbox which comprises two steps in consecutive order: (1) linear regression of potential confounding effects in the BOLD signal, and (2) temporal band-pass filtering. Step 1 implements an anatomical component-based noise correction procedure (aCompCor), and includes: noise components from cerebral white matter and cerebrospinal areas [72]; estimated subject-motion parameters and first-order derivatives [73]; identified outlier scans [74]; constant and first-order linear session effects [75]. After evaluation of denoised outputs (residual BOLD time series), the number of noise components extracted from the white matter was increased to 10. In step 2, frequencies below 0.01Hz were filtered from the BOLD signal to reduce the slow signal drift.

Longitudinal analysis of seed-based connectivity associated with GMV changes. Several studies have shown that structural and brain connectivity changes may serve as biomarkers of plasticity in adults with chronic TBI [59, 76]. In our previous study, we demonstrated that neurological music therapy was able to induced morphometric changes in chronic TBI patients [1]. In particular, our findings indicated that GMV in the right inferior frontal gyrus (IFG), among other regions, increased significantly in AB and BA groups during the intervention versus control period as well as before and after the intervention period [1]. In addition, the increase in GMV in the right IFG correlated with cognitive improvement in set shifting [1].

In the current study, we selected the statistically significant clusters in the right IFG from the intervention period analysis to assess their seed-based connectivity in resting-state, because it was the only region to significantly correlate with therapy-induced EF improvement. The seeds included two

clusters in the right IFG triangular part in standard space, that were imported into the CONN toolbox to compute seed-to-voxel functional connectivity. For more information regarding peak coordinate and size of the seeds see Table 4 in our previous study (1).

Seed-based connectivity (SBC) maps (77) between each seed and every voxel in the brain were obtained for each subject and session. SBC maps were computed as the Fisher-transformed Pearson´s bivariate correlation coefficients between a ROI BOLD timeseries and the BOLD time series of each individual voxel, using the equation:

$$r(x) = \frac{\int S(x,t)R(t)dt}{\sqrt{\int R^2(t)dtx \int S^2(x,t)dt}}$$

$$Z(x) = tanh^{-1}(r(x))$$

where S is the BOLD time series at each voxel, R is the average BOLD time series within a ROI, r is the spatial map of Pearson correlation coefficients, and Z is the SBC map of Fisher-transformed correlation coefficients for this ROI.

Connectivity pattern analysis within and between large-scale networks. To further understand functional connectivity (FC) changes induced by the neurological music therapy from the perspective of large-scale networks, we used the 8 resting-state networks (RSNs) comprising 32 regions of interest (ROIs) or nodes in the CONN toolbox, which are derived from an independent component analysis (ICA) of 498 subjects from the Human Connectome Project (77). The 8 RSNs included the default mode (DMN), salience (SAL), frontoparietal (FPN), dorsal attention (DAN), sensorimotor (SM), language (LAN), visual (VIS) and cerebellar (CER) networks. All of these networks have been widely reported in the resting-state literature.

We identified changes in both between- and within-network connectivity from those networks that have shown alterations after TBI (4) or are associated with EF, memory, and attention, which are the most commonly affected cognitive domains in TBI patients (3–6). In particular, we selected ROIs from the DMN, SAL, FPN, and DAN networks to examine the FC (a) within their nodes, and (b) between their nodes and the rest of the RSNs included in the CONN toolbox.

We used the same connectivity measure as in the SBC connectivity analysis described in the previous section: the Fisher-transformed bivariate correlation coefficient between a pair of ROIs time series (77). Each pair of ROIs (i,j) constitutes an element in the ROI-to-ROI connectivity (RRC) matrix that can be calculated as follows:

$$r(i,j) = \frac{\int Ri(x,t)Rj(t)dt}{\sqrt{\int R_i^2(t)dtx \int R_j^2(x,t)dt}}$$

$$Z(i,j) = tanh^{-1}(r(i,j))$$

where R is the BOLD time series within each ROI, r is a matrix of correlation coefficients, and Z is the RRC symmetric matrix of Fisher-transformed correlation coefficients.

Resting-state functional connectivity and cognitive performance. We investigated the link between rsFC and cognitive performance improvements after therapy. We performed Pearson's bivariate correlations between, on one hand, the SBC or ROI-to-ROI (R2R) FC changes (found significant in the main analyses), and on the other hand, neuropsychological tests measuring general EF (FAB total score), set shifting (NLT switching cost errors), and the Self-Monitor and Inhibition subscales of BRIEF-A. The chosen performance measures were those which showed a statistically significant improvement after the music therapy (described in (1) and in an article in preparation).

Quality assurance. We visually inspected all structural and functional MRI scans to ensure that patients had no significant brain atrophy. In rsMRI preprocessing, the quality of the preprocessed data was visually inspected using the quality assurance (QA) plots available in the CONN toolbox (QA normalization, registration, segmentation, and framewise displacement). In rsMRI denoising, we inspected the distribution of connectivity values for each patient and session after denoising, and adjusted the number of noise components extracted from white matter to ensure that this distribution peaked around 0.

Group-level analysis. All group-level analysis were performed in the CONN toolbox using the General Linear Model framework. In the SBC analysis, to compare pre-post intervention increases in FC, we performed a paired T-test between every voxel in the brain and the clusters that showed increased GMV after the intervention. For the AB group, the beta maps from TP1 were entered as the pre-intervention condition and the beta maps from TP2 were entered as the post-intervention condition. For the BA group, the beta maps from TP2 were entered as the pre-intervention condition and the beta maps from TP3 were entered as the post-intervention condition. We focused on the pre-post intervention effect as this was the only analysis with significant brain-behaviour relationships in our previous study (1).

In the R2R approach with RSN, we performed two group-level analyses using paired T-tests (pre- and post- intervention effect for the AB and BA groups) and mixed-model ANOVA (AB>BA×TP2>TP1 interaction). For the AB group, the denoised ROI timeseries from TP1 were entered as the pre-intervention condition, and the denoised ROI time series from TP2 were entered as the post-intervention condition. For the BA group, the denoised ROI time series from TP2 were entered as the pre-intervention condition, and the denoised ROI time series from TP3 were entered as the post-intervention condition. For the within-networks analysis, we examined the R2R connectivity of the ROIs within each of the selected 4 RSNs (DMN, SAL, FPN, and DAN). For the between-network analysis, we examined the connectivity be-

tween the ROIs of the each of these RSNs (DMN, SAL, FPN, DAN) with every other RSN included in the CONN toolbox.

The statistical parametric maps obtained were corrected for multiple comparisons based on Gaussian Random Field Theory [78]. The maps were first thresholded using a "height" threshold of $p < 0.001$ and the resulting clusters thresholded with a cluster-level False Discovery Rate (FDR)-corrected alpha threshold of $p < 0.05$ [79].

Results

The quality assurance procedure resulted in 145 rs-fMRI volumes from 23 participants (AB group, n=15; BA group, n=8). The scans were acquired at three time-points: TP1 (baseline), TP2 (3-months) and TP3 (6-months).

Seed-based resting-state functional connectivity. Previously, using voxel-based morphometry (VBM) we found that GMV in the right inferior frontal gyrus (IFG) increased significantly in both groups during the intervention period, and that this was positively correlated with enhanced set shifting ability [1]. To examine the changes in functional connectivity associated with this increase in GMV, we performed seed-based connectivity (SBC) using as sources the two clusters in the right IFG that resulted from our previous voxel-based morphometry analysis [1]. In the SBC analysis (results shown in Figure 1), a paired T-test of pre- versus post-intervention rsFC (pooling together AB and BA groups) revealed that the right IFG increased its connectivity with the right inferior parietal lobule (IPL; peak voxel x = 38, y = -46, z = 48; Figure 1A) and the left Rolandic operculum (peak voxel x = -54, y = -2, z = 4; Figure 1B) with $p < 0.05$ after FDR-correction at cluster level. There were no other significant clusters.

Between- and within-network resting-state functional connectivity. To investigate the positive effect of the neurological music therapy on the coordinated activity between large-scale resting-state networks (RSNs) that are important for high-level cognitive functions, we performed a ROI-to-ROI (R2R) analysis focusing primarily on the frontoparietal (FPN), dorsal attention (DAN), salience (SAL) and default mode (DMN) networks as source ROIs. Figure 2 shows the results of the between-network connectivity for these four RSNs, in a paired T-test of pre- versus post-intervention pooling together data from AB and BA groups (Figure 2A-C), and in the AB>BA×TP2>TP1 interaction (Figure 2D) with $p < 0.05$ FDR-corrected at the source level.

In the pre- versus post-intervention comparison (Table S1), we found that the FPN increased its temporal coupling with the sensorimotor (SM) and DAN after the music-based intervention. Likewise, when the DAN was used as a source, we observed increased between-network connectivity with several nodes of the visual (VIS) network induced by the intervention. In particular, the lateral prefrontal cortex in the left hemisphere from the FPN increased its functional connectivity with the right sensorimotor lateral node from the

SM network (Figure 2A) and the right intraparietal sulcus from the DAN network. In addition, the latter was also highly connected with the visual occipital and bilateral visual lateral nodes from the VIS network. By contrast, in the AB>BA×TP2>TP1 interaction (Table S1), we found that the spontaneous fluctuations between the medial prefrontal cortex from the DMN network and the sensorimotor superior node from the SM network were less coordinated after the intervention.

Next, we compared the connectivity within networks (Figure 3). In this case, only the pre- versus post-intervention comparison pooling together AB and BA groups (Table S2) yielded significant results with $p < 0.05$ FDR-corrected at the source level. The FPN and SAL networks showed a reduction in the functional connectivity between several of their constituents nodes. Specifically, the lateral prefrontal cortex node in the left hemisphere exhibited less coupling with its contralateral counterpart and with the posterior parietal cortex node in the left hemisphere. Similarly, within the SAL network, the supramarginal gyrus node in the left hemisphere showed a decreased coupling with its contralateral counterpart and with the anterior insula node in the right hemisphere.

Correlation between neuropsychological outcomes and resting-state functional connectivity. Next, we investigated how the changes in RSNs induced by the neurological music therapy (see above) were associated with the parallel improvement in cognitive function, specifically in general EF (FAB score), set shifting (NLT switching cost errors), and self-reported executive deficits (BRIEF-A Self-Monitor and Inhibition subscales). We found statistically significant associations between the therapy-induced cognitive improvement and within-network and between-network changes in RSNs after the intervention (Figure 4). Decreased within-network FC in the left and right lateral prefrontal cortex nodes of the FPN correlated significantly with increased FAB scores ($p = 0.040$, $r = -0.372$, Figure 4A, left) and decreased BRIEF-A Self-Monitor scores ($p = 0.013$, $r = 0.485$, Figure 4A, right). Decreased between-network FC between the DMN (medial prefrontal cortex node) and the SM (superior sensorimotor cortex node) networks correlated significantly with increased FAB scores ($p = 0.003$, $r = -0.559$, Figure 4B, left) and marginally significantly also with decreased NLT switching cost errors ($p = 0.052$, $r = 0.347$, Figure 4B, right). There were no other significant correlations. Together, these results indicate that those patients who showed a larger reduction in connectivity within the FPN and between the DMN and SM networks exhibited greater improvement in general EF (higher FAB scores) and set shifting ability (less NLT errors) as well as greater reduction in executive deficits in self-monitoring (smaller BRIEF-A Self-Monitor scores). Thus, network-specific patterns of functional connectivity induced by the music-based intervention are associated with improvement in EF.

Fig. 1. Changes in seed-based connectivity induced by the neurological music therapy intervention. A.*Top*: R IFG mask used as seed with peak coordinate x=50, y=30, y=0, overlaid on the Montreal Neurological Institute (MNI) template. A.*Bottom*: Statistical map showing the $p < 0.05$ FDR-corrected cluster in the R parietal Inf. B.*Top*: R IFG mask used as seed with peak coordinate x=54, y=22, y=12 overlaid on the MNI template. B.*Bottom*: Statistical map showing the $p < 0.05$ FDR-corrected cluster in the L Rolandic Operculum. C. Effect size plots displaying the functional connectivity (mean beta values) of each significant cluster in the pre-intervention relative to the post-intervention period. *IFG: Inferior Frontal Gyrus, Parietal Inf.: inferior parietal lobule, R: right, L: left.*

Discussion

We used resting-state functional connectivity (rsFC) methods to examine brain network changes induced by neurological music therapy after moderate and severe TBI. We conducted a ROI-to-ROI analysis with nodes from four networks [default mode (DMN), salience (SAL), frontoparietal (FPN), and dorsal attention (DAN)] as seeds and all network ROIs included in the CONN toolbox as targets. This analysis showed that music therapy strengthened network connectivity between FPN and DAN, and between these networks and the sensorimotor (SM) and visual (VIS) networks, respectively. In contrast, the music therapy reduced the connectivity between the nodes of DMN and SM network. The within-network connectivity revealed specific nodes of the FPN and SAL wherein coupling decreased in the pre- versus post-intervention comparison. Importantly, the decrease in the FPN and DMN-SM connectivity was paralleled by cognitive improvement in executive function (EF). Finally, using a seed-based connectivity analysis, we demonstrated that the right inferior frontal gyrus (IFG), in which we previously observed increased grey matter volume (GMV) induced by the music therapy (1), showed high connectivity with left frontal and right parietal regions, implicated in music processing (see Sarkamo et al. (80) for a review). These results reflect substantial functional neuroplasticity changes in resting-state networks, which taken together show a shift from a hyper-connected to a more normal state of connectivity; we believe this represents a shift in executive functioning from a compensatory increase in cognitive control and attentional mechanisms to more adaptive sensory-integrative.

Changes in cross-modal integration. Our first hypothesis (H1) was that neurological music therapy enhances the rsFC between primary sensory networks and higher-order networks as a consequence of iterated interaction between the sensory-cognitive systems recruited by music production, and perception (81). This hypothesis was conceptually inspired by the 'global workspace' (or cross-modal integration) notion empirically supported by the convergence of unimodal sensory networks into cortical hubs that contribute to perceptual integration in the human brain (82). Our results confirmed H1 as we detected an increased coupling between the FPN and SM networks. The stepwise functional connectivity performed by Sepulcre et al. (82) highlighted the dorso-lateral prefrontal cortex, which anatomically corresponds to the lateral prefrontal cortex node in the FPN, as one of the cortical hubs reached by seeds in the primary somatosensory cortex. Similarly, the intraparietal sulcus (IPS) node in the DAN was more connected with occipital nodes in the VIS network. This is not surprising given that visual streams have previously been postulated to converge in the IPS (83). We also found an increase in rsFC between the FPN and DAN, which may be related to improved regulation of perceptual attention, as recent work using meta-analytic tools has revealed (84).

Reduced connectivity in network hubs. Our second hypothesis (H2) proposed that the DMN, SAL, and FPN connectivity would be downregulated by the neurological music therapy. Our reasoning was that the repetitive engagement of cognitive control and EF during the intervention (14, 17) would counteract the increased connectivity and cognitive load associated with TBI. Brain injury has a detrimental ef-

BETWEEN NETWORKS

Fig. 2. Changes in between-network connectivity induced by the neurological music therapy intervention. Nodes are overlayed on a rendered semitransparent brain generated using CONN. Adjacency matrices display the mean post- minus pre-intervention (A–C) Fisher-transformed Z-score correlation values for each node. The bar plots (D) show the effect size of the AB>BA and TP2>TP1 interaction represented by the Fisher-transformed Z-score correlation values for each node. *FPN: frontoparietal; SM: sensorimotor; DAN: dorsal attention; VIS: visual; DMN: default mode network; FEF: frontal eye field;, IPS: intraparietal sulcus; LPFC: lateral prefrontal cortex; LAT: lateral; MED: medial; MPFC: medial prefrontal cortex; OCC: occipital; PCC: posterior cingulate cortex; PPC: posterior parietal cortex; Sup: superior; R: right, L: left.*

fect on automatic processes and increases the supervisory demand for the integration of information at all levels (85, 86), leading to the symptoms of fatigue commonly reported by these patients (87). The need to compensate for the excessive cognitive demand that arises from focal lesions and diffuse axonal injury is consistent with the hyperconnectivity hypothesis (48, 53, 88).

One prominent line of evidence in favour of the latter hypothesis comes from the observation that hyperconnectivity secondary to neurological damage, including TBI (48, 51, 52, 89), is centered around nodes with a hub connectivity profile. Network hubs can be defined in terms of their structural (90–93) or functional connectivity (82, 94–96), and are characterized by a high degree of connectivity with the rest of the brain, thus making a strong contribution to the global integration of information (97). Although functional connectivity does not reflect direct anatomical connections, graph measures derived from both methodologies show strong convergence with regards to the brain regions classified as hubs. These include regions overlapping with the nodes of the DMN, SAL, and FPN, such as the superior parietal and superior frontal cortex, the anterior and posterior cingulate cortex

as well as anterior portions of the anterior insula (82). According to the hyperconnectivity hypothesis, a major goal of the increase in functional connectivity following injury is to re-establish network communication through network hubs in order to maximize information transfer and minimize behavioural impairments (53).

Here, we found that the neurological music therapy decreased the functional connectivity within the FPN, involving a reduction in the bilateral communication between the lateral prefrontal cortices, and between the lateral prefrontal cortex and the posterior parietal cortex in the left hemisphere. Considering the hyperconnectivity hypothesis, this finding accords with recent evidence indicating that the bilateral prefrontal cortices are the subcomponents of the FPN with the highest average degree at 3 and 6 months after moderate/severe TBI (48). Several other studies reported a similar increase in connectivity in frontal regions and FPN, among mild TBI patients (41, 42, 51). Our results also showed that the anterior insula in the right hemisphere was less connected with the supramarginal gyrus in the left hemisphere and that interhemispheric connectivity in the latter was equally diminished after the intervention. This within-network connectiv-

WITHIN NETWORKS

Fig. 3. Changes in within-network connectivity induced by the neurological music therapy intervention. Nodes are overlaid on a rendered semitransparent brain generated using CONN. Adjacency matrices display the mean post- minus pre-intervention Fisher-transformed Z-score correlation values for each node. *AINS: anterior insula; IPS: intraparietal sulcus; LAT: lateral; LPFC: lateral prefrontal cortex; MPFC: medial prefrontal cortex; OCC: occipital; PPC: posterior parietal cortex; RPFC: rostral prefrontal cortex; SMG: supramarginal gyrus; Sup: superior; R: right, L: left.*

ity reduction in the SAL network, including the anterior insula, is again consistent with the hyperconnectivity reported by Hillary et al. (48). The increase in functional connectivity of the anterior insula salience network after TBI has also been supported by other longitudinal (36) and cross-sectional studies (40). Regarding the DMN, our analysis revealed a decreased connectivity between the medial prefrontal cortex

and the SM network. Given the interference that the activation of the DMN may exert on attentional switching (46), it may be the case that this reduced connectivity facilitates the sensorimotor coupling with other brain regions involved in music production and perception. The negative trend with EF would lend support to this idea (though, since this correlation was only marginally significant, it should be interpreted

Fig. 4. Within- and between-network functional connectivity changes associated with cognitive recovery induced by the neurological music therapy intervention. The scatter plots represent the bivariate Pearson correlation and shaded areas represent the 95 % CI prediction bounds. *FPN: frontoparietal; DMN: default mode network; SM: sensorimotor; MPFC: medial prefrontal cortex; LPFC: lateral prefrontal cortex; Sup: superior; R: right; L: left.*

with caution).

One clinically-relevant implication from these findings is that the music therapy effectively targeted network hubs from the FPN and SAL networks, and hence may have contributed to the shift from a hyperconnected to a normal state. Within the hyperconnectivity hypothesis framework, it has been argued that while this hyperconnectivity may be adaptive in the short-term, chronic hyperconnectivity may render network hubs vulnerable to late pathological complications due to the chronically-increased metabolic stress. Although the exact mechanisms and cognitive consequences of hyperconnectivity remain to be elucidated, this proposal has found some support from longitudinal studies examining the temporal evolution of TBI recovery (48, 88). For example, Roy et al. (88) performed a cost-efficiency analysis and found a peak in network strength in the frontal DMN and temporoparietal networks at 6-months postinjury, with some residual hyperconnectivity observable after one year but with diminished overall cost. In this context, the neurological music therapy could potentiate the cost-efficiency rebalancing in the TBI recovery trajectory. Though beyond the scope of this paper, this prediction could be directly tested by computing graph theory measures including cost, degree, as well as local and global efficiency.

Indirect support for this idea comes from our correlational re-

sults indicating improved performance in EF as a function of rsFC decrease within the FPN. The neurological music therapy was primarily designed to target a number of EFs (action planning and monitoring, inhibitory control, shifting), which are reflected by the three outcome measures (FAB, NLT, BRIEF-A) showing a correlation with the therapy-induced change in rsFC. In this context, it might be that the repetitive practice during the music intervention reduces the cognitive challenge and transfers to better executive functioning, thus reducing the need for a hyperconnected state to efficiently accomplish the task. This would fit well with the observation that the increased activity in the right prefrontal cortex and anterior insula cortex in a sample of TBI patients normalizes following practice of a working memory task (86).

Relationship between brain morphometry and rsFC changes. The simultaneous combination of both structural and rsFC measures in the context of rehabilitative strategies for TBI has only recently started to be explored (76). However, it offers an interesting possibility since both brain morphometry (98) and rsFC (99) are malleable to musical training. By cross-linking results from our previous voxel-based morphometry publication with rsFC, we were able to identify changes in GMV co-occurring with whole-brain rsFC. We focused on the right IFG, as this was the region showing a significant relationship between GMV increases and improved performance in set shifting (1). Using two right

IFG clusters from the pre- vs post-intervention comparison as seeds, we found enhanced connectivity between the right IFG and the left Rolandic operculum, which is known to be activated in the processing of music-evoked emotions (100). The right IFG also showed therapy-induced enhanced rsFC with the right inferior parietal lobule (also known as Geshwind's territory), which is particularly interesting since both regions are involved in musical syntax analysis (101–103) and have a direct anatomical connection via the anterior segment of the arcuate fasciculus (104). Therefore, we can speculate that this effect could be mediated by a concomitant increase in the underlying structural connectivity. Future tractography studies in the same TBI cohort could specifically address this issue.

Limitations and future directions. This study has some limitations, which need to be considered when evaluating its findings. Although this is the largest RCT using neurological music therapy in moderate/severe TBI to date, the final sample size is relatively modest (N=23) and may preclude the detection of small effect sizes due to lack of statistical power. Given the difficulty in finding patients fulfilling all the inclusion criteria, we were more liberal in the selection of time since injury. For this reason, even if 52% of our patients sustained TBI within 6-months or less at the time of recruitment, there is heterogeneity in our sample. While the interpretation of our findings is consistent with previous longitudinal studies examining rsFC in TBI, a direct comparison is not warranted and future RCTs should consider a stratification of patients by time since injury. Another consideration is the cross-over design: this has the advantage that patients act as their own controls, but the potential risk of a carry-over effect for the group who first participated in the intervention. This possibility seems unlikely since we ruled out such a carry-over effect in our previous publication. However, to address this issue, we restricted the group×time interaction to the two first time-points (1). Regarding rsFC, we used the resting-state networks available in the CONN toolbox that were obtained using an ICA analyses (N=498) from the Human Connectome Project. This selection, however, limited the analysis of sensory-integrative network interactions since the auditory network was not included. As we have established that the music therapy effectively enhances these interactions, future analyses will benefit from including nodes from the auditory network based on brain parcellations (93) or ICA decomposition (30). Although the human brain is a dynamic system, this study investigated how neurological music therapy affects the cross-modal integration across sensory and cognitive networks with a stationary FC approach by averaging time courses from seeds and targets. In order to gain a fine-grained view of how the intervention influences network connectivity in the time domain, future work should examine *dynamic FC* between these segregated functional networks.

Conclusion. To the best of our knowledge, this study provides the first evidence that neurological music therapy can lead to substantial functional neuroplasticity changes in resting-state networks after TBI. Our results lend support to the idea that the music intervention facilitated the integration of primary sensory information, by increasing the rsFC between multimodal and higher-level cognitive networks (FPN, DAN). The shift towards a less connected state within the FPN and SAL networks is also in line with the notion that chronic hyperconnectivy in network hubs after TBI can be maladaptive in the long-term. The relationship between improved performance in EF and lesser connectivity in the FPN and the DMN-SM networks, which might be linked to reduced interference, further supports this idea. Finally, the co-occurrence of changes in brain morphometry and rsFC connectivity, in a circuit supporting music processing, suggests a complex picture with interrelated plasticity across MRI modalities. Overall, our findings suggest that rsFC changes in brain networks can serve as sensitive biomarkers for the efficacy of music-based rehabilitation after TBI.

Bibliography

1. S. T. Siponkoski, N. Martinez-Molina, L. Kuusela, S. Laitinen, M. Holma, M. Ahlfors, P. Jordan-Kilkki, K. Ala-Kauhaluoma, S. Melkas, J. Pekkola, A. Rodriguez-Fornells, M. Laine, A. Ylinen, P. Rantanen, S. Koskinen, J. Lipsanen, and T. Sarkamo. Music therapy enhances executive functions and prefrontal structural neuroplasticity after traumatic brain injury: Evidence from a randomized controlled trial. *J Neurotrauma*, 37(4):618–634, 2020. ISSN 1557-9042 (Electronic) 0897-7151 (Linking). doi: 10.1089/neu.2019.6413.
2. A. I. R. Maas, D. K. Menon, P. D. Adelson, N. Andelic, M. J. Bell, A. Belli, P. Bragge, A. Brazinova, A. Buki, R. M. Chesnut, G. Citerio, M. Coburn, D. J. Cooper, A. T. Crowder, E. Czeiter, M. Czosnyka, R. Diaz-Arrastia, J. P. Dreier, A. C. Duhaime, A. Ercole, T. A. van Essen, V. L. Feigin, G. Gao, J. Giacino, L. E. Gonzalez-Lara, R. L. Gruen, D. Gupta, J. A. Hartings, S. Hill, J. Y. Jiang, N. Ketharanathan, E. J. O. Kompanje, L. Lanyon, S. Laureys, F. Lecky, H. Levin, H. F. Lingsma, M. Maegele, M. Majdan, G. Manley, J. Marsteller, L. Mascia, C. McFadyen, S. Mondello, V. Newcombe, A. Palotie, P. M. Parizel, W. Peul, J. Piercy, S. Polinder, L. Puybasset, T. E. Rasmussen, R. Rossaint, P. Smielewski, J. Soderberg, S. J. Stanworth, M. B. Stein, N. von Steinbuchel, W. Stewart, E. W. Steyerberg, N. Stocchetti, A. Synnot, B. Te Ao, O. Tenovuo, A. Theadom, D. Tibboel, W. Videtta, K. K. W. Wang, W. H. Williams, L. Wilson, and K. Yaffe. Traumatic brain injury: integrated approaches to improve prevention, clinical care, and research. *Lancet Neurol*, 16(12):987–1048, 2017. ISSN 1474-4422. doi: 10.1016/s1474-4422(17)30371-x.
3. D. T. Stuss. Traumatic brain injury: relation to executive dysfunction and the frontal lobes. *Curr Opin Neurol*, 24(6):584–9, 2011. ISSN 1350-7540. doi: 10.1097/WCO.0b013e32834c7eb9.
4. D. J. Sharp, G. Scott, and R. Leech. Network dysfunction after traumatic brain injury. *Nat Rev Neurol*, 10(3):156–66, 2014. ISSN 1759-4766 (Electronic) 1759-4758 (Linking). doi: 10.1038/nrneurol.2014.15.
5. T. E. Ham and D. J. Sharp. How can investigation of network function inform rehabilitation after traumatic brain injury? *Curr Opin Neurol*, 25(6):662–9, 2012. ISSN 1350-7540. doi: 10.1097/WCO.0b013e328359488f.
6. P. W. Burgess and D. T. Stuss. Fifty years of prefrontal cortex research: Impact on assessment. *J Int Neuropsychol Soc*, 23(9-10):755–767, 2017. ISSN 1355-6177. doi: 10.1017/s1355617717000704.
7. K. M. Krpan, B. Levine, D. T. Stuss, and D. R. Dawson. Executive function and coping at one-year post traumatic brain injury. *J Clin Exp Neuropsychol*, 29(1):36–46, 2007. ISSN 1380-3395 (Print) 1380-3395. doi: 10.1080/13803390500376816.
8. K. Cicerone, H. Levin, J. Malec, D. Stuss, and J. Whyte. Cognitive rehabilitation interventions for executive function: moving from bench to bedside in patients with traumatic brain injury. *J Cogn Neurosci*, 18(7):1212–22, 2006. ISSN 0898-929X (Print) 0898-929x. doi: 10.1162/jocn.2006.18.7.1212.
9. N. P. Friedman and A. Miyake. Unity and diversity of executive functions: Individual differences as a window on cognitive structure. *Cortex*, 86:186–204, 2017. ISSN 0010-9452. doi: 10.1016/j.cortex.2016.04.023.
10. S. M. Carpentier, S. Moreno, and A. R. McIntosh. Short-term music training enhances complex, distributed neural communication during music and linguistic tasks. *J Cogn Neurosci*, 28(10):1603–12, 2016. ISSN 0898-929x. doi: 10.1162/jocn_a_00988.
11. A. C. Jaschke, H. Honing, and E. J. A. Scherder. Longitudinal analysis of music education on executive functions in primary school children. *Front Neurosci*, 12:103, 2018. ISSN 1662-4548 (Print) 1662-453x. doi: 10.3389/fnins.2018.00103.
12. V. Putkinen, M. Tervaniemi, K. Saarikivi, and M. Huotilainen. Promises of formal and informal musical activities in advancing neurocognitive development throughout childhood. *Ann N Y Acad Sci*, 1337:153–62, 2015. ISSN 0077-8923. doi: 10.1111/nyas.12656.
13. S. Moreno, E. Bialystok, R. Barac, E. G. Schellenberg, N. J. Cepeda, and T. Chau. Short-term music training enhances verbal intelligence and executive function. *Psychol Sci*, 22(11):1425–33, 2011. ISSN 0956-7976. doi: 10.1177/0956797611416999.
14. M. Sachs, J. Kaplan, A. Der Sarkissian, and A. Habibi. Increased engagement of the cognitive control network associated with music training in children during an fmri stroop task. *PLoS One*, 12(10):e0187254, 2017. ISSN 1932-6203. doi: 10.1371/journal.pone.0187254.

15. A. Habibi, A. Damasio, B. Ilari, M. Elliott Sachs, and H. Damasio. Music training and child development: a review of recent findings from a longitudinal study. *Ann N Y Acad Sci*, 2018. ISSN 0077-8923. doi: 10.1111/nyas.13606.

16. L. Moradzadeh, G. Blumenthal, and M. Wiseheart. Musical training, bilingualism, and executive function: a closer look at task switching and dual-task performance. *Cogn Sci*, 39(5):992–1020, 2015. ISSN 1551-6709 (Electronic) 0364-0213 (Linking). doi: 10.1111/cogs.12183.

17. J. Zuk, C. Benjamin, A. Kenyon, and N. Gaab. Behavioral and neural correlates of executive functioning in musicians and non-musicians. *PLoS One*, 9(6):e99868, 2014. ISSN 1932-6203. doi: 10.1371/journal.pone.0099868.

18. J. A. Bugos, W. M. Perlstein, C. S. McCrae, T. S. Brophy, and P. H. Bedenbaugh. Individualized piano instruction enhances executive functioning and working memory in older adults. *Aging Ment Health*, 11(4):464–71, 2007. ISSN 1360-7863 (Print) 1360-7863. doi: 10.1080/13607860601086504.

19. J. V. Strong and B. T. Mast. The cognitive functioning of older adult instrumental musicians and non-musicians. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn*, 26(3):367–386, 2019. ISSN 1382-5585. doi: 10.1080/13825585.2018.1448356.

20. A. M. Belfi, E. Evans, J. Heskje, J. Bruss, and D. Tranel. Musical anhedonia after focal brain damage. *Neuropsychologia*, 97:29–37, 2017. ISSN 1873-3514 (Electronic) 0028-3932 (Linking). doi: 10.1016/j.neuropsychologia.2017.01.030.

21. M. H. Thaut, J. C. Gardiner, D. Holmberg, J. Horwitz, L. Kent, G. Andrews, B. Donelan, and G. R. McIntosh. Neurologic music therapy improves executive function and emotional adjustment in traumatic brain injury rehabilitation. *Ann N Y Acad Sci*, 1169:406–16, 2009. ISSN 0077-8923. doi: 10.1111/j.1749-6632.2009.04585.x.

22. C. Lynch and A. B. LaGasse. Training endogenous task shifting using music therapy: A feasibility study. *J Music Ther*, 53(3):279–307, 2016. ISSN 0022-2917. doi: 10.1093/jmt/thw008.

23. B. M. D. Vik, G. O. Skeie, E. Vikane, and K. Specht. Effects of music production on cortical plasticity within cognitive rehabilitation of patients with mild traumatic brain injury. *Brain Inj*, 32(5):634–643, 2018. ISSN 0269-9052. doi: 10.1080/02699052.2018.1431842.

24. J. C. Tomaszczyk, N. L. Green, D. Frasca, B. Colella, G. R. Turner, B. K. Christensen, and R. E. Green. Negative neuroplasticity in chronic traumatic brain injury and implications for neurorehabilitation. *Neuropsychol Rev*, 24(4):409–27, 2014. ISSN 1040-7308. doi: 10.1007/s11065-014-9273-6.

25. L. S. Miller, B. Colella, D. Mikulis, J. Maller, and R. E. Green. Environmental enrichment may protect against hippocampal atrophy in the chronic stages of traumatic brain injury. *Front Hum Neurosci*, 7:506, 2013. ISSN 1662-5161 (Print) 1662-5161. doi: 10.3389/fnhum.2013.00506.

26. P. B. de la Tremblaye, J. P. Cheng, C. O. Bondi, and A. E. Kline. Environmental enrichment, alone or in combination with various pharmacotherapies, confers marked benefits after traumatic brain injury. *Neuropharmacology*, 145(Pt A):13–24, 2019. ISSN 0028-3908. doi: 10.1016/j.neuropharm.2018.02.032.

27. T. Sarkamo, P. Ripolles, H. Vepsalainen, T. Autti, H. M. Silvennoinen, E. Salli, S. Laitinen, A. Forsblom, S. Soinila, and A. Rodriguez-Fornells. Structural changes induced by daily music listening in the recovering brain after middle cerebral artery stroke: a voxel-based morphometry study. *Front Hum Neurosci*, 8:245, 2014. ISSN 1662-5161 (Print) 1662-5161 (Linking). doi: 10.3389/fnhum.2014.00245.

28. R. M. Hutchison and S. Everling. Monkey in the middle: why non-human primates are needed to bridge the gap in resting-state investigations. *Front Neuroanat*, 6:29, 2012. ISSN 1662-5129. doi: 10.3389/fnana.2012.00029.

29. M. E. Raichle and M. A. Mintun. Brain work and brain imaging. *Annu Rev Neurosci*, 29: 449–76, 2006. ISSN 0147-006X (Print) 0147-006x. doi: 10.1146/annurev.neuro.29.051605.112819.

30. S. M. Smith, P. T. Fox, K. L. Miller, D. C. Glahn, P. M. Fox, C. E. Mackay, N. Filippini, K. E. Watkins, R. Toro, A. R. Laird, and C. F. Beckmann. Correspondence of the brain's functional architecture during activation and rest. *Proc Natl Acad Sci U S A*, 106(31): 13040–5, 2009. ISSN 0027-8424. doi: 10.1073/pnas.0905267106.

31. B. Biswal, F. Z. Yetkin, V. M. Haughton, and J. S. Hyde. Functional connectivity in the motor cortex of resting human brain using echo-planar mri. *Magn Reson Med*, 34(4): 537–41, 1995. ISSN 0740-3194 (Print) 0740-3194. doi: 10.1002/mrm.1910340409.

32. B. B. Biswal, M. Mennes, X. N. Zuo, S. Gohel, C. Kelly, S. M. Smith, C. F. Beckmann, J. S. Adelstein, R. L. Buckner, S. Colcombe, A. M. Dogonowski, M. Ernst, D. Fair, M. Hampson, M. J. Hoptman, J. S. Hyde, V. J. Kiviniemi, R. Kotter, S. J. Li, C. P. Lin, M. J. Lowe, C. Mackay, D. J. Madden, K. H. Madsen, D. S. Margulies, H. S. Mayberg, K. McMahon, C. S. Monk, S. H. Mostofsky, B. J. Nagel, J. J. Pekar, S. J. Peltier, S. E. Petersen, V. Riedl, S. A. Rombouts, B. Rypma, B. L. Schlaggar, S. Schmidt, R. D. Seidler, G. J. Siegle, C. Sorg, G. J. Teng, J. Veijola, A. Villringer, M. Walter, L. Wang, X. C. Weng, S. Whitfield-Gabrieli, P. Williamson, C. Windischberger, Y. F. Zang, H. Y. Zhang, F. X. Castellanos, and M. P. Milham. Toward discovery science of human brain function. *Proc Natl Acad Sci U S A*, 107(10):4734–9, 2010. ISSN 0027-8424. doi: 10.1073/pnas.0911855107.

33. J. H. Adams, D. Doyle, I. Ford, T. A. Gennarelli, D. I. Graham, and D. R. McLellan. Diffuse axonal injury in head injury: definition, diagnosis and grading. *Histopathology*, 15(1):49–59, 1989. ISSN 0309-0167 (Print) 0309-0167. doi: 10.1111/j.1365-2559.1989.tb03040.x.

34. P. C. Blumbergs, G. Scott, J. Manavis, H. Wainwright, D. A. Simpson, and A. J. McLean. Staining of amyloid precursor protein to study axonal damage in mild head injury. *Lancet*, 344(8929):1055–6, 1994. ISSN 0140-6736 (Print) 0140-6736. doi: 10.1016/s0140-6736(94)91712-4.

35. M. M. Mesulam. From sensation to cognition. *Brain*, 121 (Pt 6):1013–52, 1998. ISSN 0006-8950. doi: 10.1093/brain/121.6.1013.

36. V. Bonnelle, T. E. Ham, R. Leech, K. M. Kinnunen, M. A. Mehta, R. J. Greenwood, and D. J. Sharp. Salience network integrity predicts default mode network function after traumatic brain injury. *Proc Natl Acad Sci U S A*, 109(12):4690–5, 2012. ISSN 1091-6490 (Electronic) 0027-8424 (Linking). doi: 10.1073/pnas.1113455109.

37. J. C. Griffis, N. V. Metcalf, M. Corbetta, and G. L. Shulman. Structural disconnections explain brain network dysfunction after stroke. *Cell Rep*, 28(10):2527–2540.e9, 2019. doi: 10.1016/j.celrep.2019.07.100.

38. G. Deco and M. L. Kringelbach. Great expectations: using whole-brain computational connectomics for understanding neuropsychiatric disorders. *Neuron*, 84(5):892–905, 2014. ISSN 0896-6273. doi: 10.1016/j.neuron.2014.08.034.

39. M. C. Stevens, D. Lovejoy, J. Kim, H. Oakes, I. Kureshi, and S. T. Witt. Multiple resting state network functional connectivity abnormalities in mild traumatic brain injury. *Brain Imaging Behav*, 6(2):293–318, 2012. ISSN 1931-7557. doi: 10.1007/s11682-012-9157-4.

40. F. G. Hillary, J. Slocomb, E. C. Hills, N. M. Fitzpatrick, J. D. Medaglia, J. Wang, D. C. Good, and G. R. Wylie. Changes in resting connectivity during recovery from severe traumatic brain injury. *Int J Psychophysiol*, 82(1):115–23, 2011. ISSN 0167-8760. doi: 10.1016/j.ijpsycho.2011.03.011.

41. A. R. Mayer, M. V. Mannell, J. Ling, C. Gasparovic, and R. A. Yeo. Functional connectivity in mild traumatic brain injury. *Hum Brain Mapp*, 32(11):1825–35, 2011. ISSN 1097-0193 (Electronic) 1065-9471 (Linking). doi: 10.1002/hbm.21151.

42. E. Shumskaya, T. M. Andriessen, D. G. Norris, and P. E. Vos. Abnormal whole-brain functional networks in homogeneous acute mild traumatic brain injury. *Neurology*, 79(2): 175–82, 2012. ISSN 0028-3878. doi: 10.1212/WNL.0b013e31825f04fb.

43. A. A. Vakhtin, V. D. Calhoun, R. E. Jung, J. L. Prestopnik, P. A. Taylor, and C. C. Ford. Changes in intrinsic functional brain networks following blast-induced mild traumatic brain injury. *Brain Inj*, 27(11):1304–10, 2013. ISSN 0269-9052. doi: 10.3109/02699052.2013.823561.

44. D. J. Sharp, C. F. Beckmann, R. Greenwood, K. M. Kinnunen, V. Bonnelle, X. De Boissezon, J. H. Powell, S. J. Counsell, M. C. Patel, and R. Leech. Default mode network functional and structural connectivity after traumatic brain injury. *Brain*, 134(Pt 8):2233–47, 2011. ISSN 1460-2156 (Electronic) 0006-8950 (Linking). doi: 10.1093/brain/awr175.

45. Y. Zhou, M. P. Milham, Y. W. Lui, L. Miles, J. Reaume, D. K. Sodickson, R. I. Grossman, and Y. Ge. Default-mode network disruption in mild traumatic brain injury. *Radiology*, 265 (3):882–92, 2012. ISSN 0033-8419. doi: 10.1148/radiol.12120748.

46. D. J. Sharp, V. Bonnelle, X. De Boissezon, C. F. Beckmann, S. G. James, M. C. Patel, and M. A. Mehta. Distinct frontal systems for response inhibition, attentional capture, and error processing. *Proc Natl Acad Sci U S A*, 107(13):6106–11, 2010. ISSN 0027-8424. doi: 10.1073/pnas.1000175107.

47. D. Sridharan, D. J. Levitin, and V. Menon. A critical role for the right fronto-insular cortex in switching between central-executive and default-mode networks. *Proc Natl Acad Sci U S A*, 105(34):12569–74, 2008. ISSN 0027-8424. doi: 10.1073/pnas.0800005105.

48. F. G. Hillary, S. M. Rajtmajer, C. A. Roman, J. D. Medaglia, J. E. Slocomb-Dluzen, V. D. Calhoun, D. C. Good, and G. R. Wylie. The rich get richer: brain injury elicits hyperconnectivity in core subnetworks. *PLoS One*, 9(8):e104021, 2014. ISSN 1932-6203. doi: 10.1371/journal.pone.0104021.

49. F. G. Hillary, J. D. Medaglia, K. Gates, P. C. Molenaar, J. Slocomb, A. Peechatka, and D. C. Good. Examining working memory task acquisition in a disrupted neural network. *Brain*, 134(Pt 5):1555–70, 2011. ISSN 0006-8950. doi: 10.1093/brain/awr043.

50. A. Iraji, R. R. Benson, R. D. Welch, B. J. O'Neil, J. L. Woodard, S. I. Ayaz, A. Kulek, V. Mika, P. Medado, H. Soltanian-Zadeh, T. Liu, E. M. Haacke, and Z. Kou. Resting state functional connectivity in mild traumatic brain injury at the acute stage: Independent component and seed-based analyses. *J Neurotrauma*, 32(14):1031–45, 2015. ISSN 0897-7151. doi: 10.1089/neu.2014.3610.

51. R. D. Bharath, A. Munivenkatappa, S. Gohel, R. Panda, J. Saini, J. Rajeswaran, D. Shukla, I. D. Bhagavatula, and B. B. Biswal. Recovery of resting brain connectivity ensuing mild traumatic brain injury. *Front Hum Neurosci*, 9:513, 2015. ISSN 1662-5161 (Print) 1662-5161. doi: 10.3389/fnhum.2015.00513.

52. B. Johnson, K. Zhang, M. Gay, S. Horovitz, M. Hallett, W. Sebastianelli, and S. Slobounov. Alteration of brain default network in subacute phase of injury in concussed individuals: resting-state fmri study. *Neuroimage*, 59(1):511–8, 2012. ISSN 1053-8119. doi: 10.1016/j.neuroimage.2011.07.081.

53. F. G. Hillary and J. H. Grafman. Injured brains and adaptive networks: The benefits and costs of hyperconnectivity. *Trends Cogn Sci*, 21(5):385–401, 2017. ISSN 1364-6613. doi: 10.1016/j.tics.2017.03.003.

54. J. A. Mortimer, C. M. van Duijn, V. Chandra, L. Fratiglioni, A. B. Graves, A. Heyman, A. F. Jorm, E. Kokmen, K. Kondo, W. A. Rocca, and et al. Head trauma as a risk factor for alzheimer's disease: a collaborative re-analysis of case-control studies. eurodem risk factors research group. *Int J Epidemiol*, 20 Suppl 2:S28–35, 1991. ISSN 0300-5771 (Print) 0300-5771. doi: 10.1093/ije/20.supplement_2.s28.

55. P. N. Nemetz, C. Leibson, J. M. Naessens, M. Beard, E. Kokmen, J. F. Annegers, and L. T. Kurland. Traumatic brain injury and time to onset of alzheimer's disease: a population-based study. *Am J Epidemiol*, 149(1):32–40, 1999. ISSN 0002-9262 (Print) 0002-9262. doi: 10.1093/oxfordjournals.aje.a009724.

56. R. L. Buckner, J. Sepulcre, T. Talukdar, F. M. Krienen, H. Liu, T. Hedden, J. R. Andrews-Hanna, R. A. Sperling, and K. A. Johnson. Cortical hubs revealed by intrinsic functional connectivity: mapping, assessment of stability, and relation to alzheimer's disease. *J Neurosci*, 29(6):1860–73, 2009. ISSN 0270-6474. doi: 10.1523/jneurosci.5062-08.2009.

57. A. G. Vlassenko, S. N. Vaishnavi, L. Couture, D. Sacco, B. J. Shannon, R. H. Mach, J. C. Morris, M. E. Raichle, and M. A. Mintun. Spatial correlation between brain aerobic glycolysis and amyloid-beta (abeta) deposition. *Proc Natl Acad Sci U S A*, 107(41):17763–7, 2010. ISSN 0027-8424. doi: 10.1073/pnas.1010461107.

58. N. Myers, L. Pasquini, J. Gottler, T. Grimmer, K. Koch, M. Ortner, J. Neitzel, M. Muhlau, S. Forster, A. Kurz, H. Forstl, C. Zimmer, A. M. Wohlschlager, V. Riedl, A. Drzezga, and C. Sorg. Within-patient correspondence of amyloid-beta and intrinsic network connectivity in alzheimer's disease. *Brain*, 137(Pt 7):2052–64, 2014. ISSN 0006-8950. doi: 10.1093/brain/awu103.

59. K. Han, D. Martinez, S. B. Chapman, and D. C. Krawczyk. Neural correlates of reduced depressive symptoms following cognitive training for chronic traumatic brain injury. *Hum Brain Mapp*, 39(7):2955–2971, 2018. ISSN 1065-9471. doi: 10.1002/hbm.24052.

60. B. Dubois, A. Slachevsky, I. Litvan, and B. Pillon. The fab: a frontal assessment battery at bedside. *Neurology*, 55(11):1621–6, 2000. ISSN 0028-3878 (Print) 0028-3878. doi: 10.1212/wnl.55.11.1621.

61. D. Wechsler. *Wechsler Adult Intelligence Scale, 4th ed. Pearson Assessment (Finnish*

version: Psykologien kustannus oy, Helsinki, 2012). 2008.

62. D. Wechsler. *Wechsler Memory Scale, 3rd ed. San Antonio: The Psychological Corporation (Finnish version: Psykologien kustannus oy, Helsinki, 2008).* 1997.

63. N. Martin, F. Kohen, M. Kalinyak-Fliszar, A. Soveri, and M. Laine. Effects of working memory load on processing of sounds and meanings of words in aphasia. *Aphasiology*, 26(3-4):462–493, 2012. ISSN 0268-7038 (Print) 0268-7038. doi: 10.1080/02687038.2011.619516.

64. K. J. Pallesen, E. Brattico, C. J. Bailey, A. Korvenoja, J. Koivisto, A. Gjedde, and S. Carlson. Cognitive control in auditory working memory is enhanced in musicians. *PLoS One*, 5(6):e11120, 2010. ISSN 1932-6203. doi: 10.1371/journal.pone.0011120.

65. J. R. Simon and A. P. Rudell. Auditory s-r compatibility: the effect of an irrelevant cue on information processing. *J Appl Psychol*, 51(3):300–4, 1967. ISSN 0021-9010 (Print) 0021-9010. doi: 10.1037/h0020586.

66. I. H. Robertson, T. Manly, J. Andrade, B. T. Baddeley, and J. Yiend. 'oops!': performance correlates of everyday attentional failures in traumatic brain injured and normal subjects. *Neuropsychologia*, 35(6):747–58, 1997. ISSN 0028-3932 (Print) 0028-3932. doi: 10.1016/s0028-3932(97)00015-8.

67. Isquith P.K. Gioia G.A. Roth, R.M. *Behaviour Rating Inventory of executive Function - Adult Version (BRIEF-A). Lutz, FL: Psychological Assessment Resources, Inc.* 2005.

68. A. Miyake, N. P. Friedman, M. J. Emerson, A. H. Witzki, A. Howerter, and T. D. Wager. The unity and diversity of executive functions and their contributions to complex "frontal lobe" tasks: a latent variable analysis. *Cogn Psychol*, 41(1):49–100, 2000. ISSN 0010-0285 (Print) 0010-0285. doi: 10.1006/cogp.1999.0734.

69. M. Brett, A. P. Leff, C. Rorden, and J. Ashburner. Spatial normalization of brain images with focal lesions using cost function masking. *Neuroimage*, 14(2):486–500, 2001. ISSN 1053-8119 (Print) 1053-8119 (Linking). doi: 10.1006/nimg.2001.0845.

70. J. Ashburner and K. J. Friston. Unified segmentation. *Neuroimage*, 26(3):839–51, 2005. ISSN 1053-8119 (Print) 1053-8119 (Linking). doi: 10.1016/j.neuroimage.2005.02.018.

71. S. M. Andersen, S. Z. Rapcsak, and P. M. Beeson. Cost function masking during normalization of brains with focal lesions: still a necessity? *Neuroimage*, 53(1):78–84, 2010. ISSN 1095-9572 (Electronic) 1053-8119 (Linking). doi: 10.1016/j.neuroimage.2010.06.003.

72. Y. Behzadi, K. Restom, J. Liau, and T. T. Liu. A component based noise correction method (compcor) for bold and perfusion based fmri. *Neuroimage*, 37(1):90–101, 2007. ISSN 1053-8119 (Print) 1053-8119. doi: 10.1016/j.neuroimage.2007.04.042.

73. K. J. Friston, S. Williams, R. Howard, R. S. Frackowiak, and R. Turner. Movement-related effects in fmri time-series. *Magn Reson Med*, 35(3):346–55, 1996. ISSN 0740-3194 (Print) 0740-3194. doi: 10.1002/mrm.1910350312.

74. J. D. Power, A. Mitra, T. O. Laumann, A. Z. Snyder, B. L. Schlaggar, and S. E. Petersen. Methods to detect, characterize, and remove motion artifact in resting state fmri. *Neuroimage*, 84:320–41, 2014. ISSN 1053-8119. doi: 10.1016/j.neuroimage.2013.08.048.

75. S. Whitfield-Gabrieli and A. Nieto-Castanon. Conn: a functional connectivity toolbox for correlated and anticorrelated brain networks. *Brain Connect*, 2(3):125–41, 2012. ISSN 2158-0014. doi: 10.1089/brain.2012.0073.

76. K. Han, R. A. Davis, S. B. Chapman, and D. C. Krawczyk. Strategy-based reasoning training modulates cortical thickness and resting-state functional connectivity in adults with chronic traumatic brain injury. *Brain Behav*, 7(5):e00687, 2017. doi: 10.1002/brb3.687.

77. A. Nieto-Castanon. *Handbook of unctional connectivity Magnetic Resonance Imaging Methods in CONN.* Hilbert-Press, Boston, MA, 2020.

78. K. J. Worsley, S. Marrett, P. Neelin, A. C. Vandal, K. J. Friston, and A. C. Evans. A unified statistical approach for determining significant signals in images of cerebral activation. *Hum Brain Mapp*, 4(1):58–73, 1996. ISSN 1065-9471 (Print) 1065-9471. doi: 10.1002/(sici)1097-0193(1996)4:1<58::Aid-hbm4>3.0.Co;2-o.

79. J. Chumbley, K. Worsley, G. Flandin, and K. Friston. Topological fdr for neuroimaging. *Neuroimage*, 49(4):3057–64, 2010. ISSN 1053-8119. doi: 10.1016/j.neuroimage.2009.10.090.

80. T. Sarkamo, M. Tervaniemi, and M. Huotilainen. Music perception and cognition: development, neural basis, and rehabilitative use of music. *Wiley Interdisc Rev Cogn Sci*, 4(4):441–451, 2013. ISSN 1939-5078 (Print) 1939-5078 (Linking). doi: 10.1002/wcs.1237.

81. C. M. Lewis, A. Baldassarre, G. Committeri, G. L. Romani, and M. Corbetta. Learning sculpts the spontaneous activity of the resting human brain. *Proc Natl Acad Sci U S A*, 106(41):17558–63, 2009. ISSN 0027-8424 (Print). doi: 10.1073/pnas.0902455106.

82. J. Sepulcre, M. R. Sabuncu, T. B. Yeo, H. Liu, and K. A. Johnson. Stepwise connectivity of the modal cortex reveals the multimodal organization of the human brain. *J Neurosci*, 32(31):10649–61, 2012. ISSN 0270-6474. doi: 10.1523/jneurosci.0759-12.2012.

83. M. Corbetta, G. Patel, and G. L. Shulman. The reorienting system of the human brain: from environment to theory of mind. *Neuron*, 58(3):306–24, 2008. ISSN 0896-6273. doi: 10.1016/j.neuron.2008.04.017.

84. M. L. Dixon, A. De La Vega, C. Mills, J. Andrews-Hanna, R. N. Spreng, M. W. Cole, and K. Christoff. Heterogeneity within the frontoparietal control network and its relationship to the default and dorsal attention networks. *Proc Natl Acad Sci U S A*, 115(7):E1598–e1607, 2018. ISSN 0027-8424. doi: 10.1073/pnas.1715766115.

85. F. G. Hillary, H. M. Genova, N. D. Chiaravalloti, B. Rypma, and J. DeLuca. Prefrontal modulation of working memory performance in brain injury and disease. *Hum Brain Mapp*, 27(11):837–47, 2006. ISSN 1065-9471 (Print) 1065-9471. doi: 10.1002/hbm.20226.

86. J. D. Medaglia, K. S. Chiou, J. Slocomb, N. M. Fitzpatrick, B. M. Wardecker, D. Ramanathan, J. Vesek, D. C. Good, and F. G. Hillary. The less bold, the wiser: support for the latent resource hypothesis after traumatic brain injury. *Hum Brain Mapp*, 33(4):979–93, 2012. ISSN 1065-9471. doi: 10.1002/hbm.21264.

87. L. E. Nordin, M. C. Möller, P. Julin, A. Bartfai, F. Hashim, and T. Q. Li. Post mtbi fatigue is associated with abnormal brain functional connectivity. *Sci Rep*, 6, 2016. doi: 10.1038/srep21183.

88. A. Roy, R. A. Bernier, J. Wang, M. Benson, Jr. French, J. J., D. C. Good, and F. G. Hillary. The evolution of cost-efficiency in neural networks during recovery from traumatic brain injury. *PLoS One*, 12(4):e0170541, 2017. ISSN 1932-6203. doi: 10.1371/journal.pone.0170541.

89. A. Iraji, H. Chen, N. Wiseman, R. D. Welch, B. J. O'Neil, E. M. Haacke, T. Liu, and Z. Kou.

90. L. Li, X. Hu, T. M. Preuss, M. F. Glasser, F. W. Damen, Y. Qiu, and J. Rilling. Mapping putative hubs in human, chimpanzee and rhesus macaque connectomes via diffusion tractography. *Neuroimage*, 80:462–74, 2013. ISSN 1053-8119. doi: 10.1016/j.neuroimage.2013.04.024.

91. M. P. van den Heuvel and O. Sporns. Network hubs in the human brain. *Trends Cogn Sci*, 17(12):683–96, 2013. ISSN 1364-6613. doi: 10.1016/j.tics.2013.09.012.

92. E. H. Nijhuis, A. M. van Cappellen van Walsum, and D. G. Norris. Topographic hub maps of the human structural neocortical network. *PLoS One*, 8(6):e65511, 2013. ISSN 1932-6203. doi: 10.1371/journal.pone.0065511.

93. A. Zalesky, A. Fornito, I. H. Harding, L. Cocchi, M. Yucel, C. Pantelis, and E. T. Bullmore. Whole-brain anatomical networks: does the choice of nodes matter? *Neuroimage*, 50(3):970–83, 2010. ISSN 1053-8119. doi: 10.1016/j.neuroimage.2009.12.027.

94. M. W. Cole, S. Pathak, and W. Schneider. Identifying the brain's most globally connected regions. *Neuroimage*, 49(4):3132–48, 2010. ISSN 1053-8119. doi: 10.1016/j.neuroimage.2009.11.001.

95. X. N. Zuo, R. Ehmke, M. Mennes, D. Imperati, F. X. Castellanos, O. Sporns, and M. P. Milham. Network centrality in the human functional connectome. *Cereb Cortex*, 22(8):1862–75, 2012. ISSN 1047-3211. doi: 10.1093/cercor/bhr269.

96. D. Tomasi and N. D. Volkow. Association between functional connectivity hubs and brain networks. *Cereb Cortex*, 21(9):2003–13, 2011. ISSN 1047-3211. doi: 10.1093/cercor/bhq268.

97. M. P. van den Heuvel and O. Sporns. An anatomical substrate for integration among functional networks in human cortex. *J Neurosci*, 33(36):14489–500, 2013. ISSN 0270-6474. doi: 10.1523/jneurosci.2128-13.2013.

98. P. Bermudez, J. P. Lerch, A. C. Evans, and R. J. Zatorre. Neuroanatomical correlates of musicianship as revealed by cortical thickness and voxel-based morphometry. *Cereb Cortex*, 19(7):1583–96, 2009. ISSN 1460-2199 (Electronic) 1047-3211 (Linking). doi: 10.1093/cercor/bhn196.

99. Anna M. Zamorano, Ignacio Cifre, Pedro Montoya, Inmaculada Riquelme, and Boris Kleber. Insula-based networks in professional musicians: Evidence for increased functional connectivity during resting state fmri. *Human Brain Mapping*, 38(10):4834–4849, 2017. ISSN 1065-9471. doi: 10.1002/hbm.23682.

100. S. Koelsch, T. Fritz, V. Cramon DY, K. Muller, and A. D. Friederici. Investigating emotion with music: an fmri study. *Hum Brain Mapp*, 27(3):239–50, 2006. ISSN 1065-9471 (Print) 1065-9471 (Linking). doi: 10.1002/hbm.20180.

101. V. K. M. Cheung, L. Meyer, A. D. Friederici, and S. Koelsch. The right inferior frontal gyrus processes nested non-local dependencies in music. *Sci Rep*, 8(1):3822, 2018. ISSN 2045-2322 (Electronic) 2045-2322 (Linking). doi: 10.1038/s41598-018-22144-9.

102. R. Bianco, G. Novembre, P. E. Keller, S. G. Kim, F. Scharf, A. D. Friederici, A. Villringer, and D. Sammler. Neural networks for harmonic structure in music perception and action. *Neuroimage*, 142:454–464, 2016. ISSN 1095-9572 (Electronic) 1053-8119 (Linking). doi: 10.1016/j.neuroimage.2016.08.025.

103. I. Royal, D. T. Vuvan, B. R. Zendel, N. Robitaille, M. Schonwiesner, and I. Peretz. Activation in the right inferior parietal lobule reflects the representation of musical structure beyond simple pitch discrimination. *PLoS One*, 11(5):e0155291, 2016. ISSN 1932-6203. doi: 10.1371/journal.pone.0155291.

104. M. Catani and M. Thiebaut de Schotten. A diffusion tensor imaging tractography atlas for virtual in vivo dissections. *Cortex*, 44(8):1105–32, 2008. ISSN 0010-9452 (Print) 0010-9452 (Linking). doi: 10.1016/j.cortex.2008.05.004.

Compensation through functional hyperconnectivity: A longitudinal connectome assessment of mild traumatic brain injury. *Neural Plast*, 2016:4072402, 2016. ISSN 1687-5443. doi: 10.1155/2016/4072402.