# Strength of attention-sampling parietal EEG theta rhythm is linked to impaired inhibition in adult ADHD

Kristiina Juurmaa[1], Jussi Palomäki[1], and Benjamin Ultan Cowley[1,2,✉]

[1]Cognitive Science, Department of Digital Humanities, Faulty of Arts, University of Helsinki, Helsinki, Finland
[2]Educational Data Science, Faculty of Educational Sciences, University of Helsinki, Helsinki, Finland

**Attention-deficit hyperactivity disorder (ADHD) in adults is understudied, especially regarding neural mechanisms such as oscillatory control of attention sampling. We report an EEG study of such cortical oscillations, in ADHD-diagnosed adults taking a continuous performance test that measures the ability to sustain attention and inhibit impulsivity for a prolonged period of time.**

**We recorded 53 adults (28f, 25m, aged 18-60), and 18 matched healthy controls, using 128-channel EEG. We analysed features with established links to neural correlates of attention: event-related (de)synchronisation (ERS), alpha and theta frequency band activation, and stimulus-locking indices; in frontal and parietal scalp regions.**

**Test performance distinguished healthy controls from ADHD adults. The ADHD group manifested significantly less parietal 4 Hz theta ERS during correct inhibition trials, in addition to having greater sensitivity to targets in stimulus-locking measures.**

**Our results suggest that ADHD adults have impaired attention sampling in relational categorisation tasks.**

**ADHD, adult, EEG, inhibition, attention, TOVA**

**Correspondence:** *ben.cowley@helsinki.fi*

## Introduction

Attention-deficit hyperactivity disorder (ADHD) is among the most common childhood psychiatric disorders. ADHD is marked by persistent, age-inappropriate levels of inattention, hyperactivity and/or impulsivity. While the prevalence of ADHD declines with age, the condition persists into adulthood in approximately one third of those diagnosed as a child (1). There is thus a need for more research to study how the disorder manifests in adults. Here, we investigate how the neural correlates of attention differ between adults diagnosed with ADHD and healthy controls.

Both the clinical and scientific pictures of ADHD remain complex and multidimensional (2). Clinical diagnosis of ADHD operates at the level of heterogenous behavioural symptoms, subjectively assessed (3–5), with three subtypes of ADHD listed in diagnostic manuals: predominantly inattentive (ADHD-I, or ADD), predominantly hyperactive/impulsive (ADHD-HI), and combined (ADHD-C). Although diagnosis does not include behavioural attention testing, there have been efforts to develop indicative behavioural tests for ADHD (6). As yet, there remains no one-size-fits-all test that reliably classifies ADHD. Because of this, it is important to search for neurocognitive mechanisms that explain aspects of the disorder - mutual information from multiple mechanisms (e.g. behavioural and neural) could improve treatment targeting efficacy (7).

ADHD is characterised by deficits in completing cognitive tasks that challenge self-regulation of attention over extended durations. Thus, it is relevant to examine the interplay between sustained attention and inhibition, especially via the underpinning neural processes, such as regulation of perceptual processing by (dorsal or ventral) attention networks (8). High time-resolution, magneto- and electro-encephalography (M/EEG) studies have been vital in understanding the neurophysiology of ADHD by eliciting the mechanisms of attention. For example, cortical oscillations (i.e. rhythmic patterns of neural activity) have long been suggested to play a role in communication between areas of the brain (9), and have been shown to be altered in ADHD subjects (10) by measurement of event-related synchronisation.

One influential model says that ADHD is associated with deficient inhibitory control, which affects other executive functions (EF) such as sustained attention and working memory (11). The deficits in attention, inhibition, and working memory suggest that activity in the frontal cortico-striatal network and frontoparietal attention network are altered in ADHD (12), and may be linked to aberrant default mode network (DMN) suppression (13), or alpha-band oscillation suppression (14). Both volumetric alterations and changes in task-dependent BOLD (blood-oxygen level dependent) signals in prefrontal-striatal circuits support this view (15). Mowinckel et al. (16) also found causal support for these ideas from fMRI functional connectivity analysis of a decision-making task in a methylphenidate intervention. They illustrated that aberrant within-network connectivity was alleviated by treatment, but reduced sustained DMN suppression was not – implying that adults with ADHD may continue to experience debilitating effects of increased neuroenergetic demand over longer periods (attention fatigue), even when medication helps to normalise short-term performance (as predicted by the 'cognitive-energetic model' of ADHD (17)).

Sustained attention may be coordinated by patterns of synchronous oscillatory neural activation (18) – which are therefore a compelling target of study through several related measurements, including power, phase, and synchronisation.

Decline in sustained performance has been associated with

changes in the ratio of frontomedial theta (4-8 Hz) to occipital alpha (8-12 Hz) power [18]. This could possibly be due to reduced attentional sampling (at theta frequencies) and DMN suppression (at alpha frequencies), driven by neuroenergetic fatigue [17]. Errors due to attentional lapses are sometimes followed by decreases in alpha power [19], which might function to refocus attention on the task at hand and inhibit task irrelevant processes [18].

Moreover, increased frontomedial theta power has been linked to monitoring of task-relevant activities and cognitive control, both of which are implicated in sustained attention [18, 20, 21]. Frontomedial theta increase may also reflect mental fatigue [22], which consequently challenges the ability to sustain attention. Frontomedial theta power tends to increase during sustained performance tasks (for example, Wascher et al. [22]'s 4h-duration spatial stimulus-response task), correlates with increases in commission errors (incorrect responses) and reaction times (RT), and predicts post-error RT slowing [18].

Klimesch [9] proposed that reduced alpha desynchronisation is related to attentional control and inhibition, and theta synchronisation to task-specific semantic processing. Alpha oscillations tend to reduce in amplitude (due to neural desynchronisation) during effortful cognitive processing; when event-driven, this is called event-related desynchronisation of alpha (ERD). In contrast, theta oscillation amplitudes tend to increase with increasing cognitive effort [23], i.e. event-related synchronisation of theta (ERS) [24, 25].

Oscillations can become entrained (phase-locked) to predictable stimulus onsets. That is, the oscillatory phase (that facilitates the processing of subsequent stimuli) becomes more likely to coincide with the expected stimuli [26]. This entrainment can be observed already in the pre-stimulus period [10, 27]. Calderone et al. [10] suggest studying selective attention as indicated by oscillatory entrainment in clinical conditions such as ADHD. In lab conditions people with ADHD tend to be ill-prepared for incoming stimuli that may require a response [28]. Several studies have linked the entrainment of pre-stimulus alpha oscillations to fluctuations in visual awareness and sustained attention [29–35].

**Attention Testing**   Continuous Performance Tests (CPTs) are developed to measure sustained attention and inhibitory control. Among CPTs, the Test of Variables of Attention (TOVA) is considered a gold standard to detect attention problems typical for ADHD. TOVA is used on all age groups, both clinically and academically. It tests inhibition and the ability to sustain attention, with a monotonous hybrid Go/NoGo target classification task [6, 36], where subjects must respond to targets and inhibit response to non-targets – see Figure 1. TOVA software performs normative classification of test subjects' performance as "not within normal limits", "borderline" or "normal". Many studies have used TOVA to study ADHD-diagnosed individuals *behaviourally*, primarily among children (for recent work see [37, 38]). Far fewer studies have measured EEG activity during TOVA [39, 40], and of these the only one focused on adults in par-

ticular was examining substance use disorder, not ADHD.

Some previous studies have examined EEG activity during other CPTs while comparing individuals with ADHD to healthy controls. Loo et al. [41] found increased cortical arousal in the ADHD group during Conners' CPT [42, 43], indicated by attenuated frontal and parietal alpha power. This effect increased towards the end of the task. However, they did not find any differences between groups in theta power.

**Research Questions**   The aim of the present study is to examine the behavioural performance and cortical oscillations (measured with EEG) during TOVA, comparing adults with ADHD to a healthy control group. In particular, this study explores (a) how neural correlates of visual attention differ between groups throughout TOVA, and (b) how the increasing demands of sustained attention during TOVA affect oscillatory EEG and behavioural performance in adults with ADHD. We focus on the following research questions (RQs):

**RQ1**: Are there between-groups differences in neural correlates of visual attention? Specifically, we examine parietal event-related oscillatory power perturbations (ERSP) to study frequencies linked to attentional regulation, particularly alpha and theta.

**RQ2**: Does frontomedial theta power change during TOVA; is there a difference between the ADHD group and controls, or a group × time segment (first, second, or third half of the TOVA task) interaction?

**RQ3**: Does parietal or frontal alpha power change during TOVA; is there a difference between the ADHD group and controls, or a group × time segment interaction?

**RQ4**: Does pre-stimulus occipital alpha power change during TOVA; are there differences in pre-stimulus occipital alpha power between groups or between TOVA conditions (infrequent vs. frequent target mode), or a group × TOVA condition interaction?

## Methods

The measurement consisted of visual TOVA CPT with EEG recording, gathered as part of a larger project detailed in Cowley et al. [44]. An ethical approval was obtained from the Ethical Committee of the Hospital District of Helsinki and Uusimaa. Each participant gave informed consent in accordance with the Declaration of Helsinki.

**Participants.** We recruited 53 adults (25 males, age M=36.26, SD=10.22) diagnosed with either ADHD (n=44) or ADD (n=9) – we refer to both as ADHD group – and 18 adults (6 males, age M=32.78, SD=10.82) with no diagnosed neuro-cognitive deficits or ongoing medication for ADHD/ADD as a healthy control group. The healthy controls were significantly more likely to have TOVA performance within normal limits compared to ADHD adults (as detailed in the results section). All participants had normal or corrected-to-normal vision. The groups did not differ in terms of age, gender, or handedness. Due to reasons of

data quality (detailed below) a number of participants were dropped from different EEG analysis stages. The final analysed number of participants ranged from 39 to 49 in the ADHD group, and from 14 to 18 in the control group. The control group size was always at least 33% of the size of the ADHD group, often closer to 40%, and thus the statistical power of our tests is not substantially affected by this size disparity.[1]

For the current study, inclusion criteria for the ADHD group were 1) pre-existing diagnosis of ADHD/ADD, 2) no neurological diagnoses, 3) age between 18-60, 4) scores on Adult ADHD Self Report Scale (ASRS) (45) and Brown ADHD scale (BADDS) (46) indicating the presence of ADHD, and 5) an IQ score of at least 80 using WAIS IV measured by a qualified psychologist (47). No strict cut-off values were used for ASRS and BADDS to indicate the presence of ADHD/ADD. Instead, exclusion was decided by the consulting psychiatrist, who conducted structured clinical interviews with participants as per the guidelines of the Diagnostic Interview for ADHD in Adults (DIVA 2.0, (48)). Comorbidities were evaluated during the clinical interview, and exclusion criteria included outlier scores in scores of Generalized Anxiety Disorder (49), Beck Depression Inventory (50), Alcohol Use Disorders Identification Test (51), the Mood Disorder Questionnaire (52), test of prodromal symptoms of psychosis (53), and the Dissociative Experiences Scale (54). The psychiatrist made a final assessment of the balance of symptoms contributing to patients' presentation.

Within the ADHD group, there was a small difference in the ASRS hyperactivity-impulsivity scores depending on the diagnosis ($F(1,50)$=5.01, $p$<.030, $r^2$=.09). As expected, participants with ADHD-HI or ADHD-C diagnoses had higher hyperactivity-impulsivity scores (M=6.33, SD=2.56; the groups were combined) than those with an ADHD-I diagnosis (without hyperactivity-impulsivity; M=4.22, SD=2.59). There was no significant difference in the ASRS inattention scores between groups, and most analyses thus treat all ADHD subjects as one group.

### Procedure.

The behavioural and EEG data were measured in a 2.5–3 hour multi-task session, gathered in an electrically-shielded and sound-attenuated room. TOVA was administered towards the beginning of the measurement session (as suggested by Greenberg and Hughes (55, 24)). The full session included preparation (30-40 min), pre-test baseline measurement (5 min), TOVA (22 min), resting state vigilance measurement (20 min), a novel CPT (30 mins; designed by (56)) and a post-test baseline measurement (2 mins).

Participants were asked to abstain from taking their ADHD medication for 48 hours prior to the EEG measurement (washout period). They were also advised not to take any other stimulants immediately prior to the measurement (e.g.

---

[1]See www.markhw.com/blog/control-size for a simulation analysis which suggests that a control group size 30% as large as test is comfortable

**Fig. 1.** TOVA experiment protocol design. Above: number of targets and non-targets in each quarter and half, with target-to-nontarget frequency and total time. Below: individual trial structure, timings, and stimuli appearance.

coffee, cigarettes, energy drinks) and to arrive as well rested as possible to the measurement.

Karolinska sleepiness scale (KSS) (57) was employed to control for participants' sleepiness levels, as sleepiness can affect both sustained attention and EEG measurement. Participants tended to report being alert (on the scale 1-9, ADHD: M=4.60, SD=1.38, controls: M=3.73, SD=1.39; these mean scores indicated 3 "Alert", to 4 "Rather Alert", and 5 "Neither alert nor sleepy").

**TOVA.** We administered TOVA visual version 8, consisting of the test software, USB relay hardware, a low-latency microswitch hardware response button, and Synchronization Interface hardware for test-to-EEG amplifier synchrony (all products of the TOVA Company).

TOVA presents a target and a non-target stimulus, shown in Figure 1. The participant's task is to respond to targets by a button press of their dominant hand, and refrain from responding to non-targets; i.e. it has a Go/NoGo design. The participants are advised to respond as accurately and quickly as possible. A practice test of approximately 20 trials is administered before the actual test. Trials can be: correct responses, commission errors (incorrect responses), omission errors (incorrect inhibition) and correct inhibition.

TOVA has two consecutive conditions: infrequent and frequent target mode. During the first half (H1), the target stimulus appears infrequently (72 targets, and 252 non-targets, i.e. 22.5% target probability). During the second half (H2), these frequencies are reversed (252 targets and 72 non-targets, i.e. 77.5% target probability). Participants are not informed about this transition. Each stimulus is presented for 100 ms and separated from the next stimulus by a 2000 ms interstimulus interval (ISI). The anticipatory cutoff time from stimulus onset is 150 ms, meaning that responses given 0-150 ms after stimulus onset are considered invalid. The duration of TOVA is 21.6 minutes, each condition lasting 10.8 min.

***TOVA Variables.*** Analysis was performed on standardised TOVA scores, which are formed by comparing individual performance against a normed sample population (N=1596) (55). The comparison group for each participant is determined by their age group and gender.

Five standardised scores are available: 1) mean response time (RT) – mean of correct response times in milliseconds (ms); 2) response time variability (RTV) – standard deviation of mean correct RT; 3) commission errors – incorrect responses to a non-target stimulus; 4) omission errors – incorrect responses to a target stimulus; and 5) d' (d prime) – ratio of hits (correct responses) to false alarms (commission errors). One participant who showed strong evidence of symptom exaggeration (Symptom Exaggeration Index, SEI) was excluded from further analyses.

For normative comparison, TOVA is advised to be measured between 6h00 – 13h00. Our group-average measurement times were close to or within the normed hours: ADHD M=13h45, SD=2h26 (35 after 13h00); control M=12h49, SD=3h02 (7 after 13h00), and the starting times did not statistically differ between the groups.

**EEG measurement and preprocessing.** EEG was measured using Biosemi ActiveTwo equipment with 128 active electrodes, which is similar to the International 10-20 electrode placement system. Active electrode CMS (Common Mode Sense) and passive electrode DRL (Driven Right Leg) were used to create a feedback loop for amplifier reference. Electro-oculography (EOG) was recorded using bipolar montage: horizontal EOG electrodes were attached to the outer canthi of both eyes; vertical EOG electrodes were attached above and below the left eye. Electrode offsets (running average of voltage at each electrode) were kept below +/- 25mV.

During the TOVA, participants were advised to relax and sit as still as possible, avoiding excessive movements, and to fixate the middle of the screen shown as a small white dot between stimulus presentations. Three participants were excluded due to technical problems with event codes in the EEG data.

The data was preprocessed using Computational Testing Automated Preprocessing toolbox (CTAP) (58, 59), based on EEGLAB (60) for Matlab. The data was re-referenced offine to the average of the two mastoids. The data was then low-pass filtered at 45 Hz and high-pass filtered at 2 Hz. CTAP was used to detect eye blinks using the method detailed in Cowley et al. (58). Validity of eye blinks detected by CTAP were examined for each participant by visual inspection.

Each participant's continuous EEG and EOG data was decomposed using the FastICA algorithm (61). Independent Components (ICs) matching the CTAP-detected blinks were removed (58). Contaminated channels, epochs, and artifactual ICs were identified using FASTER (Fully Automated Statistical Thresholding for EEG artefact Rejection) methods (62). Data for automatically rejected channels was interpolated from adjacent electrodes. Fluctuations exceeding +/-80 microV in amplitude were discarded as bad segments. Any remaining ocular- or muscular-artifact ICs were detected and rejected manually by joint visual inspection of IC activations, scalp maps, power spectrum, and ERP image. On average 3% of channels and one IC per participant were rejected.

*Epoching for oscillatory power analyses.* For event-related analyses, the EEG data were extracted into four types of epochs: commission errors, correct responses, omission errors and correct inhibition. Each of these epochs lasted 2000 ms, starting from 1000 ms before stimulus onset to 900 ms after stimulus offset. Epochs that contained a blink (as detected by CTAP) starting from 400ms before stimulus onset to 600ms after stimulus onset, were rejected. After this, too few error trials remained to conduct statistical analyses, thus analyses were run only on correct (inhibition and response) trials. Two participants (one in each group) were completely excluded due high rate of epoch rejection. A baseline correction was applied to each epoch (from 1000 to 900 ms before stimulus onset) to counterbalance fluctuations in voltage offset across epochs and participants.

*Oscillatory power analyses.* Logarithmic mean oscillatory power was calculated using Welch's power spectral density estimation to calculate the mean log power spectrum (via EEGLAB's spectopo function). Oscillatory power was calculated for each participant within three *time-segments* of the un-epoched data: during the first 5 minutes of TOVA (infrequent mode), during the middle 5 minutes (between infrequent and frequent mode), and during the last 5 minutes (frequent mode). Power was calculated for each 1 Hz frequency bin separately within the theta band (4–7 Hz) and within the extended alpha band (8–12 Hz).

For the theta band, power from electrodes in the frontomedial region (FCz, Fz, AFz, Fpz) were averaged together. For the alpha band, power from electrodes in the frontal region (F3, F4 and Fz) and the parietal region (P3, P4, Pz) were grouped and averaged within each region. In order to minimise the problem of multiple implicit comparisons, these electrode locations were chose a priori based on previous research on parieto-occipital alpha and frontomedial theta (18, 20, 21, 41). These calculations of alpha power follow the method of Loo et al. (41).

To examine the effect of TOVA condition on pre-stimulus activity, alpha power was also calculated for the 500 ms pre-stimulus period preceding correct trials for the two TOVA conditions, H1 and H2. Among all correct trials (both correct inhibition and correct response) 102 trials were randomly chosen among H1 and H2 separately for each participant. Some participants were excluded for these specific analyses due to a low number of usable trials.

Within these correct trials, pre-stimulus power was calculated for each 1 Hz frequency bin separately within the extended alpha band (8–12 Hz). Power from electrodes in the parieto-occipital region (P3, P4, Pz, POz, Oz, O1, O2) was averaged together. The electrode locations were chosen a priori based on previous research (31, 32, 35, 63).

*Event-related Spectral Perturbations.* In addition to power calculations within different time windows, event-related spectral perturbation (ERSP) was calculated to examine event-related changes in oscillatory power dynamics. ERSP is a measure of average amplitude changes in given frequency

bands as a function of the time course of some experimental event (64). ERSPs were calculated for correct inhibition and correct response trials separately.

The number of trials were balanced to ensure that any observed effect would be due to altered neural processing, not merely due to possible differences in behavioural performance (e.g. amount of correct response trials). For each participant, 108 correct inhibition trials and 108 correct response trials were randomly selected. The same parieto-occipital electrode locations (P3, P4, Pz, POz, Oz, O1, O2) were used as for pre-stimulus alpha power calculations. Some participants were excluded from the ERSP analysis at this point, due to a low number of usable trials. The final number of available participants and trials available for statistical EEG analyses are summarised in the Appendix (Table A1).

***Response-aligned ERPs.*** Response-aligned ERP images at electrode Pz were generated for ADHD and control groups, by selecting 108 correct response trials randomly from each participant. A moving-window smoothing was applied: the control group had 1512 trials and window width 100; the ADHD group had more trials (4320) and a wider window of 285.

### Statistical analyses.

***Behavioural performance.*** Behavioural performance was analysed using a 2 x 2 two-way mixed design MANOVA, with Group (ADHD vs. control) as a between subjects factor and TOVA condition (infrequent vs. frequent) as a within-subjects factor. The DVs were mean RT, RTV, commission errors, omission errors and d' (means and standard deviations of these variables are presented in the Appendix, Table A2). These were followed up by further DV-specific analyses, which are detailed in the Results -section. We also assessed how well behavioral performance in TOVA differentiates between ADHD diagnosed adults and healthy controls.

Effect sizes for repeated measures were computed as partial eta squared. Multiple comparisons were adjusted with Bonferroni correction, keeping the alpha at .05. Possible extreme outlier values were assessed by examination of studentized residuals for values greater than +/-3. We do note, however, that excluding outlier scores in a clinical sample is not straightforward, as high behavioural variability itself has been reported as a clinical characteristic in adults with ADHD (3). In total four outlier scores were excluded from the analysis of standardised scores. Excluding them did not affect the interpretation of results, while including them would have violated the assumption of homogeneity of variance-covariance matrices. When necessary, variables with skewed distributions were log-transformed. However, group means and standard deviations are reported in original untransformed format, as these are more meaningful to interpret. The assumptions of repeated measures ANOVA were satisfied on transformed data without outliers.

***EEG data.*** Statistical analyses of EEG were run in Matlab (version 8.6), utilising EEGLAB toolbox (60) and in SPSS (version 21). Inspections of whether the data met the linear model assumptions (65) revealed moderate positive skewness and departures from normality (by Shapiro-Wilk) of alpha power. These were possible to correct by square root transformation; however the subsequent analyses produced very similar results for both original (logarithm-transformed mean power) and square root-transformed data. Therefore the results of analyses on original, logarithmic power are reported. Frontomedial theta power did not show departures from normality or skewness. Mauchly's test indicated that the assumption of sphericity had been violated for all frequencies ($p<.0005$). Degrees of freedom were corrected using Greenhouse-Geisser estimates of sphericity to counteract the inflation of Type I errors.

Three 2 x 3 mixed ANOVAs were used to determine whether there were group x time-segment interactions in: frontomedial theta power, and alpha power in parietal or frontal regions. Also, a 2 x 2 mixed ANOVA was calculated to examine if there was a group x TOVA condition (frequent vs. infrequent) interaction in parieto-occipital pre-stimulus alpha power on correct trials.

ERSP was calculated for each participant's correct response and correct inhibition trials to see if oscillatory power dynamics differed between the two groups during either type of trial. The statistical significance of the ERSPs were calculated using permutation tests (n=100) (60). False Discovery Rate (FDR) correction (66) was used to correct for multiple comparisons, maintaining the family-wise error rate at $p<.05$.

## Results

We will first describe results from our behavioural analyses, and then focus on EEG results separately for each research question (RQ); we present exploratory results when they arise.

**Behavioural performance.** Normed TOVA performance strongly distinguished diagnostic groups. Healthy controls were significantly more likely to have TOVA performance within normal limits compared to ADHD adults, as revealed by Pearson Chi-square (exact two-sided) test of independence ($\chi^2(1)=7.66$, $p<.007$).

In group-wise behavioural analysis (MANOVA), there was a significant TOVA condition × Group interaction ($F(5,61)=2.63$, $p<.032$, $\eta^2=.177$). That is, the impact that TOVA condition (H1 and H2) had on performance (as measured by the linear combination of mean RT, RT variability, commission errors, omission errors and d') depended on whether participants belonged to the ADHD group or controls. Thus, we followed up by examining the simple main effects of Group and TOVA condition. Behavioural results are illustrated in Figure 2.

First, we evaluated the difference between ADHD individuals and controls at the two TOVA conditions (H1 and H2) separately, by running two one-way MANOVAs for H1 and H2. The simple main effect of Group was not significant in

**Fig. 2.** Behavioural results. **Panel A-B**: standardised scores for commission errors and d', from TOVA first half. **Panel C**: response time variability standardised scores from TOVA second half. (Points in panels A-C shown with horizontal jitter for visibility). **Panel D-F**: rug-and-density plots show standardised scores for commission errors, d', omission errors, across both groups and TOVA halves. Red vertical lines show the data mean, 2.5%, and 97.5% quantiles.
$p < 0.05$ *, $0.01$ **, $0.001$ ***

either of these models ($Fs(5, 62) < 1.5$, $ps > .2$). We then ran univariate one-way ANOVAs separately for all DVs instead of using a multivariate approach, again separately for H1 and H2. Within H1, there was a significant between groups difference (Bonferroni-adjusted for multiple comparisons) in commission errors ($F(1,67)=4.12$, $p<.046$, $\eta^2=.059$) and in d' ($F(1,68)=5.30$, $p<.024$, $\eta^2=.074$): the ADHD group tended to produce more commission errors and had a worse d' (the ability to discriminate between targets and non-targets) than the control group. Within H2, the ADHD group had significantly greater RT variability ($F(1,68)=4.52$, $p<.037$, $\eta^2=.064$). See Table A3 in the supplementary materials for further details.

Next, we evaluated the difference between H1 and H2 (the difference in performance between the two consecutive conditions) separately for the ADHD- and control groups, by running two repeated measures MANOVAs. There was a significant simple main effect of TOVA condition in both the ADHD ($F(5,44)=8.96$, $p<.0005$, $\eta^2=.505$) and control group ($F(5,13)=7.63$, $p<.002$, $\eta^2=.746$): both groups performed worse towards the second half of TOVA. As before, we ran univariate repeated measures ANOVAs for each DV separately. Within the ADHD group, we found a significant effect of TOVA condition on commission errors ($F(1,48)=26.08$, $p<.0005$, $\eta^2=.352$), omission errors ($F(1,48)=11.15$, $p<.002$, $\eta^2=.189$) and d' ($F(1,48)=20.16$, $p<.0005$, $\eta^2=.296$): within the ADHD group performance deteriorated during the second TOVA condition across all measures. Within the control group, however, there was a significant effect of TOVA condition only on commission errors ($F(1,17)=11.58$, $p<.003$, $\eta^2=.405$), and d' ($F(1,17)=8.67$, $p<.009$, $\eta^2=.338$). See Table A4 in the supplementary materials for further details.

**EEG results.** In this section we focus on between-group analysis of EEG during non-error TOVA trials, for: parietal ERSPs (RQ 1), exploratory response-locked ERPs, and band power estimates for frontomedial theta and parieto-occipital alpha (RQs 2-4).

***Parietal ERSP on correct trials (RQ 1).*** Figure 3 shows parietal (Pz) ERSPs for correct response and inhibition trials,

for both ADHD group in row 1 and control in row 2. Between groups permutation tests (n=100) were run for each condition separately, using FDR correction for multiple comparisons (Figure 3, row 3). There were no statistically-significant differences in correct response ERSPs, but there were statistically-significant differences in correct inhibition ERSPs (Figure 3, row 3 left side, shown in brown colour).

The between-groups difference for inhibition-trials arose ~80 ms after stimulus onset and continued at least until 400 ms after stimulus onset. The frequency range of the activity was within 3-5 Hz, thus around the (lower) theta range. The strongest ERS (4dB for control group) runs from 200ms onward, consistent around 4Hz.

Figure 3 also shows that the ERS reaches a substantive power level of 1 dB (yellow colour) at different temporal offsets for each group: control ERS begins to ramp up immediately after time 0 (quickly reaching 2dB across 3-6Hz), while ADHD ERS starts some tens of ms after (and is never quite so strong).

***Exploratory ERP results.*** Given the observed group differences in target-locking of ERS, we aimed to further explore the link between behaviour and neural activation. Response-aligned ERP images, plotted for both groups at electrode Pz, are shown in Figure 4. As expected ((67)), the ADHD group has more extreme (very fast or very slow) RTs.

The control group has visibly stronger ERPs, in terms of both negative and positive power amplitudes - blue and red colours, respectively. The average maximum amplitude in the ADHD group is $\sim 4\mu V$, while in the control group it is $\sim 8\mu V$.

The ERP image clearly illustrates one particularly important group difference: control participants experience much stronger phase-resetting in early waves (P1, N2) before 200ms, and then a P3 preceding the RT by a substantial margin. In contrast, P3 in the ADHD group coincides with or follows the RT.

**Fig. 3.** ERSP plots at electrode Pz for correct inhibition and correct response trials. Time 0 represents the stimulus onset. Warm tones represent ERS and cool tones represent ERD. The scale of power perturbations goes from -4.1 to 4.1 dB. Between groups effects were tested with permutation statistics (*p*<.05) using FDR correction for multiple comparisons. The statistically significant differences in the time frequency matrix are shown in brown in the bottom row.

***Exploratory phase analysis.*** Given the suggestions in ERSP and ERP data of control group's stronger temporal coherence between target onset and neural responses, we investigated by plotting phase-aligned ERPs and inter-trial coherence (for methods and figures see supplementary materials). Both these analyses focused on 10Hz, not only because it is the canonical value of peak alpha, but also the timing of TOVA stimuli is 10Hz, i.e. stimuli are shown for 100ms per trial.

Supplementary Figure 5/S1 shows ERP images at Pz, aligned to the pre-stimulus (-80ms) phase of 10Hz alpha, for randomly selected correct trials per-group and per-half. The figure confirms that the control group had much stronger phase resetting with higher amplitude early waves.

Supplementary Figure 6/S2 shows the 10Hz alpha inter-trial coherence (ITC) calculated per-group and per-half at Pz. Both group have significant 10Hz ITC peaking at

around 200ms and 400ms, but it is markedly stronger for control group, by around 50%. For ADHD group, peak ITC droppped from H1 to H2, and the temporal extent of significant-ITC spread out, to begin earlier and end later. In contrast, for controls both H1 and H2 had almost equal profiles of significant ITC.

***Oscillatory power across time (RQs 2-4).*** There was no significant group×time interaction, or a significant main effect of group in frontomedial theta power (RQ 2). There was however a significant main effect of time ($F(8,58)=3.52$, $p<.002$, $\eta^2=.327$) on frontomedial theta in both groups. Pairwise comparisons revealed that only power at 7 Hz between time 1 and time 3 was significant (Bonferroni adjusted $p<.050$).

There was a significant main effect of time on alpha power (RQ 3), as measured from both frontal ($F(10,56)=5.77$, $p<.0005$, $\eta^2=.508$) and parietal electrodes ($F(10,56)=4.68$,

**Fig. 4.** Response-aligned ERP images at Pz during correct response trials. Each row is a single trial from -1s to +1s with respect to stimulus onset, with colour-coded amplitude from -8.1 to 8.1 $\mu V$ (blue and red colors, respectively), and time of response shown by the black sigmoidal curve.
We can see that the stimulus-locked early waves (i.e. vertically-stacked amplitudes at a fixed lag from 0), and the response-locked waves, are both much stronger for control than ADHD group. The P3 wave of the control group also clearly precedes the response, in contrast to the ADHD group.

$p$<.0005, $\eta^2$=.455). That is, power in the extended alpha band (8-12 Hz) increased with time in both groups. There was no significant group×time interaction in either electrode site.

In terms of parieto-occipital pre-stimulus alpha (RQ 4), there was no significant group×time interaction, nor a significant main effect of group on correct trials. There was however a significant main effect of time ($F$(5,47)=3.4, $p$=.004, $\eta^2$=302).

Tables of the complete set of results for the oscillatory band-power analyses and pairwise comparisons are presented in the supplementary materials (Tables A5 and A6).

***Sub-group differences.*** Since we did not find the expected between-group differences in oscillatory power, we split the ADHD group into *three* diagnostic groups for more detailed analyses: ADHD-HI/C, ADHD-I, and control.

There was a significant but small main effect of group on parietal alpha power ($F$(10,122)=2.35, $p$<.014, $\eta^2$=.162) in addition to the significant effect of time segment ($F$(10,55)=5.82, $p$<.0005, $\eta^2$=.514). Pairwise comparisons revealed that all time comparisons were significant with the exception of time 2 versus time 3 at 8 Hz and at 9 Hz (Table

A5). The between groups difference lay between control and ADHD-I groups at 10 Hz (Bonferroni adjusted $p$<.046) and at 11 Hz (Bonferroni adjusted $p$<.034). In these frequency bands the ADHD-I group's oscillatory power was twice as high (10 Hz M=9.77, SD=6.31 and 11 Hz M=8.81, SD=6.07) as the control group's (10 Hz M=4.86, SD=3.76 and 11 Hz M=4.13, SD=3.42). This effect was not due to an outlier participant in the ADHD-I group, as assessed by studentized residuals exceeding ±3. This small effect was only observed for logarithmic power, not normalised (additionally square root transformed) power.

## Discussion

The present study explored cortical oscillations during sustained attention in adults with ADHD and in a healthy control group, measuring EEG while they performed the TOVA task.

**Group performance results.** In terms of behavioural performance, both groups performed worse as the test progressed, from the low response-demand of the first TOVA condition (H1 – testing inattention, infrequent targets) to the higher response-demand of the second TOVA condition (H2 – testing inhibition, frequent targets). However the pattern of

performance within and across test-halves differed between groups.

The ADHD group was affected by each TOVA condition in line with expected deficits of the diagnostic subtypes. During H1, testing inattention, the ADHD group had more commission errors and a lower d' than the control group. During H2, testing inhibition, the ADHD group had more variable RTs than the control group.

Analysing performance between halves, both groups were affected by the change from H1 to H2, reflected in decreased d' (possibly due to reduced vigilance) and increased commission errors (possibly due to fatigue). However, for the ADHD group the amount of omission errors also increased during H2, possibly due to mind wandering as suggested by the Cognitive-Energetic model of ADHD (17, 28, 68, 69).

**EEG-related research questions.** We positively answered RQ1: Are there between-groups differences in neural correlates of visual attention? Indeed, both groups manifested ERS in the theta range on correct inhibition trials, but the amount of parietal synchronisation at 3-5 Hz was significantly higher for the control group. Landau et al. (70) examined parietal theta in the context of visual discrimination-task performance, establishing a precise relationship between two-target spatial attention and gamma-band activity phase-modulated by a parietal 4 Hz source. More recently, Spyropoulos et al. (8) showed that (in macaque V4) parietal theta plays a role in modulating gamma-frequency coding of visual input, and in thus mediating visual attention. Evidence shows that sustained attention is a rhythmic sampling process occurring at a base rate of 8 hz when monitoring a single stimulus; this sampling rhythm decreases to 4 hz when there are two targets to monitor, and keeps decreasing with increasing number of targets (71). Observation of these effects in human electrophysiology indicates that they could be found also in scalp EEG, if generated strongly. TOVA sequentially presents two spatially-distinct (but otherwise identical) stimuli, thus providing ideal conditions to generate strong ∼4 Hz parietal theta ERS, exactly as observed.

Further observations of parietal neural correlates of attention focused on the stronger phase-resetting to stimuli of the control group. At electrode Pz, controls showed ERS more promptly after stimulus onset (Figure 3); they showed early ERP waves with roughly double the amplitude of the ADHD group (Figures 4, 5/S1); and their ITC profiles were both more focal and 50% stronger at peak (Figure 6/S2). Taken together, the evidence shows that ADHD participants were more weakly tuned to the periodic temporal profile of the TOVA trials, and thus had diminished capacity to predict the onset of the next stimulus (as expected from prior findings (28)).

Regarding the questions on band-power, RQs 2-4, results were mixed. RQ2 asked whether frontomedial theta power changed during TOVA, and our results were (surprisingly) negative, on balance. Frontomedial theta power did not increase during the increased demands on sustained attention

in TOVA, with the exception of the increase at 7 Hz from first to last 5-minute time segment (see supplementary materials). No between-groups effect in frontomedial theta power was observed. This is in line with the results of Loo et al. (41).

RQ3 asked if alpha power at parietal or frontal locations changed during TOVA. Power in the extended alpha band (8-12 Hz) increased during TOVA in both groups, indicating decreased cortical arousal. Perhaps heightened demand on sustained attention resulted in fatigue and reduced alpha suppression capacity. This interpretation is in line with the finding that both groups performed worse during the second half of TOVA.

Again we found a null between-groups result; in particular, in contrast to Loo et al. (41), the ADHD group did not exhibit relatively lower parietal alpha power (more cortical arousal) compared with controls. Indeed, when we performed exploratory subgroup analysis, we found that the ADHD-I (predominantly inattentive type) subgroup's parietal 10-11 Hz alpha power was twice that of the control group. This implies that these participants exerted less cognitive control and had diminished alpha suppression during TOVA, reflected as lower cortical arousal (i.e. higher alpha power). It is of note that differences were found only at 10-11Hz, precisely in time with the speed of presentation of TOVA stimuli. The fact this occurred parietally is interesting, as it may relate to the observation of weaker (attention-sampling) parietal theta.

Finally, RQ4 asked if pre-stimulus occipital alpha power changed during TOVA, which was answered positively. On correct trials, pre-stimulus occipital alpha power increased towards the end of TOVA – potentially a reflection of reduced alpha suppression due to fatigue. However, this effect was not different between groups.

Overall, the deterioration of performance during TOVA appears a normal consequence of fatigue, brought on by paying sustained attention over an extended period of time.

**Interpretation of results.** *Synthesising all observed results*, we propose an interpretation based on deficient rhythmic attention-sampling in a cortical area responsible for relational processing, the posterior parietal cortex. This may be inferred from the nature of the TOVA task itself, as follows.

The TOVA task is temporally regular, meaning that every trial is predictable. For studying ADHD, this provides an advantage since the ADHD behavioural and neural responses can be compared to responses of a healthy control group which are facilitated by predictability. Thus, our particular observations point toward the ADHD behavioural deficit arising from reduced neural entrainment to task, possibly due to weaker endogenous attention sampling. The latter seems to be a parietal rather than frontal issue.

Specifically, based on the theory that attention continuously samples attended locations (at some fraction of an 8 Hz base rate (70)), our results suggest that it is the weaker parietal theta activation (reflected as reduced parietal theta ERS) that

causes reduced phase resetting – i.e. reduced resources to monitor stimuli leads to weaker modulation of ongoing oscillations by those stimuli (or their predicted occurence). This consequently leads to weaker early ERPs (see Figure 5/S1) and smaller peak ITC (see Figure 6/S2). Thus, ADHD may have poorer TOVA performance because of weaker modulation of attentional sampling for stimuli requiring inhibition, reflected (for example) as significantly higher commission error rate in H1 (the half with frequent non-targets).

Further, TOVA is a relational classification task: Go/NoGo (respond/inhibit) to targets/non-targets, which share the same stimulus properties (smaller black rectangles within larger white squares), at two different spatial locations (above-below fixation). This relationship between target types permits a *unidimensional encoding*. Based on this property of the task, we can *hypothesise a cognitive mechanism* that could generate our above results, as follows.

Summerfield et al. (72) recently described a theory for the role of posterior parietal cortex (PPC, i.e. the dorsal visual stream, or the 'what' stream) in processing relational structure in the environment. In brief, Summerfield et al. (72) suggest that primate PPC supports learning and processing of relational structure, for example in visual scenes. They build on work establishing that PPC, in particular lateral intraparietal area (LIP, Brodman's areas 39/40), provides spatially-selective coding for regions of ego-centric space that subserve functions including decision-making and top-down spatial attention (73). PPC neurons *also* code for abstract categories (74), in a scalar manner wherein unidimensional relations are coded asynchronously – if A is coded by many neurons, then B may be coded by relatively fewer.

Since TOVA stimuli consist exactly of a unidimensional relational structure, every trial requires recruitment of the posterior parietal cortex to judge the presented relationship and trigger appropriate action. The observed lack of between-group difference in frontomedial theta suggests that an executive function deficit is not at work, and reinforces the implication that the ADHD group was deficient in relational processing. This hypothesis refines the interpretation of the ERSP result: that not only does our ADHD group show reduced theta-rhythmic attention sampling, but they do so in the specific function of sampling relationships in the world.

The question then arises, why was there no significant difference in between-groups ERSP to response trials? Here, the control group's ERSP was still visibly stronger (than ADHD), so the trend remains. We can thus draw again on the Summerfield *et al* theory, and the scalar coding of unidimensional relations. In our data, the matching interpretation would be that ERS to response trials is simply weaker due to having fewer coding neurons, and thus any between-group differences get washed out due to lower signal-to-noise ratio.

Alternative explanations are of course possible. For one thing, primate data (75) supporting the above theory shows that individuals code category classes *arbitrarily*, and the group-wise aggregate in our data should then 'even out' – i.e.

higher firing for targets in some participants, and vice versa in others. If this is the case, our results would imply that inhibition trials differed in some other way than PPC coding. One option is that ADHD theta ERS for inhibition trials was spread uniformly across frequencies and thus not strong enough at the monitoring frequency 4Hz, because ADHD participants were not sufficiently focused on attending the correct spatial location for inhibition. This contrasts with the response trials because responses have action affordances, which can leverage premotor cortex coding to supplement PPC category judgement. It also goes to the hyperactive-impulsive aspect of ADHD, plus our observation that ADHD group had more commission errors (inability to inhibit responses) in H1 than controls.

**Limitations.** The present study is among the first of its kind combining TOVA with an EEG measurement in adults (40) – and the first to focus on ADHD. As such, there were a number of limitations, both in the sense of constraints and issues.

*Constraints.* A number of aspects were suboptimal but were constrained by the fixed design of TOVA task, and size of sample locally available to meet the recruitment criteria.

For example, the small size of some of the TOVA norming groups, such as males aged 30-39 (n=4), might have increased the number of false positives. In the present study 19 participants (of which 15 with ADHD) belonged to this cohort. The low sample sensitivity and specificity highlight the issue of behavioural heterogeneity among people diagnosed with ADHD, and further motivates our sub-group analysis which found elevated parietal 10Hz alpha among the ADHD-I group (compared to controls).

In terms of constraints of the task, TOVA does not aim to induce a large number of error trials, which limited the types of analyses available. For example, we could not subtract correct response trials from commission error trials, or *vice versa*. As a result, the ERSP plot for correct response trials (Figure 3) contains not only cognitive processing but also motor processing related to the response.

Finally, in relation to the oscillatory analyses of time-dependency, it should be noted that the time-segment variable is confounded by the two TOVA conditions H1 and H2, the order of which are not counter-balanced in the TOVA software.

*Issues.* In order to recruit sufficient N the present study had a sample with a wide range of ages. There are well-documented age-related changes in tonic alpha and theta oscillations; for example, the age-related evolution of alpha and theta frequency spectra is nonlinear and may lead in opposing directions for alpha and for theta (9). However, given that our analyses were conducted within (not between) frequency bands, and the age range of ADHD and control groups were balanced, we do not believe this issue alters our interpretation of results.

We did not control for individual differences in alpha peak frequency (iAPF), that is, the single frequency in the alpha-

band showing the highest power per individual. Counting iAPF as the central frequency of alpha, instead of 10 Hz, can be used to recalibrate all frequency bands and potentially account for clinical variation in EEG spectra (76). Individual differences in frequency spectrum may be more prominent than any age-related differences, particularly in the alpha and theta bands (9). On the other hand, for ADHD population such variation may be more prominent in children than adults.

The neuropathological heterogeneity of ADHD implies that distinct neurocognitive subtypes of ADHD may exist, and yet not map to symptomatic subgroups. Due to this, one criticism of ADHD studies has been that group-wise comparisons between ADHD participants and healthy controls are likely to yield small, hard-to-replicate effects (4). While the present study compared groups based on their diagnostic status and did find robust behavioural differences, also clustering participants as a function of both behavioural performance and cortical oscillations may find meaningful distinctions in test responses. This type of analysis-by-outcome (as opposed to analysis-by-diagnosis) might shed light on how oscillatory dynamics are related to performance, and whether this relationship is altered in ADHD as compared to healthy controls. In the context of the present study, we did find strong differences between diagnostic groups, especially in event-related spectral and stimulus-locking analyses.

**Conclusion.** In conclusion, this first-of-its-kind study examined the combination of TOVA CPT with EEG measurement for assessing neural correlates of ADHD in adults, an understudied question. We focused on participants' diagnostic status and its relation to differences in oscillatory markers of sustained and relational attention during TOVA.

We showed that the ADHD group showed lower theta synchronisation during correct inhibition trials, and demonstrated reduced sensitivity to stimulus timing across several parietal measures. **The interpretation of these results is notable, as it points to deficient attention sampling in relational categorisation tasks** (72). This result contributes to the overall understanding of the neuropathology of ADHD in adults.

Clinically, combining EEG measurement with TOVA has the potential to explain why differences in behavioural performance in TOVA arise. Identifying subgroups among ADHD diagnosed adults could eventually be used as an additional grouping variable to predict treatment outcomes.

**ACKNOWLEDGEMENTS**

All authors co-wrote and approved the manuscript. KJ gathered and analysed the data, and created figures. BC conceived and designed the experiment, analysed data, and created figures. JP contributed to text and figures.

Authors wish to thank Mona Moisala and Marko Repo for data gathering; Otto Lappi, Alina Leminen and Eeva Palomäki for comments to the manuscript.

# Bibliography

1. JM Swanson, JA Sergeant, E Taylor, EJS Sonuga-Barke, PS Jensen, and DP Cantwell. Attention-deficit hyperactivity disorder and hyperkinetic disorder. *The Lancet*, 351(9100): 429–433, 1998.

2. Peter R Killeen, Rosemary Tannock, and Terje Sagvolden. The four causes of adhd: a framework. In *Behavioral neuroscience of attention deficit hyperactivity disorder and its treatment*, pages 391–425. Springer, 2011.

3. F Xavier Castellanos, Edmund JS Sonuga-Barke, Anouk Scheres, Adriana Di Martino, Christopher Hyde, and Judith R Walters. Varieties of attention-deficit/hyperactivity disorder-related intra-individual variability. *Biological psychiatry*, 57(11):1416–1423, 2005.

4. Joel T Nigg, Erik G Willcutt, Alysa E Doyle, and Edmund JS Sonuga-Barke. Causal heterogeneity in attention-deficit/hyperactivity disorder: do we need neuropsychologically impaired subtypes? *Biological psychiatry*, 57(11):1224–1230, 2005.

5. Erik G Willcutt, Alysa E Doyle, Joel T Nigg, Stephen V Faraone, and Bruce F Pennington. Validity of the executive function theory of attention-deficit/hyperactivity disorder: a meta-analytic review. *Biological psychiatry*, 57(11):1336–1346, 2005.

6. Esther Strauss, Elisabeth MS Sherman, Otfried Spreen, et al. *A compendium of neuropsychological tests: Administration, norms, and commentary*, pages 645–651. American Chemical Society, 2006.

7. Martijn Arns, Jay Gunkelman, Marinus Breteler, and Desirée Spronk. EEG phenotypes predict treatment outcome to stimulants in children with ADHD. *Journal of integrative neuroscience*, 7(3):421–38, 2008. ISSN 0219-6352.

8. Georgios Spyropoulos, Conrado Arturo Bosman, and Pascal Fries. A theta rhythm in macaque visual cortex and its attentional modulation. *Proceedings of the National Academy of Sciences*, 115(24):E5614–E5623, 2018. doi: 10.1073/pnas.1719433115.

9. Wolfgang Klimesch. Eeg alpha and theta oscillations reflect cognitive and memory performance: a review and analysis. *Brain research reviews*, 29(2-3):169–195, 1999.

10. Daniel J Calderone, Peter Lakatos, Pamela D Butler, and F Xavier Castellanos. Entrainment of neural oscillations as a modifiable substrate of attention. *Trends in cognitive sciences*, 18 (6):300–309, 2014.

11. Russell A Barkley. Behavioral inhibition, sustained attention, and executive functions: constructing a unifying theory of adhd. *Psychological bulletin*, 121(1):65, 1997.

12. Sandra K Loo, Lorie A Humphrey, Terttu Tapio, Irma K Moilanen, JAMES J McGOUGH, JAMES T McCRACKEN, May H Yang, Jeff Dang, Anja Taanila, Hanna Ebeling, et al. Executive functioning among finnish adolescents with attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 46(12):1594–1604, 2007.

13. Edmund JS Sonuga-Barke and F Xavier Castellanos. Spontaneous attentional fluctuations in impaired states and pathological conditions: a neurobiological hypothesis. *Neuroscience & Biobehavioral Reviews*, 31(7):977–986, 2007.

14. Agatha Lenartowicz, Ali Mazaheri, Ole Jensen, and Sandra K. Loo. Aberrant Modulation of Brain Oscillatory Activity and Attentional Impairment in Attention-Deficit/Hyperactivity Disorder. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 3(1):19–29, January 2018. doi: 10.1016/j.bpsc.2017.09.009.

15. F Xavier Castellanos and Erika Proal. Large-scale brain systems in adhd: beyond the prefrontal–striatal model. *Trends in cognitive sciences*, 16(1):17–26, 2012.

16. Athanasia M. Mowinckel, Dag Alnæs, Mads L. Pedersen, Sigurd Ziegler, Mats Fredriksen, Tobias Kaufmann, Edmund Sonuga-Barke, Tor Endestad, Lars T. Westlye, and Guido Biele. Increased default-mode variability is related to reduced task-performance and is evident in adults with ADHD. *NeuroImage: Clinical*, 16:369–382, 2017. ISSN 2213-1582. doi: 10.1016/j.nicl.2017.03.008.

17. Peter R Killeen. Absent without leave; a neuroenergetic theory of mind wandering. *Frontiers in psychology*, 4:373, 2013.

18. Michael S Clayton, Nick Yeung, and Roi Cohen Kadosh. The roles of cortical oscillations in sustained attention. *Trends in cognitive sciences*, 19(4):188–195, 2015.

19. Joram van Driel, K Richard Ridderinkhof, and Michael X Cohen. Not all errors are alike: theta and alpha eeg dynamics relate to differences in error-processing dynamics. *Journal of Neuroscience*, 32(47):16795–16806, 2012.

20. James F Cavanagh, Michael X Cohen, and John JB Allen. Prelude to and resolution of an error: Eeg phase synchrony reveals cognitive control dynamics during action monitoring. *Journal of Neuroscience*, 29(1):98–105, 2009.

21. James F Cavanagh and Michael J Frank. Frontal theta as a mechanism for cognitive control. *Trends in cognitive sciences*, 18(8):414–421, 2014.

22. Edmund Wascher, Björn Rasch, Jessica Sänger, Sven Hoffmann, Daniel Schneider, Gerhard Rinkenauer, Herbert Heuer, and Ingmar Gutberlet. Frontal theta activity reflects distinct aspects of mental fatigue. *Biological psychology*, 96:57–65, 2014.

23. Paul Sauseng and Wolfgang Klimesch. What does phase information of oscillatory brain activity tell us about cognitive processes? *Neuroscience & Biobehavioral Reviews*, 32(5): 1001–1013, 2008.

24. Wolfgang Klimesch. Alpha-band oscillations, attention, and controlled access to stored information. *Trends in cognitive sciences*, 16(12):606–617, 2012.

25. Wolfgang Klimesch, Paul Sauseng, and Simon Hanslmayr. Eeg alpha oscillations: the inhibition–timing hypothesis. *Brain research reviews*, 53(1):63–88, 2007.

26. Luc H Arnal and Anne-Lise Giraud. Cortical oscillations and sensory predictions. *Trends in cognitive sciences*, 16(7):390–398, 2012.

27. Peter Lakatos, George Karmos, Ashesh D Mehta, Istvan Ulbert, and Charles E Schroeder. Entrainment of neuronal oscillations as a mechanism of attentional selection. *science*, 320 (5872):110–113, 2008.

28. Vivienne A Russell, Robert D Oades, Rosemary Tannock, Peter R Killeen, Judith G Auerbach, Espen B Johansen, and Terje Sagvolden. Response variability in attention-deficit/hyperactivity disorder: a neuronal and glial energetics hypothesis. *Behavioral and Brain Functions*, 2(1):30, 2006.

29. Niko A Busch and Rufin VanRullen. Spontaneous eeg oscillations reveal periodic sampling of visual attention. *Proceedings of the National Academy of Sciences*, 107(37):16048–16053, 2010.

30. Jordan P Hamm, Kara A Dyckman, Jennifer E McDowell, and Brett A Clementz. Pre-cue fronto-occipital alpha phase and distributed cortical oscillations predict failures of cognitive control. *Journal of Neuroscience*, 32(20):7034–7041, 2012.

31. Simon Hanslmayr, Alp Aslan, Tobias Staudigl, Wolfgang Klimesch, Christoph S Herrmann, and Karl-Heinz Bäuml. Prestimulus oscillations predict visual perception performance between and within subjects. *Neuroimage*, 37(4):1465–1473, 2007.

32. James Stuart Peter Macdonald, Santosh Mathan, and Nick Yeung. Trial-by-trial variations in subjective attentional state are reflected in ongoing prestimulus eeg alpha oscillations. *Frontiers in psychology*, 2:82, 2011.

33. Gregor Thut, Annika Nietzel, Stephan A Brandt, and Alvaro Pascual-Leone. $\alpha$-band electroencephalographic activity over occipital cortex indexes visuospatial attention bias and predicts visual target detection. *Journal of Neuroscience*, 26(37):9494–9502, 2006.

34. Rufin VanRullen, Niko Busch, Jan Drewes, and Julien Dubois. Ongoing eeg phase as a trial-by-trial predictor of perceptual and attentional variability. *Frontiers in psychology*, 2:60, 2011.

35. Kyle E Mathewson, Gabriele Gratton, Monica Fabiani, Diane M Beck, and Tony Ro. To see or not to see: prestimulus $\alpha$ phase predicts visual awareness. *Journal of Neuroscience*, 29 (9):2725–2732, 2009.

36. R. A. Leark, L. K. Greenberg, C. L. Kindschi, T. R. Dupuy, and S. J. Hughes. Test of Variables of Attention: Professional Manual., 2007.

37. Celestino Rodríguez, Paloma González-Castro, Marisol Cueli, Debora Areces, and Julio A. González-Pienda. Attention Deficit/Hyperactivity Disorder (ADHD) Diagnosis: An Activation-Executive Model. *Frontiers in Psychology*, 7, September 2016. doi: 10.3389/fpsyg.2016.01406.

38. Paloma González-Castro, Marisol Cueli, Celestino Rodríguez, Trinidad García, and Luis Álvarez. Efficacy of Neurofeedback Versus Pharmacological Support in Subjects with ADHD. *Applied Psychophysiology and Biofeedback*, 41(1):17–25, March 2016. doi: 10.1007/s10484-015-9299-4.

39. Islam F. Halawa, Basma B. El Sayed, Omnia R. Amin, Nagwa A. Meguid, and Ann A. Abdel Kader. Frontal theta/beta ratio changes during TOVA in Egyptian ADHD children. *Neurosciences (Riyadh, Saudi Arabia)*, 22(4), 2017. ISSN 1319-6138. doi: 10.17712/nsj.2017.4.20170067.

40. Julian R. Keith, Lobsang Rapgay, Don Theodore, Jeffrey M. Schwartz, and Jae L. Ross. An assessment of an automated EEG biofeedback system for attention deficits in a substance use disorders residential treatment setting. *Psychology of Addictive Behaviors: Journal of the Society of Psychologists in Addictive Behaviors*, 29(1):17–25, March 2015. doi: 10.1037/adb0000016.

41. Sandra K Loo, T Sigi Hale, James Macion, Grant Hanada, James J McGough, James T McCracken, and Susan L Smalley. Cortical activity patterns in adhd during arousal, activation and sustained attention. *Neuropsychologia*, 47(10):2114–2119, 2009.

42. CK Conners. The continuous performance test (cpt): Use as a diagnostic tool and measure of treatment outcome. *American Psychological Association*, 1994.

43. C Keith Conners, MHS Staff, V Connelly, S Campbell, M MacLean, and J Barnes. Conners' continuous performance test ii (cpt ii v. 5). *Multi-Health Syst Inc*, 29:175–96, 2000.

44. Benjamin Ultan Cowley, Édua Holmström, Kristiina Juurmaa, Levas Kovarskis, and Christina M Krause. Computer enabled neuroplasticity treatment: a clinical trial of a novel design for neurofeedback therapy in adult adhd. *Frontiers in Human Neuroscience*, 10:205, 2016.

45. Ronald C Kessler, Lenard Adler, Minnie Ames, Olga Demler, Steve Faraone, EVA Hiripi, Mary J Howes, Robert Jin, Kristina Secnik, Thomas Spencer, et al. The world health organization adult adhd self-report scale (asrs): a short screening scale for use in the general population. *Psychological medicine*, 35(2):245–256, 2005.

46. Thomas E Brown. *Brown Attention-Deficit Disorder Scales (Brown ADD Scales): For Adolescents and Adults*. Psychological Corporation, 1996.

47. David Wechsler. *WAIS-IV technical and interpretive manual*. Pearson, 2008.

48. JJS Kooij and MH Francken. Diagnostic interview for adhd in adults 2.0 (diva 2.0). *Adult ADHD: Diagnostic assessment and treatment*, pages 33–99, 2010.

49. Robert L Spitzer, Kurt Kroenke, Janet BW Williams, and Bernd Löwe. A brief measure for assessing generalized anxiety disorder: the gad-7. *Archives of internal medicine*, 166(10): 1092–1097, 2006.

50. Aaron T Beck, Robert A Steer, and Gregory K Brown. *BDI-II: Beck depression inventory*. Pearson, 1996.

51. John B Saunders, Olaf G Aasland, Thomas F Babor, Juan R De la Fuente, and Marcus Grant. Development of the alcohol use disorders identification test (audit): Who collaborative project on early detection of persons with harmful alcohol consumption-ii. *Addiction*, 88 (6):791–804, 1993.

52. Robert MA Hirschfeld, Janet BW Williams, Robert L Spitzer, Joseph R Calabrese, Laurie Flynn, Paul E Keck Jr, Lydia Lewis, Susan L McElroy, Robert M Post, Daniel J Rapport, et al. Development and validation of a screening instrument for bipolar spectrum disorder: the mood disorder questionnaire. *American Journal of Psychiatry*, 157(11):1873–1875, 2000.

53. M Heinimaa, RKR Salokangas, T Ristkari, M Plathin, J Huttunen, T Ilonen, T Suomela, J Korkeila, and TH McGlashan. Prod-screen–a screen for prodromal symptoms of psychosis. *International Journal of Methods in Psychiatric Research*, 12(2):92–104, 2003.

54. Michael R Liebowitz. Dissociative experiences scale. *The American journal of psychiatry*, 1992.

55. Kindschi C. L. Dupuy T. R. Greenberg, L. K. and S. J. Hughes. Test of Variables of Attention: Clinical Manual., 2016.

56. Benjamin Ultan Cowley. Studying the integrated functional cognitive basis of sustained attention with a primed subjective-illusory-contour attention task. *Scientific reports*, 8, 2018.

57. Torbjörn Åkerstedt and Mats Gillberg. Subjective and objective sleepiness in the active individual. *International Journal of Neuroscience*, 52(1-2):29–37, 1990.

58. Benjamin U Cowley, Jussi Korpela, and Jari Torniainen. Computational testing for automated preprocessing: a matlab toolbox to enable large scale electroencephalography data processing. *PeerJ Computer Science*, 3:e108, 2017.

59. Benjamin U Cowley and Jussi Korpela. Computational testing for automated preprocessing 2: practical demonstration of a system for scientific data-processing workflow management for high-volume eeg. *Frontiers in neuroscience*, 12:236, 2018.

60. Arnaud Delorme and Scott Makeig. Eeglab: an open source toolbox for analysis of single-trial eeg dynamics including independent component analysis. *Journal of neuroscience methods*, 134(1):9–21, 2004.

61. Aapo Hyvarinen. Fast and robust fixed-point algorithms for independent component analysis. *IEEE transactions on Neural Networks*, 10(3):626–634, 1999.

62. Hugh Nolan, Robert Whelan, and Richard B Reilly. Faster: fully automated statistical thresholding for eeg artifact rejection. *Journal of neuroscience methods*, 192(1):152–162, 2010.

63. Simon Hanslmayr, Wolfgang Klimesch, Paul Sauseng, Walter Gruber, Michael Doppelmayr, Roman Freunberger, and Thomas Pecherstorfer. Visual discrimination performance is related to decreased alpha amplitude but increased phase locking. *Neuroscience letters*, 375 (1):64–68, 2005.

64. Scott Makeig, Stefan Debener, Julie Onton, and Arnaud Delorme. Mining event-related brain dynamics. *Trends in cognitive sciences*, 8(5):204–210, 2004.

65. Steven J Luck. *An introduction to the event-related potential technique*. MIT press, 2014.

66. Yoav Benjamini and Yosef Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*, 57(1):289–300, 1995.

67. Harald Baayen and Petar Milin. Analyzing Reaction Times. *International Journal of Psychological Research*, 3:12–28, 2010.

68. Joseph Sergeant. The cognitive-energetic model: an empirical approach to attention-deficit hyperactivity disorder. *Neuroscience & Biobehavioral Reviews*, 24(1):7–12, 2000.

69. Joseph A Sergeant. Modeling attention-deficit/hyperactivity disorder: a critical appraisal of the cognitive-energetic model. *Biological psychiatry*, 57(11):1248–1255, 2005.

70. Ayelet Nina Landau, Helene Marianne Schreyer, Stan van Pelt, and Pascal Fries. Distributed attention is implemented through theta-rhythmic gamma modulation. *Current Biology*, 25 (17):2332 – 2337, 2015. ISSN 0960-9822. doi: 10.1016/j.cub.2015.07.048.

71. Alex O. Holcombe and Wei-Ying Chen. Splitting attention reduces temporal resolution from 7 Hz for tracking one object to <3 Hz when tracking three. *Journal of Vision*, 13(1):12–12, Jan 2013. ISSN 1534-7362. doi: 10.1167/13.1.12.

72. Christopher Summerfield, Fabrice Luyckx, and Hannah Sheahan. Structure learning and the parietal cortex, Jul 2019.

73. David J. Freedman and Guilhem Ibos. An Integrative Framework for Sensory, Motor, and Cognitive Functions of the Posterior Parietal Cortex. *Neuron*, 97(6):1219–1234, March 2018. ISSN 0896-6273. doi: 10.1016/j.neuron.2018.01.044.

74. David J. Freedman and John A. Assad. Neuronal Mechanisms of Visual Categorization: An Abstract View on Decision Making. *Annual Review of Neuroscience*, 39:129–147, July 2016. doi: 10.1146/annurev-neuro-071714-033919.

75. Jamie K. Fitzgerald, David J. Freedman, Alessandra Fanini, Sharath Bennur, Joshua I. Gold, and John A. Assad. Biased Associative Representations in Parietal Cortex. *Neuron*, 77(1): 180–191, January 2013. doi: 10.1016/j.neuron.2012.11.014.

76. Marieke M Lansbergen, Martijn Arns, Martine van Dongen-Boomsma, Desirée Spronk, and Jan K Buitelaar. The increase in theta/beta ratio on resting-state eeg in boys with attention-deficit/hyperactivity disorder is mediated by slow alpha peak frequency. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 35(1):47–52, 2011.