A Bayesian computational model reveals a failure to adapt interoceptive precision

estimates across depression, anxiety, eating, and substance use disorders

Ryan Smith[1], Rayus Kuplicki[1], Justin Feinstein[1,2], Katherine L. Forthman, Jennifer L. Stewart[1,2],

Martin P. Paulus[1,2], Tulsa 1000 investigators[†], and Sahib S. Khalsa[1,2]

[1]Laureate Institute for Brain Research, Tulsa, OK, USA

[2]Oxley College of Health Sciences, The University of Tulsa, Tulsa, OK.

[†]The Tulsa 1000 Investigators include the following contributors: Robin L. Aupperle,

Ph.D.; Jerzy Bodurka, Ph.D.; Jonathan B. Savitz, Ph.D.; and Teresa A. Victor, Ph.D.

Corresponding author:

Ryan Smith

Laureate Institute for Brain Research

6655 S Yale Ave, Tulsa, OK 74136, USA

Email: rsmith@laureateinstitute.org

Abstract

Recent neurocomputational theories have hypothesized that abnormalities in prior beliefs and/or

the precision-weighting of afferent interoceptive signals may facilitate the transdiagnostic

emergence of psychopathology. Specifically, it has been suggested that, in certain psychiatric

disorders, interoceptive processing mechanisms either over-weight prior beliefs or under-weight

signals from the viscera (or both), leading to a failure to accurately update beliefs about the body.

However, this has not been directly tested empirically. To evaluate the potential roles of prior

beliefs and interoceptive precision in this context, we fitted behavior in a transdiagnostic patient

population on an interoceptive awareness (heartbeat tapping) task to a Bayesian computational

model. Modeling revealed that, during an interoceptive perturbation condition (inspiratory

breath-holding during heartbeat tapping), healthy individuals (N=52) assigned greater precision

to ascending cardiac signals than individuals with symptoms of anxiety (N=15), depression

(N=69), co-morbid depression/anxiety (N=153), substance use disorders (N=131), and eating

disorders (N=14) – who failed to increase their precision estimates from resting levels. In

contrast, we did not find strong evidence for differences in prior beliefs. These results provide

the first empirical computational modeling evidence of a selective dysfunction in adaptive

interoceptive processing in psychiatric conditions, and lay the groundwork for future studies

examining how reduced interoceptive precision influences body regulation and interoceptively-

guided decision-making.

Author Summary

Interoception is the process by which the nervous system senses the internal state of the body. It

provides the brain with important information to adaptively guide the regulation of both internal

body states and behavior. Interoceptive dysfunction is thought to play a role in multiple

psychiatric disorders. Theoretical models propose that the computational mechanisms of

interoceptive dysfunction are caused by overly precise prior beliefs about body states

("hyperprecise priors") or underestimates of the reliability of the information carried by

ascending signals from the body ("low sensory precision"). Our empirical approach tested for

evidence of these mechanisms across several psychiatric disorders, using a computational model

of perception during performance of a heartbeat perception task. We found evidence of low

sensory precision within individuals with anxiety, depression, eating disorders, and/or substance

use disorders, relative to healthy individuals. This difference occurred only during a breath-

holding condition designed to enhance heartbeat signals. We did not find evidence for

hyperprecise priors in the patient groups. The data from this study support the argument for

computational mechanisms of interoceptive dysfunction across several psychiatric disorders, and

suggests that these conditions may be characterized by an inability to adjust sensory precision

when signals from the body change.

Keywords: Interoception; Active Inference; Depression; Anxiety; Substance Use; Eating
Disorders; Precision; Priors; Bayesian Perception; Computational Modeling

Introduction

Interoception plays an important role in a number of psychiatric disorders. Interoceptive

dysfunction has been observed in depression, anxiety, eating, and substance use disorders,

among others (reviewed in (1)). For example, depressed patients exhibit reduced accuracy when

asked to count their own heartbeats (2-4), and counting accuracy is negatively correlated with

depressive symptoms (5). Several studies have also shown heightened interoceptive sensations in

panic disorder under high arousal states (reviewed in (6)). Other studies have reported evidence

of blunted neural responses during interoceptive processing in substance users (7), and that

patients with eating disorders show stronger expectation effects on interoception during

modulations of arousal (8). However, the mechanisms underlying these potential interoceptive

dysfunctions in psychiatric disorders have not yet been demonstrated in empirical work.

In recent years, a growing body of theoretical work within neuroscience and psychiatry has

begun to highlight plausible neurocomputational accounts of interoceptive processing (9-15), as

well as accounts of exteroceptive (16, 17), cognitive (18-26), emotional (27, 28), and motor

control (29-31) functions that plausibly interact with interoception. In some cases, empirical

studies have also found support for computational models and computational abnormalities in

relation to specific psychiatric symptoms/disorders (26, 32-34). With respect to sensory

processing, which has focused to a large degree on neural models of approximate probabilistic

(Bayesian) inference, empirical results in this area have come almost exclusively from the

exteroceptive domain. And while recent work has begun to test qualitative predictions of

Bayesian models of interoception (35), no study to our knowledge has yet explicitly fit

computational models to brain or behavioral responses during experimental modulations of

interoception in individuals with psychiatric disorders.

Drawing on this theoretical literature, we and others (9, 13-15, 36-38) have previously proposed

that symptoms of multiple psychiatric conditions could be explained by faulty interoceptive

computational processes (and associated visceromotor control processes), due to an inappropriate

weighting of prior beliefs and sensory evidence. Although these models of computational

dysfunction in interoception could have broad explanatory power, the putative mechanisms they

propose have yet to be empirically tested using formal computational models. In this paper, we

use a formal Bayesian computational model of perception to examine whether there is evidence

for the transdiagnostic dysfunction of interoceptive processing suggested in these previous

proposals within the cardiovascular system.

Historically, the most common empirical paradigms for studying interoception have focused on

the perception of cardiac signals. Common heartbeat perception tasks include those involving

heartbeat counting (39), heartbeat tapping (40), and other means of assessing cardiac

interoception (41-45). While each approach has certain limitations (46-50), one consistent

finding is that cardiac interoceptive awareness is quite poor in the majority of participants tested

during resting conditions – where only roughly 35% of individuals appear to accurately perceive

their own heartbeats (6). In contrast, when visceral states are perturbed, cardiac perception

becomes more accurate. This is particularly true for interoceptive accuracy under conditions of

heightened cardiorespiratory arousal (51-53).

Active inference models, and Bayesian predictive processing models more generally, have been

the main computational framework within which interoceptive dysfunction has been discussed

(13, 38, 54-56). Several authors have suggested that one important transdiagnostic factor within

mental disorders may be an inability of the brain to update its model of the body in the face of

interoceptive prediction errors (i.e., mismatches between expected and received afferent

interoceptive signals from the body). In predictive coding and active inference models, this kind

of aberrant belief updating is thought to come about through a dysfunctional "precision

weighting" mechanism, which governs the relative influence of prior beliefs and afferent bodily

signals in determining perception (and informing visceral regulation). Simply put, it is suggested

that, across multiple mental health conditions, the brain may treat afferent bodily signals (and

associated prediction errors) as though they are not reliable indicators of bodily states during

interoceptive inference – leading perception to be insufficiently constrained by true visceral

states and primarily determined by (in many cases maladaptive) prior beliefs. Misestimating the

state of the body could in turn promote a number of transdiagnostic symptoms. For example,

interoceptive feelings are intimately tied to emotions (57-59); poor body perception (e.g., high

uncertainty about internal physiological conditions and a resulting inability to efficiently regulate

them) could thus maintain unpleasant emotional states. Chronic underestimates of available

metabolic resources may contribute to apathy and anhedonia (56), and overestimates of the

evidence uncomfortable bodily sensations provide for physical threat (e.g., a heart attack) may

contribute to anxiety and panic (60, 61).

It's important to emphasize, however, that computational models often include several additional

parameters. Aside from the precision-weighting of sensory signals, individuals can also have

differences in (for example) prior beliefs about what they will perceive (assumed in any

Bayesian model of perception; e.g., predictive coding (62)), and differences in how quickly they

update those prior beliefs over repeated observations (i.e., "learning rate"; i.e., which can be

dependent upon the relative precisions of sensory signals and prior beliefs). Formal

computational models are often necessary to distinguish which parameters show differences

between individuals and best explain differences in perception. Thus, there are multiple

computational mechanisms that could account for individual differences in interoception in

clinical populations. One major goal in computational psychiatry is to "computationally

phenotype" patients by identifying which sets of a parameter values best account for their neural

and behavioral responses (including self-reported perceptual experience) and use this

information to guide treatment (20, 22, 25).

In the present study, we apply a novel computational phenotyping approach, using a Bayesian

model of perception, to identify the computational parameters that best explain behavior on a

cardiac perception (heartbeat tapping) task performed by a transdiagnostic clinical sample of

individuals with psychiatric disorders as well as a healthy comparison (HC) sample. Given

previous findings that performance on cardiac perception tasks can be significantly altered by

different task instructions ((63); i.e., potentially by adjusting prior beliefs), we chose to assess

task performance under two resting conditions in which participants were instructed to apply

different confidence thresholds: 1) a condition where guessing was allowed and 2) a condition in

which they should not guess (i.e., they should only tap a key if they were sure they actually felt a

heartbeat). Given the notably poorer cardiac perception of most human beings at rest compared

to during altered arousal states (6), in addition to the resting conditions we also chose to assess

individual differences in cardiac perception during a non-invasive interoceptive perturbation (a

breath-hold) condition (which also included the no-guessing instruction) that was expected to

increase the precision of the afferent cardiac signal and improve cardiac perception above floor

values in a greater number of individuals (i.e., we expected that cardiac perception would be

generally poor during resting conditions, and that the breath-hold condition would result in

improved performance on average). Thus, comparing the guessing and no-guessing conditions

allowed assessment of the effects of altering prior beliefs with different task instructions, and

comparison of the no-guessing a breath-hold conditions allowed assessment of changes in

afferent signal precision under identical (i.e., no-guessing) task instructions. Our primary aims

were to 1) demonstrate the sensitivity of our novel computational approach in measuring the

condition-specific precision-weighting of interoceptive signals and prior beliefs across a

transdiagnostic sample of individuals with depression, anxiety, substance use disorders, and/or

eating disorders, 2) test the hypothesis (as previously proposed; e.g., (10, 38, 56)) that these

patient groups would show lower interoceptive precision weightings than HCs, more precise

prior beliefs than HCs, or both, and 3) establish whether prior beliefs and/or interoceptive

precision is abnormal in general (relative to healthy participants) or selectively within resting or

interoceptive perturbation conditions.

<div align="center">

Methods

</div>

Participants

Data were collected from 500 participants (153 male) as a part of the Tulsa 1000 (T1000)

project, a naturalistic longitudinal study that recruited subjects based on the dimensional NIMH

Research Domain Criteria framework (a full description of this pre-planned project can be found

in (64)). Individuals aged 18-55 years were screened on the basis of dimensional

psychopathology scores. Inclusion was based on the following measures: Patient Health

Questionnaire (PHQ-9; (65)) ≥ 10, Overall Anxiety Severity and Impairment Scale (OASIS;

(66)) ≥ 8, Drug Abuse Screening Test (DAST-10; (67)) score > 2, and/or Eating Disorder Screen

(SCOFF; (68)) score ≥ 2 (for screening measure scores, see **Table 1**). HCs who did not show

elevated symptoms or psychiatric diagnoses were also included. Participants were excluded if

they (i) tested positive for drugs of abuse, (ii) met criteria for psychotic, bipolar, or obsessive-

compulsive disorders, or reported (iii) history of moderate-to-severe traumatic brain injury,

neurological disorders, or severe or unstable medical conditions, (iv) active suicidal intent or

plan, or (v) change in psychotropic medication status within 6 weeks. Full inclusion/exclusion

criteria are described in (64). The study was approved by the Western Institutional Review

Board. All participants provided written informed consent prior to completion of the study

protocol, in accordance with the Declaration of Helsinki, and were compensated for

participation. ClinicalTrials.gov identifier: #NCT02450240.

Similar to previous studies of the T1000 cohort (69), participants were grouped based on DSM-

IV or DSM-5 diagnosis using the Mini International Neuropsychiatric Inventory 6 or 7 (MINI;

(70)), and included those with major depressive disorder only (DEP), an anxiety disorder only

(ANX; social anxiety, generalized anxiety, panic, or posttraumatic stress disorder), comorbid

MDD and anxiety disorder (DEP+ANX), substance use disorders (SUDs; recreational drugs

excluding alcohol and nicotine; with or without comorbid anxiety and mood disorders), eating

disorders (EDs; with or without comorbid anxiety and mood disorders), and HCs with no mental

health diagnoses. In this study we examined all groups from the first 500 participants of the

T1000 project (recruited from January 5, 2015 to February 22, 2017).

Table 1. Mean (and standard deviation) for clinical and demographic variables.

Individual difference variable*	Healthy comparison (N = 52)	Anxiety (N = 15)	Depression (N = 69)	Depression + Anxiety (N = 153)	Eating Disorder (N = 14)	Substance use disorder (N = 131)	p-value
Age	32.04 (11.08)	36.42 (10.01)	37.12 (11.83)	35.28 (11.23)	27.40 (9.83)	34.03 (8.88)	0.012
Sex	50% male	33% male	30% male	26% male	14% male	45% male	0.001
PHQ-9	0.83 (1.29)	7.47 (5.80)	13.48 (4.73)	13.10 (5.21)	12.93 (7.98)	6.68 (5.91)	<0.001
OASIS	1.37 (1.89)	10.60 (2.16)	7.55 (3.34)	10.51 (3.12)	9.50 (4.93)	5.94 (4.77)	<0.001
DAST-10	0.10 (0.30)	0.33 (0.49)	0.72 (1.66)	0.58 (1.13)	1.23 (2.62)	7.55 (2.05)	<0.001
PTT	0.20 (0.02)	0.20 (0.01)	0.19 (0.01)	0.20 (0.02)	0.20 (0.02)	0.20 (0.02)	0.261

| BMI | 27.59 (5.54) | 27.05 (6.02) | 28.51 (5.47) | 28.77 (5.49) | 22.25 (4.43) | 28.23 (4.56) | 0.001 |

191 *PHQ-9 = Patient Health Questionnaire 9; OASIS = Overall Anxiety Sensitivity and Impairment Scale;

192 DAST-10 = Drug Abuse Screening Test; PTT = median pulse transit time; BMI = Body Mass Index.

Heartbeat perception task

As part of the T1000 project, participants completed a large number of assessments, self-report

measures, and behavioral tasks (detailed in (64)). Here we focus on data from a cardiac

perception task on which we have previously reported (i.e., on a subset of the participants

reported here, with analyses unrelated to computational modeling (71, 72)), wherein participants

were asked to behaviorally indicate the times at which they felt their heartbeat. The utilization of

the heartbeat tapping measure as an index of perception was based on a previously developed

heartbeat tapping task (40); for a more recent example, see (73)). The task was repeated under

multiple conditions designed to assess the influence of cognitive strategy and physiological

perturbation on performance. In the initial task condition, participants were simply instructed to

close their eyes and press down on a key when they felt their heartbeat, to try to mirror their

heartbeat as closely as possible, and even if they weren't sure they should take their best guess

(the "guessing" condition). Participants completed this (and each other) task condition over a

period of 60 seconds. In the second task condition, all instructions were identical except that they

were told to only press the key when they actually feel their heartbeat, and if they do not feel

their heartbeat then they should not press the key (the "no-guessing" condition). In other words,

unlike the first time they completed the task, they were specifically instructed not to guess if they

didn't feel anything, which could be understood as altering prior beliefs associated with

confidence thresholds (i.e., higher confidence was required to choose to tap than in the guessing

condition, where this change in prior confidence thresholds could differ between individuals).

Finally, in the perturbation condition, participants were again instructed not to guess but were

also asked to first empty their lungs of all air and then inhale as deeply as possible and hold it for

as long as they could tolerate (up to the length of the one-minute trial) while reporting their

perceived heartbeat sensations. This third condition (the "breath-hold" condition) was used in an

attempt to putatively increase the strength of the afferent cardiac signal by increasing

physiological arousal. We expected that cardiac perception would be poor in the guessing

condition, that tapping would be more conservative in the no-guessing condition, and that the

breath-hold condition would result in improved performance on average. As a control condition,

we also included an identical task where participants were instructed to tap every time they heard

a 1000Hz auditory tone presented for 100ms (78 tones, randomly jittered by +/- 10% and

presented in a pattern following a sine curve with a frequency of 13 cycles/minute, mimicking

the range of respiratory sinus arrhythmia during a normal breathing range of 13 breaths per

minute). This was completed between the first (guessing) and second (no-guessing) heartbeat

tapping conditions.

Directly after completing each task condition, individuals were asked the following using a

visual analogue scale:

"How accurate was your performance?"

"How difficult was the previous task?"

"How intensely did you feel your heartbeat?"

Each scale had anchors of "not at all" and "extremely" on the two ends. Numerical scores could

range from 0 to 100.

Computational model

To model behavior on the heartbeat tapping task, we first divided each task time series into

intervals corresponding to the periods of time directly before and after each heartbeat. Potentially

perceivable heartbeats were based on the timing of the peak of the electrocardiogram (EKG) R-

wave (signaling electrical depolarization of the atrioventricular neurons of the heart) + 200

milliseconds (ms). This 200 ms interval was considered a reasonable estimate of participants'

pulse transit time (PTT) according previous estimates for the ear PTT (74). We also measured

the average PTT of each participant during a separate resting-state period, defined as the distance

between the peak of the EKG R-wave and the onset of the peak of the PPG waveform (signaling

mechanical transmission of the systolic pressure wave to the earlobe). These average PTT values

were used as covariates in some analyses (described below), and the group-level average PTT

value was imputed for individuals where PPG data quality was poor. The length of each

heartbeat interval (i.e., the "before-beat interval" and "after-beat interval") depended on the heart

rate. For example, if two heartbeats were 1 second apart, the "after-beat interval" would include

the first 500 ms after the initial beat and the "before-beat interval" would correspond to the 2^{nd}

500 ms. The after-beat intervals were considered the time periods in which the systole (heart

muscle contraction) signal was present and in which a tap should be chosen if it was felt. The

before-beat intervals were treated as the time periods where the diastole (heart muscle relaxation)

signal was present and in which tapping should not occur (i.e., assuming taps are chosen in

response to detecting a systole; e.g., as supported by (75)). This allowed us to formulate each

interval as a "trial" in which either a tap or no tap could be chosen and whether a systole or

diastole signal was present (see **Figure 1**).

Figure 1. Bayesian approach used to model interoceptive awareness on the heartbeat tapping task. The generative model is here depicted graphically, such that arrows indicate dependencies between variables. Associated vectors/matrices are also shown. At each time point (t), observations (o) depend on hidden states (s), where this relationship is specified by the **A** matrix, and those states depend on previous states (as specified by the **B** matrix, or the initial states specified by the **D** vector). This model represents a simplified version of a commonly used active inference formulation of partially observable Markov decision processes (for more details regarding the structure and mathematics describing these models, see (76-78). In this model, the observations were systole/diastole, and the hidden states included beliefs about the presence or absence of a heartbeat. Selection of the tap vs. no tap actions were sampled from the posterior distribution over states ($\bar{s}$) – that is, a higher posterior probability of a heartbeat state ($P(HB)$) corresponded to a higher probability of choosing to tap, and a higher posterior probability of the no heartbeat state ($P(nHB)$) corresponded to a higher probability of choosing not to tap. The model parameters we estimated corresponded to: 1) interoceptive precision (IP) – the precision of the mapping from systole/diastole to beliefs about heartbeat/no heartbeat in the **A** matrix, which can be associated with the weight assigned to sensory prediction errors; and 2) prior beliefs for the presence of a heartbeat (pHB). Because minimal precision corresponds to an IP value of .5, and both higher and lower values indicate that taps will more reliably track systoles (albeit in an anticipatory or reactive manner), our ultimate measure of precision subtracted 0.5 from raw IP values and then took their absolute value. The raw IP values were then used to assess for group differences in the tendency to tap before vs. after each systole. We also compared this model to an analogous model that included learning (see main text). On each trial, beliefs about the probability of a heartbeat (corresponding to the probability of choosing to tap) relied on Bayesian inference as implemented in the "heartbeat perception" equations shown at the bottom of the figure. Note that, by convention in the active inference models from which our model was derived, the dot product (·) applied to matrices here indicates transposed matrix multiplication. Observed diastoles and systoles were taken from ECG traces, after dividing the time series into periods before and after each systole (exemplified in the upper left portion of the figure; see text for details). Each time period in which a diastole or systole could be present was treated as a separate trial, in which the participant started in the 'start' state and then

updated beliefs about hidden states based on observation of a diastole or systole. For this reason, the

pHB parameter in the transition matrix (**B**) only specifies the probability of transitioning from the 'start'

state to the heartbeat vs. no heartbeat states, and the heartbeat vs. no heartbeat states simply have

identity mappings (i.e., a given trial cannot transition between these two states).

To model interoception and task behavior, we used a Bayesian model of perception derived from

a Markov decision process (MDP) formulation of active inference that has been used in previous

work; for more details about the structure and mathematics of this class of (discrete state space)

models, see (77, 79, 80). See **Figure 1** for a graphical depiction of our model and the associated

vectors and matrices. Observations (*o*) in the model were categorical and included systole,

diastole, and a "start" observation. Hidden states (*s*) in the model, which were inferred based on

observations, were also categorical and included a heartbeat state, a no heartbeat state, and a

"start" state. Each trial in the model corresponded to a window in the EKG time series in which

either a systole or diastole signal was present. Each trial formally had two timesteps (*t*=1 and

t=2). At *t*=1, the participant always formally began in the "start" state and made the associated

"start" observation. At *t*=2, the participant either observed a systole or a diastole and inferred

whether they had transitioned from the "start" state into the heartbeat state or the no heartbeat

state. In other words, they inferred a posterior distribution over states $P(\bar{s}_{t=2})$ that assigned a

probability to the heartbeat state and to the no heartbeat state, where this posterior distribution

was informed by 1) prior beliefs about the probability of transitioning from the "start" state to the

heartbeat state vs. the no heartbeat state, $P(s_{t=2}|s_{t=1})$, and 2) beliefs about the likelihood of

observing a systole or diastole given a heartbeat or no heartbeat state, $P(o_t|s_t)$.

A vector **D** encoded prior beliefs over initial states, $P(s_{t=1})$, which specified that the participant

always started the trial in the "start" state with a probability of 1 (see vector in upper left portion

of the model depiction in **Figure 1**). A matrix **B** encoded the probability that each state would

transition into any other state:

$$\boldsymbol{B} = P(s_{t=2}|s_{t=1}) = \begin{bmatrix} 0 & 0 & 0 \\ 1 - pHB & 1 & 0 \\ pHB & 0 & 1 \end{bmatrix}$$

Here, columns indicate (from left to right) the "start" state, the no heartbeat state, and the

heartbeat state at time $t=1$, and rows (from top to bottom) indicate the "start" state, the no

heartbeat state, and the heartbeat state at time $t=2$. The probability of transitioning from the

"start" state to a heartbeat state vs. a no heartbeat state was encoded by a parameter pHB, where

values above 0.5 indicate prior beliefs that transitions from the "start" state to the heartbeat state

are more likely (e.g., expecting a faster heart rate), and values below 0.5 indicate prior beliefs

that transitions from the "start" state to the heartbeat state are less likely (e.g., expecting a slower

heart rate). Note that the second and third columns simply indicate that, once entering a heartbeat

or no heartbeat state, this does not subsequently change within the trial (i.e., as subsequent

systole/diastole observations are modelled as subsequent trials).

A matrix **A** encoded the probability of observations given states:

$$\boldsymbol{A} = P(o_t|s_t) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & IP & 1 - IP \\ 0 & 1 - IP & IP \end{bmatrix}$$

Here, columns indicate (from left to right) the "start" state, the no heartbeat state, and the

heartbeat state, and rows (from top to bottom) indicate the "start" observation, the diastole

observation, and the systole observation. The probability of observing a systole or diastole if a

heartbeat or no heartbeat state were present was encoded by an "interoceptive precision"

parameter (IP). A value of 0.5 for IP indicates minimal precision – that is, that the probability of

observing a systole or diastole is 0.5 when in either the heartbeat or no heartbeat state. In

contrast, a value approaching 1 indicates high precision – that is, that the probability of

observing a systole is high when in a heartbeat state and low when in a no heartbeat state (and

vice-versa when observing a diastole). Importantly, however, IP values approaching 0 also

indicate high precision, in that observing a systole vs. diastole still provides strong evidence for a

heartbeat vs. a no heartbeat state (but where the probabilistic relationships are reversed).

Behaviorally, IP values approaching 0 would indicate that a participant consistently tapped the

button during the task in a way that reliably anticipated each upcoming systole. It is important to

note here that 'precision' in discrete state-space models, such as the one used here, takes on a

formally different meaning than in continuous state-space models, as can be seen in the

likelihood ($\mathbf{A}$) matrix specified above. While precision refers to inverse variance in continuous

state-space models, in discrete models it more simply reflects how peaked vs. flat the

probabilistic mapping is between states and observations.

Belief updating in the perception model was based on the following equations at times $t = 1$ and

$t = 2$, respectively:

$$\bar{s}_{t=1} = \sigma\left(\frac{1}{2}(\ln \mathbf{D} + \ln \mathbf{B} \cdot s_{t+1}) + \ln \mathbf{A} \cdot o_t\right)$$

$$\bar{s}_{t=2} = \sigma(\ln \boldsymbol{B} \, s_{t-1} + \ln \boldsymbol{A} \cdot o_t)$$

Because time $t = 1$ in each trial always included the "start" observation with a fully precise prior

belief in $\mathbf{D}$ for being in the "start" state, the posterior belief $\bar{s}_{t=1}$ always corresponded to a fully

precise belief of being in the "start" state. The equation for time $t = 2$ corresponds to Bayesian

inference, in which prior beliefs ($\ln\mathbf{B}s_{t-1}$) are integrated with the likelihood distribution and the

systole or diastole observation at the second timepoint ($\ln\boldsymbol{A}\ \cdot o_t$), and then converted into a

proper probability distribution via a softmax (normalized exponential) function $\boldsymbol{\sigma}(\cdot)$ – leading to

a posterior distribution over heartbeat states ($\bar{s}_{t=2}$). Note that, by convention in the active

inference literature, the dot notation ($\cdot$) here indicates a matrix transpose, meaning that $\boldsymbol{A}\cdot o_t =$

$\boldsymbol{A}^T o_t$.

Our response model formally included two actions, the choice to tap or not tap. This model made

the assumption that the probability of choosing to tap vs. not tap corresponded to the posterior

probability assigned to the heartbeat vs. no heartbeat state at time $t=2$ in each trial:

360
$$P(tap) = P(\bar{s}_{t=2} = heartbeat).$$

In other words, tapping behaviors were sampled from the posterior distribution over heartbeat vs.

no heartbeat states, such that choices to tap became more likely as the posterior probability of a

heartbeat state approached 1 and choices not to tap became more likely as the posterior

probability of a heartbeat state approached 0. No further parameters were included in the

response model to account for behavioral stochasticity. This is because, in the context of the

present task, parameters encoding randomness in behavior cannot be distinguished from IP, as

both effectively control the precision of the posterior distribution from which tapping actions are

sampled in response to the systole/diastole signal. As described further below, we instead took

further analysis steps at the group level to account for the potential influence of individual

differences in motor stochasticity on IP and pHB estimates.

Table 2. Description of computational model elements

Model variable	General Definition	Model-specific specification
t	Timepoint within a trial	There were 2 timepoints in each trial (i.e., for each EKG time window). At $t = 1$, the participant was modelled as waiting to infer the presence or absence of a heartbeat. At $t = 2$, either a systole or diastole observation was presented (depending on whether the EKG time window for that trial contained a systole or diastole), and a posterior probability of the presence vs. absence or a heartbeat was inferred.
o_t	Observable outcomes at time t	Outcomes: 1. Start 2. Systole 3. Diastole
s_t	Hidden states at time t	Hidden states: 1. Start 2. Heartbeat 3. No Heartbeat
A matrix $p(o_t \mid s_t)$	A matrix encoding beliefs about the relationship between hidden states and observable outcomes (i.e., the probability that specific outcomes will be observed given specific hidden states).	Encodes beliefs about the relationship between heartbeat vs. no heartbeat states and diastole vs. systole observations. The precision of the relationship between heartbeat/no heartbeat states and diastole/systole observations is controlled by a parameter IP, which specifies how much evidence systole provides for a heartbeat and how much evidence diastole provides for no heartbeat.
B matrix $p(s_{t+1} \mid s_t)$	A matrix encoding beliefs about how hidden states will evolve over time (transition probabilities).	Encodes the prior belief that either a heartbeat or no heartbeat would occur on each trial, as controlled by a parameter pHB.
D vector $p(s_{t=1})$	A matrix encoding beliefs about (a probability distribution over) initial hidden states.	Ensures the individual always begins in an initial starting state.

In this task, it is unclear whether or not learning over time (e.g., from paying attention to one's

heartbeat) contributes to task performance. To assess this using Bayesian model comparison, we

compared evidence for the "perception only" model described above with evidence for a model

that included learning to update prior beliefs over time. Learning within this model involves

updating beliefs about the probability of feeling a heartbeat vs. no heartbeat at each time point,

based on how frequently one believes they have felt their heartbeat in the past. Essentially, every

time a heartbeat is felt, prior beliefs favoring feeling a heartbeat go up, and every time no

heartbeat is felt this (relative) belief goes back down. Formally, this corresponds to updating the

concentration parameters of Dirichlet (Dir) priors associated with the **B** matrix (b) that specify

beliefs about state transitions. At $t = 0$:

383
$$P(\mathbf{B}) = Dir(b)$$

384
$$b(heartbeats) = P(s_{t+1}|s_t) = \begin{bmatrix} 0 & 0 & 0 \\ 1 - pHB & 1 & 0 \\ pHB & 0 & 1 \end{bmatrix} \times b_0$$

385
$$b_t = b_{t-1} + \sum_t s_t \otimes s_{t-1}$$

Here $\otimes$ indicates the cross-product, and b_0 is a scalar on the prior value for concentration

parameters, where its value prior to learning encodes (inverse) sensitivity to information, such

that higher values will reduce the rate at which prior beliefs are updated over time with new

observations. In the learning model, b_0 was also estimated for each individual to capture the

possibility of different learning rates for updating prior beliefs (this could also be thought of as

differences in a kind of interoceptive "belief rigidity").

Thus, the final parameters estimated for each participant included the IP, pHB, and b_0

parameters. Our approach to parameter estimation used Bayesian inference at two levels (81).

First, each participant's responses were modeled using the Bayesian model of perception

described above. We then used a commonly used Bayesian optimization algorithm (called

Variational Bayes) to estimate each participant's parameter values that maximized the likelihood

of their responses (under the assumption that a higher/lower probability assigned to feeling a

heartbeat corresponded to a higher/lower probability of choosing to tap), as described in (25).

We optimized these parameters for each model using this likelihood and variational Laplace

(82), implemented within the spm_nlsi_Newton.m parameter estimation routine available within

the freely available SPM12 software package (Wellcome Trust Centre for Neuroimaging,

London, UK, http://www.fil.ion.ucl.ac.uk/spm). This estimation approach has the advantage of

preventing overfitting, due to the greater cost it assigns to moving parameters farther from their

prior values. Estimating parameters required setting prior means and prior variances for each

parameter. The prior variance was set to a high precision value of 1/2 for each parameter (i.e.,

deterring overfitting), and the prior means were set as follows: IP = .5, pHB = .5, and b_0 = 1. Our

decision for selecting these priors was motivated in part by initial simulations confirming that

parameter values were recoverable under these prior values. The IP and pHB prior values were

further chosen to minimize estimate bias, as pHB = .5 assumes flat prior beliefs, and IP = .5 does

not bias estimates in favor of values assuming anticipatory vs. reactive strategies. The b_0 prior of

1 is equivalent to this parameter having no effect on the model. After fitting parameters for each

model, we then performed Bayesian model comparison (based on (83, 84)) to determine the best

model. We then used classical inference to test for the effects of group differences in parameter

estimates (i.e., posterior mean estimates) for the best model, using a standard summary statistic

approach.

Before using these parameters in further analyses, however, the "raw" IP parameter values

(IP_{raw}) were transformed to correctly capture 2 distinct constructs of interest. First, because

IP_{raw} values both above and below .5 indicate higher precision (i.e., values below approaching 0

indicate reliable anticipatory tapping, whereas values approaching 1 indicate reliable tapping

after a systole), our ultimate measure of precision was recalculated by centering IP_{raw} on 0 and

taking its absolute value as follows:

$$IP = |IP_{raw} - 0.5|$$

This means that IP has a minimum value of 0 and a maximum value of 0.5. The IP_{raw} values

were then instead used to assess individual differences in the tendency to tap in an anticipatory or

reactive (AvR) fashion:

$$AvR = IP_{raw}$$

Higher AvR values (> 0.5) thus indicated a stronger tendency to reactively tap in response to a

heartbeat as opposed to tapping in an anticipatory fashion (< 0.5).

Physiological measurements

Electrocardiography was used to assess the objective timing of participants' heartbeats

throughout the task. A BIOPAC MP150 was used to collect a three lead EKG signal and the

pulse oximeter signal, using a pulse plethysmography (PPG) device attached to the ear lobe.

Response times were collected using a task implemented in PsychoPy, with data collection

synchronized via a parallel port interface.

EKG and response data were scored using in-house developed MATLAB code. As described

above in relation to modelling, each participant's pulse transit time (PTT) was estimated as the

median delay between R wave and the corresponding inflection in the PPG signal.

Supplementary materials display an example participant's EKG and PPG trace to illustrate

how this was calculated (also supporting the 200ms delay assumption used in modelling and in

individuals without usable PPG).

Quality Control and Final Sample Sizes

Prior to performing our analyses, several participants were removed due to quality control

checks: 19 individuals were removed due to "cheating" (i.e., video revealed they were taking

their pulse while performing the task); 3 individuals didn't complete the task; and 13 individuals

had poor EKG across all trials that didn't allow reliable identification of heartbeat timing. An

additional 31 individuals were removed due to being outliers when performing the tone task

(using Iterative Grubb's with $p < 0.01$) – assumed to reflect inappropriate engagement during the

task (e.g., being inattentive, tapping rapidly without listening to the tones, etc.). This resulted in

434 participants, including 52 HCs, 15 ANX, 69 DEP, 153 co-morbid ANX/DEP, 131 SUDs,

and 14 eating disorders. In the three heartbeat tapping conditions, a few other participants were

removed due to poor EKG, leading to the following participant numbers in each condition: 433

(guessing), 431 (no-guessing), and 427 (breath-hold). **Table 3** reports the final number of

participants included for model-based analyses in each condition as well as descriptive statistics

for all task-related measures.

Statistical analysis

Correlational analyses were first conducted to examine the relationships between parameters

across conditions. For purposes of parameter validation, we ran further correlational analyses to

examine the relationships between each parameter and task-specific measures, including the self-

report ratings of difficulty, confidence, and heartbeat intensity collected after each trial.

Linear mixed effects models (LMEs; described further below) were conducted (using the lme4

package in R) to identify possible group differences (i.e., between HCs and the five patient

groups) for each parameter and in how they differed between conditions (i.e., guessing, no-

guessing, breath-hold), while accounting for individual differences in age, sex, BMI, median

PTT, number of heartbeats (and its interaction with group and condition), and medication status

(i.e., one analysis per parameter). Inclusion of the peripheral physiological variables was done to

rule out the possibility that differences in parameters between groups/conditions was due to

differences in peripheral physiology between groups/conditions. We also included sensory

precision estimates for the tone condition in the models in order to account for any variability in

tapping behavior due to motor stochasticity (e.g., differences in reaction times). This was based

on the assumption that, because the sensory signal in the tone condition is highly precise, any

variability in precision would be better explained by random influences on behavior as opposed

to perception. Because the first half of the T1000 sample was pre-specified as an exploratory

sample, we report relationships at $p < .05$ as a potential basis for a priori hypotheses verification

in the second (confirmatory) half of the T1000 dataset in subsequent work. Although our

analyses are exploratory, we note that a Bonferroni corrected threshold for multiple comparisons

with three parameters is $p < .017$ ($\alpha=0.05$).

For each parameter we ran LMEs (with the covariates mentioned above) that compared the three

task conditions by group. Given our transdiagnostic focus, we first divided participants into two

groups comprised of healthy participants and those diagnosed with one or more psychiatric

disorders (transdiagnostic group). We subsequently divided those with psychiatric diagnoses into

the 5 distinct diagnostic groups to assess the potential for diagnosis-specific differences.

To test the relative evidence for models with vs. without group and condition effects, we ran a

JZS Bayes factor analyses with default prior scales in R (85, 86) comparing null models (with

only an effect of subject) to the space of models including all combinations of main effects and

interactions for group and condition. We also compared evidence for models with

transdiagnostic vs. condition-specific effects.

Results

Participant characteristics

Complete information on sample size, demographics, and symptom screening measures is

provided in **Table 1**. Separate ANOVAs showed significant differences between groups in age

($F(5,428) = 2.98$, $p = .01$) and body mass index (BMI; $F(5,413) = 4.04$, $p = .001$). A chi-squared

analysis also showed significant differences in the proportion of males to females between

groups (chi-squared $= 19.43$, $df = 5$, $p = .002$). Therefore, in our analyses of model parameters,

we also confirm our results after controlling for these other factors.

Interoceptive Perturbation Validation

As expected, across all participants self-reported heartbeat intensity, confidence in task

performance, and task difficulty differed significantly between the three heartbeat tapping

conditions in separate LMEs (intensity: $F(2858) = 79.09$, $p < .001$; confidence: $F(2,858) =$

36.23, $p < .001$; difficulty: $F(2,858) = 31.8$, $p < .001$), reflecting a greater perceived intensity of

heartbeat sensations and greater confidence during the breath-hold perturbation than in the other

2 conditions, as well as lower difficulty in the breath-hold condition than the no-guessing

condition (and greater difficulty in the no-guessing than guessing condition; $p < .001$ for all post-

hoc comparisons). An LME analysis of heart rate revealed a significant difference in the number

of heartbeats between conditions ($F(2,856) = 15.08$, $p < .001$), reflecting a faster heart rate in the

breath-hold condition than in the no-guessing condition ($p < .001$) and guessing condition ($p =$

.001)

Table 3. Summary statistics for all task-related variables.

Panel A.

Guessing Condition	**Healthy comparisons (N = 52)**	**Anxiety (N = 15)**	**Depression (N = 69)**	**Depression + Anxiety (N = 153)**	**Eating Disorder (N = 14)**	**Substance use disorder (N = 130)**	**p-value***
Number of taps	67.92 (29.58)	58.40 (28.40)	67.65 (31.97)	62.02 (32.58)	65.86 (27.86)	65.50 (30.82)	0.701
Number of heartbeats	66.67 (9.48)	67.13 (9.55)	70.23 (10.90)	70.34 (10.86)	67.21 (8.71)	71.23 (10.47)	0.1
Self-reported difficulty	56.87 (22.70)	53.20 (26.69)	57.55 (28.12)	53.18 (25.84)	71.57 (17.78)	48.24 (24.65)	0.009
Self-reported confidence	32.67 (19.27)	41.27 (21.54)	39.52 (22.67)	39.29 (21.20)	25.14 (21.36)	42.20 (22.39)	0.022
Self-reported intensity	20.71 (16.72)	26.73 (21.51)	26.26 (21.52)	28.27 (23.25)	15.36 (18.64)	35.94 (24.65)	<0.001
Counting accuracy	0.69 (0.33)	0.67 (0.30)	0.65 (0.29)	0.63 (0.30)	0.68 (0.30)	0.67 (0.29)	0.755

509   *p-values correspond to the results of ANOVAs comparing the groups (i.e., not including all task
510   conditions within the analysis as reported in the main text).

512   *Panel B.*

No-Guessing Condition	**Healthy comparisons (N = 52)**	**Anxiety (N = 15)**	**Depression (N = 68)**	**Depression + Anxiety (N = 153)**	**Eating Disorder (N = 14)**	**Substance use disorder (N = 129)**	**p-value**
Number of taps	18.73 (22.69)	22.13 (24.04)	23.43 (24.22)	24.10 (24.26)	18.29 (21.74)	31.47 (25.46)	0.015
Number of heartbeats	65.88 (10.22)	66.73 (11.07)	69.97 (10.45)	69.78 (11.10)	66.14 (8.43)	70.69 (10.14)	0.066
Self-reported difficulty	67.83 (27.94)	59.73 (35.14)	68.00 (32.72)	62.88 (31.35)	69.86 (26.68)	55.14 (30.75)	0.039
Self-reported confidence	39.94 (30.32)	36.80 (35.54)	41.15 (31.36)	41.41 (29.49)	31.93 (32.16)	42.29 (28.27)	0.858
Self-reported intensity	18.15 (19.63)	16.80 (22.11)	25.82 (26.38)	25.31 (25.67)	13.50 (13.51)	33.64 (27.53)	0.001
Counting accuracy	0.28 (0.29)	0.31 (0.33)	0.32 (0.31)	0.33 (0.31)	0.28 (0.32)	0.43 (0.32)	0.028

Panel C.

Breath Hold Condition	Healthy comparisons (N = 52)	Anxiety (N = 15)	Depression (N = 68)	Depression + Anxiety (N = 151)	Eating Disorder (N = 14)	Substance use disorder (N = 127)	p-value
Number of taps	27.12 (21.47)	30.60 (42.93)	24.75 (21.07)	26.46 (23.58)	17.71 (16.80)	34.17 (25.05)	0.027
Number of heartbeats	69.40 (9.80)	66.87 (12.01)	71.26 (10.48)	70.59 (11.20)	66.29 (6.29)	72.96 (10.75)	0.061
Self-reported difficulty	46.44 (27.08)	54.20 (23.24)	56.53 (25.74)	53.50 (28.32)	55.64 (25.67)	45.86 (29.08)	0.07
Self-reported confidence	47.79 (26.20)	41.27 (26.06)	50.78 (26.49)	52.42 (27.42)	36.50 (33.37)	53.04 (25.55)	0.158
Self-reported intensity	38.19 (28.40)	26.40 (23.61)	41.85 (29.60)	41.09 (29.99)	32.71 (30.63)	46.81 (28.88)	0.074
Counting accuracy	0.39 (0.29)	0.28 (0.26)	0.34 (0.27)	0.35 (0.29)	0.28 (0.29)	0.44 (0.30)	0.03

Panel D.

Tone Condition	Healthy comparisons (N = 52)	Anxiety (N = 15)	Depression (N = 69)	Depression + Anxiety (N = 153)	Eating Disorder (N = 14)	Substance use disorder (N = 131)	p-value
Number of taps	77.90 (1.00)	77.40 (1.12)	77.71 (1.03)	77.82 (1.17)	77.57 (0.94)	78.06 (0.97)	0.072
Number of heartbeats	65.63 (11.10)	67.27 (9.81)	70.74 (9.94)	71.37 (11.14)	66.86 (8.16)	71.30 (10.62)	0.01
Self-reported difficulty	18.04 (15.84)	29.20 (21.12)	22.14 (19.52)	24.15 (22.00)	35.21 (22.33)	25.77 (22.40)	0.065
Self-reported confidence	77.56 (14.26)	73.93 (17.84)	78.84 (16.60)	74.22 (19.59)	59.79 (24.84)	75.16 (18.56)	0.017
Self-reported intensity	85.58 (13.76)	75.67 (15.19)	86.20 (15.09)	82.86 (17.83)	87.79 (12.46)	81.47 (16.51)	0.102
Counting accuracy	0.99 (0.01)	0.99 (0.01)	0.99 (0.01)	0.99 (0.01)	0.99 (0.01)	0.99 (0.01)	0.922

Bayesian Model Comparison

When comparing models (based on (83, 84)), there was more evidence for the "perception only"

model than for the model that included learning prior beliefs for the no-guessing and breath-hold

conditions (protected exceedance probability = 1), whereas there was not clear evidence favoring

a single model for the guessing condition (protected exceedance probabilities = .46 vs. .54,

slightly favoring the learning model). The learning model was favored in the tone condition

(protected exceedance probability = 1). No group differences were observed when comparing

model fits between groups.

For consistency/comparability, we use the "perception only" model parameters to compare

conditions in our analyses below, as this model best explained heartbeat tapping behavior

overall. The accuracy of this model – defined as the percentage of choices to tap/not tap that

matched the highest probability action in the model (e.g., a tap occurring when the highest

probability percept in the model was a heartbeat) – was 74% across all conditions; by condition,

model accuracy was: guessing condition = 67% (SD = 12%); tone condition = 67% (SD = 12%);

no-guessing condition = 85% (SD = 14%); breath-hold condition = 84% (SD = 13%).

Relationship between parameters

Parameter values for each group and condition are listed in **Table 4**. Across conditions, all

parameters were normally distributed (skew < 2). As shown in **Figure 2**, correlations between IP

across task conditions were generally low. Correlations between pHB estimates across conditions

were moderate, most notably between the no-guessing and breath-hold conditions (which also

included the no-guessing instruction). The tendency to tap in an anticipatory vs. reactionary

manner (AvR) showed no relationships across conditions. Correlations between IP and pHB (or

pTone) within each condition were also low ($0 \leq r \leq .27$), as were correlations between these

parameters and AvR ($-.28 < r < .01$; see **supplementary materials**).

Figure 2. Pearson correlations between model parameters across task conditions across all participants. For reference, correlations at p < .001 (uncorrected) are marked with red asterisks.

Parameter face validity

Figure 3 shows the correlations, including some significant relationships ($p \leq .001$, uncorrected), across all participants between model parameters in each condition and several task-relevant variables. IP showed positive relationships with self-reported heartbeat intensity ratings in both the no-guessing and breath hold conditions. Additionally, pHB was lower in those self-reporting greater difficulty in the no-guessing condition, and higher in those self-reporting higher confidence and higher heartbeat intensity in both the no-guessing and breath hold conditions. Across heartbeat tapping conditions, model parameters were also weakly (IP) to strongly (pHB) related to the traditional counting accuracy measure (39).

In the breath-hold condition, t-tests revealed significantly greater IP in males than females ($t(251) = 2.30, p = .02$).

Figure 3. Exploratory Pearson correlations between model parameters and self-report and other task-
relevant variables for each task condition across all participants. IP = interoceptive precision parameter,
pHB = prior belief for heartbeat parameter, pT = prior belief for tone parameter, AvR = anticipate vs.
react strategy parameter, PTT = median pulse transit time, #HBs = number of heartbeats during the task
condition, BMI = body mass index. For reference, correlations at p < .001 (uncorrected) are marked with
red asterisks.

Group by condition interactions in model parameters

567 **Table 4**. Mean (and standard deviation) for model parameters by group and condition.

Individual difference variable	Healthy comparisons	Anxiety	Depression	Depression + Anxiety	Eating disorder	Substance use disorder	p-value*
Sensory precision							
Guessing	0.04 (0.05)	0.04 (0.02)	0.04 (0.03)	0.04 (0.04)	0.05 (0.05)	0.04 (0.04)	0.815
No-Guessing	0.04 (0.05)	0.04 (0.04)	0.03 (0.04)	0.03 (0.03)	0.06 (0.06)	0.04 (0.05)	0.09
Breath-hold	0.07 (0.07)	0.03 (0.03)	0.04 (0.04)	0.04 (0.04)	0.03 (0.03)	0.04 (0.04)	<0.001
Tone	0.18 (0.12)	0.12 (0.11)	0.18 (0.12)	0.14 (0.11)	0.21 (0.10)	0.16 (0.11)	0.052
Prior beliefs							
Guessing	0.44 (0.16)	0.37 (0.16)	0.41 (0.17)	0.38 (0.18)	0.43 (0.18)	0.40 (0.16)	0.331

No-Guessing	0.15 (0.13)	0.15 (0.12)	0.17 (0.13)	0.17 (0.14)	0.14 (0.12)	0.21 (0.14)	0.05
Breath-hold	0.18 (0.11)	0.19 (0.19)	0.17 (0.11)	0.18 (0.13)	0.14 (0.11)	0.22 (0.13)	0.056
Tone	0.50 (0.01)	0.50 (0.01)	0.50 (0.01)	0.50 (0.01)	0.50 (0.01)	0.50 (0.01)	0.139
Anticipate vs. React							
Guessing	0.49 (0.06)	0.50 (0.04)	0.49 (0.05)	0.49 (0.06)	0.54 (0.07)	0.50 (0.06)	0.042
No-Guessing	0.50 (0.07)	0.52 (0.06)	0.49 (0.05)	0.50 (0.05)	0.50 (0.08)	0.50 (0.06)	0.683
Breath-hold	0.49 (0.10)	0.49 (0.04)	0.51 (0.06)	0.50 (0.06)	0.48 (0.04)	0.50 (0.05)	0.529
Tone	0.50 (0.22)	0.45 (0.16)	0.52 (0.21)	0.49 (0.18)	0.37 (0.20)	0.46 (0.19)	0.088

*p-values correspond to the results of exploratory ANOVAs comparing the groups (i.e., not including all
task conditions within the analysis as reported in the main text).

In this section, we present LME analyses assessing the main effects of task condition for each

parameter, and then subsequent LMEs that exclude the tone condition, in which we assess the

main effects of group and task condition, and their interaction, while accounting for several

covariates that account for potential confounding effects of other demographic, behavioral, and

peripheral physiological variables (see methods). In LMEs including BMI as a covariate, 15

participants were removed due to lack of available data (2 HCs, 1 ANX, 3 DEP, 1 DEP+ANX, 7

SUDs, and 1 ED).

Due to our transdiagnostic focus, we examined the transdiagnostic vs. condition-specific nature

of these effects both when dividing participants into two groups (healthy comparisons vs.

patients transdiagnostically) and six groups (healthy comparisons and each of the five patient

groups separately). We then perform Bayes factor analyses to compare evidence for the space of

possible models with vs. without group or task effects, including models with transdiagnostic

effects, and models with condition-specific effects (see methods). As this is a pre-defined

exploratory sample, we report post-hoc comparisons as significant with an uncorrected threshold

of $p < .05$.

Interoceptive Precision. An initial LME revealed that sensory precision (IP or auditory

precision in the tone condition) was significantly different between conditions ($F(3,1292) =$

339.4, $p < .001$; see **Figure 4**), reflecting greater values in the tone condition than in the three

heartbeat tapping conditions (post-hoc comparisons indicated $p < .001$ in all cases). Subsequently

focusing only on IP (excluding the tone condition), and grouping participants transdiagnostically

(i.e., based on the presence vs. absence of a diagnosis), a further LME revealed a significant

group by condition interaction ($F(2,839) = 10.30$, $p < .001$), but no main effects of group

($F(1,607) = 0..54$, $p = .46$) or condition ($F(2,885) = 1.35$, $p = .26$). Post-hoc comparisons

indicated that the interaction was driven by: 1) the fact that IP significantly increased in HCs in

the breath-hold condition when compared to the two resting conditions ($p < .001$ for each), 2)

that IP in HCs during breath-hold was significantly greater than IP in all three conditions in the

transdiagnostic patient sample ($p < .001$ in all cases), and 3) that there were no significant

differences between conditions in the patient sample. This LME also revealed no significant

effect of age, sex, precision within the tone condition, BMI, median PTT, medication status,

number of heartbeats, and the interaction between number of heartbeats and both group and

condition (*Fs* between .02 and 1.7, *ps* between .19 and .88*).

A further LME including each diagnostic group separately confirmed the same group by

condition interaction ($F(10,821) = 3.21$, $p < .001$). Here, post-hoc comparisons showed that IP in

the breath-hold condition was significantly greater in HCs when compared to each clinical group

(ANX, $p = .002$; DEP, $p < .001$; DEP+ANX, p $p < .001$, ED, $p = .007$, SUDs, $p < .001$). No other

clinical group showed a significant difference between conditions, with the exception of

DEP+ANX, which showed lower IP in the no-guessing condition than in the guessing condition

($p = .03$). DEP also showed lower IP than ED in the no-guessing condition ($p = .03$).

In the Bayesian analyses, the transdiagnostic model (HCs vs. all patients) that included main

effects of group, condition, and their interaction, was most strongly favored over the null model

(Bayes factor = 142.5), while the disorder-specific model was not favored over the null model

(Bayes factor = .18). The transdiagnostic model was also strongly favored over the diagnosis-

specific model (Bayes factor = 796.7). See **Figure 5** for plots of IP values by group and

condition.

Prior expectations. An LME revealed that prior expectations (pHB or priors in the tone

condition) were significantly different between conditions ($F(3,1291) = 861.5$, $p < .001$; see

Figure 4), reflecting greater values in the tone condition than in the three heartbeat tapping

conditions, and greater values in the guessing condition than in the other two heartbeat tapping

conditions (post-hoc comparisons indicated $p < .001$ in all cases). Subsequently focusing only on

pHB (excluding the tone condition), and grouping participants transdiagnostically, a further

LME revealed a main effect of condition ($F(2,863) = 27.09$, $p < .001$), and a group by condition

interaction ($F(2,842) = 4.38$, $p = .01$), but no main effect of group ($F(1,722) = .03$, $p = .86$). Post-

hoc comparisons again indicated greater pHB in the guessing condition than in the other two

conditions ($p < .001$). However, post-hoc comparisons did not show any significant differences

between groups by condition to account for the interaction. This LME also revealed a negative

association between pHB and age ($F(1,408) = 8.84$, $p = .003$) and a heart rate by condition

interaction ($F(2,865) = 3.56$, $p = .03$), but no significant effect of sex, tone precision, BMI,

median PTT, medication status, number of heartbeats, or the interaction between number of

heartbeats and group (*Fs* between .02 and 3.49, *ps* between .06 and .90*)*. The heart rate by

condition interaction reflected a stronger negative correlation between heart rate and pHB in the

guessing condition (r = -.13) than in the no-guessing and breath-hold conditions (r = -.04 and

.03, respectively).

A further LME with diagnosis-specific groups confirmed the same group by condition

interaction ($F(10,821)$ = 2.21, p = .02). Here, post-hoc comparisons suggested this interaction

was driven primarily by the SUD group, who showed higher pHB values in the breath-hold

condition than the DEP, DEP+ANX, and ED groups (p = .048, .03, and .04, respectively), and

also showed higher pHB values in the no-guessing condition than the DEP+ANX, ED, and HC

groups (p = .03, .046, and .01, respectively). HCs also showed higher pHB values in the guessing

condition than the DEP+ANX group (p = .03).

In the Bayesian analyses, the most strongly favored model over the null model only included a

main effect of condition (Bayes factor = 1.07E+135). This model was also favored over 1) a

model including both main effects and the transdiagnostic group by condition interaction (Bayes

factor = 4.8), and 2) an analogous model with the diagnosis-specific group by condition

interaction (Bayes factor = 63.21). See **Supplementary materials** for plots of pHB values by

group and condition.

Anticipating vs. reacting. An initial LME showed no significant differences in AvR between

task conditions ($F(3,1292)$ = 2.01, p = .11; see **Figure 4**). Detailed results of further LMEs and

Bayes factor analyses for AvR are presented in **Supplementary materials**. No clear evidence

was found for differences between HCs and the clinical groups by condition, although some

potential relationships with heart rate and medication status were noted. Bayesian analyses

strongly favored the null model over all other models. See **Supplementary materials** for plots

of AvR values by group and condition.

Comparison to existing measures

To examine the ability of traditional measures to capture similar group differences, we ran

analogous LMEs using the traditional heartbeat counting task formula for interoceptive accuracy

(39). When using the transdiagnostic grouping, analyses did not find a main effect of group

($F(1,688) = .01$, $p = .92$) or condition ($F(2,868) = 2.4$, $p = .09$), but did detect an interaction

between group and condition ($F(2,842) = 3.37$, $p = .03$). There was negative association with age

($F(1,408) = 4.59$, $p = .03$) and PTT ($F(1,412) = 4.83$, $p = .03$), and no effect of any other

covariate (Fs between .0003 and 1.09, ps between .30 and .99). Post-hoc comparisons suggested

the group by condition interaction was driven by greater counting accuracy in the guessing

condition than the other two conditions in both groups ($p < .001$ in all cases), and greater

counting accuracy in the breath-hold condition than the no-guessing condition in HCs but not in

the transdiagnostic group ($p = .045$ and $p = .55$, respectively). When using the diagnosis-specific

grouping, there was no main effect of group ($F(5,620) = .23$, $p = .95$) or group by condition

interaction ($F(10,821) = 1.46$, $p = .15$). A Bayes factor analysis showed that the most highly

favored model (relative to a null model) only included an effect of condition (Bayes factor =

1.7E+62), reflecting higher counting accuracy in the guessing condition than in the other two

heartbeat tapping conditions ($p < .001$; also see **Table 3**). It was also strongly favored relative to

the second-best model including an effect of group and condition (Bayes factor = 7.4).

To assess potential group differences in the effect of task condition on self-reported experience

and physiology, we also carried out analogous LMEs assessing confidence, intensity, and

difficulty, as well as heart rate. These results are reported in the **supplementary materials**. No

group by condition interactions in these variables were observed mirroring our IP results.

Figure 4. Bottom: Bar plots illustrating means and standard errors for model parameters by task condition across all participants. Prior beliefs for heartbeats were lower in the no-guessing and breath-hold conditions. Sensory precision (i.e., interoceptive precision for the heartbeat or auditory precision for the tone condition) was much greater in the tone condition. This was expected given the unambiguous nature of this signal relative to the heartbeat signal. There were no significant differences in sensory precision between the heartbeat conditions. The Anticipate vs. React values revealed no mean differences between conditions. Top: For more complete data characterization, we also show raincloud plots depicting the same results in terms of individual datapoints, boxplots (median and upper/lower quartiles), and distributions. These illustrate that, for the Anticipate vs. React parameter, nearly equally sized clusters of participants appeared to adopt more anticipatory (<.5) vs. reactive (>.5) strategies in the tone condition, and that prior beliefs remained unbiased (.5; with little variance) in the tone condition relative to the heartbeat tapping conditions.

Figure 5. Bottom: Mean and standard error for interoceptive precision estimates by condition and clinical group. Interoceptive precision (IP) was significantly greater in healthy comparisons than all other groups in the breath-hold condition, and healthy comparisons showed a significant increase in IP from the guessing and no guessing condition to the breath-hold condition that was absent in the other groups. Top: For more complete data characterization, we also show raincloud plots depicting the same results in terms of individual datapoints, boxplots (median and upper/lower quartiles), and probability densities.

Associations with symptom severity measures and interoceptive awareness scales

Given the heterogeneity in our clinical sample, we ran subsequent exploratory correlational

analyses with continuous scores on the clinical measures gathered, excluding HCs, to assess

whether model parameters might provide additional information about symptom severity. We

note a few weak relationships at uncorrected levels in **supplementary materials**, but none

survive correction for multiple comparisons.

The T1000 dataset also includes self-report measures commonly used in interoception research,

including the Multidimensional Assessment of Interoceptive Awareness (MAIA; (87)), the

Toronto Alexithymia Scale (TAS-20; (88)), and the Anxiety Sensitivity Index (ASI; (89)). For

708   the interested reader, we also show exploratory correlation matrices between model parameters

709   and these common measures within **supplementary materials**. While a couple of relationships

710   were significant at uncorrected levels, the strength of these relationships was low.

Discussion

This investigation aimed to examine whether a novel Bayesian computational model of

perception could provide a principled approach to empirically characterizing interoceptive

processing dysfunctions that have previously been proposed. Specifically, we used behavior

during a heartbeat perception task in conjunction with this model to estimate quantitative

differences in the prior beliefs and sensory precision estimates that individuals implicitly apply

to afferent interoceptive (cardiac) signals – in both healthy individuals and a transdiagnostic

sample of individuals with depression, anxiety, substance use, and/or eating disorder symptoms.

We observed several relationships in the expected directions between model parameters and

other task-related variables that supported the construct validity of our model parameters.

Of greatest interest, we found evidence, using both frequentist and Bayesian analyses, that an

interoceptive (breath-hold) perturbation increased the precision estimates assigned to cardiac

signals in healthy individuals, but that this perturbation had no effect on interoceptive precision

in individuals with a range of psychiatric disorders, including depression, anxiety, substance use,

and/or eating disorders (**Figure 5**). Notably, Bayesian analyses found evidence against

differences in interoceptive precision between the different clinical groups, suggesting that low

precision during perturbation represents a transdiagnostic feature across these psychiatric

conditions. Bayesian analyses also found evidence against differences in prior expectations

between healthy participants and psychiatric patients, suggesting that differences in sensory

precision, and not prior beliefs, best account for interoceptive differences in the clinical groups

(although frequentist analyses suggested that substance users may show less attenuated prior

expectations than some other groups in the no-guessing and breath-hold conditions). Model

comparison further suggested that individuals did not update prior beliefs over time during the

task, which is perhaps unsurprising given the low cardiac awareness commonly seen in previous

studies (6). That is, individuals may not have had sufficient signal to learn from (note that, in

contrast, model comparison supported the presence of learning in our auditory control condition

with a clear signal). We expand on these points below.

Parameter validity/sensitivity

IP showed positive relationships with self-reported heartbeat intensity ratings (no-guessing and

breath-hold conditions) – as would be expected in the context of more precise cardiac signals. In

the no-guessing and breath-hold conditions, pHB was lower than in the guessing condition. In

other words, participants appear to have successfully adjusted their prior beliefs to comply with

the no-guessing instructions. Further, consistent with the role of prior beliefs in perception, pHB

was also lower in those reporting greater difficulty (no-guessing condition) and higher in those

reporting greater confidence and higher heartbeat intensity (no-guessing and breath-hold

conditions). Each of these results support the notion that our parameters tracked theoretically

meaningful individual differences in perception/behavior and that our approach can disentangle

the effects of sensory precision from those of prior beliefs and anticipatory vs. reactive strategies.

Further, while model parameters had some shared variance with traditional accuracy measures

(e.g., higher counting accuracy was weakly associated with higher interoceptive precision), they

mainly captured unique variance that was not tracked by standard measures. Further, no group

differences analogous to those seen with model parameters were found using the traditional

interoceptive accuracy measure (i.e., across all participants, counting accuracy showed an

opposite pattern, being highest in the guessing condition). This highlights the unique ability of

this computational method to uncover differences in perceptual decision making across different

psychiatric subtypes.

It is also worth noting the strong positive correlation we observed between pHB and counting

accuracy in the no-guessing and breath-hold conditions, suggesting that heartbeat counting

accuracy primarily reflects prior beliefs (as previously proposed in (38)). In the present task this

is explained by the fact that higher pHB values led to a greater number of taps and the fact that

the average number of taps in these conditions was low (i.e., because of the no-guessing

instruction). Thus, those who tapped more often (i.e., due to stronger priors, independent of

precision) approached the actual number of heartbeats and therefore had higher counting

accuracy scores. This highlights one specific way in which, in the context of restrictions on

guessing, counting accuracy may be most closely associated with prior beliefs.

Differences in Interoceptive Precision

Our primary results were that: 1) IP was significantly higher in the breath-hold condition in

healthy comparisons than in each of the clinical groups; and 2) there was a group by condition

interaction, demonstrating that the interoceptive perturbation (breath-hold) increased IP (relative

to resting conditions) in the healthy participants, whereas this perturbation had no effect in any of

the clinical groups. These group differences were not diagnosis-specific and were not accounted

for by any other demographic (e.g., age, sex) or physiological variables (e.g., pulse transit time,

changes in heart rate), suggesting it was not a result of group-specific changes in peripheral

physiology between conditions. Thus, as the clinical groups had comparable IP to healthy

participants during resting conditions, it appears that interoceptive processing differences in

psychiatric populations may not be present across all physiological contexts. Instead,

interoceptive processing differences may manifest selectively when in altered physiological

states (e.g., such as the high arousal states linked with anxiety).

The finding that IP was reduced during the breath-hold across psychiatric groups included in this

study may be of clinical interest. First, multiple neurocognitive (55, 90, 91) and computational

(11, 12, 15, 27, 92) theories of emotion, and associated empirical findings (e.g.,(42, 93-97)),

suggest that interoceptive awareness may be an important transdiagnostic factor in promoting

emotional awareness. As low emotional awareness has been linked to multiple psychiatric and

systemic medical conditions (reviewed in (98, 99)), reduced IP during altered interoceptive states

(e.g., affective arousal) could contribute to low emotional awareness and its maladaptive

consequences irrespective of diagnostic category. Second, visceral regulation might be expected

to be less effective in the absence of precise feedback signals from the body, which could relate

to visceral dysregulation in psychiatric conditions (e.g., see (56)). For example, to adaptively

regulate visceral states, visceromotor control regions plausibly require reliable feedback signals

from the body to determine whether descending visceromotor commands have successfully led

to the intended adjustments in internal bodily states (10, 13, 36). Thus, the low IP observed

across multiple psychiatric conditions during the breath-hold could potentially contribute to

visceral dysregulation in other contexts in which interoceptive perturbations occur, such as

highly arousing, negatively valenced states (e.g. panic anxiety, irritability/anger etc.).

These results build on previous bodies of work suggesting associations between psychiatric

disorders and interoceptive processing deficits (38, 56, 100-103). For example, previous cardiac

perception studies have shown that depressed patients exhibit reduced accuracy on a heartbeat

counting task (2-4), and that performance is negatively correlated with depressive symptoms (5)

as well as associated with both lower positivity and poorer decision-making (2); although, the

limitations of heartbeat counting tasks should be kept in mind when interpreting such findings

(47, 49, 63, 104). While the literature on interoceptive dysfunction is mixed for anxiety disorders

broadly, it is well-stablished in panic disorder (reviewed in (6)). A couple recent studies have

also reported evidence of differences in interoceptive processing in both SUDs (e.g., blunted

brain responses (7)) and eating disorders (e.g., stronger effects of prior beliefs on perception

during low arousal (8)). The computational framework within which our findings were observed

is also in line with several recent proposals about the role of interoceptive inference in guiding

(predictive) autonomic control and the potential breakdown of this mechanism within different

psychiatric conditions (10-15, 36, 38); however, in contrast to previous emphasis on altered prior

beliefs in these proposals, our results more selectively support the existence of deficits in

adjusting precision estimates for afferent interoceptive signals – and do not provide strong

support for the presence of altered priors.

That said, the neurobiological mechanisms promoting reduced IP in mental disorders during

interoceptive perturbation remains unclear. As IP increased during the interoceptive perturbation

in HCs, but not in the clinical groups, it could be that altered brain processes in individuals with

certain psychiatric disorders fail to update IP estimates during states of acute bodily arousal. The

neural process theory associated with active inference and related Bayesian models suggests an

inability to adjust IP estimates would correspond to reductions in synaptic plasticity in response

to changes in patterns of interoceptive prediction-errors (10, 24, 77, 79, 105), most plausibly

within neural networks supporting interoception and visceromotor control (e.g., insula and

anterior cingulate (15, 106)). This could be tested using our model/task in conjunction with

neuroimaging. Alternatively, IP estimates could be accurate, and afferent interoceptive signals

may in fact be conveyed with less fidelity (i.e., greater noise) to the brain in the context of

psychiatric conditions. For example, such differences could be due to altered signaling in

interoceptive sensory neurons, which could in principle be affected by many factors

(genetic/epigenetic influences, early adversity and related socio-environmental factors, and/or

effects of disease-related chronic stress, among others). Future research will need to investigate

these different possibilities.

Prior expectations and task strategy

Unlike the results for sensory precision described above, our results did not provide strong

support for group differences in prior expectations or the tendency to tap in an anticipatory vs.

reactive manner. While frequentist analyses suggested a few potential group differences in prior

beliefs in the no-guessing and breath-hold conditions, Bayesian analyses found strong evidence

favoring a model without group differences over a model with transdiagnostic or diagnosis-

specific effects. This suggests that baseline expectations for heart rate do not differ in the

psychiatric populations examined here, and that these priors can be adjusted in response to

instructions in the same manner as in healthy participants. The lack of group differences with

respect to strategy might also be seen as supporting a similar conclusion in that one might expect

greater tendencies to tap in an anticipatory manner if basing behavior on prior expectations.

Thus, further study of differences in the ability to adjust precision estimates may be more

promising for improving our understanding of interoceptive dysfunction in psychiatric

populations. However, the finding in frequentist analyses that pHB in substance users was less

attenuated relative to some other groups in the no-guessing and breath-hold conditions could

suggest that less sensory evidence is required to generate changes in interoceptive feelings within

these individuals (consistent with the greater self-reported heartbeat intensity and lower self-

reported difficulty in this group relative to the other groups suggested by our supplementary

analyses; see **Table 3** and **Supplementary Materials**). This possibility is worth further

investigation in future studies.

Strengths, Limitations, and Conclusion

This study has several major strengths. First is the novel application of a computational model to

behavior on an interoceptive awareness task, which allowed for model comparison (e.g.,

allowing us to rule out learning effects) and parameter estimates to disentangle distinct

computational mechanisms (e.g., the role of prior beliefs vs. interoceptive precision). A second

strength is the application of this model to interoceptive processing in individuals with

psychiatric disorders, which to our knowledge, has never been reported. While accounting for

other influences, this approach allowed us to test a theoretical prediction – that afferent

interoceptive signals (and resulting prediction-errors) are assigned low precision estimates in

those with psychiatric disorders, leading those signals to be under-weighted during interoceptive

inference (although, as discussed above, other interpretations are possible). While an important

first step, a further test of this hypothesis would require replication in a confirmatory sample,

something we plan to investigate in the second cohort of participants in the Tulsa 1000 project

(for some confirmatory work in an independent sample of healthy individuals, see (107)).

Additionally, combining this approach with neuroimaging measures would help to establish the

expected relationship between IP and neural activity within interoceptive cortices.

Computational approaches derived from Active inference models may be especially useful in this

regard, as they allow for simulation of predicted neuronal responses (79).

Some limitations of our modeling approach are also important to keep in mind. First, we were

required to make specific choices about model structure. For example, while priors during

estimation were fixed at plausible values (justification explained within the methods section),

other values might have been chosen. Also, we chose not to include a temperature parameter in

our action model. While this was possible, in simple perceptual tasks like ours that lack reward

feedback, any variability in the precision of action selection would be correlated with sensory

precision. Therefore, we chose a simpler model that treated behavior as a direct readout of

posterior beliefs in perception. To account for potential differences in motor stochasticity, we

then controlled for sensory precision in the tone condition, where, because the sensory signal is

highly precise, between-subject differences in precision would most plausibly reflect between-

subject differences in variability of motor responses. Here it is also worth highlighting that, if

motor effects (such as reaction time) were stable within-subjects but varied between subjects,

this would not influence precision estimates because the temporal relationship between systoles

and taps would still be clear. However, the novelty of our approach entails that it should be

replicated in future studies.

Another limitation is that, while we did compare models with vs. without learning, we did not

compare our model to other existing approaches. The heartbeat tapping task was not well-suited

for the several alternatives we considered. For example, it did not have a sufficient number of

trials to be suited for traditional signal detection approaches, which would be most comparable to

our model-based precision measure (108). Computational models based on reinforcement

learning were also inappropriate as the task did not include planning or learning from reward,

and instead dealt mainly with uncertainty in perception. One alternative approach we might have

taken would have been to model the task with a hierarchical Gaussian filter (109), which has

similar Bayesian foundations and can estimate the relative precisions of sensory input and prior

beliefs (i.e., as opposed to the separate estimates of each in our model), which could be useful to

examine in future work.

Our modelling approach also required making choices about how to include EKG signals as

observations and relate them to behavior. Here this involved discretizing timepoints where

responses were considered co-occurrent (or not) with diastole vs. systole; but other such

discretizations could have been chosen. That said, given the relationships we observed between

parameters and other task measures, our choices appear to have led to estimates that track

meaningful individual differences in task behavior. The task measurement conditions also had

certain limitations. For example, many individuals had low IP values – reflecting the low cardiac

awareness commonly seen at rest in previous studies (6) – which may have limited the variability

necessary to assess relationships with other variables. Finally, there were relatively lower sample

sizes in the eating disorders and anxiety disorders groups. Thus, studies looking at these groups

with larger sample sizes will be necessary to afford stronger confidence in our results for these

groups.

In summary, this study 1) demonstrated the sensitivity of individual difference measures

(parameter estimates) derived from a novel Bayesian computational model, with a focus on

estimating the precision weighting of interoceptive signals across a transdiagnostic sample of

individuals with psychiatric disorders, and 2) tested—and found evidence supporting—the

hypothesis that individuals with psychiatric disorders fail to update the precision-weighting of

afferent interoceptive signals during homeostatic perturbations. While the underlying

neurophysiological mechanisms leading to this difference remain unidentified, these results point

to a potential origin of visceral dysregulation (and perhaps its influence on maladaptive

behavior) across multiple psychiatric conditions. This represents an important step towards a

primary goal of computational psychiatry – computationally phenotyping individuals with

psychiatric disorders with the tools of computational neuroscience in hopes of using this

information to guide the development of precision medicine interventions.

Software Note: All model simulations were implemented using standard routines
(**spm_MDP_VB_X.m**) that are available as Matlab code in the latest version of SPM academic
software: http://www.fil.ion.ucl.ac.uk/spm/. The specific code used for our model can be found
in supplementary materials.

Funding: This work has been supported in part by The William K. Warren Foundation, the
National Institute of Mental Health, K23MH112949 (SSK), and the National Institute of General
Medical Sciences Center Grant, P20GM121312 (MPP, SSK, JF, JLS).

Conflict of Interest: None of the authors have any conflicts of interest to disclose.

Supplementary Files

Supplementary materials. This file includes results of supplementary analyses as well as
supplementary figures, as referred to in the main text.

Supplementary code. This file contains the MATLAB code used for modelling task behavior.

Study data. This file includes all data used in the analyses reported in the manuscript.

References

1.   Khalsa SS, Adolphs R, Cameron OG, Critchley HD, Davenport PW, Feinstein JS, et al.
Interoception and Mental Health: A Roadmap. Biol Psychiatry Cogn Neurosci Neuroimaging.
2018;3(6):501-13.
2.   Furman D, Waugh C, Bhattacharjee K, Thompson R, Gotlib I. Interoceptive awareness, positive
affect, and decision making in major depressive disorder. Journal of affective disorders. 2013;151:780-5.
3.   Terhaar J, Viola FC, Bär K-J, Debener S. Heartbeat evoked potentials mirror altered body
perception in depressed patients. Clinical neurophysiology. 2012;123:1950-7.
4.   Dunn BD, Stefanovitch I, Evans D, Oliver C, Hawkins A, Dalgleish T. Can you feel the beat?
Interoceptive awareness is an interactive function of anxiety- and depression-specific symptom
dimensions. Behav Res Ther. 2010;48(11):1133-8.
5.   Pollatos O, Traut-Mattausch E, Schandry R. Differential effects of anxiety and depression on
interoceptive accuracy. Depression and anxiety. 2009;26:167-73.
6.   Khalsa SS, Lapidus RC. Can Interoception Improve the Pragmatic Search for Biomarkers in
Psychiatry? Front Psychiatry. 2016;7:121.

7.   Stewart JL, Khalsa SS, Kuplicki R, Puhl M, Investigators T, Paulus MP. Interoceptive attention in
opioid and stimulant use disorder. Addict Biol. 2019:e12831.
8.   Khalsa SS, Craske MG, Li W, Vangala S, Strober M, Feusner JD. Altered interoceptive awareness
in anorexia nervosa: effects of meal anticipation, consumption and bodily arousal. International Journal
of Eating Disorders. 2015;48(7):889-97.
9.   Allen M, Levy A, Parr T, Friston KJ. In the Body's Eye: The Computational Anatomy of
Interoceptive Inference. bioRxiv. 2019:603928.
10.   Owens AP, Allen M, Ondobaka S, Friston KJ. Interoceptive inference: From computational
neuroscience to clinic. Neurosci Biobehav Rev. 2018;90:174-83.
11.   Seth A. Interoceptive inference, emotion, and the embodied self. Trends in cognitive sciences.
2013;17:565-73.
12.   Seth AK, Critchley HD. Extending predictive processing to the body: Emotion as interoceptive
inference. Behavioral and brain sciences. 2013;36:47-8.
13.   Stephan K, Manjaly Z, Mathys C, Weber L, Paliwal S, Gard T, et al. Allostatic Self-efficacy: A
Metacognitive Theory of Dyshomeostasis-Induced Fatigue and Depression. Frontiers in human
neuroscience. 2016;10:550.
14.   Petzschner F, Weber L, Gard T, Stephan K. Computational Psychosomatics and Computational
Psychiatry: Toward a Joint Framework for Differential Diagnosis. Biological Psychiatry. 2017;82:421-30.
15.   Barrett L, Simmons W. Interoceptive predictions in the brain. Nature reviews Neuroscience.
2015;16:419-29.
16.   Bastos A, Usrey W, Adams R, Mangun G, Fries P, Friston K. Canonical microcircuits for predictive
coding. Neuron. 2012;76:695-711.
17.   Friston K. A theory of cortical responses. Philosophical transactions of the Royal Society of
London Series B, Biological sciences. 2005;360:815-36.
18.   Chen C, Takahashi T, Nakagawa S, Inoue T, Kusumi I. Reinforcement learning in depression: A
review of computational research. Neurosci Biobehav Rev. 2015;55:247-67.
19.   Clark J, Watson S, Friston K. What is mood? A computational perspective. Psychological
Medicine. 2018:1-8.
20.   Friston K, Stephan K, Montague R, Dolan R. Computational psychiatry: the brain as a phantastic
organ. The lancet Psychiatry. 2014;1:148-58.
21.   Huys Q, Maia T, Frank M. Computational psychiatry as a bridge from neuroscience to clinical
applications. Nature Neuroscience. 2016;19:404-13.
22.   Montague P, Dolan R, Friston K, Dayan P. Computational psychiatry. Trends in Cognitive
Sciences. 2012;16:72-80.
23.   Moutoussis M, Shahar N, Hauser T, Dolan R. Computation in Psychotherapy, or How
Computational Psychiatry Can Aid Learning-Based Psychological Therapies. Computational Psychiatry.
2017:1-21.
24.   Parr T, Friston K. The Anatomy of Inference: Generative Models and Brain Structure. Frontiers in
Computational Neuroscience. 2018;12:90.
25.   Schwartenbeck P, Friston K. Computational Phenotyping in Psychiatry: A Worked Example.
eNeuro. 2016;3:ENEURO.0049-16.2016.
26.   Sharp PB, Eldar E. Computational models of anxiety: nascent efforts and future directions.
Current Directions in Psychological Science. 2019;28(2):170-6.
27.   Smith R, Parr T, Friston KJ. Simulating Emotions: An Active Inference Model of Emotional State
Inference and Emotion Concept Learning. Front Psychol. 2019;10:2844.
28.   Smith R, Lane RD, Parr T, Friston KJ. Neurocomputational mechanisms underlying emotional
awareness: Insights afforded by deep active inference and their potential clinical relevance. Neurosci
Biobehav Rev. 2019;107:473-91.

29.   Adams R, Shipp S, Friston K. Predictions not commands: active inference in the motor system.
Brain Structure and Function. 2013;218:611-43.
30.   Adams R, Perrinet L, Friston K. Smooth Pursuit and Visual Occlusion: Active Inference and
Oculomotor Control in Schizophrenia. PLoS ONE. 2012;7:e47502.
31.   Edwards M, Adams R, Brown H, Pareés I, Friston K. A Bayesian account of 'hysteria'. Brain.
2012;135:3495-512.
32.   Schwartenbeck P, FitzGerald T, Mathys C, Dolan R, Friston K. The Dopaminergic Midbrain
Encodes the Expected Certainty about Desired Outcomes. Cerebral Cortex. 2015;25:3434-45.
33.   Iglesias S, Mathys C, Brodersen K, Kasper L, Piccirelli M, den Ouden H, et al. Hierarchical
Prediction Errors in Midbrain and Basal Forebrain during Sensory Learning. Neuron. 2013;80:519-30.
34.   Diaconescu A, Mathys C, Weber L, Kasper L, Mauer J, Stephan K. Hierarchical prediction errors in
midbrain and septum during social learning. Social Cognitive and Affective Neuroscience.
2017;12:nsw171.
35.   Petzschner FH, Weber LA, Wellstein KV, Paolini G, Do CT, Stephan KE. Focus of attention
modulates the heartbeat evoked potential. Neuroimage. 2019;186:595-606.
36.   Smith R, Thayer JF, Khalsa SS, Lane RD. The hierarchical basis of neurovisceral integration.
Neurosci Biobehav Rev. 2017;75:274-96.
37.   Owens A, Friston K, Low D, Mathias C, Critchley H. Investigating the relationship between
cardiac interoception and autonomic cardiac control using a predictive coding framework. Autonomic
Neuroscience. 2018;210:65-71.
38.   Paulus MP, Feinstein JS, Khalsa SS. An Active Inference Approach to Interoceptive
Psychopathology. Annu Rev Clin Psychol. 2019;15:97-122.
39.   Schandry R. Heart beat perception and emotional experience. Psychophysiology.
1981;18(4):483-8.
40.   Ludwick-Rosenthal R, Neufeld RW. Heart beat interoception: a study of individual differences.
Int J Psychophysiol. 1985;3(1):57-65.
41.   Garfinkel S, Seth A, Barrett A, Suzuki K, Critchley H. Knowing your own heart: Distinguishing
interoceptive accuracy from interoceptive awareness. Biological Psychology. 2014;104:65-74.
42.   Simmons W, Avery J, Barcalow J, Bodurka J, Drevets W, Bellgowan P. Keeping the body in mind:
insula functional organization and functional connectivity integrate interoceptive, exteroceptive, and
emotional awareness. Human brain mapping. 2013;34:2944-58.
43.   Brener J, Kluvitse C. Heartbeat detection: judgments of the simultaneity of external stimuli and
heartbeats. Psychophysiology. 1988;25(5):554-61.
44.   Katkin ES, Morell MA, Goldband S, Bernstein GL, Wise JA. Individual differences in heartbeat
discrimination. Psychophysiology. 1982;19(2):160-6.
45.   Whitehead WE, Drescher VM, Heiman P, Blackwell B. Realtion of heart rate control to heartbeat
perception. Biofeedback Self Regul. 1977;2(4):317-92.
46.   Phillips GC, Jones GE, Rieger EJ, Snell JB. Effects of the presentation of false heart-rate feedback
on the performance of two common heartbeat-detection tasks. Psychophysiology. 1999;36(4):504-10.
47.   Ring C, Brener J, Knapp K, Mailloux J. Effects of heartbeat feedback on beliefs about heart rate
and heartbeat counting: a cautionary tale about interoceptive awareness. Biol Psychol. 2015;104:193-8.
48.   Windmann S, Schonecke OW, Frohlig G, Maldener G. Dissociating beliefs about heart rates and
actual heart rates in patients with cardiac pacemakers. Psychophysiology. 1999;36(3):339-42.
49.   Ring C, Brener J. Heartbeat counting is unrelated to heartbeat detection: A comparison of
methods to quantify interoception. Psychophysiology. 2018;55(9):e13084.
50.   Murphy J, Millgate E, Geary H, Ichijo E, Coll MP, Brewer R, et al. Knowledge of resting heart rate
mediates the relationship between intelligence and the heartbeat counting task. Biol Psychol.
2018;133:1-3.

51.   Khalsa SS, Rudrauf D, Sandesara C, Olshansky B, Tranel D. Bolus isoproterenol infusions provide
a reliable method for assessing interoceptive awareness. Int J Psychophysiol. 2009;72(1):34-45.
52.   Hassanpour MS, Yan L, Wang DJ, Lapidus RC, Arevian AC, Simmons WK, et al. How the heart
speaks to the brain: neural activity during cardiorespiratory interoceptive stimulation. Philos Trans R Soc
Lond B Biol Sci. 2016;371(1708).
53.   Schandry R, Bestler M, Montoya P. On the relation between cardiodynamics and heartbeat
perception. Psychophysiology. 1993;30(5):467-74.
54.   Huang H, Thompson W, Paulus MP. Computational dysfunctions in anxiety: Failure to
differentiate signal from noise. Biological psychiatry. 2017;82(6):440-6.
55.   Barrett L. How emotions are made: The secret life of the brain. 2017.
56.   Barrett LF, Quigley KS, Hamilton P. An active inference theory of allostasis and interoception in
depression. Philos Trans R Soc Lond B Biol Sci. 2016;371(1708).
57.   Critchley H, Garfinkel S. Interoception and emotion. Current Opinion in Psychology. 2017;17:7-
14.
58.   Smith R, Lane RD. The neural basis of one's own conscious and unconscious emotional states.
Neurosci Biobehav Rev. 2015;57:1-29.
59.   Craig AD. How do you feel--now? The anterior insula and human awareness. Nature Reviews
Neuroscience. 2009;10:59-70.
60.   Domschke K, Stevens S, Pfleiderer B, Gerlach AL. Interoceptive sensitivity in anxiety and anxiety
disorders: an overview and integration of neurobiological findings. Clin Psychol Rev. 2010;30(1):1-11.
61.   Yoris A, Esteves S, Couto B, Melloni M, Kichic R, Cetkovich M, et al. The roles of interoceptive
sensitivity and metacognitive interoception in panic. Behav Brain Funct. 2015;11:14.
62.   Bogacz R. A tutorial on the free-energy framework for modelling perception and learning.
Journal of mathematical psychology. 2017;76:198-211.
63.   Desmedt O, Luminet O, Corneille O. The heartbeat counting task largely involves non-
interoceptive processes: Evidence from both the original and an adapted counting task. Biol Psychol.
2018;138:185-8.
64.   Victor TA, Khalsa SS, Simmons WK, Feinstein JS, Savitz J, Aupperle RL, et al. Tulsa 1000: a
naturalistic study protocol for multilevel assessment and outcome prediction in a large psychiatric
sample. BMJ open. 2018;8(1):e016620.
65.   Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. J
Gen Intern Med. 2001;16(9):606-13.
66.   Norman SB, Cissell SH, Means-Christensen AJ, Stein MB. Development and validation of an
Overall Anxiety Severity And Impairment Scale (OASIS). Depress Anxiety. 2006;23(4):245-9.
67.   Staley D, el-Guebaly N. Psychometric properties of the Drug Abuse Screening Test in a
psychiatric patient population. Addict Behav. 1990;15(3):257-64.
68.   Morgan JF, Reid F, Lacey JH. The SCOFF questionnaire: a new screening tool for eating disorders.
West J Med. 2000;172(3):164-5.
69.   Aupperle RL, Paulus MP, Kuplicki R, Touthang J, Victor T, Yeh HW, et al. Web-Based Graphic
Representation of the Life Course of Mental Health: Cross-Sectional Study Across the Spectrum of
Mood, Anxiety, Eating, and Substance Use Disorders. JMIR Ment Health. 2020;7(1):e16919.
70.   Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-
International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured
diagnostic psychiatric interview for DSM-IV and ICD-10. The Journal of clinical psychiatry. 1998;59 Suppl
20:22-33;quiz 4-57.
71.   Smith R, Feinstein J, Kuplicki R, Forthman KL, Stewart JL, Paulus MP, et al. Perceptual
Insensitivity to the Modulation of Interoceptive Signals in Depression, Anxiety, and Substance Use
Disorders. PsyArXiv. 2020.

72. DeVille DC, Kuplicki R, Stewart JL, Tulsa I, Aupperle RL, Bodurka J, et al. Diminished responses to
bodily threat and blunted interoception in suicide attempters. Elife. 2020;9.
73. Canales-Johnson A, Silva C, Huepe D, Rivera-Rei Á, Noreika V, Garcia MdC, et al. Auditory
Feedback Differentially Modulates Behavioral and Neural Markers of Objective and Subjective
Performance When Tapping to Your Heartbeat. Cerebral Cortex. 2015;25(11):4490-503.
74. Allen J, Murray A. Age-related changes in the characteristics of the photoplethysmographic
pulse shape at various body sites. Physiol Meas. 2003;24(2):297-307.
75. Ring C, Brener J. The temporal locations of heartbeat sensations. Psychophysiology.
1992;29(5):535-45.
76. Friston K, Lin M, Frith C, Pezzulo G, Hobson J, Ondobaka S. Active Inference, Curiosity and
Insight. Neural Computation. 2017;29:2633-83.
77. Friston K, Parr T, de Vries B. The graphical brain: Belief propagation and active inference.
Network Neuroscience. 2017;1:381-414.
78. Da Costa L, Parr T, Sajid N, Veselic S, Neacsu V, Friston K. ACTIVE INFERENCE ON DISCRETE
STATE-SPACES – A SYNTHESIS. arXiv. 2020:2001.07203v2 [q-bio.NC]
79. Friston K, FitzGerald T, Rigoli F, Schwartenbeck P, Pezzulo G. Active Inference: A Process Theory.
Neural Computation. 2017;29:1-49.
80. Parr T, Friston K. Working memory, attention, and salience in active inference. Scientific Reports.
2017;7:14678.
81. Daunizeau J, den Ouden HE, Pessiglione M, Kiebel SJ, Stephan KE, Friston KJ. Observing the
observer (I): meta-bayesian models of learning and decision-making. PLoS One. 2010;5(12):e15554.
82. Friston K, Mattout J, Trujillo-Barreto N, Ashburner J, Penny W. Variational free energy and the
Laplace approximation. Neuroimage. 2007;34(1):220-34.
83. Rigoux L, Stephan KE, Friston KJ, Daunizeau J. Bayesian model selection for group studies -
revisited. Neuroimage. 2014;84:971-85.
84. Stephan KE, Penny WD, Daunizeau J, Moran RJ, Friston KJ. Bayesian model selection for group
studies. Neuroimage. 2009;46(4):1004-17.
85. Morey RD, Rouder JN. BayesFactor (Version 0.9.10-2)[Computer software]. 2015.
86. Rouder JN, Morey RD, Speckman PL, Province JM. Default Bayes Factors for ANOVA Designs.
Journal of Mathematical Psychology. 2012;56:356-74.
87. Mehling WE, Price C, Daubenmier JJ, Acree M, Bartmess E, Stewart A. The Multidimensional
Assessment of Interoceptive Awareness (MAIA). PLoS One. 2012;7(11):e48230.
88. Parker J, Taylor G, Bagby R. The 20-Item Toronto Alexithymia Scale: III. Reliability and factorial
validity in a community population. Journal of Psychosomatic Research. 2003;55:269-75.
89. Sandin B, Chorot P, McNally RJ. Anxiety sensitivity index: normative data and its differentiation
from trait anxiety. Behav Res Ther. 2001;39(2):213-9.
90. Smith R, Killgore WDS, Lane RD. The structure of emotional experience and its relation to trait
emotional awareness: A theoretical review. Emotion. 2018;18(5):670-92.
91. LeDoux JE, Brown R. A higher-order theory of emotional consciousness. Proc Natl Acad Sci U S A.
2017;114(10):E2016-E25.
92. Smith R, Lane RD, Parr T, Friston KJ. Neurocomputational mechanisms underlying emotional
awareness: insights afforded by deep active inference and their potential clinical relevance. Neurosci
Biobehav Rev. 2019.
93. Terasawa Y, Moriguchi Y, Tochizawa S, Umeda S. Interoceptive sensitivity predicts sensitivity to
the emotions of others. Cognition & emotion. 2014;28:1435-48.
94. Gu X, Hof P, Friston K, Fan J. Anterior insular cortex and emotional awareness. The Journal of
comparative neurology. 2013;521:3371-88.

95. Zaki J, Davis JI, Ochsner K. Overlapping activity in anterior insula during interoception and
emotional experience. NeuroImage. 2012;62:493-9.
96. Herbert B, Herbert C, Pollatos O. On the Relationship Between Interoceptive Awareness and
Alexithymia: Is Interoceptive Awareness Related to Emotional Awareness? Journal of Personality.
2011;79:1149-75.
97. Smith R, Lane RD, Sanova A, Alkozei A, Smith C, Killgore WDS. Common and Unique Neural
Systems Underlying the Working Memory Maintenance of Emotional vs. Bodily Reactions to Affective
Stimuli: The Moderating Role of Trait Emotional Awareness. Front Hum Neurosci. 2018;12:370.
98. Lane RD, Weihs KL, Herring A, Hishaw A, Smith R. Affective agnosia: Expansion of the alexithymia
construct and a new opportunity to integrate and extend Freud's legacy. Neurosci Biobehav Rev.
2015;55:594-611.
99. Kashdan T, Barrett L, McKnight P. Unpacking Emotion Differentiation: Transforming Unpleasant
Experience by Perceiving Distinctions in Negativity. Current Directions in Psychological Science.
2015;24:10-6.
100. Paulus MP, Stein MB. Interoception in anxiety and depression. Brain Struct Funct. 2010;214(5-
6):451-63.
101. Paulus MP, Stewart JL. Interoception and drug addiction. Neuropharmacology. 2014;76 Pt
B:342-50.
102. Paulus MP, Tapert SF, Schulteis G. The role of interoception and alliesthesia in addiction.
Pharmacol Biochem Behav. 2009;94(1):1-7.
103. Verdejo-Garcia A, Clark L, Dunn BD. The role of interoception in addiction: a critical review.
Neurosci Biobehav Rev. 2012;36(8):1857-69.
104. Ring C, Brener J. Influence of beliefs about heart rate and actual heart rate on heartbeat
counting. Psychophysiology. 1996;33(5):541-6.
105. Friston K, FitzGerald T, Rigoli F, Schwartenbeck P, O Doherty J, Pezzulo G. Active inference and
learning. Neuroscience and biobehavioral reviews. 2016;68:862-79.
106. Kleckner I, Zhang J, Touroutoglou A, Chanes L, Xia C, Simmons W, et al. Evidence for a Large-
Scale Brain System Supporting Allostasis and Interoception in Humans. Nature human behaviour.
2017;1:0069.
107. Smith R, Kuplicki R, Teed A, Upshaw V, Khalsa SS. Confirmatory evidence that healthy individuals
can adaptively adjust prior expectations and interoceptive precision estimates. 1st International
Workshop on Active Inference (IWAI), ECML/PKDD conference 2020 Published on bioRxiv ahead of issue:
Springer; 2020. p. 2020.08.31.275594.
108. Kleckner IR, Wormwood JB, Simmons WK, Barrett LF, Quigley KS. Methodological
recommendations for a heartbeat detection-based measure of interoceptive sensitivity.
Psychophysiology. 2015;52(11):1432-40.
109. Mathys CD, Lomakina EI, Daunizeau J, Iglesias S, Brodersen KH, Friston KJ, et al. Uncertainty in
perception and the Hierarchical Gaussian Filter. Front Hum Neurosci. 2014;8:825.

1177