An active inference model reveals a failure to adapt interoceptive precision estimates across

depression, anxiety, eating, and substance use disorders

Ryan Smith[1], Rayus Kuplicki[1], Justin Feinstein[1,2], Katherine L. Forthman, Jennifer L. Stewart[1,2],

Martin P. Paulus[1,2], Tulsa 1000 investigators[†], and Sahib S. Khalsa[1,2]

[1]Laureate Institute for Brain Research, Tulsa, OK, USA

[2]Oxley College of Health Sciences, The University of Tulsa, Tulsa, OK.

[†]The Tulsa 1000 Investigators include the following contributors: Robin L. Aupperle,

Ph.D.; Jerzy Bodurka, Ph.D.; Jonathan B. Savitz, Ph.D.; and Teresa A. Victor, Ph.D.

Corresponding author:

Ryan Smith

Laureate Institute for Brain Research

6655 S Yale Ave, Tulsa, OK 74136, USA

Email: rsmith@laureateinstitute.org

Abstract

Recent neurocomputational theories have hypothesized that abnormalities in prior expectations and/or the precision-weighting of afferent interoceptive signals (i.e. the degree to which afferent bodily signals contribute to interoceptive perceptual inference), may facilitate the transdiagnostic emergence of psychopathology. Specifically, it has been suggested that, in certain psychiatric disorders, interoceptive processing mechanisms either over-weight prior expectations or under-weight signals from the viscera (or both), leading to a failure to accurately update beliefs about the body (potentially resulting in visceral dysregulation among other maladaptive phenomena). However, this has not been directly tested empirically. To evaluate the potential roles of prior expectations and interoceptive precision in this context, we fitted behavior in a transdiagnostic patient population on an interoceptive awareness (heartbeat tapping) task to a Bayesian computational model based on active inference (i.e., in which interoceptive inference can be cast as prediction-error minimization). Modeling revealed that, during an interoceptive perturbation condition (inspiratory breath-holding during heartbeat tapping), healthy individuals (N=52) assigned greater precision to ascending cardiac signals than individuals with symptoms of anxiety (N=15), depression (N=69), co-morbid depression/anxiety (N=153), substance use disorders (N=131), and eating disorders (N=14) – who failed to increase their precision estimates from resting levels. In contrast, no differences were found in prior expectations. These results provide the first empirical computational modeling evidence of a selective dysfunction in adaptive interoceptive processing in psychiatric conditions, and lay the groundwork for future studies examining how reduced interoceptive precision influences body regulation and interoceptively-guided decision-making.

Author Summary

Interoception is the process by which the nervous system senses the internal state of the body. It

provides the brain with important information to adaptively guide the regulation of both internal

body states and behavior. Interoceptive dysfunction is thought to play a role in multiple

psychiatric disorders. Theoretical models propose that the computational mechanisms of

interoceptive dysfunction are caused by overly precise prior expectations about body states

("hyperprecise priors") or underestimates of the reliability of the information carried by

ascending signals from the body ("low sensory precision"). Our empirical approach tested for

evidence of these mechanisms across several psychiatric disorders, using a computational model

of perception during performance of a heartbeat perception task. We found evidence of low

sensory precision within individuals with anxiety, depression, eating disorders, and/or substance

use disorders, relative to healthy individuals. This difference occurred only during a breath-

holding condition designed to enhance heartbeat signals. We did not find evidence for

hyperprecise priors in the patient groups. The data from this study support the argument for

computational mechanisms of interoceptive dysfunction across several psychiatric disorders, and

suggests that these conditions may be characterized by an inability to adjust sensory precision

when signals from the body change.

Keywords: Interoception; Active Inference; Depression; Anxiety; Substance Use; Eating
Disorders; Precision; Prior expectations; Bayesian Perception; Computational Modeling

Introduction

Interoception plays an important role in a number of psychiatric disorders. Interoceptive dysfunction has been observed in depression, anxiety, eating, and substance use disorders, among others (reviewed in (1)). For example, depressed patients exhibit reduced accuracy when asked to count their own heartbeats (2-4), and counting accuracy is negatively correlated with depressive symptoms (5). Several studies have also shown heightened interoceptive sensations in panic disorder under high arousal states (reviewed in (6)). Other studies have reported evidence of blunted neural responses during interoceptive processing in substance users (7), and that patients with eating disorders show stronger expectation effects on interoception during modulations of arousal (8). However, a well-established, principled approach for identifying a generative explanatory model of interoceptive dysfunctions in psychiatric disorders has not yet emerged.

In recent years, a growing body of theoretical work within neuroscience and psychiatry has begun to highlight plausible neurocomputational accounts of interoceptive processing (9-15), as well as accounts of exteroceptive (16, 17), cognitive (18-26), emotional (27, 28), and motor control (29-31) functions that plausibly interact with interoception. In some cases, empirical studies have also found support for computational models and computational abnormalities in relation to specific psychiatric symptoms/disorders (26, 32-34). With respect to sensory processing, which has focused to a large degree on neural models of approximate probabilistic (Bayesian) inference, empirical results in this area have come almost exclusively from the exteroceptive domain. And while recent work has begun to test qualitative predictions of Bayesian models of interoception (35), no study to our knowledge has yet explicitly fit

computational models to brain or behavioral responses during experimental modulations of

interoception in individuals with psychiatric disorders.

Drawing on this theoretical literature, we and others (9, 13-15, 36-38) have previously proposed

that symptoms of multiple psychiatric conditions could be explained by faulty interoceptive

computational processes (and associated visceromotor control processes), due to an inappropriate

weighting of prior expectations and sensory evidence. Although these models of computational

dysfunction in interoception could have broad explanatory power, the putative mechanisms they

propose have yet to be empirically tested using formal computational models. In this paper, we

use a formal Bayesian computational model of perception to examine whether there is evidence

for the transdiagnostic dysfunction of interoceptive processing suggested in these previous

proposals within the cardiovascular system.

Historically, the most common empirical paradigms for studying interoception have focused on

the perception of cardiac signals. Common heartbeat perception tasks include those involving

heartbeat counting (39), heartbeat tapping (40), and other means of assessing cardiac

interoception (41-45). While each approach has certain limitations (46-50), one consistent

finding is that cardiac interoceptive awareness is quite poor in the majority of participants tested

during resting conditions – where only roughly 35% of individuals appear to accurately perceive

their own heartbeats (6). In contrast, when visceral states are perturbed, cardiac perception

becomes more accurate. This is particularly true for interoceptive accuracy under conditions of

heightened cardiorespiratory arousal (51-53).

Active inference models, and Bayesian predictive processing models more generally, have been

the main computational framework within which interoceptive dysfunction has been discussed

(13, 38, 54-56). Several authors have suggested that one important transdiagnostic factor within

mental disorders may be an inability of the brain to update its model of the body in the face of

interoceptive prediction errors (i.e., mismatches between expected and received afferent

interoceptive signals from the body). In predictive coding and active inference models, this kind

of aberrant belief updating is thought to come about through a dysfunctional "precision

weighting" mechanism, which governs the relative influence of prior expectations and afferent

bodily signals in determining perception (and informing visceral regulation). Simply put, it is

suggested that, across multiple mental health conditions, the brain may treat afferent bodily

signals (and associated prediction errors) as though they are not reliable indicators of bodily

states during interoceptive inference – leading perception to be insufficiently constrained by true

visceral states and primarily determined by (in many cases maladaptive) prior expectations.

Misestimating the state of the body could in turn promote a number of transdiagnostic

symptoms. For example, interoceptive feelings are intimately tied to emotions (57-59); poor

body perception (e.g., high uncertainty about internal physiological conditions and a resulting

inability to efficiently regulate them) could thus maintain unpleasant emotional states. Chronic

underestimates of available metabolic resources may contribute to apathy and anhedonia (56),

and overestimates of the evidence uncomfortable bodily sensations provide for physical threat

(e.g., a heart attack) may contribute to anxiety and panic (60, 61).

It's important to emphasize, however, that computational models often include several additional

parameters. Aside from the precision-weighting of sensory signals, individuals can also have

distinct differences in (for example) prior expectations about what they will perceive (assumed in

any Bayesian model of perception; e.g., predictive coding (62)), and differences in how quickly

they update those prior expectations over repeated observations (i.e., "learning rate"). Formal

computational models are often necessary to distinguish which parameters show differences

between individuals and best explain differences in perception. Thus, there are multiple

computational mechanisms that could account for individual differences in interoception in

clinical populations. One major goal in computational psychiatry is to "computationally

phenotype" patients by identifying which sets of a parameter values best account for their neural

and behavioral responses (including self-reported perceptual experience) and use this

information to guide treatment (20, 22, 25).

In the present study, we apply a novel computational phenotyping approach, using a Bayesian

(active inference) model of perception, to identify the computational parameters that best explain

behavior on a cardiac perception (heartbeat tapping) task performed by a transdiagnostic clinical

sample of individuals with psychiatric disorders as well as a healthy comparison (HC) sample.

Given the notably poor cardiac perception of most human beings at rest (6), we chose to assess

individual differences in cardiac interoceptive precision in the context of normal physiological

baseline states as well as during a non-invasive interoceptive perturbation (a breath-hold)

condition that was expected to improve cardiac perception above floor values in a greater

number of individuals (i.e., we expected that cardiac perception would be generally poor during

resting conditions, and that the breath-hold condition would result in improved performance on

average). Our primary aims were to 1) demonstrate the sensitivity of our novel computational

approach in measuring the precision weighting of interoceptive signals and prior expectations

across a transdiagnostic sample of individuals with depression, anxiety, substance use disorders,

and/or eating disorders, 2) test the hypothesis (as previously proposed; e.g., (10, 38, 56)) that

these patient groups would show lower interoceptive precision weightings than HCs, more

precise prior expectations than HCs, or both, and 3) explore whether prior expectations and/or

interoceptive precision is abnormal in general or selectively within resting or interoceptive

perturbation conditions

Methods

Participants

Data were collected from 500 participants (153 male) as a part of the Tulsa 1000 (T1000) project

(63), a naturalistic longitudinal study that recruited subjects based on the dimensional NIMH

Research Domain Criteria framework. Individuals aged 18-55 years were screened on the basis

of dimensional psychopathology scores. Inclusion was based on the following measures: Patient

Health Questionnaire (PHQ-9; (64)) $\geq$ 10, Overall Anxiety Severity and Impairment Scale

(OASIS; (65)) $\geq$ 8, Drug Abuse Screening Test (DAST-10; (66)) score > 2, and/or Eating

Disorder Screen (SCOFF; (67)) score $\geq$ 2 (for screening measure scores, see **Table 1**). HCs who

did not show elevated symptoms or psychiatric diagnoses were also included. Participants were

excluded if they (i) tested positive for drugs of abuse, (ii) met criteria for psychotic, bipolar, or

obsessive-compulsive disorders, or reported (iii) history of moderate-to-severe traumatic brain

injury, neurological disorders, or severe or unstable medical conditions, (iv) active suicidal intent

or plan, or (v) change in psychotropic medication status within 6 weeks. Full inclusion/exclusion

criteria are described in (63). The study was approved by the Western Institutional Review

Board. All participants provided written informed consent prior to completion of the study

protocol, in accordance with the Declaration of Helsinki, and were compensated for

participation. ClinicalTrials.gov identifier: #NCT02450240.

Similar to previous studies of the T1000 cohort (68), participants were grouped based on DSM-

IV or DSM-5 diagnosis using the Mini International Neuropsychiatric Inventory 6 or 7 (MINI;

(69)), and included those with major depressive disorder only (DEP), an anxiety disorder only

(ANX; social anxiety, generalized anxiety, panic, or posttraumatic stress disorder), comorbid

MDD and anxiety disorder (DEP+ANX), substance use disorders (SUDs; recreational drugs

excluding alcohol and nicotine; with or without comorbid anxiety and mood disorders), eating

disorders (EDs; with or without comorbid anxiety and mood disorders), and HCs with no mental

health diagnoses. In this study we examined all groups from the first 500 participants of the

T1000 project (recruited from January 5, 2015 to February 22, 2017).

Table 1. Mean (and standard deviation) for clinical and demographic variables

Individual difference variable*	Healthy comparisons (N = 52)	Anxiety (N = 15)	Depression (N = 69)	Depression + Anxiety (N = 153)	Eating Disorder (N = 14)	Substance use disorder (N = 131)	p
Age	32.04 (11.08) `	36.42 (10.01)	37.12 (11.83)	35.28 (11.23)	27.40 (9.83)	34.03 (8.88)	0.012
Sex	50% male	33% male	30% male	26% male	14% male	45% male	0.001
PHQ-9	0.83 (1.29)	7.47 (5.80)	13.48 (4.73)	13.10 (5.21)	12.93 (7.98)	6.68 (5.91)	<0.001
OASIS	1.37 (1.89)	10.60 (2.16)	7.55 (3.34)	10.51 (3.12)	9.50 (4.93)	5.94 (4.77)	<0.001
DAST-10	0.10 (0.30)	0.33 (0.49)	0.72 (1.66)	0.58 (1.13)	1.23 (2.62)	7.55 (2.05)	<0.001
PTT	0.20 (0.02)	0.20 (0.01)	0.19 (0.01)	0.20 (0.02)	0.20 (0.02)	0.20 (0.02)	0.261
BMI	27.59 (5.54)	27.05 (6.02)	28.51 (5.47)	28.77 (5.49)	22.25 (4.43)	28.23 (4.56)	0.001

*PHQ-9 = Patient Health Questionnaire 9; OASIS = Overall Anxiety Sensitivity and Impairment Scale;
DAST-10 = Drug Abuse Screening Test; PTT = median pulse transit time; BMI = Body Mass Index

Heartbeat perception task

As part of the T1000 project, participants completed a large number of assessments, self-report

measures, and behavioral tasks (detailed in (63)). Here we focus on data from a cardiac

perception task on which we have previously reported (i.e., on a subset of the participants

reported here, with analyses unrelated to computational modeling (70, 71)), wherein participants

were asked to behaviorally indicate the times at which they felt their heartbeat. The utilization of

the heartbeat tapping measure as an index of perception was based on a previously developed

heartbeat tapping task (40); for a more recent example, see (72)). The task was repeated under

multiple conditions designed to assess the influence of cognitive strategy and physiological

perturbation on performance. In the initial task condition, participants were simply instructed to

close their eyes and press down on a key when they felt their heartbeat, to try to mirror their

heartbeat as closely as possible, and even if they weren't sure they should take their best guess

(the "guessing" condition). Participants completed this (and each other) task condition over a

period of 60 seconds. In the second task condition, all instructions were identical except that they

were told to only press the key when they actually feel their heartbeat, and if they do not feel

their heartbeat then they should not press the key (the "no-guessing" condition). In other words,

unlike the first time they completed the task, they were specifically instructed not to guess if they

didn't feel anything. Finally, in the perturbation condition, participants were again instructed not

to guess but were also asked to first empty their lungs of all air and then inhale as deeply as

possible and hold it for as long as they could tolerate (up to the length of the one-minute trial)

while reporting their perceived heartbeat sensations. This third condition (the "breath-hold"

condition) was used in an attempt to putatively increase the strength of the afferent cardiac signal

by increasing physiological arousal. We expected that cardiac perception would be poor in the

guessing condition, that tapping would be more conservative in the no-guessing condition, and

that the breath-hold condition would result in improved performance on average. As a control

condition, we also included an identical task where participants were instructed to tap every time

they heard a 1000Hz auditory tone presented for 100ms (78 tones, randomly jittered by +/- 10%

and presented in a pattern following a sine curve with a frequency of 13 cycles/minute,

mimicking the range of respiratory sinus arrhythmia during a normal breathing range of 13

breaths per minute). This was completed between the first (guessing) and second (no-guessing)

heartbeat tapping conditions.

Directly after completing each task condition, individuals were asked the following using a

visual analogue scale:

"How accurate was your performance?"

"How difficult was the previous task?"

"How intensely did you feel your heartbeat?"

Each scale had anchors of "not at all" and "extremely" on the two ends. Numerical scores could

range from 0 to 100.

Computational model

To model behavior on the heartbeat tapping task, we first divided each task time series into

intervals corresponding to the periods of time directly before and after each heartbeat. Potentially

perceivable heartbeats were based on the timing of the peak of the electrocardiogram (EKG) R-

wave (signaling electrical depolarization of the atrioventricular neurons of the heart) + 200

milliseconds (ms). This 200 ms interval was considered a reasonable estimate of participants'

pulse transit time (PTT) according previous estimates for the ear PTT (73). We also measured

the average PTT of each participant, defined as the distance between the peak of the EKG R-

wave and the onset of the peak of the PPG waveform (signaling mechanical transmission of the

systolic pressure wave to the earlobe). The length of each heartbeat interval (i.e., the "before-beat

interval" and "after-beat interval") depended on the heart rate. For example, if two heartbeats

were 1 second apart, the "after-beat interval" would include the first 500 ms after the initial beat

and the "before-beat interval" would correspond to the 2nd 500 ms. The after-beat intervals were

considered the time periods in which the systole (heart muscle contraction) signal was present

and in which a tap should be chosen if it was felt. The before-beat intervals were treated as the

time periods where the diastole (heart muscle relaxation) signal was present and in which tapping

should not occur (i.e., assuming taps are chosen in response to detecting a systole; e.g., as

supported by (74)). This allowed us to formulate each interval as a "trial" in which either a tap or

no tap could be chosen and whether a systole or diastole signal was present (see **Figure 1**).

Figure 1. Bayesian approach used to model interoceptive awareness on the heartbeat tapping task. The generative model is here depicted graphically, such that arrows indicate dependencies between variables. Associated vectors/matrices are also shown. At each time point (t), observations (o) depend on hidden states (s), where this relationship is specified by the **A** matrix, and those states depend on previous states (as specified by the **B** matrix, or the initial states specified by the **D** vector). This model represents a simplified version of a commonly used active inference formulation of partially observable Markov decision processes, but which does not explicitly model action selection (for more details regarding the structure and mathematics describing these models, see (75-77). In our model, the observations were systole/diastole, and the hidden states included beliefs about the presence or absence of a heartbeat. For simplicity, the probability of choosing to tap is here assumed to correspond to the posterior distribution over states ($\bar{s}$) – that is, the relative confidence in the presence vs. absence of a heartbeat: $P(HB)$ and $P(nHB)$, respectively. The model parameters we estimated corresponded to: 1) interoceptive precision (IP) – the precision of the mapping from systole/diastole to beliefs about heartbeat/no heartbeat in the **A** matrix, which can be associated with the weight assigned to sensory prediction errors; and 2) prior expectations for the presence of a heartbeat (pHB). Because minimal precision corresponds to an IP value of .5, and both higher and lower values indicate that taps will more reliably track systoles (albeit in an anticipatory or reactive manner), our ultimate measure of precision subtracted 0.5 from raw IP values and then took their absolute value. The raw IP values were then used to assess for group differences in the tendency to tap before vs. after each systole. We also compared this model to an analogous model that included learning (see main text). On each trial, beliefs about the probability of a heartbeat (corresponding to the probability of choosing to tap) relied on Bayesian inference as implemented in the "heartbeat perception" equations shown at the bottom of the figure. Note that, by convention in active inference, the dot product (·) applied to matrices here indicates transposed matrix multiplication. Observed diastoles and systoles were taken from ECG traces, after dividing the time series into periods before and after each systole (exemplified in the right portion of the figure; see text for details).

To model behavior, we used a Bayesian model of perception (see **Figure 1**) derived from a

Markov decision process (MDP) formulation of active inference that has been used in previous

work; for more details about the structure and mathematics of this class of models, see (76, 78,

79). Unlike the full MDP model, however, which includes both a perception model and an action

model, here we only explicitly included a generative model of perception. This model specified

the random variables (see **Table 2**), and their dynamics over time, associated with each time

period ("trial") in the task in which a systole either did or did not occur. Observations (*o*) in the

model formally included systole, diastole, and a "start" observation. Perceptual states (*s*)

included either feeling one's heartbeat or not, as well as a "start" state. Here, a trial formally

included two timesteps: 1) a "start" time point, followed by 2) the possibility of either a systole

or diastole.

A vector **D** encoding prior expectations over initial states, $P(s_{t=1})$, specified that the participant

always started the trial in the start state with perfect confidence. The relationships between states

and observations at a time (*t*) within a trial are described by a set of matrices. The **A** matrix

encodes how perceptual states are informed by observations, $P(o_t|s_t)$ – here corresponding to

the probability of receiving an afferent cardiac signal reflecting systole/diastole if a heartbeat did

or did not occur. The precision of this matrix was controlled by an "interoceptive precision" (IP)

parameter. A value of 0.5 would indicate minimal precision (i.e., leading the individual to simply

"guess" whether he/she experienced a heartbeat), whereas a value approaching 0 or 1 indicates

high precision (i.e., leading the individual to more consistently respond before, or consistently

respond after, a heartbeat has occurred). The **B** matrix encodes the probability that one state will

transition into another, $P(s_{t+1}|s_t)$ – here corresponding to the probability of transitioning from

the "start" state to a "heartbeat state" vs. a "no heartbeat" state. This probability was controlled

by a parameter pHB, where values above .5 indicate prior expectations favoring feeling a

heartbeat (e.g., expecting a faster heart rate), and values less than .5 indicate stronger

expectations not to feel a heartbeat (e.g., expecting slower heart rate). Both IP and pHB were

estimated for each participant based on their tapping behavior, as described below.

Table 2. *Description of active inference model elements*

Model variable	General Definition	Model-specific specification
o_t	Observable outcomes at time t	Outcome modalities: 1. Diastole vs. systole
s_t	Hidden states at time t	Hidden state factors: 1. Feel heartbeat vs. no heartbeat
A matrix $p(o_t \mid s_t)$	A matrix encoding beliefs about the relationship between hidden states and observable outcomes (i.e., the probability that specific outcomes will be observed given specific hidden states).	Encodes beliefs about the relationship between felt heartbeats and diastole vs. systole. The precision of the relationship between heartbeats and diastole/systole was controlled by a parameter IP, which specified how much evidence systole provides for a heartbeat and how much evidence diastole provides for the absence

		of a heartbeat.
B matrix $p(s_{t+1} \mid s_t)$	A matrix encoding beliefs about how hidden states will evolve over time (transition probabilities).	Encodes the prior expectation that either a heartbeat or no heartbeat would occur on each trial, as controlled by a parameter pHB.
D vector $p(s_1)$	A matrix encoding beliefs about (a probability distribution over) initial hidden states.	Ensures the individual always begins in an initial starting state.

In this task, it is unclear whether or not learning over time (e.g., from paying attention to one's

heartbeat) contributes to task performance. To assess this using Bayesian model comparison, we

compared evidence for the "perception only" model described above with evidence for a model

that included learning to update prior expectations over time. Learning within this model

involves updating beliefs about the probability of feeling a heartbeat vs. no heartbeat at each time

point, based on how frequently one believes they have felt their heartbeat in the past. Essentially,

every time a heartbeat is felt, prior expectations favoring feeling a heartbeat go up, and every

time no heartbeat is felt this (relative) expectation goes back down. Formally, this corresponds to

updating the concentration parameters of Dirichlet (Dir) priors associated with the **B** matrix (*b*)

that specify beliefs about state transitions. At $t = 0$:

312
$$P(\boldsymbol{B}) = Dir(b)$$

313
$$b(heartbeats) = P(s_{t+1}|s_t) = \begin{bmatrix} 0 & 0 & 0 \\ 1 - pHB & 1 & 0 \\ pHB & 0 & 1 \end{bmatrix} \times b_0$$

314
$$b_t = b_{t-1} + \sum_t s_t \otimes s_{t-1}$$

Here $\otimes$ indicates the cross-product, and b_0 is a scalar on the prior value for concentration

parameters, where its value prior to learning encodes (inverse) sensitivity to information, such

that higher values will reduce the rate at which prior expectations are updated over time with

new observations. In the learning model, b_0 was also estimated for each individual to capture the

possibility of different learning rates for updating prior expectations (this could also be thought

of as differences in a kind of interoceptive "belief rigidity").

Thus, the final parameters estimated for each participant included the IP, pHB, and b_0

parameters. Our approach to parameter estimation used Bayesian inference at two levels (80).

First, each participant's responses were modeled using the Bayesian model of perception

described above. We then used a commonly used Bayesian optimization algorithm (called

Variational Bayes) to estimate each participant's parameter values that maximized the likelihood

of their responses (under the assumption that a higher/lower probability assigned to feeling a

heartbeat corresponded to a higher/lower probability of choosing to tap), as described in (25).

We optimized these parameters for each model using this likelihood and variational Laplace

(81), implemented within the spm_nlsi_Newton.m parameter estimation routine available within

the freely available SPM12 software package (Wellcome Trust Centre for Neuroimaging,

London, UK, http://www.fil.ion.ucl.ac.uk/spm). This estimation approach has the advantage of

preventing overfitting, due to the greater cost it assigns to moving parameters farther from their

prior values. Estimating parameters required setting prior means and prior variances for each

parameter. The prior variance was set to a high precision value of 1/2 for each parameter (i.e.,

deterring overfitting), and the prior means were set as follows: IP = .5, pHB = .5, and b_0 = 1. Our

decision for selecting these priors was motivated in part by initial simulations confirming that

parameter values were recoverable under these prior values. The IP and pHB prior values were

further chosen to minimize estimate bias, as pHB = .5 assumes flat prior expectations, and IP =

.5 does not bias estimates in favor of values assuming anticipatory vs. reactive strategies. The b_0

prior of 1 is equivalent to this parameter having no effect on the model. After fitting parameters

for each model, we then performed Bayesian model comparison (based on (82, 83)) to determine

the best model. We then used classical inference to test for the effects of group differences in

parameter estimates for the best model, using a standard summary statistic approach.

Before using these parameters in further analyses, however, the "raw" IP parameter values

(IP_{raw}) were transformed to correctly capture 2 distinct constructs of interest. First, because

IP_{raw} values both above and below .5 indicate higher precision (i.e., values below approaching 0

indicate reliable anticipatory tapping, whereas values approaching 1 indicate reliable tapping

after a systole), our ultimate measure of precision was recalculated by centering IP_{raw} on 0 and

taking its absolute value as follows:

350
$$IP = |IP_{raw} - 0.5|$$

351   This means that IP has a minimum value of 0 and a maximum value of 0.5. The IP_{raw} values

352   were then instead used to assess individual differences in the tendency to tap in an anticipatory or

353   reactive (*AvR*) fashion:

354
$$AvR = IP_{raw}$$

355 Higher AvR values (> 0.5) thus indicated a stronger tendency to reactively tap in response to a

356 heartbeat as opposed to tapping in an anticipatory fashion (< 0.5).

Physiological measurements

Electrocardiography was used to assess the objective timing of participants' heartbeats

throughout the task. A BIOPAC MP150 was used to collect a three lead EKG signal and the

pulse oximeter signal, using a pulse plethysmography (PPG) device attached to the ear lobe.

Response times were collected using a task implemented in PsychoPy, with data collection

synchronized via a parallel port interface.

EKG and response data were scored using in-house developed MATLAB code. As described

above in relation to modelling, each participant's pulse transit time (PTT) was estimated as the

median delay between R wave and the corresponding inflection in the PPG signal.

Supplementary materials display an example participant's EKG and PPG trace to illustrate

how this was calculated (also supporting the 200ms delay assumption used in modelling and in

individuals without usable PPG).

Quality Control and Final Sample Sizes

Prior to performing our analyses, several participants were removed due to quality control

checks: 19 individuals were removed due to "cheating" (i.e., video revealed they were taking

their pulse while performing the task); 3 individuals didn't complete the task; and 13 individuals

had poor EKG across all trials that didn't allow reliable identification of heartbeat timing. An

additional 31 individuals were removed due to being outliers when performing the tone task

(using Iterative Grubb's with p < 0.01) – assumed to reflect inappropriate engagement during the

task (e.g., being inattentive, tapping rapidly without listening to the tones, etc.). This resulted in

434 participants, including 52 HCs, 15 ANX, 69 DEP, 153 co-morbid ANX/DEP, 131 SUDs,

and 14 eating disorders. In the three heartbeat tapping conditions, a few other participants were

removed due to poor EKG, leading to the following participant numbers in each condition: 433

(guessing), 431 (no-guessing), and 427 (breath-hold). **Table 3** reports the final number of

participants included for model-based analyses in each condition as well as descriptive statistics

for all task-related measures.

Statistical analysis

Correlational analyses were first conducted to examine the relationships between parameters

across conditions. For purposes of parameter validation, we ran further correlational analyses to

examine the relationships between each parameter and task-specific measures, including the self-

report ratings of difficulty, confidence, and heartbeat intensity collected after each trial.

Analyses of variance (ANOVAs) and covariance (ANCOVAs) were conducted to identify

possible group differences (i.e., between HCs and the five patient groups) in each parameter and

in how they differed across trials (i.e., guessing, no-guessing, breath-hold), while accounting for

individual differences in age, sex, BMI, median PTT, number of heartbeats (and its interaction

with condition), IP for the tone condition (to rule out potential effects of reaction time), and

medication status (i.e., one analysis per parameter). Because the first half of the T1000 sample

was intentionally designed as an exploratory sample, we report relationships at $p < .05$ as a

potential basis for a priori hypotheses verification in the second (confirmatory) half of the T1000

dataset in subsequent work. Although our analyses are exploratory, we note that a Bonferroni

corrected threshold for multiple comparisons with three parameters is $p < .017$ (α=0.05).

Results

Participant characteristics

Complete information on sample size, demographics, and symptom screening measures is

provided in **Table 1**. Separate ANOVAs showed significant differences between groups in age

($F(5,428) = 2.98$, $p = .01$) and body mass index (BMI; $F(5,413) = 4.04$, $p = .001$). A chi-squared

analysis also showed significant differences in the proportion of males to females between

groups (chi-squared $= 19.43$, df $= 5$, $p = .002$). Therefore, in our analyses of model parameters,

we also confirm our results after controlling for these other factors.

Interoceptive Perturbation Validation

As expected, across all participants self-reported heartbeat intensity, confidence in task

performance, and task difficulty differed significantly between the three heartbeat tapping

conditions in separate ANOVAs (intensity: $F(2,1288) = 42.67$, $p < .001$; confidence: $F(2,1288)$

$= 24.94$, $p < .001$; difficulty: $F(2,1288) = 18.26$, $p < .001$), reflecting a greater perceived

intensity of heartbeat sensations and greater confidence during the breath-hold perturbation than

in the other 2 conditions, as well as lower difficulty in the breath-hold condition than the no-

guessing condition (and greater difficulty in the no-guessing than guessing condition; Tukey

post-hoc comparisons: $p < .001$ for all). Analysis of heart rate revealed a marginal difference in

the number of heartbeats between conditions ($F(2,1381) = 2.49$, $p = .08$), reflecting a faster heart

rate in the breath-hold than no-guessing conditions (Tukey post-hoc comparison: $p = .07$).

Table 3. Summary statistics for all task-related variables

Panel A.

Guessing Condition	Healthy comparisons (N = 52)	Anxiety (N = 15)	Depression (N = 69)	Depression + Anxiety (N = 153)	Eating Disorder (N = 14)	Substance use disorder (N = 130)	p-value*
Number of taps	67.92 (29.58)	58.40 (28.40)	67.65 (31.97)	62.02 (32.58)	65.86 (27.86)	65.50 (30.82)	0.701
Number of heartbeats	66.67 (9.48)	67.13 (9.55)	70.23 (10.90)	70.34 (10.86)	67.21 (8.71)	71.23 (10.47)	0.1
Self-reported difficulty	56.87 (22.70)	53.20 (26.69)	57.55 (28.12)	53.18 (25.84)	71.57 (17.78)	48.24 (24.65)	0.009
Self-reported confidence	32.67 (19.27)	41.27 (21.54)	39.52 (22.67)	39.29 (21.20)	25.14 (21.36)	42.20 (22.39)	0.022
Self-reported intensity	20.71 (16.72)	26.73 (21.51)	26.26 (21.52)	28.27 (23.25)	15.36 (18.64)	35.94 (24.65)	<0.001
Counting accuracy	0.69 (0.33)	0.67 (0.30)	0.65 (0.29)	0.63 (0.30)	0.68 (0.30)	0.67 (0.29)	0.755

419   *p-values correspond to the results of ANOVAs comparing the groups (i.e., not including all task
420   conditions within the analysis as reported in the main text).

422   *Panel B.*

No-Guessing Condition	Healthy comparisons (N = 52)	Anxiety (N = 15)	Depression (N = 68)	Depression + Anxiety (N = 153)	Eating Disorder (N = 14)	Substance use disorder (N = 129)	p-value
Number of taps	18.73 (22.69)	22.13 (24.04)	23.43 (24.22)	24.10 (24.26)	18.29 (21.74)	31.47 (25.46)	0.015
Number of heartbeats	65.88 (10.22)	66.73 (11.07)	69.97 (10.45)	69.78 (11.10)	66.14 (8.43)	70.69 (10.14)	0.066
Self-reported difficulty	67.83 (27.94)	59.73 (35.14)	68.00 (32.72)	62.88 (31.35)	69.86 (26.68)	55.14 (30.75)	0.039
Self-reported confidence	39.94 (30.32)	36.80 (35.54)	41.15 (31.36)	41.41 (29.49)	31.93 (32.16)	42.29 (28.27)	0.858
Self-reported intensity	18.15 (19.63)	16.80 (22.11)	25.82 (26.38)	25.31 (25.67)	13.50 (13.51)	33.64 (27.53)	0.001
Counting accuracy	0.28 (0.29)	0.31 (0.33)	0.32 (0.31)	0.33 (0.31)	0.28 (0.32)	0.43 (0.32)	0.028

424   *Panel C.*

Breath Hold Condition	Healthy comparisons (N = 52)	Anxiety (N = 15)	Depression (N = 68)	Depression + Anxiety (N = 151)	Eating Disorder (N = 14)	Substance use disorder (N = 127)	p-value
Number of taps	27.12 (21.47)	30.60 (42.93)	24.75 (21.07)	26.46 (23.58)	17.71 (16.80)	34.17 (25.05)	0.027

Number of heartbeats	69.40 (9.80)	66.87 (12.01)	71.26 (10.48)	70.59 (11.20)	66.29 (6.29)	72.96 (10.75)	0.061
Self-reported difficulty	46.44 (27.08)	54.20 (23.24)	56.53 (25.74)	53.50 (28.32)	55.64 (25.67)	45.86 (29.08)	0.07
Self-reported confidence	47.79 (26.20)	41.27 (26.06)	50.78 (26.49)	52.42 (27.42)	36.50 (33.37)	53.04 (25.55)	0.158
Self-reported intensity	38.19 (28.40)	26.40 (23.61)	41.85 (29.60)	41.09 (29.99)	32.71 (30.63)	46.81 (28.88)	0.074
Counting accuracy	0.39 (0.29)	0.28 (0.26)	0.34 (0.27)	0.35 (0.29)	0.28 (0.29)	0.44 (0.30)	0.03

426 *Panel D.*

Tone Condition	**Healthy comparisons (N = 52)**	**Anxiety (N = 15)**	**Depression (N = 69)**	**Depression + Anxiety (N = 153)**	**Eating Disorder (N = 14)**	**Substance use disorder (N = 131)**	**p-value**
Number of taps	77.90 (1.00)	77.40 (1.12)	77.71 (1.03)	77.82 (1.17)	77.57 (0.94)	78.06 (0.97)	0.072
Number of heartbeats	65.63 (11.10)	67.27 (9.81)	70.74 (9.94)	71.37 (11.14)	66.86 (8.16)	71.30 (10.62)	0.01
Self-reported difficulty	18.04 (15.84)	29.20 (21.12)	22.14 (19.52)	24.15 (22.00)	35.21 (22.33)	25.77 (22.40)	0.065
Self-reported confidence	77.56 (14.26)	73.93 (17.84)	78.84 (16.60)	74.22 (19.59)	59.79 (24.84)	75.16 (18.56)	0.017
Self-reported intensity	85.58 (13.76)	75.67 (15.19)	86.20 (15.09)	82.86 (17.83)	87.79 (12.46)	81.47 (16.51)	0.102
Counting accuracy	0.99 (0.01)	0.99 (0.01)	0.99 (0.01)	0.99 (0.01)	0.99 (0.01)	0.99 (0.01)	0.922

Bayesian Model Comparison

When comparing models (based on (82, 83)), the "perception only" model was better than the

model that included learning prior expectations for the no-guessing and breath-hold conditions

(protected exceedance probability = 1), whereas there was not a significantly better model for the

guessing condition (protected exceedance probabilities = .46 vs. .54, slightly favoring the

learning model). The learning model was better in the tone condition (protected exceedance

probability = 1). No group differences were observed when comparing model fits between

groups.

For consistency/comparability, we use the "perception only" model parameters to compare

conditions in our analyses below, as this model best explained heartbeat tapping behavior

overall. The accuracy of this model – defined as the percentage of choices to tap/not tap that

matched the highest probability action in the model (e.g., a tap occurring when the highest

probability percept in the model was a heartbeat) – was 74% across all conditions; by condition,

model accuracy was: guessing condition = 67% (SD = 12%); tone condition = 67% (SD = 12%);

no-guessing condition = 85% (SD = 14%); breath-hold condition = 84% (SD = 13%).

Relationship between parameters

As shown in **Figure 2**, correlations between IP across task conditions were generally low.

Correlations between pHB estimates across conditions were moderate, most notably between the

no-guessing and breath-hold conditions (which also included the no-guessing instruction). The

tendency to tap in an anticipatory vs. reactionary manner (AvR) showed no relationships across

conditions. Correlations between IP and pHB (or pTone) within each condition were also low (0

$\leq$ r $\leq$ .27), as were correlations between these parameters and AvR (-.28 < r < .01; see

supplementary materials).

Figure 2. Pearson correlations between model parameters across task conditions across all participants. For reference, correlations at p < .001 (uncorrected) are marked with red asterisks.

Parameter face validity in terms of

Figure 3 shows the correlations, including some significant relationships ($p \leq .001$, uncorrected), across all participants between model parameters in each condition and several task-relevant variables. IP showed positive relationships with self-reported heartbeat intensity ratings in both the no-guessing and breath hold conditions. Additionally, pHB was lower in those self-reporting greater difficulty in the no-guessing condition, and higher in those self-reporting higher confidence and higher heartbeat intensity in both the no-guessing and breath hold conditions. Across heartbeat tapping conditions, model parameters were also weakly (IP) to strongly (pHB) related to the traditional counting accuracy measure (39).

In the breath-hold condition, t-tests revealed significantly greater IP in males than females $(t(251) = 2.30, p = .02)$.

Figure 3. Exploratory Pearson correlations between model parameters and self-report and other task-relevant variables for each task condition across all participants. IP = interoceptive precision parameter, pHB = prior expectation for heartbeat parameter, pT = prior expectation for tone parameter, AvR = anticipate vs. react strategy parameter, PTT = median pulse transit time, #HBs = number of heartbeats during the task condition, BMI = body mass index. For reference, correlations at p < .001 (uncorrected) are marked with red asterisks.

Group by condition interactions in model parameters

Parameter values for each group and condition are listed in **Table 4**. Across conditions, all

parameters were normally distributed (skew < 2).

Table 4. Mean (and standard deviation) for model parameters by group and condition

Individual difference variable	Healthy comparisons	Anxiety	Depression	Depression + Anxiety	Eating disorder	Substance use disorder	p-value*
Sensory precision							
Guessing	0.04 (0.05)	0.04 (0.02)	0.04 (0.03)	0.04 (0.04)	0.05 (0.05)	0.04 (0.04)	0.815

No-Guessing	0.04 (0.05)	0.04 (0.04)	0.03 (0.04)	0.03 (0.03)	0.06 (0.06)	0.04 (0.05)	0.09
Breath-hold	0.07 (0.07)	0.03 (0.03)	0.04 (0.04)	0.04 (0.04)	0.03 (0.03)	0.04 (0.04)	<0.001
Tone	0.18 (0.12)	0.12 (0.11)	0.18 (0.12)	0.14 (0.11)	0.21 (0.10)	0.16 (0.11)	0.052
Prior expectations							
Guessing	0.44 (0.16)	0.37 (0.16)	0.41 (0.17)	0.38 (0.18)	0.43 (0.18)	0.40 (0.16)	0.331
No-Guessing	0.15 (0.13)	0.15 (0.12)	0.17 (0.13)	0.17 (0.14)	0.14 (0.12)	0.21 (0.14)	0.05
Breath-hold	0.18 (0.11)	0.19 (0.19)	0.17 (0.11)	0.18 (0.13)	0.14 (0.11)	0.22 (0.13)	0.056
Tone	0.50 (0.01)	0.50 (0.01)	0.50 (0.01)	0.50 (0.01)	0.50 (0.01)	0.50 (0.01)	0.139
Anticipate vs. React							
Guessing	0.49 (0.06)	0.50 (0.04)	0.49 (0.05)	0.49 (0.06)	0.54 (0.07)	0.50 (0.06)	0.042
No-Guessing	0.50 (0.07)	0.52 (0.06)	0.49 (0.05)	0.50 (0.05)	0.50 (0.08)	0.50 (0.06)	0.683
Breath-hold	0.49 (0.10)	0.49 (0.04)	0.51 (0.06)	0.50 (0.06)	0.48 (0.04)	0.50 (0.05)	0.529
Tone	0.50 (0.22)	0.45 (0.16)	0.52 (0.21)	0.49 (0.18)	0.37 (0.20)	0.46 (0.19)	0.088

*p-values correspond to the results of ANOVAs comparing the groups (i.e., not including all task

conditions within the analysis as reported in the main text).

An ANOVA revealed that sensory precision (IP or auditory precision in the tone condition) was

significantly different between conditions ($F(3,1721) = 332$, $p < .001$; see **Figure 4**), reflecting

greater values in the tone condition than in the three heartbeat tapping conditions (post-hoc

Tukey comparisons indicated $p < .001$ in all cases). A subsequent ANOVA focused only on IP

(excluding the tone condition), including condition, clinical group, and their interaction revealed

a main effect of group ($F(5,1273) = 2.35$, $p = .04$) and a group by condition interaction

($F(10,1273) = 3.10$, $p < .001$); post-hoc Tukey comparisons indicated greater IP in HCs than

DEP/ANX ($p = .05$) and DEP ($p = .04$), and that the interaction was driven by the following: 1)

IP was significantly greater in the breath-hold condition than in the guessing ($p = .005$) and no-

guessing ($p = .02$) conditions in HCs, but not in any of the other clinical groups, and 2) HCs had

greater IP than ANX, DEP, DEP+ANX, and SUDs in the breath-hold condition ($p = .04$, $p = $

.008, p = .001, and $p < .001$, respectively). We subsequently confirmed that the group by

condition interaction remained significant ($F(10,1114) = 2.7$, $p = .003$) when also including age,

sex, BMI, median PTT, precision within the tone condition, number of heartbeats (and it's

interaction with condition), and medication status within the model. No significant effect on IP

was observed for any of these other variables, with the exception of a negative association with

age ($F(1,1114) = 5.02$, $p = .03$). The main effect of group was only present at a trend level after

accounting for these additional variables ($F(2,1114) = 2.7$, $p = .07$). See **Figure 5** for plots of IP

values by group and condition.

pHB was significantly different between the different task conditions ($F(3,1721) = 704.48$, $p <$

.001), indicating a lower prior expectation to feel a heartbeat in the no-guessing and breath-hold

condition (which retained the no-guessing instruction) than in the guessing condition, as well as

a higher prior expectation to hear a tone in the tone condition than to feel a heartbeat in the three

interoception conditions (post-hoc Tukey comparisons indicated p < .001 in all cases). No effect

of group, or group by condition interactions were observed for pHB in a subsequent

interoception-focused model excluding the tone condition. See **supplementary materials** for

plots of pHB values by group and condition.

No difference was observed between conditions in the tendency to anticipate vs. react (AvR) to

the tone/heartbeat using an ANOVA. There was also no effect of group, or condition by group

interactions for AvR in a subsequent interoception-focused model excluding the tone condition.

See **supplementary materials** for plots of AvR values by group and condition.

Comparison to existing measures

To examine the ability of traditional measures to capture similar group differences, we ran

analogous analyses using the traditional heartbeat counting task formula for interoceptive

accuracy (39). An ANOVA revealed a main effect of trial condition ($F(2,1273) = 139.00$, $p <$

.001) and group ($F(5,1273) = 3.57$, $p = .03$) on counting accuracy, but no group by condition

interaction. Tukey post-hoc comparisons revealed that counting accuracy was greater in the

guessing condition than in the no-guessing and breath-hold conditions (both $p < .001$), and that

accuracy was higher in SUDs than in DEP and DEP/ANX ($p = .04$ and $.004$, respectively). These

effects remained significant after accounting for the other variables we controlled for above.

To assess potential group differences in the effect of task condition on self-reported experience

and physiology, we also carried analogous analyses assessing confidence, intensity, and

difficulty, as well as heart rate. These results are reported in the **supplementary materials**. No

group by condition interactions in these variables were observed mirroring our IP results.

Figure 4. Bottom: Bar plots illustrating means and standard errors for model parameters by task
condition across all participants. Prior expectations for heartbeats were lower in the no-guessing and
breath-hold conditions. Sensory precision (i.e., interoceptive precision for the heartbeat or auditory
precision for the tone condition) was much greater in the tone condition. This was expected given the

unambiguous nature of this signal relative to the heartbeat signal. There were no significant differences
in sensory precision between the heartbeat conditions. The Anticipate vs. React values revealed no
mean differences between conditions. Top: For more complete data characterization, we also show
raincloud plots depicting the same results in terms of individual datapoints, boxplots (median and
upper/lower quartiles), and distributions. These illustrate that, for the Anticipate vs. React parameter,
nearly equally sized clusters of participants appeared to adopt more anticipatory (<.5) vs. reactive (>.5)
strategies in the tone condition, and that prior expectations remained unbiased (.5; with little variance)
in the tone condition relative to the heartbeat tapping conditions.

Figure 5. Bottom: Mean and standard error for interoceptive precision estimates by condition and
clinical group. Interoceptive precision (IP) was significantly greater in healthy comparisons than all other
groups in the breath-hold condition (*p < 0.05; with the exception of the eating disorders group, likely
due to the small sample size for this group), and healthy comparisons showed a significant increase in IP
from the guessing to breath-hold conditions that was absent in the other groups. Top: For more
complete data characterization, we also show raincloud plots depicting the same results in terms of
individual datapoints, boxplots (median and upper/lower quartiles), and distributions.

Associations with symptom severity measures and interoceptive awareness scales

Given the heterogeneity in our clinical sample, we ran subsequent exploratory correlational

analyses with continuous scores on the clinical measures gathered, excluding HCs, to assess

whether model parameters might provide additional information about symptom severity. We

note a few weak relationships at uncorrected levels in **supplementary materials**, but none

survive correction for multiple comparisons.

The T1000 dataset also includes self-report measures commonly used in interoception research,

including the Multidimensional Assessment of Interoceptive Awareness (MAIA; (84)), the

Toronto Alexithymia Scale (TAS-20; (85)), and the Anxiety Sensitivity Index (ASI; (86)). For

the interested reader, we also show exploratory correlation matrices between model parameters

and these common measures within **supplementary materials**. While a couple of relationships

were significant at uncorrected levels, the strength of these relationships was low.

Discussion

This investigation aimed to examine whether a novel computational (active inference) model of

perception could provide a principled approach to empirically characterizing interoceptive

processing dysfunctions that have previously been proposed. Specifically, we used behavior

during a heartbeat perception task in conjunction with this model to estimate quantitative

differences in the prior expectations and sensory precision estimates that individuals implicitly

apply to afferent interoceptive (cardiac) signals – in both healthy individuals and a

transdiagnostic sample of individuals with depression, anxiety, substance use, and/or eating

disorder symptoms. We observed several relationships in the expected directions between model

parameters and other task-related variables that supported the construct validity of our model

parameters. Of greatest interest, we found evidence that an interoceptive (breath-hold)

perturbation increased the precision estimates assigned to cardiac signals in healthy individuals,

but that this perturbation had no effect on interoceptive precision in individuals with depression,

anxiety, substance use, and/or eating disorders. In contrast, no significant differences in prior

expectations were observed, suggesting sensory precision, and not prior expectations, best

accounts for interoceptive differences in the clinical groups. Model comparison also suggested

that individuals did not update prior expectations over time during the task, which is perhaps

unsurprising given the low cardiac awareness commonly seen in previous studies (6). That is,

individuals may not have had sufficient signal to learn from (note that, in contrast, model

comparison supported the presence of learning in our auditory control condition with a clear

signal). We expand on these points below.

Parameter validity/sensitivity

IP showed positive relationships with self-reported heartbeat intensity ratings (no-guessing and

breath-hold conditions) – as would be expected in the context of more precise cardiac signals. In

the no-guessing and breath-hold conditions, pHB was lower than in the guessing condition. In

other words, participants appear to have successfully adjusted their prior expectations to comply

with the no-guessing instructions. Further, consistent with the role of prior expectations in

perception, pHB was also lower in those reporting greater difficulty (no-guessing condition) and

higher in those reporting greater confidence and higher heartbeat intensity (no-guessing and

breath-hold conditions). Each of these results support the notion that our parameters tracked

theoretically meaningful individual differences in perception/behavior and that our approach can

disentangle the effects of sensory precision from those of prior expectations and anticipatory vs.

reactive strategies.

Further, while model parameters had some shared variance with traditional accuracy measures

(e.g., higher counting accuracy was weakly associated with higher interoceptive precision), they

mainly captured unique variance that was not tracked by standard measures. Further, no group

differences analogous to those seen with model parameters were found using the traditional

interoceptive accuracy measure (i.e., across all participants, counting accuracy showed an

opposite pattern, being highest in the guessing condition). This highlights the unique ability of

this computational method to uncover differences in perceptual decision making across different

psychiatric subtypes.

It is also worth noting the strong positive correlation we observed between pHB and counting

accuracy in the no-guessing and breath-hold conditions, suggesting that heartbeat counting

accuracy primarily reflects prior expectations (as previously proposed in (38)). In the present

task this is explained by the fact that higher pHB values led to a greater number of taps and the

fact that the average number of taps in these conditions was low (i.e., because of the no-guessing

instruction). Thus, those who tapped more often approached the actual number of heartbeats and

therefore had higher counting accuracy scores. This highlights one specific way in which, in the

context of restrictions on guessing, counting accuracy may be most closely associated with prior

expectations.

Differences in Interoceptive Precision

Our primary results were that: 1) IP was higher in the healthy comparisons than in the clinical

groups (in general for those with depression/anxiety, and for all clinical groups but eating

disorders in the breath-hold condition; note that IP values in eating disorders were numerically

comparable to the other clinical groups, but did not reach significance due to small sample size);

and 2) there was a group by condition interaction, demonstrating that the interoceptive

perturbation (breath-hold) increased IP (relative to resting conditions) in the healthy participants,

whereas this perturbation had no effect in any of the clinical groups. These group differences

were not accounted for by any other demographic (e.g., age, sex) or physiological variables (e.g.,

pulse transit time, changes in heart rate). The hypothesized finding that IP was reduced across

psychiatric groups included in this study may be of clinical interest. First, multiple

neurocognitive (55, 87, 88) and computational (11, 12, 15, 27, 89) theories of emotion, and

associated empirical findings (e.g.,(42, 90-94)), suggest that interoceptive awareness may be an

important transdiagnostic factor in promoting emotional awareness. As low emotional awareness

has been linked to multiple psychiatric and systemic medical conditions (reviewed in (95, 96)),

reduced IP could contribute to low emotional awareness and its maladaptive consequences

irrespective of diagnostic category. Second, visceral regulation might be expected to be less

effective in the absence of precise feedback signals from the body, which could relate to visceral

dysregulation in psychiatric conditions (e.g., see (56)). For example, to adaptively regulate

visceral states, visceromotor control regions plausibly require reliable feedback signals from the

body to determine whether descending visceromotor commands have successfully led to the

intended adjustments in internal bodily states (10, 13, 36). Thus, the low IP observed across

multiple psychiatric conditions could potentially contribute to visceral dysregulation in other

contexts in which interoceptive perturbations occur, such as highly arousing, negatively valenced

states (e.g. panic anxiety, irritability/anger etc.).

These results build on previous bodies of work suggesting associations between psychiatric

disorders and interoceptive processing deficits (38, 56, 97-100). For example, previous cardiac

perception studies have shown that depressed patients exhibit reduced accuracy on a heartbeat

counting task (2-4), and that performance is negatively correlated with depressive symptoms (5)

as well as associated with both lower positivity and poorer decision-making (2); although, the

limitations of heartbeat counting tasks should be kept in mind when interpreting such findings

(47, 49, 101, 102). While the literature on interoceptive dysfunction is mixed for anxiety

disorders broadly, it is well-stablished in panic disorder (reviewed in (6)). A couple recent

studies have also reported evidence of differences in interoceptive processing in both SUDs (e.g.,

blunted brain responses (7)) and eating disorders (e.g., stronger expectation effects on perception

during low arousal (8)). The computational framework within which our findings were observed

is also in line with several recent proposals about the role of interoceptive inference in guiding

(predictive) autonomic control and the potential breakdown of this mechanism within different

psychiatric conditions (10-15, 36, 38); however, in contrast to previous emphasis on altered prior

expectations in these proposals, our results more selectively support the existence of deficits in

adjusting precision estimates for afferent interoceptive signals – and do not suggest the presence

of altered priors.

That said, the neurobiological mechanisms promoting reduced IP in mental disorders during

interoceptive perturbation remains unclear. As IP increased during the interoceptive perturbation

in HCs, but not in the clinical groups, it could be that altered brain processes in individuals with

certain psychiatric disorders fail to update IP estimates during states of acute bodily arousal. The

neural process theory associated with active inference suggests an inability to adjust IP estimates

would correspond to reductions in synaptic plasticity in response to changes in patterns of

interoceptive prediction-errors (10, 24, 76, 78, 103), most plausibly within neural networks

supporting interoception and visceromotor control (e.g., insula and anterior cingulate (15, 104)).

This could be tested using our model/task in conjunction with neuroimaging. Alternatively, IP

estimates could be accurate, and afferent interoceptive signals may in fact be conveyed with less

fidelity (i.e., greater noise) to the brain in the context of psychiatric conditions. For example,

such differences could be due to altered signaling in interoceptive sensory axons, which could in

principle be affected by many factors (genetic/epigenetic influences, early adversity and related

socio-environmental factors, and/or effects of disease-related chronic stress, among others).

Future research will need to investigate these different possibilities.

Strengths, Limitations, and Conclusion

This study has several major strengths. First is the novel application of a computational (active

inference) model to behavior on an interoceptive awareness task, which allowed for model

comparison (e.g., allowing us to rule out learning effects) and parameter estimates to disentangle

distinct computational mechanisms (e.g., the role of prior expectations vs. interoceptive

precision). A second strength is the application of this model to interoceptive processing in

individuals with psychiatric disorders, which to our knowledge, has never been reported. While

accounting for other influences, this approach allowed us to test a theoretical prediction – that

afferent interoceptive signals (and resulting prediction-errors) are assigned low precision

estimates in those with psychiatric disorders, leading those signals to be under-weighted during

interoceptive inference (although, as discussed above, other interpretations are possible). While

an important first step, a further test of this hypothesis would require replication in a

confirmatory sample, something we plan to investigate in the second cohort of participants in the

Tulsa 1000 project. Additionally, combining this approach with neuroimaging measures would

help to establish the expected relationship between IP and neural activity within interoceptive

cortices. Active inference models may be especially useful in this regard, as they allow for

simulation of predicted neuronal responses (78).

Some limitations of our modeling approach are also important to keep in mind. First, we were

required to make specific choices about model structure. For example, while priors during

estimation were fixed at plausible values (justification explained within the methods section),

other values might have been chosen. Also, we chose not to include an explicit action model.

While this was possible, in simple perceptual tasks like ours that lack reward feedback, any

variability in the precision of action selection would be correlated with sensory precision;

693 therefore, we chose a simpler model that treated behavior as a direct readout of confidence in

694 perception (and controlled for sensory precision in the tone condition to account for individual

695 differences in motor stochasticity). However, the novelty of our approach entails that it should be

696 replicated in future studies.

697 Another limitation is that, while we did compare models with vs. without learning, we did not

698 compare our perceptual model to other existing approaches. The heartbeat tapping task was not

699 well-suited for the several alternatives we considered. For example, it did not have a sufficient

700 number of trials to be suited for traditional signal detection approaches, which would be most

701 comparable to our model-based precision measure (105). Computational models based on

702 reinforcement learning were also inappropriate as the task did not include planning or learning

703 from reward, and instead dealt mainly with uncertainty in perception. Consequently, we

704 employed an active inference model that only includes perceptual inference. This involved

705 discretizing timepoints where responses were considered co-occurrent (or not) with diastole vs.

706 systole; but other such discretizations could have been chosen. That said, given the relationships

707 we observed between parameters and other task measures, our choices appear to have led to

708 estimates that track meaningful individual differences in task behavior. The task measurement

709 conditions also had certain limitations. For example, many individuals had low IP values –

710 reflecting the low cardiac awareness commonly seen at rest in previous studies (6) – which may

711 have limited the variability necessary to assess relationships with other variables. Finally, low

712 power, due to a relatively lower sample size in the eating disorders and anxiety disorders groups,

713 may have prevented us from detecting some effects in these groups.

714 In summary, this study 1) demonstrated the sensitivity of individual difference measures

715 (parameter estimates) derived from a novel active inference-based computational model, with a

focus on estimating the precision weighting of interoceptive signals across a transdiagnostic

sample of individuals with psychiatric disorders, and 2) tested—and found evidence

supporting—the hypothesis that individuals with psychiatric disorders fail to update the precision

weighting of afferent interoceptive signals during homeostatic perturbations. While the

underlying neurophysiological mechanisms leading to this difference remain unidentified, these

results point to a potential origin of visceral dysregulation (and perhaps its influence on

maladaptive behavior) across multiple psychiatric conditions. This represents an important step

towards a primary goal of computational psychiatry – computationally phenotyping individuals

with psychiatric disorders with the tools of computational neuroscience in hopes of using this

information to guide the development of precision medicine interventions.

Software Note: All model simulations were implemented using standard routines
(**spm_MDP_VB_X.m**) that are available as Matlab code in the latest version of SPM academic
software: http://www.fil.ion.ucl.ac.uk/spm/.

Funding: This work has been supported in part by The William K. Warren Foundation, the
National Institute of Mental Health, K23MH112949 (SSK), and the National Institute of General
Medical Sciences Center Grant, P20GM121312 (MPP, SSK, JF, JLS).

Conflict of Interest: None of the authors have any conflicts of interest to disclose.

<u>Supplementary Files</u>

Supplementary materials. This file includes results of supplementary analyses as well as
supplementary figures, as referred to in the main text.

Study data. This file includes all data used in the analyses reported in the manuscript.

References

1.   Khalsa SS, Adolphs R, Cameron OG, Critchley HD, Davenport PW, Feinstein JS, et al.
Interoception and Mental Health: A Roadmap. Biol Psychiatry Cogn Neurosci Neuroimaging.
2018;3(6):501-13.
2.   Furman D, Waugh C, Bhattacharjee K, Thompson R, Gotlib I. Interoceptive awareness, positive
affect, and decision making in major depressive disorder. Journal of affective disorders. 2013;151:780-5.
3.   Terhaar J, Viola FC, Bär K-J, Debener S. Heartbeat evoked potentials mirror altered body
perception in depressed patients. Clinical neurophysiology. 2012;123:1950-7.
4.   Dunn BD, Stefanovitch I, Evans D, Oliver C, Hawkins A, Dalgleish T. Can you feel the beat?
Interoceptive awareness is an interactive function of anxiety- and depression-specific symptom
dimensions. Behav Res Ther. 2010;48(11):1133-8.
5.   Pollatos O, Traut-Mattausch E, Schandry R. Differential effects of anxiety and depression on
interoceptive accuracy. Depression and anxiety. 2009;26:167-73.
6.   Khalsa SS, Lapidus RC. Can Interoception Improve the Pragmatic Search for Biomarkers in
Psychiatry? Front Psychiatry. 2016;7:121.
7.   Stewart JL, Khalsa SS, Kuplicki R, Puhl M, Investigators T, Paulus MP. Interoceptive attention in
opioid and stimulant use disorder. Addict Biol. 2019:e12831.
8.   Khalsa SS, Craske MG, Li W, Vangala S, Strober M, Feusner JD. Altered interoceptive awareness
in anorexia nervosa: effects of meal anticipation, consumption and bodily arousal. International Journal
of Eating Disorders. 2015;48(7):889-97.
9.   Allen M, Levy A, Parr T, Friston KJ. In the Body's Eye: The Computational Anatomy of
Interoceptive Inference. bioRxiv. 2019:603928.
10.   Owens AP, Allen M, Ondobaka S, Friston KJ. Interoceptive inference: From computational
neuroscience to clinic. Neurosci Biobehav Rev. 2018;90:174-83.
11.   Seth A. Interoceptive inference, emotion, and the embodied self. Trends in cognitive sciences.
2013;17:565-73.
12.   Seth AK, Critchley HD. Extending predictive processing to the body: Emotion as interoceptive
inference. Behavioral and brain sciences. 2013;36:47-8.
13.   Stephan K, Manjaly Z, Mathys C, Weber L, Paliwal S, Gard T, et al. Allostatic Self-efficacy: A
Metacognitive Theory of Dyshomeostasis-Induced Fatigue and Depression. Frontiers in human
neuroscience. 2016;10:550.
14.   Petzschner F, Weber L, Gard T, Stephan K. Computational Psychosomatics and Computational
Psychiatry: Toward a Joint Framework for Differential Diagnosis. Biological Psychiatry. 2017;82:421-30.
15.   Barrett L, Simmons W. Interoceptive predictions in the brain. Nature reviews Neuroscience.
2015;16:419-29.
16.   Bastos A, Usrey W, Adams R, Mangun G, Fries P, Friston K. Canonical microcircuits for predictive
coding. Neuron. 2012;76:695-711.
17.   Friston K. A theory of cortical responses. Philosophical transactions of the Royal Society of
London Series B, Biological sciences. 2005;360:815-36.
18.   Chen C, Takahashi T, Nakagawa S, Inoue T, Kusumi I. Reinforcement learning in depression: A
review of computational research. Neurosci Biobehav Rev. 2015;55:247-67.
19.   Clark J, Watson S, Friston K. What is mood? A computational perspective. Psychological
Medicine. 2018:1-8.
20.   Friston K, Stephan K, Montague R, Dolan R. Computational psychiatry: the brain as a phantastic
organ. The lancet Psychiatry. 2014;1:148-58.
21.   Huys Q, Maia T, Frank M. Computational psychiatry as a bridge from neuroscience to clinical
applications. Nature Neuroscience. 2016;19:404-13.
22.   Montague P, Dolan R, Friston K, Dayan P. Computational psychiatry. Trends in Cognitive
Sciences. 2012;16:72-80.

23.   Moutoussis M, Shahar N, Hauser T, Dolan R. Computation in Psychotherapy, or How
Computational Psychiatry Can Aid Learning-Based Psychological Therapies. Computational Psychiatry.
2017:1-21.
24.   Parr T, Friston K. The Anatomy of Inference: Generative Models and Brain Structure. Frontiers in
Computational Neuroscience. 2018;12:90.
25.   Schwartenbeck P, Friston K. Computational Phenotyping in Psychiatry: A Worked Example.
eNeuro. 2016;3:ENEURO.0049-16.2016.
26.   Sharp PB, Eldar E. Computational models of anxiety: nascent efforts and future directions.
Current Directions in Psychological Science. 2019;28(2):170-6.
27.   Smith R, Parr T, Friston KJ. Simulating Emotions: An Active Inference Model of Emotional State
Inference and Emotion Concept Learning. Front Psychol. 2019;10:2844.
28.   Smith R, Lane RD, Parr T, Friston KJ. Neurocomputational mechanisms underlying emotional
awareness: Insights afforded by deep active inference and their potential clinical relevance. Neurosci
Biobehav Rev. 2019;107:473-91.
29.   Adams R, Shipp S, Friston K. Predictions not commands: active inference in the motor system.
Brain Structure and Function. 2013;218:611-43.
30.   Adams R, Perrinet L, Friston K. Smooth Pursuit and Visual Occlusion: Active Inference and
Oculomotor Control in Schizophrenia. PLoS ONE. 2012;7:e47502.
31.   Edwards M, Adams R, Brown H, Pareés I, Friston K. A Bayesian account of 'hysteria'. Brain.
2012;135:3495-512.
32.   Schwartenbeck P, FitzGerald T, Mathys C, Dolan R, Friston K. The Dopaminergic Midbrain
Encodes the Expected Certainty about Desired Outcomes. Cerebral Cortex. 2015;25:3434-45.
33.   Iglesias S, Mathys C, Brodersen K, Kasper L, Piccirelli M, den Ouden H, et al. Hierarchical
Prediction Errors in Midbrain and Basal Forebrain during Sensory Learning. Neuron. 2013;80:519-30.
34.   Diaconescu A, Mathys C, Weber L, Kasper L, Mauer J, Stephan K. Hierarchical prediction errors in
midbrain and septum during social learning. Social Cognitive and Affective Neuroscience.
2017;12:nsw171.
35.   Petzschner FH, Weber LA, Wellstein KV, Paolini G, Do CT, Stephan KE. Focus of attention
modulates the heartbeat evoked potential. Neuroimage. 2019;186:595-606.
36.   Smith R, Thayer JF, Khalsa SS, Lane RD. The hierarchical basis of neurovisceral integration.
Neurosci Biobehav Rev. 2017;75:274-96.
37.   Owens A, Friston K, Low D, Mathias C, Critchley H. Investigating the relationship between
cardiac interoception and autonomic cardiac control using a predictive coding framework. Autonomic
Neuroscience. 2018;210:65-71.
38.   Paulus MP, Feinstein JS, Khalsa SS. An Active Inference Approach to Interoceptive
Psychopathology. Annu Rev Clin Psychol. 2019;15:97-122.
39.   Schandry R. Heart beat perception and emotional experience. Psychophysiology.
1981;18(4):483-8.
40.   Ludwick-Rosenthal R, Neufeld RW. Heart beat interoception: a study of individual differences.
Int J Psychophysiol. 1985;3(1):57-65.
41.   Garfinkel S, Seth A, Barrett A, Suzuki K, Critchley H. Knowing your own heart: Distinguishing
interoceptive accuracy from interoceptive awareness. Biological Psychology. 2014;104:65-74.
42.   Simmons W, Avery J, Barcalow J, Bodurka J, Drevets W, Bellgowan P. Keeping the body in mind:
insula functional organization and functional connectivity integrate interoceptive, exteroceptive, and
emotional awareness. Human brain mapping. 2013;34:2944-58.
43.   Brener J, Kluvitse C. Heartbeat detection: judgments of the simultaneity of external stimuli and
heartbeats. Psychophysiology. 1988;25(5):554-61.

44.   Katkin ES, Morell MA, Goldband S, Bernstein GL, Wise JA. Individual differences in heartbeat
discrimination. Psychophysiology. 1982;19(2):160-6.
45.   Whitehead WE, Drescher VM, Heiman P, Blackwell B. Realtion of heart rate control to heartbeat
perception. Biofeedback Self Regul. 1977;2(4):317-92.
46.   Phillips GC, Jones GE, Rieger EJ, Snell JB. Effects of the presentation of false heart-rate feedback
on the performance of two common heartbeat-detection tasks. Psychophysiology. 1999;36(4):504-10.
47.   Ring C, Brener J, Knapp K, Mailloux J. Effects of heartbeat feedback on beliefs about heart rate
and heartbeat counting: a cautionary tale about interoceptive awareness. Biol Psychol. 2015;104:193-8.
48.   Windmann S, Schonecke OW, Frohlig G, Maldener G. Dissociating beliefs about heart rates and
actual heart rates in patients with cardiac pacemakers. Psychophysiology. 1999;36(3):339-42.
49.   Ring C, Brener J. Heartbeat counting is unrelated to heartbeat detection: A comparison of
methods to quantify interoception. Psychophysiology. 2018;55(9):e13084.
50.   Murphy J, Millgate E, Geary H, Ichijo E, Coll MP, Brewer R, et al. Knowledge of resting heart rate
mediates the relationship between intelligence and the heartbeat counting task. Biol Psychol.
2018;133:1-3.
51.   Khalsa SS, Rudrauf D, Sandesara C, Olshansky B, Tranel D. Bolus isoproterenol infusions provide
a reliable method for assessing interoceptive awareness. Int J Psychophysiol. 2009;72(1):34-45.
52.   Hassanpour MS, Yan L, Wang DJ, Lapidus RC, Arevian AC, Simmons WK, et al. How the heart
speaks to the brain: neural activity during cardiorespiratory interoceptive stimulation. Philos Trans R Soc
Lond B Biol Sci. 2016;371(1708).
53.   Schandry R, Bestler M, Montoya P. On the relation between cardiodynamics and heartbeat
perception. Psychophysiology. 1993;30(5):467-74.
54.   Huang H, Thompson W, Paulus MP. Computational dysfunctions in anxiety: Failure to
differentiate signal from noise. Biological psychiatry. 2017;82(6):440-6.
55.   Barrett L. How emotions are made: The secret life of the brain. 2017.
56.   Barrett LF, Quigley KS, Hamilton P. An active inference theory of allostasis and interoception in
depression. Philos Trans R Soc Lond B Biol Sci. 2016;371(1708).
57.   Critchley H, Garfinkel S. Interoception and emotion. Current Opinion in Psychology. 2017;17:7-
14.
58.   Smith R, Lane RD. The neural basis of one's own conscious and unconscious emotional states.
Neurosci Biobehav Rev. 2015;57:1-29.
59.   Craig AD. How do you feel--now? The anterior insula and human awareness. Nature Reviews
Neuroscience. 2009;10:59-70.
60.   Domschke K, Stevens S, Pfleiderer B, Gerlach AL. Interoceptive sensitivity in anxiety and anxiety
disorders: an overview and integration of neurobiological findings. Clin Psychol Rev. 2010;30(1):1-11.
61.   Yoris A, Esteves S, Couto B, Melloni M, Kichic R, Cetkovich M, et al. The roles of interoceptive
sensitivity and metacognitive interoception in panic. Behav Brain Funct. 2015;11:14.
62.   Bogacz R. A tutorial on the free-energy framework for modelling perception and learning.
Journal of mathematical psychology. 2017;76:198-211.
63.   Victor TA, Khalsa SS, Simmons WK, Feinstein JS, Savitz J, Aupperle RL, et al. Tulsa 1000: a
naturalistic study protocol for multilevel assessment and outcome prediction in a large psychiatric
sample. BMJ open. 2018;8(1):e016620.
64.   Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. J
Gen Intern Med. 2001;16(9):606-13.
65.   Norman SB, Cissell SH, Means-Christensen AJ, Stein MB. Development and validation of an
Overall Anxiety Severity And Impairment Scale (OASIS). Depress Anxiety. 2006;23(4):245-9.
66.   Staley D, el-Guebaly N. Psychometric properties of the Drug Abuse Screening Test in a
psychiatric patient population. Addict Behav. 1990;15(3):257-64.

67.   Morgan JF, Reid F, Lacey JH. The SCOFF questionnaire: a new screening tool for eating disorders.
West J Med. 2000;172(3):164-5.
68.   Aupperle RL, Paulus MP, Kuplicki R, Touthang J, Victor T, Yeh HW, et al. Web-Based Graphic
Representation of the Life Course of Mental Health: Cross-Sectional Study Across the Spectrum of
Mood, Anxiety, Eating, and Substance Use Disorders. JMIR Ment Health. 2020;7(1):e16919.
69.   Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-
International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured
diagnostic psychiatric interview for DSM-IV and ICD-10. The Journal of clinical psychiatry. 1998;59 Suppl
20:22-33;quiz 4-57.
70.   Smith R, Feinstein J, Kuplicki R, Forthman KL, Stewart JL, Paulus MP, et al. Perceptual
Insensitivity to the Modulation of Interoceptive Signals in Depression, Anxiety, and Substance Use
Disorders. PsyArXiv. 2020.
71.   DeVille DC, Kuplicki R, Stewart JL, Tulsa I, Aupperle RL, Bodurka J, et al. Diminished responses to
bodily threat and blunted interoception in suicide attempters. Elife. 2020;9.
72.   Canales-Johnson A, Silva C, Huepe D, Rivera-Rei Á, Noreika V, Garcia MdC, et al. Auditory
Feedback Differentially Modulates Behavioral and Neural Markers of Objective and Subjective
Performance When Tapping to Your Heartbeat. Cerebral Cortex. 2015;25(11):4490-503.
73.   Allen J, Murray A. Age-related changes in the characteristics of the photoplethysmographic
pulse shape at various body sites. Physiol Meas. 2003;24(2):297-307.
74.   Ring C, Brener J. The temporal locations of heartbeat sensations. Psychophysiology.
1992;29(5):535-45.
75.   Friston K, Lin M, Frith C, Pezzulo G, Hobson J, Ondobaka S. Active Inference, Curiosity and
Insight. Neural Computation. 2017;29:2633-83.
76.   Friston K, Parr T, de Vries B. The graphical brain: Belief propagation and active inference.
Network Neuroscience. 2017;1:381-414.
77.   Da Costa L, Parr T, Sajid N, Veselic S, Neacsu V, Friston K. ACTIVE INFERENCE ON DISCRETE
STATE-SPACES – A SYNTHESIS. arXiv. 2020:2001.07203v2 [q-bio.NC]
78.   Friston K, FitzGerald T, Rigoli F, Schwartenbeck P, Pezzulo G. Active Inference: A Process Theory.
Neural Computation. 2017;29:1-49.
79.   Parr T, Friston K. Working memory, attention, and salience in active inference. Scientific Reports.
2017;7:14678.
80.   Daunizeau J, den Ouden HE, Pessiglione M, Kiebel SJ, Stephan KE, Friston KJ. Observing the
observer (I): meta-bayesian models of learning and decision-making. PLoS One. 2010;5(12):e15554.
81.   Friston K, Mattout J, Trujillo-Barreto N, Ashburner J, Penny W. Variational free energy and the
Laplace approximation. Neuroimage. 2007;34(1):220-34.
82.   Rigoux L, Stephan KE, Friston KJ, Daunizeau J. Bayesian model selection for group studies -
revisited. Neuroimage. 2014;84:971-85.
83.   Stephan KE, Penny WD, Daunizeau J, Moran RJ, Friston KJ. Bayesian model selection for group
studies. Neuroimage. 2009;46(4):1004-17.
84.   Mehling WE, Price C, Daubenmier JJ, Acree M, Bartmess E, Stewart A. The Multidimensional
Assessment of Interoceptive Awareness (MAIA). PLoS One. 2012;7(11):e48230.
85.   Parker J, Taylor G, Bagby R. The 20-Item Toronto Alexithymia Scale: III. Reliability and factorial
validity in a community population. Journal of Psychosomatic Research. 2003;55:269-75.
86.   Sandin B, Chorot P, McNally RJ. Anxiety sensitivity index: normative data and its differentiation
from trait anxiety. Behav Res Ther. 2001;39(2):213-9.
87.   Smith R, Killgore WDS, Lane RD. The structure of emotional experience and its relation to trait
emotional awareness: A theoretical review. Emotion. 2018;18(5):670-92.

88.   LeDoux JE, Brown R. A higher-order theory of emotional consciousness. Proc Natl Acad Sci U S A. 2017;114(10):E2016-E25.

89.   Smith R, Lane RD, Parr T, Friston KJ. Neurocomputational mechanisms underlying emotional awareness: insights afforded by deep active inference and their potential clinical relevance. Neurosci Biobehav Rev. 2019.

90.   Terasawa Y, Moriguchi Y, Tochizawa S, Umeda S. Interoceptive sensitivity predicts sensitivity to the emotions of others. Cognition & emotion. 2014;28:1435-48.

91.   Gu X, Hof P, Friston K, Fan J. Anterior insular cortex and emotional awareness. The Journal of comparative neurology. 2013;521:3371-88.

92.   Zaki J, Davis JI, Ochsner K. Overlapping activity in anterior insula during interoception and emotional experience. NeuroImage. 2012;62:493-9.

93.   Herbert B, Herbert C, Pollatos O. On the Relationship Between Interoceptive Awareness and Alexithymia: Is Interoceptive Awareness Related to Emotional Awareness? Journal of Personality. 2011;79:1149-75.

94.   Smith R, Lane RD, Sanova A, Alkozei A, Smith C, Killgore WDS. Common and Unique Neural Systems Underlying the Working Memory Maintenance of Emotional vs. Bodily Reactions to Affective Stimuli: The Moderating Role of Trait Emotional Awareness. Front Hum Neurosci. 2018;12:370.

95.   Lane RD, Weihs KL, Herring A, Hishaw A, Smith R. Affective agnosia: Expansion of the alexithymia construct and a new opportunity to integrate and extend Freud's legacy. Neurosci Biobehav Rev. 2015;55:594-611.

96.   Kashdan T, Barrett L, McKnight P. Unpacking Emotion Differentiation: Transforming Unpleasant Experience by Perceiving Distinctions in Negativity. Current Directions in Psychological Science. 2015;24:10-6.

97.   Paulus MP, Stein MB. Interoception in anxiety and depression. Brain Struct Funct. 2010;214(5-6):451-63.

98.   Paulus MP, Stewart JL. Interoception and drug addiction. Neuropharmacology. 2014;76 Pt B:342-50.

99.   Paulus MP, Tapert SF, Schulteis G. The role of interoception and alliesthesia in addiction. Pharmacol Biochem Behav. 2009;94(1):1-7.

100.   Verdejo-Garcia A, Clark L, Dunn BD. The role of interoception in addiction: a critical review. Neurosci Biobehav Rev. 2012;36(8):1857-69.

101.   Ring C, Brener J. Influence of beliefs about heart rate and actual heart rate on heartbeat counting. Psychophysiology. 1996;33(5):541-6.

102.   Desmedt O, Luminet O, Corneille O. The heartbeat counting task largely involves non-interoceptive processes: Evidence from both the original and an adapted counting task. Biol Psychol. 2018;138:185-8.

103.   Friston K, FitzGerald T, Rigoli F, Schwartenbeck P, O Doherty J, Pezzulo G. Active inference and learning. Neuroscience and biobehavioral reviews. 2016;68:862-79.

104.   Kleckner I, Zhang J, Touroutoglou A, Chanes L, Xia C, Simmons W, et al. Evidence for a Large-Scale Brain System Supporting Allostasis and Interoception in Humans. Nature human behaviour. 2017;1:0069.

105.   Kleckner IR, Wormwood JB, Simmons WK, Barrett LF, Quigley KS. Methodological recommendations for a heartbeat detection-based measure of interoceptive sensitivity. Psychophysiology. 2015;52(11):1432-40.

Supplementary material

<u>Supplementary Analyses</u>

Group by condition interactions in other task-relevant variables

To assess potential group differences in the effect of task condition on self-reported experience

and physiology, we also carried analogous analyses assessing confidence, intensity, and

difficulty, as well as heart rate.

Heart beat intensity. A main effect of both condition ($F(2,1273) = 44.14$, $p < .001$) and group

($F(5,1273) = 9.9$, $p < .001$) was found on self-reported heart beat intensity, but no group by

condition interaction; post-hoc Tukey comparisons indicated greater intensity in the breath-hold

than guessing or no-guessing conditions (ps $< .001$), and that SUDs reported greater intensity

than HCs ($p < .001$), ANX ($p = .002$), DEP ($p = .01$), DEP/ANX ($p < .001$), and eating disorders

($p < .001$).

Confidence. A main effect of both condition ($F(2,1273) = 25.11$, $p < .001$) and group ($F(5,1273)$

$= 3.25$, $p = .006$) was found on self-reported confidence, but no group by condition interaction;

post-hoc Tukey comparisons indicated greater intensity in the breath-hold than guessing or no-

guessing conditions (ps $< .001$), and that eating disorders reported lower confidence than SUDs

and DEP/ANX ($p = .008$ and $.02$, respectively).

Difficulty. A main effect of both condition ($F(2,1273) = 18.62$, $p < .001$) and group ($F(5,1273)$

$= 5.94$, $p < .001$) was found on self-reported difficulty, but no group by condition interaction;

post-hoc Tukey comparisons indicated greater difficulty in the no-guessing condition than in the

guessing or breath-hold conditions ($ps < .001$), and that SUDs reported less difficulty than DEP,

DEP/ANX, and eating disorders ($ps < .001$, .007, and .007, respectively).

Heart rate. A main effect of group ($F(5,1273) = 5.60$, $p < .001$) was found on heart rate, as well

as a marginal effect of condition ($F(2,1273) = 2.55$, $p = .08$), but no group by condition

interaction; post-hoc Tukey comparisons indicated marginally faster heart rate in the breath-hold

condition than in the no-guessing condition ($p = .07$), and that SUDs and DEP/ANX had faster

heart rate than HCs ($ps < .001$ and .04, respectively), and that SUDs had faster heart rate than

eating disorders ($p = .04$).

Associations with symptom severity

Given the heterogeneity in our clinical sample, we ran subsequent exploratory correlational

analyses with continuous scores on the clinical measures gathered, excluding the healthy

comparisons, to assess whether model parameters in the heartbeat tapping conditions might

provide additional information about symptom severity. The only notable (but weak)

relationships observed were as follows. In the no-guessing condition, IP was negatively

associated with both depression (PHQ; $r = -.13$, $p = .01$) and anxiety (OASIS; $r = -.11$, $p = .03$)

severity, and positively associated with substance use severity (DAST; $r = .10$, $p = .04$); pHB

was also positively associated with substance use severity (DAST; $r = .17$, $p = .001$). In the

breath-hold condition, pHB was positively associated with substance use severity (DAST; $r =$

.16, $p = .002$). Higher anxiety (OASIS) was also associated with a slightly more anticipatory

response pattern (AvR; $r = -.12$, $p = .02$).

1020

1021

1022 **Supplementary Figures**

1023

1024
1025 *Figure S1. Top: Example EKG trace segment from one participant (red). Bottom: Simultaneous PPG trace from the same*
1026 *participant. Each vertical black line denotes the upswing in the PPG signal after each EKG R-spike. Each cyan line indicates the*
1027 *length of the delay (i.e., the pulse transit time; PTT) between each R-spike and each subsequent upswing in PPG, the median of*
1028 *which was calculated for each participant. These showed that a delay of 200ms was a good estimate of PTT, which is what was*
1029 *assumed for computational modelling.*

1030

Figure S2. Correlations between model parameters by condition across all participants. For reference, correlations at p < .05 are marked with red asterisks.

Figure S3. Bottom: Mean and standard error for prior expectation estimates by condition and clinical group. Prior expectations (pHB) were significantly higher in the guessing condition, but were not significantly different between groups in the analyses reported in the main text (although marginally greater pHB in the no-guessing and breath-hold conditions can be seen in the substance use group relative to the other groups). Top: Raincloud plots showing the same results in terms of individual datapoints, boxplots, and distributions.

Figure S4. Bottom: Mean and standard error for Anticipate vs. React (AvR) parameter estimates by condition and clinical group. AvR was not significantly different between groups in the analyses reported in the main text. Top: Raincloud plots showing the same results in terms of individual datapoints, boxplots, and distributions.

Figure S5. Correlations between model parameters by condition across all participants. For reference, correlations at p < .05 (uncorrected) are marked with red asterisks. IP = interoceptive precision, pHB = prior expectation for heartbeat, AvR = anticipatory vs. reactive tapping strategy parameter.

$$P(s_{t=1})$$

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \begin{matrix} \text{Start} \\ \text{No Heartbeat} \\ \text{Heartbeat} \end{matrix}$$

$$P(s_{t=2})$$

$$\begin{bmatrix} 0 \\ P(nHB) \\ P(HB) \end{bmatrix}$$

$$s_t$$

$$\begin{matrix} & \text{Start} & & \\ \text{Start} & 0 & 0 & 0 \\ & 1-pHB & 1 & 0 \\ & pHB & 0 & 1 \end{matrix} \begin{matrix} \text{Start} \\ s_{t+1} \\ \\ \end{matrix}$$

$$B = P(s_{t+1}|s_t)$$

D → s_t → **B** → s_t

s_t → **A** → o_t

s_t → **A** → o_t

$$\begin{matrix} \text{Start} \\ \text{Diastole} \\ \text{Systole} \end{matrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

No Heartbeat   Heartbeat

$$\begin{matrix} & \text{Start} & & \\ 1 & 0 & 0 \\ 0 & IP & 1-IP \\ 0 & 1-IP & IP \end{matrix} \begin{matrix} \text{Start} \\ \text{Diastole} \\ \text{Systole} \end{matrix}$$

$$A = P(o_t|s_t)$$

Diastole   Systole

Heartbeat perception:

$$\bar{s}_{t=1} = \boldsymbol{\sigma}\left(\frac{1}{2}(\ln \mathbf{D} + \ln \mathbf{B}\cdot s_{t+1}) + \ln \mathbf{A}\cdot o_t\right) \qquad \bar{s}_{t=2} = \boldsymbol{\sigma}(\ln \mathbf{B}\, s_{t-1} + \ln \mathbf{A}\cdot o_t)$$

IP = interoceptive precision
pHB = prior expectation for heartbeat

Blue = diastole
Purple = systole
Light Red = no tap
Dark Green = tap

Green vertical lines = R-wave + 200ms
Red vertical lines = tap location

| | Anticipate vs. React | Prior Expectation for Heartbeat or Tone | Sensory Precision |

Interoceptive Precision Estimates by Group and Trial Type