FAST: a Flexible, Accurate and Speedy Test for COVID-19

Linjiajie Fang[1], Bing-Yi Jing[2], Shen Ling[3], and Qing Yang[4]

Abstract— As the COVID-19 pandemic continues worldwide, there is an urgent need to detect infected patients as quickly and accurately as possible. Group testing proposed by Technion [1][2] could improve efficiency greatly. However, the false negative rate (FNR) would be doubled. Using USA as an example, group testing would have over 70,000 false negatives, compared to 35,000 false negatives by individual testing.

In this paper, we propose a Flexible, Accurate and Speedy Test (FAST), which is faster and more accurate than any existing tests. FAST first forms small close contact subgroups, e.g. families and friends. It then pools subgroups to form larger groups before RT-PCR test is done. FAST needs a similar number of tests to Technion's method, but sharply reduces the FNR to a negligible level. For example, FAST brings down the number of false negatives in USA to just 2000, and it is seven times faster than individual testing.

I. INTRODUCTION

Reverse Transcription Polymerase Chain Reaction (RT-PCR) is one of the main diagnostic methods for early detection of coronavirus, SARS-CoV-2. However, RT-PCR is a technical intensive testing method, which requires professional technicians to perform in some specialized laboratories [3]. The overload at testing facilities would hinder the process of virus tests for a large population. As a result, there is an urgent need to improve the efficiency of overall testing. Researchers at Technion - Israel Institute of Technology and Rambam Health Care Campus introduced a faster way [1][2] to test coronavirus by pooling samples (Fig.1). They also suggested a positive pool up to the size of 32 or even 64 containing a single infected sample could still be confidently identified [1][2]. Such idea of applying group testing to reduce the number of necessary tests can be traced back to Dorfman [4].

Despite the high efficiency of Technion's proposal, one major drawback is that its false negative rate will double that of individual testing. Taking USA as an example, as in late May, there were about 1.7 million confirmed COVID-19 cases [5]. Assuming the false negative rate of RT-PCR test is 2%, individual test will estimate that around 35,000 infected patients as healthy, while the number would be 70,000 using Technion's method.

To overcome this problem and maintain the high efficiency, we propose a *Flexible, Accurate and Speedy Test*

Fig. 1: Group testing suggested by Technion. The pools of samples are tested and positive pools are later tested individually.

(FAST) (Fig.2). We first form small close contact subgroups of people e.g. families, colleagues, passengers from the same flight. These subgroups are then pooled together to form larger groups. Each group will be tested twice. If both results are negative (- -), the group is regarded as safe and testing stops. Otherwise, if either test is positive (+ +, + -, - +), testing will be conducted twice for each subgroup. Then the above "stop/continue" rule applies to the subgroups and also down to the individuals. A detailed example is shown in Fig.2.

According to the prevalence in USA, FAST pools an average number of 32 individuals in each group. The number of performed tests is less than 14% of the number of population size. In the meantime, comparing with Technion's strategy, FAST drastically lowers the false negatives from over 70,000 to about 2,000. In terms of estimating false positives of universal testing in USA, the individual test would wrongly diagnose 300,000 healthy people as positive, which is a huge pressure for medical systems. Technion's strategy would reduce this number to 22,500, yet FAST could lower this number to 3,600 in total.

The reasons why FAST can improve both efficiency and accuracy can be partially explained by its unique design, which can be characterised as follows.

- First, at each level (group, subgroup, and individual), two tests will be carried out. This can greatly decrease false negative rate of each single test to second order. However, the downside is that the procedure doubles both the number of tests and the false positive rate.
- Secondly, to overcome the above problems, FAST adopts two-layer grouping methods, which pools subgroups to a larger group. This two-layer structure can largely reduce the number of required tests and at the same time sharply reduce false positive rate.
- Finally, to further enhance efficiency, FAST pools close

[1]Mr. Linjiajie Fang is a PhD student in Data Science and Analytics, Hong Kong University of Science and Technology Email: lfangad@connect.ust.hk

[2]Prof. Bing-Yi Jing is a Professor in Mathematics, Hong Kong University of Science and Technology Email: majing@ust.hk (Corresponding Author)

[3]Ms. Shen Ling is a PhD candidate in Mathematics and Statistics, Dalhousie University, NS, Canada Email: lisaling@dal.ca

[4]Ms. Qing Yang is a PhD candidate in Mathematics, Hong Kong University of Science and Technology Email: qyangad@connect.ust.hk

Fig. 2: An example of performing FAST when each group containing four close contact subgroups. FAST performs two RT-PCR tests for each test simultaneously, and either one of the two tests shows positive, such group, subgroup or individual would be labeled positive. FAST first tests groups of samples in the first step, subgroups in a positive group will be tested one-by-one. Finally, samples in a positive subgroup will be re-tested individually.

contact subgroups to a larger group. Such close contact groups are common in real life, e.g., families, friends, colleagues. FAST is flexible in that it allows these subgroups to have different sizes.

Interestingly, we discover a simple relationship between the optimal pooling size and the prevalence i.e. $k_{opt} \approx 1/\sqrt{\text{prevalence}}$, which can be used to determine the optimal pooling size in practice easily. More details will be described in RESULTS.

All the results below concern the COVID-19 cases in USA as it is currently the worse affected country. However, we point out that FAST is very flexible and easily applied to other different countries/regions at different periods. FAST can be potentially more useful in some less developed countries/regions as it needs much fewer tests and is more accurate. Also, the COVID-19 may stay with us for quite some time, and thus testing will be a regular event in many parts of the world. FAST will be more useful as the prevalence becomes smaller as time moves on.

II. RESULTS

Efficiency comparisons Fig.3 compares the performance of individual test, Technion's strategy and FAST on the simulation set of 10,000 families. We assume the prevalence $\rho = 0.005$ and the infection rate within close-contact-subgroup $\beta = 0.7$. Here β is defined as the probability that one individual is infected by a virus carrier in the same subgroup.

Fig.3(a) shows that optimal pooling sizes of FAST and Technion's strategy are 15 and 32 respectively. And both FAST and Technion's strategies are over seven times efficient as the individual test. We evaluate the efficiency of testing strategies by calculating persons per test, i.e. the size of population divided by the total number of tests performed.

Dramatically lower FNR Fig.3(b) shows the results of false negative rates under different pooling sizes. Technion's strategy doubles the false negative rate from 2% to 4%. FAST dramatically lowers the false negative rate to nearly 0.0012. Such improvement is significant in reality: Taking USA as an example, 1.7 million cases were confirmed in late May. Individual test would miss 35,000 infected patients while Technion's strategy would double such number to 70,000. If FAST is applied, the number would sharply drop to around 2,000.

Low and stable FPR Fig.3(c) shows the results of false positive rates under different pooling sizes. By assuming the false positive rate to be 0.001 for RT-PCR test, the simulation shows that FAST has stable and almost zero false positive rate. Although the false positive rate of Technion's strategy is also far less than individual test, it is higher than FAST and nearly linear to the pooling size, which suggests avoiding pooling too much samples.

However, FAST can pool a much larger number of samples without increasing false positive rate. If the universal virus testing is applied to USA with around 300 million residents, and the optimal pooling sizes are applied, individual tests would wrongly diagnose 300,000 healthy people as infected. Technion's strategy would reduce this number to 22,500, yet FAST can lower this number to 3,600 in total.

Effects of infection rate within close-contact-subgroup Fig.4 compares the performance of Technion's strategy and FAST with optimal pooling size under different infection rate $\beta \in [0, 1]$. Here we fix the prevalence $\rho = 0.005$. Fig.4(a) shows that FAST always pools more samples than Technion's

Fig. 3: The performance of FAST, Technion's strategy and individual tests on 50 simulation sets. Each simulation set contains 10,000 families with distribution of household sizes following data in USA[6]. We fixed individual prevalence $\rho = 0.005$ and infection rate within close-contact-subgroup $\beta = 0.7$. We evaluate the efficiency and accuracy along different pooling sizes. (a): The efficiency versus pool size. (b): The false negative rate versus pool size. (c): The false positive rate changing pool size.

Fig. 4: The best performance of FAST and Technion's strategy under different infection rate within close-contact-subgroup. The prevalence and infection rate are fixed to be the same as Fig.2. (a): The optimal pooling size of FAST and Technion's strategy with respect to changing β. (b): The best efficiency and relative efficiency of FAST and Technion's strategy with respect to changing β.

Fig. 5: The best performance of FAST and Technion's strategy under different prevalence. The prevalence and infection rate are fixed to be the same as Fig.2.(a): The optimal pooling size of two methods with respect to changing ρ. (b): The efficiency and relative efficiency of two methods with respect to changing ρ.

strategy and the pooling size is linear to β. Fig.4(b) shows that when $\beta \geq 0.72$, FAST will be more efficient than Technion's strategy. Since SARS-CoV-2 spreads very easily and sustainably between people [7], it is reasonable to speculate that FAST would be more efficient in reality.

Different prevalence levels Fig.5 compares the performance of two methods under different prevalence. We fix the infection rate within close-contact-subgroup $\beta = 0.7$ and change the prevalence from 0.01% to 1%. Fig.5(a) and Fig.5(b) show that optimal pooling sizes and efficiencies of both two group testings decrease when the prevalence increases. Fig.5(b) shows that the relative efficiencies of both methods do not change much with prevalence. And the smaller the prevalence, the more efficient the two group testings should be.

Estimation of the optimal pooling size. Interestingly, we discover a simple relationship between the optional pooling size and the prevalence level. The significance of these simple estimations is that they can be used in reality to guide the implementation of grouping testing without tedious computation. Here we introduce the results without proofs:

1. Given the prevalence ρ, the optimal pooling size k_{opt} of individuals could be estimated by $k_{opt} \approx 1/\sqrt{\rho}$.

For example, when the prevalence is 0.005, it is estimated that the optimal pooling size of Technion's strategy is $1/\sqrt{0.005} \approx 14$. It is very close to the simulation result which optimally pools 15 samples. This can also be used to estimate the optimal pooling size of sub-pools:

2. Suppose the probability of a sub-pool containing infections is ρ^*, the optimal pooling size k_{opt}^* of sub-pools could be estimated by $k_{opt}^* \approx 1/\sqrt{\rho^*}$.

For FAST, $\rho^* = \frac{m}{1+(m-1)\beta}\rho$ with known average close contact subgroup size m and infection rate β. In our simulation $m = 2.5$ [6], $\beta = 0.7$, $\rho = 0.005$, thus $\rho^* \approx 0.0061$, we can quickly compute the optimal pooling size to be about $1/\sqrt{0.0061} \approx 13$ families or equivalently 32 people, which is the same as our simulation result. In our experiments, such estimation is effective with an error of at most one sample for $\rho \leq 0.4$.

Fig. 6: The theoretical efficiency of one-layer group testing method with repeated tests when the prevalence level is high.

III. DISCUSSION

How accurate can group testing achieve? If repetition is not adopted, two-layer grouping, which applies to the groups of subgroups, is much more efficient than one-layer grouping. However, compared with individual tests, one-layer grouping (like Technion's method) doubles FNR, and its FPR is nearly linear to the group size. Two-layer grouping triples FNR, and their FPR is nearly linear to the subgroup size. These facts direct us to avoid overmuch number of layers and prevent pooling too many individual samples in the last layer of group testing strategy.

To alleviate the problem of multiplying FNR without loss of much efficiency, FAST adopts a two-layer grouping structure and performs repeated tests, whose FNR is controlled to be at the level of second order and FPR is stable along different pooling sizes. Although repetition intuitively increases the total number of tests, by virtue of two-layer structure and infection information, FAST maintains comparable efficiency as Technion's strategy.

Necessity of group testing When the prevalence becomes very high, both efficiency and accuracy of group testing will be greatly reduced. Thus we have to consider if it is necessary to implement group testing when the level of prevalence is severe. Fig.6 provides theoretical efficiency of naive grouping testing with repeated tests, which suggests that when prevalence is over 10%, the number of tests of grouping patients would outnumber the individual tests. It could be a loss of the original intention for designing group testing. For areas with very severe epidemics, small pooling size will be appreciated or just remain as simple as individual testing. However, to the best of our knowledge, prevalence in most areas for COVID-19 is about 1% [8] or even lower, so the necessity of group testing is still evident for COVID-19 pandemic.

Determine close contact group The concept of close contact group can be freely generalized to any clique of close contacts, not only families. For example, the residential relations, roommates staying in the same dormitory unit. Furthermore, when applying group testing strategies in airport to diagnose the virus status of commuters, passengers sitting in front and back rows on the same plane can be pooled in the same close contact group.

Unlike Technion's strategy, which pools a fixed number of individuals, FAST pools samples by a fixed number of close contact subgroups. Since the size of close contact subgroups is flexible, FAST pools different number of samples from group to group, which concerns more about the contact information than restrictions on specific pooling size. In this paper, the contact between groups or subgroups is ignored. However, in reality, the contact between people is much more complicated. The contact information of patients plays an important role to distinguish high risk susceptible patients from low risk ones. A flexible grouping strategy based on contact network might help cluster the infected patients to certain pools and largely reduce the number of tests.

Flexibility of FAST strategy Not limited to USA, FAST can be flexibly adopted to different nations in different disease stage. With appropriate settings including prevalence and infected rate within close-contact-subgroup, FAST can generate the optimal pooling size with the corresponding best efficiency to achieve. Under the guidance of FAST, the false negative rate and the false positive rate can be maintained at a dramatically low level with good performance in efficiency in some range of parameters.

"Predictive" group testing In some areas, government would propose universal testing among all the population, where millions of individuals could be involved. In order to perform a more efficient testing, it is prioritized to test people who have high probability of getting infected. Before testing, we can make prediction of the risk level for someone based on their contact information, symptoms and residential areas. Such risk prediction is usually non-uniform among different groups of people. In the future, we hope to develop methods to predict the risk of people and build testing strategies based on different predictions.

IV. METHODS

Simulation Setup To fairly evaluate the performance of FAST, we set up our simulations with some practical assumptions. We assume 0.005 prevalence according to the number of infections in USA up to 29th May 2020 [5], and the distribution of family sizes follows data of the household in America [6]. That is, we choose the number of family members from $\{1, 2, 3, 4, 5, 6, 7\}$ with probabilities equal to $\{0.28, 0.35, 0.15, 0.13, 0.06, 0.02, 0.01\}$ respectively. As for the accuracy of each RT-PCR test, we assume FNR=0.02 and FPR=0.001, following the same configuration proposed in Hanel etc.(2020)[9].

In Fig.3 we assume a infection rate within close-contact-subgroup i.e. $\beta = 0.7$, in order to model the realistic case for COVID-19. In our data preparation, we first generate 10,000 families with fraction of infection equal to 0.0061, then we generate the number of members and infections of each family by the household distribution and the infection rate to make sure the overall individual prevalence is 0.005. For each pooling size, we simulate 50 replicates and evaluate the average performance.

In Fig.4, we follow the same configuration in Fig.3 except for changing infection rate β from 0 to 1 with a gap equal to 0.1. For each β, we search for the optimal pooling size

by evaluating strategies with pooling sizes less than 500 and choose the size with the best efficiency. Similarly, in Fig.5 we fix $\beta = 0.7$ and change prevalence from 10^{-4} to 10^{-2} with a gap of log-scale equal to 0.2 in order to evaluate the best performance on different prevalence levels. To avoid the variance when searching for the best efficiency, we simulate 10 replicates of simulation set containing 50,000 families and the average performance is compared.

V. CONCLUSION

In this paper, we introduce FAST, a *Flexible, Accurate and Speedy Test*, based on the design of group testing, adoption of close contact information and necessary repetition. Simulations concerning the cases in USA demonstrate that FAST can be over 7 times efficient as the individual test, and greatly reduce the false negative rate and the false positive rate to be far below the level of naive group testing. We also introduce a simple and very close estimation of the optimal pooling size $k_{opt} \approx 1/\sqrt{\text{prevalence}}$, which can be flexibly implemented in reality.

VI. CODE AVAILABILITY

Our code is available on https://github.com/Fang-Lin93/FAST_group_testing

REFERENCES

[1] N. Jeffay, "To ease global virus test bottleneck israeli scientists suggest pooling samples pooling method for accelerated testing of covid-19, url https://www.timesofisrael.com/to-ease-global-virus-test-bottleneck-israeli-scientists-suggest-pooling-samples/," 2020.

[2] "Pooling method for accelerated testing of covid-19, url https://www.technion.ac.il/en/2020/03/pooling-method-for-accelerated- testing-of-covid-19/," 2020.

[3] T. Nolan, R. E. Hands, and S. A. Bustin, "Quantification of mrna using real-time rt-pcr," *Nature protocols*, vol. 1, no. 3, p. 1559, 2006.

[4] R. Dorfman, "The detection of defective members of large populations," *The Annals of Mathematical Statistics*, vol. 14, no. 4, pp. 436–440, 1943.

[5] "Coronavirus disease 2019 (covid-19): Cases in the u.s., url cdc.gov/coronavirus/2019-ncov/cases-updates/cases-in-us.html," 2020.

[6] E. Duffin, "Distribution of households in the united states from 1970 to 2019, by household size, url https://www.statista.com/statistics/242189/distribution-of-households-in-the-us -by-household-size/," Nov. 2019.

[7] "Coronavirus disease 2019, url cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/how -covid-spreads.html," 2020.

[8] S. Lohse, T. Pfuhl, B. Berkó-Göttel, J. Rissland, T. Geißler, B. Gärtner, S. L. Becker, S. Schneitler, and S. Smola, "Pooling of samples for testing for sars-cov-2 in asymptomatic people," *The Lancet Infectious Diseases*, 2020.

[9] R. Hanel and S. Thurner, "Boosting test-efficiency by pooled testing strategies for sars-cov-2," *arXiv preprint arXiv:2003.09944*, 2020.

VII. ACKNOWLEDGEMENTS

Fang, Jing and Yang's research is partially supported by the Hong Kong RGC Grants GRF 16304419 and GRF 16305616.

VIII. AUTHOR CONTRIBUTIONS

L.J.J.F. developed the solutions of the model and executed the simulation and theoretical analysis. S.L. prepared the Figure1 and Figure 2. B.Y.J. conceived the work and revised the contents and forms of the manuscript. Q.Y. wrote the main manuscript and did the data analysis.

IX. COMPETING INTERESTS

The author declares no competing interest.