Pollen Explains Flu-Like and COVID-19 Seasonality

Martijn J. Hoogeveen[1], Eric C.M. van Gorp[2], and Ellen K. Hoogeveen[3].

1. Department Technical Sciences & Environment, Open University, The Netherlands.

2. Department of Viroscience and department of Infectious Diseases, Erasmus Medical Centre,

Rotterdam, The Netherlands.

3. Department of Internal Medicine, Jeroen Bosch Hospital, Den Bosch, The Netherlands.

Corresponding author:

Martijn J. Hoogeveen

De Liesbosch 12d, 3439LC Nieuwegein, The Netherlands

martijn.hoogeveen@icecat.com

Abstract

Current models for flu-like epidemics insufficiently explain multi-cycle seasonality. Meteorological

factors alone, including associated behavior, do not predict seasonality, given substantial climate

differences between countries that are subject to flu-like epidemics or COVID-19. Pollen is

documented to be allergenic, plays a role in immuno-activation, and seems to create a bio-aerosol

lowering the reproduction number of flu-like viruses. Therefore, we hypothesize that pollen may

explain the seasonality of flu-like epidemics including COVID-19 in conjunction with meteorological

variables.

We tested the Pollen-Flu Seasonality Theory for 2016-2020 flu-like seasons, including COVID-19, in

The Netherlands with its 17 million inhabitants. We combined changes in flu-like incidence per

100K/Dutch citizens (code: ILI) with weekly pollen concentrations and meteorological data. Finally, a

discrete, predictive model is tested using pollen and meteorological threshold values displaying

inhibitory effects on flu-like incidence.

We found a highly significant inverse correlation of r(224)= -0.38 (p < 0.00001) between pollen and

changes in flu-like incidence corrected for incubation period. The correlation was stronger after

taking into account incubation time, which satisfies the temporality criteria. We found that our

predictive model has the highest inverse correlation with changes in flu-like incidence of r(222) = -

0.48 (p < 0.001) when average thresholds of 610 total pollen grains/m3, 120 allergenic pollen

grains/m3, and a solar radiation of 510 J/cm2 are passed. The passing of at least the pollen

thresholds, preludes the beginning and end of flu-like seasons. Solar radiation is a co-inhibitor of flu-

like incidence, temperature makes no difference, and higher relative humidity associates even with flu-

like incidence increases.

We conclude that pollen is a predictor of the inverse seasonality of flu-like epidemics including

COVID-19, and that solar radiation is a co-inhibitor. The observed seasonality of COVID-19 during

Spring, suggests that COVID-19 may revive in The Netherlands after week 33, preceded by the

relative absence of pollen, and follows pollen-flu seasonality patterns.

Introduction

Current models for flu-like epidemics insufficiently explain multi-cycle seasonality. Meteorological

factors alone do not fully explain the seasonality of flu-like epidemics (Tamerius et al., 2011) or

COVID-19 (Yao et al, 2020). Pollen is documented to be allergenic (Klemens et al, 2007;

Rosenwasser, 2011; Howarth, 2000), plays a role in immuno-activation (Brandelius et al, 2020), and

there are some indications that pollen might be antiviral (Wachsman et al, 2000; Ghanem et al., 2015)

and anti-influenza (Chen, 2016). Interestingly, "other allergic diseases" are absent as a co-morbid

condition of COVID-19 according to Zhang et al. (2020), although more confirmation is needed

before conclusions can be drawn. One more indication for a pollen effect might be from a controlled

experiment indicating that flowering (and foliage) plants in hospital rooms lead to positive

physiological responses and a reduced recovery time for post-surgery patients (Park & Mattson, 2008).

At the other hand, it is reported that pollen might suppress certain immune parameters in non-allergic

subjects and is found to be correlated to rhinovirus-positive cases (Gilles et al, 2020), but not to other

flu-like virus positive cases (Nivel.nl, 2020).

Recently, we identified pollen bio-aerosol as a discrete seasonal factor in inhibiting flu-like epidemics

for the period 2016 to 2019 in The Netherlands (Hoogeveen, 2020). In this epidemiological study, we

found strong inverse correlations between allergenic pollen concentrations and hay fever on the one

hand, and flu-like incidence on the other hand. The study was based on the persistent observation that

pollen and flu season predictably alternate each other in moderate climate zones, and the absence of

sufficient meteorological explanations (Tamerius et al., 2011). We further observed that the passing of

pollen threshold values of around 100 allergenic pollen grains/m^3, reliably mark the onset and decline

of moderate flu-like epidemic lifecycles, and thus might be used as predictor. Such a concentration of

allergenic pollen makes sense as in an overview of real-life studies, clinical pollen threshold values are

observed between 1 and 400 grains/m^3, whereby the first symptoms are typically observed in the range

of 1 – 50 grains/m^3 (De Weger et al, 2013), depending on country, period, and vegetation of study, and

probably the susceptibility of subjects to allergens.

The seasonality of respiratory viral infections has already been recognized for thousands of years in

temperate regions (Moriyama et al., 2020). More in detail, virologists observed that the cold, and flu-

like epidemics (e.g., influenza and corona caused) "go away in May" in the Northern Hemisphere,

while emerging in the Southern Hemisphere with its opposite seasonality, to re-emerge in the Northern

Hemisphere during its next Autumn and Winter in a slightly mutated form. Furthermore, all new flu-

like pandemics since 1889 typically emerged in the Northern Hemisphere at the tail-end of respective

flu-seasons (Fox et al., 2017), whereby the current COVID-19 pandemic is clearly no exception. The

emergence of COVID-19 and other pandemics at the tail-end of flu season makes sense. It takes time

for a spontaneous new cross-over virus with a sufficiently high reproduction number (Ro) – for SARS-

CoV-2 it is estimated to be initially around 3 (Liu et al., 2020) - to develop from patient 0 to a full-

fledged pandemic during flu season on the Northern Hemisphere. Chances for the Northern

Hemisphere with its larger populations are higher to be the initial breeding ground for a new flu-like

pandemic than the Southern Hemisphere. Fox et al. showed further that most flu-like pandemics are

multi-wave, whereby the initial wave at the tail-end of flu season is typically short-lived. This gives

rise to the suspicion that COVID-19 is subject to such multi-wave seasonality as well (Kissler et al.,

2020) and has a short waive at the tail end of the 2019/2020 flu-like season in the Northern

Hemisphere.

Numerous studies try to explain flu-like seasonality from meteorological factors such as sun light

including UV radiation (Schuit et al., 2020), temperature and humidity (Chong et al., 2020; Shaman et

al, 2011). However, Postnikov (2016) concluded that ambient temperature is not a good predictor for

influenza seasonality in The Netherlands, and inconsistent correlations are found for the relation

between COVID-19 and temperature as well (Toseupu et al, 2020; Xie & Zhu, 2020; Ma et al 2020;

Qi et al, 2020). Also findings about the relation between humidity and influenza (Soebiyanto et al.,

2014), and humidity and COVID-19 (Ahmadi et al, 2020; Ma et al, 2020; Qi et al, 2020) are equally

inconsistent. Although UV light is detrimental for the flu-like virus aerosol under laboratory

conditions, associated with immuno-activation (Abhimanyu & Coussens, 2017; Tan & Ruegiger,

2020), and circadian rhythms regulating lung immunity (Nosal et al., 2020), the onset of flu season,

halfway August in The Netherlands, coincides with an annual peak in hot, sunny days and is still in the

middle of the Summer season. According to Yao et al. (2020) also for the decrease of COVID-19

infections, nor high UV nor high temperature are good predictors. The contradictory findings related

to COVID-19 understandingly based on the analysis of a limited part of the year and disease cycle,

might be partly due to sub-seasonal bias and unstandardized data collection methods. With sub-

seasonal bias, we mean that if only a part of a season or cycle is analyzed, different and thus biased

conclusions can be drawn.

Nevertheless, these meteorological variables are known factors in flowering, and pollen maturation

and dispersion. Meteorological variables such as increased solar radiation and temperature – among

others the absence of frost - are not only triggering flowering and pollen maturation, but also affect the

pollen bio-aerosol formation: dry and warm conditions stimulate pollen to be airborne. Rain, to the

opposite, makes pollen less airborne, and cools the bio-aerosol down. Very high humidity levels (RH

98%) are even detrimental for pollen (Guarnieri, 2006). The RH 98% effect on pollen, could thus

provide an alternative explanation of why flu-like incidence in tropical countries is higher during rainy

season, and reduced during the rest of the year.

We hypothesize that there is an inverse effect of pollen bio-aerosol on flu-like incidence including

COVID-19 (See Figure 1), whereby pollen are known to be triggered and influenced by

meteorological variables, which can then jointly explain the seasonality of flu-like incidence. The

indirect explanation of the pollen effect is based on the fact that pollen bio-aerosol and UV light

exposure lead to immuno-activation, and sometimes allergic symptoms, which *might* reduce the

effectiveness of flu-like viruses. The indirect pollen effect is explained by the spread of pollen bio-

aerosol under sunny and dry conditions, which *might* absorb viral bio-aerosol, and *might* as well

become an agent that leads to an increased exposure of flu-like viruses to virus-degrading solar

radiation including the UV spectrum, and - possibly – anti-viral properties of pollen.

To further understand the impact of pollen as an environmental factor influencing the life cycle of flu-

like epidemics, the objective of this study is to determine the correlations of pollen and meteorological

variables with (changes in) flu-like incidence, develop and test a discrete predictive model that

combines pollen and meteorological co-inhibitors, and infer whether COVID-19 in the tail-end of the

2019/2020 flu-like season is able to defy flu-like seasonality or not. Our main hypothesis therefore is

that pollen are the missing link, jointly explaining with certain meteorological variables, flu-like

seasonality including COVID-19, and that a compound threshold based factor - combining detected

flu-inhibitors - is a good unified predictor of such seasonality.

Figure 1: a simplified theoretic model explaining pollen-flu inverse seasonality, whereby pollen might

have a direct (I) "antiviral", indirect allergenic (II) and/or immune triggering (III) function,

inhibiting flu-like epidemic incidence in combination with meteorological conditions and triggers.

Methods

To study the relation between pollen and flu-like incidence in The Netherlands, we used the public

datasets of Elkerliek Hospital (Elkerliek.nl) about the weekly allergenic, low-allergenic and total

pollen concentrations in The Netherlands in grains/m^3, whereby for 42 types of pollen particles the

numbers are counted and averaged per day per 1 m^3 of air. The common Burkard spore trap is used

through which a controlled amount of air is ingested. The applied classification and analysis method is

conform to the standard by EACCI (European Academy of Allergology and Clinical Immunology)

and the EAN (European Allergy Network). Allergenic pollen includes nine types of particles that are

classified as moderate (*Corylus*, *Alnus*, *Rumex*, *Plantago* and *Cedrus Libani*), strong (*Betula* and

Artemisia), or very strong allergenic (*Poaceae* and *Ambrosia*). Additionally, we included low-

allergenic pollen concentrations in addition to the allergenic ones as it should not matter for the bio-

aerosol filter function (see Figure 1) whether pollen is allergenic or not. Low-allergenic pollen

includes the other 33 particles that are classified as non-allergenic to low-allergenic (*Cupressaceae,*

Ulmus, Populus, Fraxinus, Salix, Carpinus, Hippophae, Fagus, Quercus, Aesculus, Juglans, Acer,

Platanus, Pinus, Ilex, Sambucus, Tilia, Ligustrum, Juncaceae, Cyperaceae, Ericaceae, Rosaceae,

Asteraceae, Ranunculaceae, Apiaceae, Brassicaceae, Urtica, Chenopodiaceae, Fabaceae, Humulus,

Filipendula, and *Indet*). Total pollen concentration is the sum of the average allergenic and low-

allergenic pollen concentrations. One advantage of the total pollen metric is that there are hardly 0

values (only 3 out of 266), and we do not need to limit ourselves to just parts of the seasonal cycle

which could import sub-seasonal bias in our research. We assume that long distance pollen transport is

also accounted for as foreign pollen will be counted as well by the pollen measuring station that works

all year round.

Further, we use the data from the Dutch State Institute for Public Health (RIVM.nl) gathered by Nivel

(Nivel.nl) about weekly flu-like incidence (WHO code "ILI" - Influenza Like Illnesses) reports at the

primary medical care, per 100,000 citizens in The Netherlands. Primary medical care is the day-to-

day, first line healthcare given by local health care practices to their registered clients as typical for

The Netherlands with a population of currently 17.4 million. The reports relate to a positive RIVM

laboratory test for ILI after a medical practitioner diagnosed ILI after a consult, whether that leads to

hospitalization or not. The ILI metric is according to a standardized WHO method as ILI data are

gathered and compared globally. ILI is defined by WHO as a combination of a measured fever of ≥ 38

°C, and cough, with an onset within the last 10 days. The flu-like incidence metric is a weekly average

based on a representative group of 40 primary care units, and calculated using the number of

influenza-like reports per primary care unit divided by the number of patients registered at that unit,

averaged for all primary care units, and next extrapolated to the complete population. The datasets run

from week 1 of 2016 till week 18 of 2020 (n = 226 data points) to include the recent COVID-19

pandemic in the tail-end of the 2019/2020 flu-like season. To underpin the relative importance of

COVID-19: SARS-CoV-2 is detected in the Netherlands since week 9, 2020. According to figures of

Nivel.nl (2020; see Figure 2), from week 13 on, SARS-CoV-2 is the outcome of the (vast) majority of

positive tests for patients at primary care with flu-like complaints, whereby at week 18 even 100% of

positive tests indicate SARS-CoV-2 (other tested viruses are five Influenza A and B subtypes, RSV,

Rhinovirus and Enterovirus). A small bump in rhinovirus positive cases is seen after week 18 in 2020,

so not influencing our ILI statistics till week 18, at least not for 2020.

Figure 2: overview flu-like virus positive cases in 2020 show that from week 13 on SARS-CoV-2 is

dominant. The Rhinovirus cases bump is outside the timeline of our research. (source: Nivel.nl, 2020)

Further, we included meteorological datasets form the Royal Dutch Meteorological Institute

(KNMI.nl), including average relative humidity/day, average temperature/day and global solar

radiation in J/cm^2 per day as an indicator of UV radiation, from its centrally located De Bilt weather

station. Next, we calculated the weekly averages for the same periods as used in the other datasets. De

Bilt is traditionally chosen as it provides an approximation of modal meteorological parameters in The

Netherlands, which is a small country. Furthermore, all major population centra in The Netherlands

are within a radius of only 60 kilometer from De Bilt which covers around 70% of the Dutch

population. So, we assume in this study that the measurements of De Bilt are sufficiently

representative for the meteorological conditions that the Dutch population experiences on average.

To test allergenic versus low-allergenic pollen assumptions, against hay fever and pre-covid-19 flu-

like incidence, we make use of the hay fever index. The hay fever index is defined as turnover for hay

fever medication as reported by all Dutch pharmacies to the Dutch Central Bureau of Statistics

(CBS.nl) based on respective ATC codes (especially R01A/R01AC). We use a dataset from week 1 of

2016 till week 10 of 2019 (n=166 data points), as no further data has been made available. For the

interpretation of findings, we assume for The Netherlands a prevalence of allergic rhinitis that is more

or less similar to that in Western Europe being around 23% and frequently undiagnosed (Bauchau &

Durham, 2004). Further, it can be noted that the prevalence of allergic diseases in general in the

Netherlands is around 52% (Van de Ven et al, 2006).

Datasets are complete, except that there are three weekly pollen concentration measurements missing

(1.3%) because of a malfunctioning monitoring station, namely week 26 of 2016, week 21 of 2017 and

week 22 of 2019. These missings appear to be completely at random. We imputed missing values to

avoid bias and maintain power. We used a four weeks surrounding average to estimate the three

missing data points to avoid breaking lines in visuals. We checked that these missing data have no

material impact on the results by comparing these averages with the data of previous years for similar

periods, and by observing if removal from statistical tests has any effect on outcomes and conclusions.

Regarding the incidence of flu-like symptoms, we calculated the weekly change compared to the

previous week ($\Delta ILI = ILI_t - ILI_{t-1}$) to get an indication of the flu-like epidemic life cycle progression,

whereby a decline is interpreted as Ro<1 and an increase as Ro>1 (Ro is the reproduction number of

flu-like viruses). Further, to cover, in one time-series metric, for changes in flu-like incidence as well

as for an incubation period of up to two weeks, we calculated a three weeks moving average (3WMA)

of changes in flu-like incidence, of which two weeks are forward looking:

$$(\Delta ILI_{3WMA} = (\Delta ILI_t + \Delta ILI_{t+1} + \Delta ILI_{t+2})/3)$$

Whenever we use the term incubation time, we also mean to include reporting delay (estimated to be

around 4.5 days). We don't assume delay effects for meteorological variables or pollen concentrations,

so don't calculate moving areas for other time series.

Compared to our previous study (Hoogeveen, 2020) there is an overlap in datasets of less than 25%,

given the extension in time, the addition of meteorological datasets and non-allergenic pollen, and the

introduction of newly calculated variables such as total pollen concentration, ΔILI_{3WMA}, the compound

predictor and log10 transformations.

We formulated the following statistical null hypotheses for falsification:

$H1_0$: there are no inverse correlations for total pollen concentrations with flu-like incidence (corrected

for incubation period).

$H2_0$: there are no inverse correlations between pollen and *changes* in flu-like incidence (ΔILI or

corrected for incubation time: ΔILI_{3WMA}).

$H3_0$: there is no predictive significance of a discrete model's compound value, based on thresholds for

pollen and meteorological co-inhibitors, related to changes in flu-like incidence (ΔILI_{3WMA}).

To understand the role of meteorological variables, to check if - in our datasets - meteorological

variables show their well-established effects on pollen as assumed, and to select co-inhibitors:

$H4_0$: meteorological variables – solar radiation, temperature and relative humidity – have no effect on

pollen and/or flu-like incidence change (ΔILI_{3WMA}).

Low-allergenic pollen are known to have sometimes a little allergenic effect. To understand how to

interpret adding none-to-low-allergenic pollen to the total pollen metric, we want to verify their effects

on the hay fever index:

$H5_0$: low-allergenic pollen has no effect on hay fever and (changes in) flu-like incidence.

Note that except H5, all hypotheses are related to potential causality: the temporal sequentiality

(temporality) of the respective independent variables, and flu-like incidence corrected for incubation

period. Whenever we refer to temporality, we mean to indicate that the datasets behave *as if* there is

causality, in the understanding that statistics alone cannot proof causality in uncontrolled settings.

Statistical analyses

Variables are presented with their mean and standard deviation (SD).

We calculated correlation coefficients to test the hypotheses, as datasets show normal distributions, to

assess the strength and direction of relationships. We use the full datasets, to avoid sub-seasonal bias,

and by extending the number of years, the distortions by incidental and uncontrolled events are

supposed to be minimized. Only to verify the assumed, well-documented relation between Winter and

Spring temperature and pollen maturing, we used a sub-seasonal dataset as will be explained in the

Results section.

Next, linear regression (F-test) on identified inhibitors and interactions is used *descriptively* to

determine whether the relation can (statistically) be described as linear, and to determine the equation

using estimates and intercept values, and produce probability, significance level, F-value, and the

Multiple R squared correlation to understand the predictive power of the respective inhibitor. Standard

deviations and errors, and degrees of freedom (DF) are used as input for calculating the 95%

probability interval. We report in the text the outcome of statistical tests in APA style, adapted to

journal requirements. For relations that appear non-linear - logarithmic or exponential – we use the

log10 function to transform the data if that makes the relation appear linear, before re-applying linear

regression.

Finally, we created a simple, discrete model resulting in one compound value using selected flu-like

inhibitors, to determine the optimal average threshold values for these inhibitors which have the

highest joint correlation with changes in flu-like incidence (ILI_{3wMA}). We apply linear regression (F-

test) to understand the predictive power of the compound value, and determine the linear equation

when significant. By constructing one compound independent variable, we cover for collinearity or

interaction effects between joined co-inhibitors. In our analysis, we base the compound value on three

selected thresholds. For example, when one threshold value is passed this leads to a compound value =

1, and when all three threshold values are passed this leads to a compound value = 3. Therefore, the

values of the compound value are in the range of [0, 3].

263 The compound value equation can be expressed as follows, whereby iv = the respective independent

264 variable that acts as inhibitor of flu-like incidence and k relates to the respective calculated threshold

265 value. For each respective threshold passed (iv > k), +1 is added to the compound value (CV):

266 $$CV = \sum_{n=1}^{3} (iv_n > k_n) \rightarrow 1 \ \wedge \ (iv_n \leq k_n) \rightarrow 0$$

267 For three selected co-inhibitors, this takes in Excel the form of CV = IF($IV_1 > K_1$;1;0) + IF($IV_2 > K_2$;1;0)

268 + IF($IV_1 > K_3$;1;0), whereby we use a threshold value for solar radiation (k_r), and both pollen threshold

269 values for allergenic (k_{ap}) and total pollen (k_p) for K_1, K_2 and K_3 respectively, as will be shown in the

270 Results section.

271 It is outside of the scope of this research to verify the underlying datasets of Elkerliek Ziekenhuis,

272 RIVM/Nivel, CBS, and KNMI by examining the validity and reliability of data collection methods as

273 these institutes have well-established and internationally standardized protocols for data collection and

274 verification.

275 All regression analyses are done using the statistical package R version 3.5.

277 **Results**

278 The mean value and SD for all weekly pollen concentrations are 732 grains/m^3 (SD = 1368), for

279 allergenic pollen 349 grains/m^3 (SD = 987 grains/m^3), for low-allergenic pollen 383 grains/m^3 (SD =

280 626 grains/m^3) and for flu-like incidence 47 incidence/100K citizens per week (SD = 40.2

281 incidence/100K citizens).

Figure 3: Total pollen concentrations vs flu-like incidence in The Netherlands, whereby especially

passing the 610 total pollen grains/m³ threshold marks the seasonal switches around week 10 (±5

weeks) and week 33 (±2 weeks). For the interpretation of the relation an incubation time of up to 2

weeks should be taken into account, and one should consider the change in flu-like incidence (Ro >1

of Ro <1).

When further inspecting the datasets regarding pollen concentrations and flu-like incidence reported

by primary medical care in The Netherlands, it is clear that there are continuous pollen bursts (Figure

3), whereby only a few of these pollen bursts are classified as more allergenic (Figure 6). These bursts

of pollen, allergenic or low-allergenic, typically coincide with and precede a decline of flu-like

incidence.

In our previous study, we found a significant inverse correlation between allergenic pollen

concentrations and changes in flu-like incidence (ΔILI), but no significant inverse correlations

between allergenic pollen concentrations and flu-like incidence due to the short and extreme bursts of

these allergenic pollen (Hoogeveen, 2020). However, the addition of low-allergenic pollen reveals

highly significant negative correlations (r(224)= -0.19, p < 0.01) with flu-like incidence, which

become stronger when correcting for two weeks incubation time (r(222)= -0.31, p < 0.00001). The

correlation for total pollen and flu-like incidence is highly significant when taking into account

incubation time: r(222) = -0.19, p < 0.01. We can thus reject the null-hypotheses $H1_0$ in favor of the

alternative hypothesis that there is a negative correlation between total pollen and flu-like incidence,

including the first cycle of the COVID-19 pandemic, when taking into account incubation time.

Further, we can reject $H5_0$ in favor of our assumption that it makes sense to include non-allergenic

pollen concentrations in our study as well.

That the correlations become stronger when taking into account incubation time, implies temporality

(see Table 1). Furthermore, we can also observe from Figure 3 that flu-like incidence starts to decline

after the first pollen bursts. And that flu-like incidence starts to increase sharply *after* pollen

concentrations become very low or close to zero. This is a qualitative indication of temporality.

Further, we can notice that the first COVID-19 cycle behaves according to pollen-flu seasonality, at

least does not defy it.

	Total pollen		Low-allergenic pollen		DF
Flu-like incidence	-0.05	n.s.	-0.19	**	226
+2week incubation	-0.19	**	-0.31	***	224

Table 1. Correlation coefficients between flu-like incidence and total pollen and low-allergenic pollen.

*(n.s.: not significant, * p < 0.05, ** p < 0.01, *** p < 0.001)*

When testing the impact on ΔILI, the weekly *changes* in medical flu-like incidence (M = -0.25

incidence per 100K/citizens, SD = 15.4 incidence per 100K/citizens), the extended dataset till 2020,

including COVID-19, shows a strong and highly significant inverse correlation with allergenic pollen

(r(223) = -0.23, p = 0.000413). For total pollen, including low-allergenic pollen, the correlation is even

somewhat stronger (r(223) = -0.27, p = 0.000059). Therefore, we can falsify the null-hypothesis ($H2_0$)

that there are no inverse correlations between the weekly pollen concentrations – total or only

allergenic - and weekly changes in flu-like incidence (ΔILI), including the period covering the first

cycle of the COVID-19 pandemic. These inverse correlations provide thus further support for the

alternative hypothesis that the presence of an elevated level of pollen has an inhibiting effect on flu-

like incidence, and starts to immediately influence the direction and course of the epidemic life cycle.

Also during the COVID-19 dominated period of the last 9 weeks, it appears that flu-like incidence

behaves according to the expected pollen-flu seasonality. This strengthens the idea that COVID-19

itself might be seasonal as well, like all other flu-like pandemics since the end of the 19[th] century. Also

when looking at other data from RIVM.nl about COVID-19 hospitalizations, we cannot conclude that

COVID-19 breaks through the seasonal barrier. For example, new COVID-19 hospitalizations went

down from a peak of 611 on March 27 to just 33 on May 3, the last day of week 18.

Figure 4: scatter diagram showing the inverse relation between changes in flu-like incidence (ILI$_{3WMA}$)

and log10(pollen concentrations in grains/m^3).

Using the three weeks moving average (ΔILI$_{3WMA}$) of changes in flu-like incidence (M = -0.26 per

100K/citizens, SD = 8.9 per 100K/citizens), the correlation coefficients become stronger and are again

all highly significant for both allergenic pollen (r(224) = -0.34, p < 0.00001) and total pollen

concentration (r(224) = -0.38, p < 0.00001). We can thus also reject the null-hypothesis (H2$_0$) that

there is no inverse relation between pollen and ΔILI$_{3WMA}$, 3 weeks moving average of changes in flu-

like incidence including incubation time. As these correlations (see also Figure 4) are stronger than if
not corrected for incubation period, it is a further indication of temporality, and as they are stronger
now with the 2019/2020 flu-like season included, it appears to support the idea that COVID-19 is
subject to pollen induced flu-seasonality as well.

Linear regression analysis shows that there is a highly significant inhibitory effect of pollen on flu-like
incidence change (ΔILI_{3WMA}) of $F(1, 222) = 37.1$, $p < 0.001$ (see Table 2 line 1), as a further basis for
using total pollen concentration as a predictor. A Log10 transformation of pollen to compensate for
visual non-linearity leads to a similar outcome: $F(1, 222) = 45.60$, $p < 0.001$ (see Table, line 4). At
least visually, it gives a good fit (see Figure 4).

Δ Flu-change (ΔILI_{3WMA})	Estimate	95%CI	Intercept	Multiple R sq.	F-stat on DF	P<
1. Total pollen per 100/m³ incr.	-0.253	-0.334 to -0.171	1.53	0.14	37.1 (1, 222)	0.001
2. Solar radiation per 100 J/cm² incr.	-0.312	-0.475 to -0.153	2.98	0.06	14.43 (1, 222)	0.001
3. Compound predictor per incr. of 1	-3.88	-4.82 to -2.94	4.95	0.23	65.59 (1, 222)	0.001
4. Log10 (Total pollen per 100/ m³) incr.	-3.67	-4.21 to -3.13	7.54	0.17	45.60 (1, 222)	0.001

Table 2: Summary of univariate regression analyses of pollen (1), solar radiation (2) and our
compound pollen/solar radiation predictor (3) on changes in flu-like incidence (ΔILI_{3WMA}) showing all
highly significant (p < 0.001) results, but the correlation for solar radiation is weak (0.06) and the
compound predictor the strongest (0.23). The log10 (4) analysis is a check on non-linearity of pollen
concentrations in relation to ΔILI_{3WMA}, leading to the same conclusion.

Flu-like incidence per 100K p.w.	Estimate	95%CI	Intercept	Multiple R sq.	F-stat on DF	P<
Hay fever index per 1% incr.	-0.12677	-0.181 to -0.0723	63.81	0.11	20.74 (1, 164)	0.001

Δ Hay Fever Index (2016=100)	Estimate	95%CI	Intercept	Multiple R sq.	F-stat on DF	P<
Total pollen per 100/m³ incr.	5.3049	3.922 to 6.688	67.78	0.26	56.52 (1, 164)	0.001

Log10 (Flu-like incidence per 100K)	Estimate	95%CI	Intercept	Multiple R sq.	F-stat on DF	P<
Log10(hay fever index)	-0.444	-0,379 to -0,510	2.360	0.22	45.96 (1, 164)	0.001

Table 3: Summary of univariate regression analyses of total pollen concentration on hay fever (p <
0.001) and hay fever on flu-like incidence/100K citizens per week (p < 0.001), whereby pollen leads to
an increase in hay fever, which in turn is associated with a decrease in flu-like incidence. To
compensate for non-linearity, the regression of log10(hay fever) on log10(flu-like incidence) is added,
with the same highly significant outcome.

Previously, we established a highly significant inverse correlation of $r(164)= -0.34$ ($p = .00001$)
between the hay fever index ($M = 101$, $SD = 115.7$) and flu-like medical incidence (Hoogeveen,
2020). Univariate regression analyses show that there is a highly significant positive effect of total
pollen on hay fever incidence, which in turn has a highly significant inhibitory effect on flu-like
incidence (see Table 3). In line with the correlation between pollen and flu-like incidence, the
correlation between *total* pollen concentration and hay fever is stronger ($r(164)= 0.51$, $p < 0.00001$)
than those for allergenic and low-allergenic pollen individually. This confirms that we can best use
total pollen concentration as predictor.

Low-allergenic pollen has as well a highly significant effect on hay fever ($r(164)= 0.48$, $p < 0.00001$).
We can thus reject the null-hypothesis $H5_0$ in favor of the alternative hypothesis that also low-
allergenic pollen has a positive effect on hay fever. This might imply that pollen classified as none-to-
low-allergenic might still be responsible for certain allergic effects, and not just the more allergenic
pollen. Therefore, trying to use low-allergenic pollen to discriminate effects outside the allergenic path
regarding the immune system might be challenging.

377   *Figure 5: inverse correlation between hay fever and flu-like incidence*

378   The nature of the relation between hay fever and flu-like incidence, might be statistically described as

379   linear, but could be better described as logarithmic (Figure 5). In the context of this study, we interpret

380   it as a further indication that it could be described as a thresholds based switching pattern as well,

381   conforming the thresholds-based approach we take in our compound value as calculated below.

Figure 6: Both allergenic and low-allergenic pollen are positively correlated to hay fever.

The expected effects of relative humidity ($r(224) = -0.46$, $p < 0.001$), temperature ($r(224) = 0.10$ n.s.) and solar radiation ($r(224) = 0.35$, $p < 0.001$) on pollen are found. So, there is indeed more pollen with sunny, warmer and dry weather. We can reject the null-hypothesis that meteorological variables ($H4_0$) have no effect on pollen, for solar radiation ($M = 1047$ J/cm2, $SD = 709$ J/cm2) and relative humidity ($M = 79\%$, $SD = 8.3\%$), whereby relative humidity is reducing the amount of aerosol pollen. The relation with temperature ($M = 10.8$ ℃, $SD = 5.8$ ℃), is only highly significant during the start of pollen season – between week 1 and till week 20 - as it is known to shift the start of the pollen calendar ($r(98) = 0.51$, $p < .00001$). For other meteorological variables such a sub-seasonal analysis is not leading to materially different outcomes.

Counter to findings in other studies, relative humidity is positively associated to changes in flu-like incidence (ΔILI_{3WMA}) in The Netherlands ($r(224) = 0.34$, $p < 0.00001$). Dutch flu season is cold and humid, and on rainy days the effect of pollen and solar radiation are reduced. Although temperature

strongly correlates with flu-like incidence ($r(226) = -0.71$, $p < 0.001$), it has a negligible effect on Δ

ILI, weekly changes in flu-like incidence ($r(224) = -0.02$ n.s.), and it doesn't change with incubation

time. Therefore, it seems unlikely that temperature has a direct effect on aerosol flu-like viruses and

the life cycle of a flu-like epidemic. In line with this, temperature is also not a good marker for the

onset or the end of flu season as the end of flu season (Ro<1) can coincide with an average

temperature of close to 0 ºC and the start of flu season (Ro>1) can coincide with temperatures as high

as 17 ºC in The Netherlands.

Of the meteorological variables, only for solar radiation there is a highly significant inverse correlation

with changes in flu-like incidence (ΔILI_{3wMA}): ($r(224) = -0.25$, $p = 0.000156$).

Thus, from the meteorological variables, for solar radiation and relative humidity the null-hypothesis

($H4_0$) can also be rejected for that they have no effect on the flu-like epidemic lifecycle. But, of these

two only solar radiation is a flu-like inhibitor in line with its positive effect on pollen concentration,

and association with immune-activation and UV-effect on viruses.

A univariate linear regression also shows the highly significant negative correlation for solar radiation

on flu-like incidence change (ΔILI_{3wMA}) ($F(1, 222) = 14.43$, $p < 0.001$ (see Table 2 line 2). As the

correlation is weak (Multiple R-squared = .06), we interpret solar radiation as a co-inhibitor in relation

to pollen, that has stand-alone a too weak effect to explain flu-like seasonality.

Taking into account all these findings, we developed a discrete, compound model in which we take the

changes in flu-like incidence (ΔILI_{3wMA}), a threshold value for solar radiation (k_r), and both pollen

threshold values for allergenic (k_{ap}) and total pollen (k_p). We found that the compound model (M = 1.4

thresholds passed, SD = 1.1 thresholds passed) has the highest inverse correlation ($r(222) = -0.48$, $p <$

0.001) for the following threshold values: k_r: 510 J/cm^2, k_{ap}: 120 allergenic pollen grains/m^3, and k_p:

610 total pollen grains/m^3. In line with the previous outcomes, inclusion of relative humidity, low-

allergenic pollen or temperature did not improve the correlation strength of this model. As they also

did not show significant interaction effects with pollen, also such interactions are not meaningful to

consider for the model.

In each of the observed years the now (re)defined pollen thresholds are being passed in week 10 ($\pm$ 5

weeks) depending on meteorological conditions controlling the pollen calendar, which coincides also

with reaching flu-like peaks, and again in week 33 ($\pm$ 2 weeks) marking the start of the new flu-like

season.

There is a highly significant inverse relation of our compound thresholds based predictor value with

flu-like incidence change (ΔILI_{3WMA}) of $F(1, 222) = 65.59$, $p < 0.001$ and a Multiple R-squared

correlation of 0.2281 (see Table 2 line 3). This confirms the usefulness of a discrete, pollen and solar

radiation thresholds based model as a predictor of switches in flu-like seasonality, whereby the effect

of pollen is stronger than that of solar radiation. As a consequence we can reject the null-hypothesis

($H3_0$) that this compound pollen/solar radiation value has no predictive significance for flu-like

seasonality.

Discussion

We first discuss the possible implications of the results for our theoretic model and alternative

explanations. Next, we discuss our methods: alternatives or ways to improve them.

Theoretic model

We found highly significant inverse relations between pollen and solar radiation on the one hand, and

(changes in) flu-like incidence on the other hand: a higher number of pollen or an increase in solar

radiation, is related to a decline in flu-like incidence. The inverse correlation with pollen becomes

stronger when including the 2019/2020 period, which is increasingly COVID-19 dominated during the

last 9 weeks. This might be cautiously interpreted as an early indication that also COVID-19 is subject

to pollen-flu seasonality as all previous pandemics since the end of the 19[th] century before. However,

social distancing is likely to have contributed to flattening both the flu-like epidemic and COVID-19

pandemic curves at the tail-end of the 2019/2020 flu-like season: the Dutch government has applied

hygienic measures since March 9, 2020, and a mild form of a lockdown with social distancing from

March 11 on. Such behavioral policies will need to be included in the theoretic model in addition to

pollen and meteorological variables to understand the relative importance of social distancing versus seasonality.

As also non-to-low-allergenic pollen – we called them low-allergenic pollen – has a positive effect on the hay fever index, we could not use it as an independent variable to discriminate between the immuno-activation explanation and the ("anti-viral") bio-aerosol filter explanation in our model (see Figure 1).

The highly significant inverse correlation between hay fever and flu-like incidence can be interpreted in a number of ways, which are not mutually exclusive. A) Allergic rhinitis symptoms might make it more difficult for flu-like viruses to find their way to the lung cells that are vulnerable to it, such as ACE-2 receptor positive cells in case of SARS-CoV-2 (Wan et al., 2020). B) The anti-histamine hay fever medication possibly suppresses flu-like symptoms as well. Further, given the flower experiment mentioned in the introduction, it might make sense to look beyond allergies, and interpret immuno-activation in the broadest sense to include any auto-repair function of the human body.

The only meteorological variable that has a co-inhibitive effect on changes in flu-like incidence, solar radiation, has a stimulating effect on aerosol pollen formation, and is responsible for melatonin-induced immuno-activation. Relative humidity reduces pollen aerosol formation, and correlates positively with flu-like incidence. We did not specifically look at precipitation, but it might make sense as it reduces pollen dissemination.

In our study we showed that temperature, except for influencing the onset of pollen season, has no predictive value for changes in flu-like incidence, and therefore its inverse correlation with flu-like incidence might be interpreted in a number of ways: a) as spurious: the common causal factor is solar radiation, or b) as a stressor that has immediate effects on the functioning of the immune system of already infected persons. If we talk about the influence of meteorological variables, we assume associated behavioral aspects being covered, which are sometimes summarized as seasonal behavior, but this independent variable might have a cultural dimension that needs to be understood better.

We showed that a compound value, based on threshold values for pollen and solar radiation, results in

a stronger correlation with the flu-like lifecycle than the individual inhibitors. This model could form

an empirical basis for understanding flu-like seasonality, its Ro and reliably predicting the start and

end of each flu-like cycle. As behavior, in the form of hygiene and social distancing, is also widely

seen as an inhibitor, it might be of value to include this factor in our compound value as well, which

probably leads to an even stronger predictor for the evolution of the reproduction number Ro of flu-

like epidemics, although this might be beyond explaining the seasonality effect itself.

As long as, for COVID-19, the level of herd immunity (Fine et al., 2011) is still below required

thresholds for ending pandemics (Plans-Rubio, 2012), it might make sense to include indications of

herd immunity levels in the theoretic model as well.

Methods

In general, statistical research cannot proof causal relationships in uncontrolled environments, even if

datasets seem to behave *as if* there is causality. Such statistics, however, can provide indications and

identify reliable predictors, help filter out bad ideas, and be the inspiration for testable hypotheses that

can be verified in laboratory and other fully controlled experiments. Regarding statistical methods,

other approaches for time series analysis and hypothesis testing could be applied, non-linear tests

instead of our log10 transformations on logarithmic data or our discrete switching model, but in the

end only replicational studies in other countries and controlled experiments are the best way to

confront and verify our observations and hypotheses.

Regarding meteorological data, it might be useful to look if including more weather stations can help

to better approximate the weather conditions the Dutch population is experiencing on average. It might

be useful to distinguish patterns per province, and it might be of importance for understanding wind as

the main vector for dispersal of pollen in The Netherlands with its maritime and temperate climate. It

might also be good to look for the effects of climate change on pollen maturation (Frei & Gassner,

2008) as the climate in The Netherlands is affected.

Regarding the pollen dataset, it might be of interest when particle counts would include more pollen

types than currently covered, to get a further understanding of the magnitude and impact of airborne

pollen material, classified as allergenic or not. Such an expansion of pollen counting methodology

would probably require an update of the EAN protocols.

Regarding the hay fever index, there are probably lag effects, but the data collection method for this

finance based dataset, did not include assessment of validity and reliability, and thus there is no

estimate of such a lag effect. It might be good to include alternative datasets such as search engine

based trend analysis to be able to generate a complete dataset for a whole period of study, which can

be validated separately.

The ILI dataset is now based on a sample of 40 representative local primary care units. It might be an

improvement if the sample would be based on patients visiting *any* primary care unit in The

Netherlands to reduce the need for extrapolations, which would likely improve the reliability of the

dataset. Further, if it is confirmed that rhinovirus positive cases are positively correlated to pollen

concentrations, it might make sense to remove these from the ILI metrics, as it seems that this effect

cannot be generalized to other ILI viruses. The use of ΔILI_{3WMA} including incubation time, seems to be

more elegant than multiple tests with ΔILI. At the same time it could be argued that this metric could

better become a 2 weeks forward-looking moving average, as it is unlikely that in the first week any

effects can be noticed given the average reporting delays and incubation period.

We could more explicitly include behavioral variables (Gozzi et al, 2020), such as social distancing,

lockdowns and hygiene, in the current method, for example by rating lockdown regimes on a Likert-

type scale [1, 5] from no lockdown to a complete lockdown. Although seasonal behavior might be

implicitly covered by the meteorological variables, it might still make sense to model them more

explicit as there might be cultural patterns – such as holidays or seasonal celebrations – that need to be

checked for.

Although pollution is not an inhibitor of flu-like incidence (Coccia, 2020), it still might interact with

pollen. A more complete theoretic model, controlling for the (interactions with) pollution, can give

more insight in how to interpret the findings of this or similar studies.

It will require further research to test the findings, threshold values and predictive model for flu-like

seasonality in other countries with different climates.

Conclusion

We conclude that pollen and solar radiation both have highly significant inverse correlations with

changes in flu-like incidence. The inverse correlation of pollen with (changes in) flu-like incidence

becomes stronger when incubation time is included. A compound variable – based on the thresholds

for total pollen, allergenic pollen and solar radiation – shows the strongest correlation with changes in

flu-like incidence, and appears to be useful as predictor for switches in flu-like seasonality.

COVID-19 dominates the tail-end of the flu-like season The Netherlands. And, COVID-19 appears to

be subject to flu-like seasonality as well like other pandemics before. This gives rise to the hypothesis

that COVID-19 might be multicycle as well, and might thus resurface (Ro>1) from week 33 on,

together with other flu-like viruses, when pollen season is over in the Northern Hemisphere, and will

gain in force during Autumn and Winter when also solar radiation is reduced.

Controlled experiments are needed to understand the interaction between pollen and viral bio-aerosol,

and whether immuno-activation by pollen is indeed a causal factor in reducing the spread of flu-like

viruses and if confirmed, how that works. Understanding this might lead to new therapeutic

approaches.

Acknowledgements: thanks to Sowjanya Putrevu, data scientist at Icecat, for her voluntary support

with executing statistical tests.

Literature

Abhimanyu O.P. & Coussens A.K. (2017). The role of UV radiation and vitamin D in the seasonality

and outcomes of infectious disease. Photochemical & Photobiological Studies, issue 3.

Ahmadi M., Sharifi A.,Dorosti S., Ghoushchi S.J., &Ghanbari N. Investigation of effective

climatology parameters on COVID-19 outbreak in Iran. Science of The Total Environment

Volume 729, 10 August 2020, 138705. https://doi.org/10.1016/j.scitotenv.2020.138705

Bauchau V. & Durham S.R. (2004). Prevalence and rate of diagnosis of allergic rhinitis in Europe.

European Respiratory Journal, 24: 758-764; DOI: 10.1183/09031936.04.00013904

Brandelius A., Andersson M. & Uller L. (2020). Topical dsRNA challenges may induce

overexpression of airway antiviral cytokines in symptomatic allergic disease. A pilot in vivo study in

nasal airways. Respiratory Medicine, vol 108(12) 1816-1819

https://doi.org/10.1016/j.rmed.2014.10.010

Chen X., Dai G.H., Ren Z.M., Tong Y.L., Yang F., & Zhu Y.Q. (2016). Identification of Dietetically

Absorbed Rapeseed (Brassica campestris L.) Bee Pollen MicroRNAs in Serum of Mice. Hindawi

Publishing Corporation, BioMed Research International, Article ID 5413849.

http://dx.doi.org/10.1155/2016/5413849

Chong K.C., Liang J., Jia K.M., Kobayashi N., Wang M.W., Wei L., Lau S.Y.F., & Sumi A. Latitudes

mediate the association between influenza activity and meteorological factors: A nationwide

modelling analysis in 45 Japanese prefectures from 2000 to 2018. Science of the Total Environment.

Vol 703, February 2020. https://doi.org/10.1016/j.scitotenv.2019.134727

Coccia M. (2020). Diffusion of COVID-19 outbreaks: the interaction between air pollution-to-human

and human-to-human transmission dynamics in hinterland regions with cold weather and low average

wind speed. Working Paper CocciaLab n. 48/2020, CNR - National Research Council of Italy.

De Weger L.A. et al. (2013). Impact of Pollen. In: Sofiev M., Bergmann KC. (eds) Allergenic Pollen.

Springer, Dordrecht.

Fine P., Eames K. & Heymann D.L., "Herd Immunity": A Rough Guide, Clinical Infectious Diseases,

Vol 52(7), April 2011, 911–916, https://doi.org/10.1093/cid/cir007.

Fox S.J., Miller J.C., & Meyers L.A. (2017). Seasonality in risk of pandemic influenza emergence.

PLoS Comput Biol. Oct; 13(10): e1005749. doi: 10.1371/journal.pcbi.1005749

Ghanem K.Z., Ramadan M.M., Ghanem H.Z., Fadel M. Improving the production of unsaturated fatty

acid esters and flavonoids from date palm pollen and their effects as anti-breast-cancer and antiviral

agents: An in-vitro study. J Arab Soc Med Res 2015;10:47-55.

Gozzi N., Perrotta D., Paolotti D., and Perra N. (2020). Towards a data-driven characterization of

behavioral changes induced by the seasonal flu. Cornell University. arXiv:2002.00671 [physics.soc-

ph].

Guarnieri, M, Speranza, A., Nepi, M. et al. (2006). Ripe pollen carbohydrate changes in Trachycarpus

fortunei: the effect of relative humidity. Sex Plant Reprod 19, 117. https://doi.org/10.1007/s00497-

006-0027-3

Frei, T. & Gassner, E. Climate change and its impact on birch pollen quantities and the start of the

pollen season an example from Switzerland for the period 1969–2006. Int J Biometeorol 52, 667

(2008). https://doi.org/10.1007/s00484-008-0159-2

Gilles, S, Blume, C, Wimmer, M, et al. Pollen exposure weakens innate defense against respiratory

viruses. Allergy. 2019; 75: 576– 587. https://doi.org/10.1111/all.14047

Hoogeveen M.J. (2020. Pollen likely seasonal factor in inhibiting flu-like epidemics. A Dutch study

into the inverse relation between pollen counts, hay fever and flu-like incidence 2016–2019, Science

of The Total Environment, Vol. 727, 2020, https://doi.org/10.1016/j.scitotenv.2020.138543.

Howarth, P.H. (2000). Allergic rhinitis: not purely a histamine-related disease. European Journal of

Allergy and Clinical Immunology, Vol 55(s64), 7-16.

Klemens, C., Rasp, G., Jund, F., Hilgert, E., Devens, C., Pfrogner, E., Kramer, M.F. Mediators and

cytokines in allergic and viral-triggered rhinitis. Allergy & Asthma Proceedings . Jul/Aug 2007, Vol

28(4), 434-441.

Kissler S.M., Tedijanto C., Goldstein E., Grad Y.H. & Lipsitch M. (2020). Projecting the transmission

dynamics of SARS-CoV-2 through the post-pandemic period. Science 10.1126/science.abb5793.

Liu Y., Gayle A.A., Wilder-Smith A., Rocklöv J. The reproductive number of COVID-19 is higher

compared to SARS coronavirus, Journal of Travel Medicine, Vol 27(2), March 2020,

https://doi.org/10.1093/jtm/taaa021.

Ma Y., Zhao Y., Liu J., He X., Wang B., Fu S., Yan J., Niu J., Zhou J. & Luo B. Effects of

temperature variation and humidity on the death of COVID-19 in Wuhan, China. Science of The Total

Environment Volume 724, 1 July 2020, 138226. https://doi.org/10.1016/j.scitotenv.2020.138226

Moriyama M., Hugentobler W.J. & Iwasaki A.. Seasonality of Respiratory Viral Infections. Annual

Review of Virology 2020 7:1. https://doi.org/10.1146/annurev-virology-012420-022445.

Nivel.nl (2020). Nivel zorgregistraties eerste lijn – Surveillance – wekelijks bulletin over symptomen

en aandoeningen op basis van gegevens van huisartsen, version June 17, 2020.

https://www.nivel.nl/sites/default/files/algemene-content/NivelSurveillance_2020_wk24b.pdf

Nosal C., Ehlers A. & Haspel J.A.. Why Lungs Keep Time: Circadian Rhythms and Lung Immunity.

Annual Review of Physiology 2020 82:1, 391-412. https://doi.org/10.1146/annurev-physiol-021119-

034602.

Park, S., & Mattson, R. H. (2008). Effects of Flowering and Foliage Plants in Hospital Rooms on

Patients Recovering from Abdominal Surgery, HortTechnology hortte, 18(4), 563-568. Retrieved Jun

20, 2020, from https://journals.ashs.org/horttech/view/journals/horttech/18/4/article-p563.xml

Plans-Rubió P.. The vaccination coverage required to establish herd immunity against influenza

viruses. Preventive Medicine, Vol 55(1), July 2012, 72-77.

https://doi.org/10.1016/j.ypmed.2012.02.015.

Postnikov, E. (2016). Dynamical prediction of flu seasonality driven by ambient temperature:

influenza vs. common cold. Eur. Phys. J. B 89, 13. https://doi.org/10.1140/epjb/e2015-50845-7

Rosenwasser, L. J. (2011). Current Understanding of the Pathophysiology of Allergic Rhinitis.

Immunology and Allergy Clinics, Vol 31(3), p 433 – 439.

Qi H., Xiao S., Shi R., Ward M.P., Chen Y., Tu W., Su Q., Wang W., Wang X. & Zhang Z. COVID-

19 transmission in Mainland China is associated with temperature and humidity: A time-series

analysis. Science of The Total Environment Volume 728, 1 August 2020, 138778.

https://doi.org/10.1016/j.scitotenv.2020.138778

Schuit M., Gardner S., Wood S., Bower K., Williams G., Freeburger D., Dabisch P. The Influence of

Simulated Sunlight on the Inactivation of Influenza Virus in Aerosols, The Journal of Infectious

Diseases, Vol 221(3), February 2020, p 372–378, https://doi.org/10.1093/infdis/jiz582.

Shaman J., Goldstein E. & Lipsitch M. (2011) Absolute Humidity and Pandemic Versus Epidemic

Influenza, American Journal of Epidemiology, Vol 173(2), January 2011, p 127–135,

https://doi.org/10.1093/aje/kwq347

Soebiyanto R.P., Clara W., Jara J., Castillo L., Sorto O.R., Marinero S., Barnett de Antinori M.E.,

McCracken J.P., Widdowson M.A., 9 Azziz-Baumgartner E., & Kiang R.K. (2014). The Role of

Temperature and Humidity on Seasonal Influenza in Tropical Areas: Guatemala, El Salvador and

Panama, 2008–2013. PLoS One 9(6): e100659. Published online 2014 Jun 23. doi:

10.1371/journal.pone.0100659

Tamerius, J., Nelson, M. I., Zhou, S. Z., Viboud, C., Miller, M. A., & Alonso, W. J. (2011). Global

influenza seasonality: reconciling patterns across temperate and tropical regions. Environmental health

perspectives, 119(4), 439–445. https://doi.org/10.1289/ehp.1002383.

Tan D.X. & Ruegiger Hardeland R. (2020). Potential utility of melatonin in deadly infectious diseases

related to the overreaction of innate immune response and destructive inflammation: focus on COVID-

19. Melatonin in deadly virus infection. Melatonin Research Vol 3(1). DOI:

https://doi.org/10.32794/mr11250052

Tosepu R., Gunawan J., Effendy D.S. Ahmad L.O.A.I., Lestari H., Bahar H. & Asfiang P. Correlation

between weather and Covid-19 pandemic in Jakarta, Indonesia. Science of The Total Environment,

Volume 725, 10 July 2020, 138436. https://doi.org/10.1016/j.scitotenv.2020.138436

Van De Ven M.O.M., Van Den Eijnden R.J.J.M., Engels R.C.M.E. Atopic diseases and related risk

factors among Dutch adolescents, *European Journal of Public Health*, Vol 16(5), October 2006, p

549–558, https://doi.org/10.1093/eurpub/ckl022.

Wachsman M.B., Lopez E.M.F., Ramirez J.A., Galagovksy L.R., & Coto C.E. (2000). Antiviral effect

of brassinosteroids against herpes virus and arenaviruses. Antiviral Chemistry & Chemotherapy

11:71–77.

Wan, Y. et al. (2020). J. Virol. Receptor Recognition by the Novel Coronavirus from Wuhan: an

Analysis Based on Decade-Long Structural Studies of SARS Coronavirus. [Epub ahead of print].

https://jvi.asm.org/content/94/7/e00127-20.long

Xie J. & Zhu Y. Association between ambient temperature and COVID-19 infection in 122 cities from

China. Science of The Total Environment Volume 724, 1 July 2020, 138201.

https://doi.org/10.1016/j.scitotenv.2020.138201

Yao Y., Pan J.H., Liu Z.X., Meng X., Wang W.D., Kan H.D. & Wang W.B. (2020). No Association of

COVID-19 transmission with temperature or UV radiation in Chinese cities. European Respiratory

Journal 2020; DOI: 10.1183/13993003.00517-2020

Zhang, J-J, Dong, X, Cao, Y-Y, et al. Clinical characteristics of 140 patients infected with SARS-

CoV-2 in Wuhan, China. Allergy. 2020; 00: 1– 12. https://doi.org/10.1111/all.14238