1 *UV-C irradiation is highly effective in inactivating SARS-CoV-2 replication*
2

Mara Biasin[1], Andrea Bianco[2], Giovanni Pareschi[2], Adalberto Cavalleri[3], Claudia Cavatorta[3], Claudio

Fenizia[1,6], Paola Galli[2], Luigi Lessio[4], Manuela Lualdi[5], Enrico Tombetti[1], Alessandro Ambrosi[6], Edoardo Maria

Alberto Redaelli[2], Irma Saulle[1,7], Daria Trabattoni[1], Alessio Zanutta[2], Mario Clerici[7,8,*]

[1] *Department of Biomedical and Clinical Sciences L. Sacco, University of Milano, Milano, Italy.*
[2]*Italian National Institute for Astrophysics (INAF) – Brera Astronomical Observatory, Merate, Italy.*
[3] *Epidemiology and Prevention Unit, IRCCS Foundation, Istituto Nazionale dei Tumori, Milan, Italy.*
[4] *Italian National Institute for Astrophysics (INAF) – Padova Astronomical Observatory, Padova, Italy.*
[5] *Department of Imaging Diagnostic and Radioterapy, IRCCS Foundation, Istituto Nazionale dei Tumori,*
Milan, Italy.
[6] *University Life and health San Raffaele, Milan, Italy.*
[7] *Department of Pathophysiology and Transplantation, University of Milano, Milano, Italy.*
[8]*Don C. Gnocchi Foundation, IRCCS Foundation, Milano, Italy.*
16

The potential virucidal effects of UV-C irradiation on SARS-CoV-2 were experimentally evaluated for

different illumination doses and virus concentrations (1000, 5, 0.05 MOI). At a virus density comparable to

that observed in SARS-CoV-2 infection, an UV-C dose of just 3.7 mJ/cm^2 was sufficient to achieve a more

than 3-log inactivation without any sign of viral replication. Moreover, a complete inactivation at all viral

concentrations was observed with 16.9 mJ/cm^2. These results could explain the epidemiological trends of

COVID-19 and are important for the development of novel sterilizing methods to contain SARS-CoV-2

infection.

Introduction

The COVID-19 pandemic caused by SARS-CoV-2 virus[1] has had an enormous, as yet barely understood,

impact on health and economic outlook at the global level[2]. The identification of effective microbicide

approaches is of paramount importance in order to limit further viral spread, as the virus can be

transmitted via aerosol[3,4] and can survive for hours outside the body[5–7]. Non-contact disinfection

technologies are highly desirable, and UV radiation, in particular UV-C (200 – 280 nm) has been suggested

to be able to inactivate different viruses, including SARS-CoV[8–12]. The interaction of UV-C radiations with

viruses has been extensively studied[13–15], and direct absorption of the UV-C photon by the nucleic acid basis

and/or capsid proteins leading to the generation of photoproducts that inactivate the virus was suggested

* e-mail: mario.clerici@unimi.it

to be one of the main UV-C-associated virucidal mechanisms[16,17]. Some models have been proposed to

correlate the nucleic acid structure with the required dose to inactivate the virus, but a reliable model is

still unavailable[18]. This is also due to the fact that UV-C measurements were conducted using different

viruses and diverse experimental conditions[19–22]. This led to an extremely wide range of values for the same

virus and, e.g., in the case of SARS-CoV-1 values reported in the literature range from a few mJ/cm^2 to

hundreds mJ/cm$^{2\ 19,22,23}$. Likewise, recent papers reported values for UVC inactivation ranging from 3 to

1000 mJ/cm$^{2\ 24–27}$. A better understanding of the effects of UV-C on SARS-CoV-2 , which take into account

all the key factors involved in the experimental setting (including culture medium, SARS-CoV-2

concentration, UV-C irradiance, time of exposure, and UV-C absorbance) will allow to replicate the results

in other laboratory with different devices. Moreover, as recent evidences suggest that UV light from

sunlight are efficient in inactivating the virus[28], these measurements will be relevant for the setting up of

further experiment considering the role of UV-A and UV-B on SARS-CoV-2 replication.

Results and discussion

Herein, we report the effect of monochromatic UV-C (254 nm) on SARS-CoV-2, showing that virus

inactivation can be easily achieved. Experiments were conducted using a custom-designed low-pressure

mercury lamp system, which has been spectral-calibrated providing an average intensity of 1.082 mW/cm^2

over the illumination area (see the details reported in the Method section). Three different illumination

exposure times, corresponding to 3.7, 16.9 and 84.4 mJ/cm^2, were administered to SARS-CoV-2 either at a

multiplicity of infection (MOI) of 0.05, 5, 1000. The first concentration is equivalent to the low-level

contamination observed in closed environments (e.g. hospital rooms), the second one corresponds to the

average concentration found in the sputum of COVID-19 infected patients, and the third one is a very large

concentration, corresponding to that observed in terminally diseased COVID-19 patients[29] . After UV-C

exposure, viral replication was assessed by culture-polymerase chain reaction (C-RT-PCR) targeting two

regions (N1 and N2) of the SARS-CoV-2 nucleocapsid gene, as well as by analyzing SARS-CoV-2-induced

cytopathic effect. Analyses were performed in the culture supernatant of infected cells at three different

time points (24, 48 and 72 hours for SARS-CoV-2 at MOI 1000 and 5; 24, 48 hours and 6 days for SARS-CoV-

2 at MOI 0.05), as well as on cell lysates at the end of cellular culture (72 hours: MOI 1000 and 5; 6 days:

MOI 0.05). This approach allows to follow the kinetic of viral growth and to verify whether the used dose is

sufficient to completely inactivate the virus over time. This is useful from a practical point of view, when

UV-C devices are used to disinfect surfaces and the environment.

The effect of the UV-C exposure on SARS-CoV-2 replication was extremely evident and independent from

the MOI employed; dose-response and time-dependent curves were observed. Figure 1, 2 and 3 report for

different MOI the number of SARS-CoV-2 copies for the three concentrations as a function of the UV-C dose

and time, quantified on a standard curve from a plasmid control. The corresponding normalised curves of

the virus copies are reported in the same figures.

70
Figure 1. Viral replication of UV-irradiated SARS-CoV-2 (0.05 MOI) virus in in vitro VeroE6 cells. Vero E6 cells were infected with

UV-C irradiated SARS-CoV-2 virus at a MOI of 0.05. Culture supernatants were harvested at the indicated times (24, 48 hours and 6

days) and virus titers were measured (Panel A, B) by absolute copy number quantification (Real-Time PCR). Viral replication was

assessed even on cell lysate harvested at the end of cell cultures (6 days) (Panel C). All cell culture conditions were seeded in

duplicate. Panel D reports the plots of the measured virus copies normalized at the untreated sample in the different conditions.

For descriptive purposes, mean values and whiskers representing the observed half-ranges are shown.

Viral replication was not observed at the lowest viral concentration (0.05 MOI) in either untreated or in UV-

C-irradiated samples in the initial 48 hours (Figure 1). However, 6 days after infection, viral replication was

distinctly evident in the UV-C unexposed condition, but was completely absent following UV-C irradiation

even at 3.7 mJ/cm^2 both in cell culture supernatants (Figure1, panel A and B) and in cell lysate (Figure 1,

panel C). A two-way ANOVA analysing the effect of UV-C dose and time of incubation failed to identify a

significant effect of the UV exposure on viral replication. This is due to the fact that at very low MOI

relevant increases in N1 and N2 copy numbers were detectable only in a single condition -at six days in the

absence of UV-C exposure- thus hampering the statistical power of the analysis.

87
Figure 2. Viral replication of UV-irradiated SARS-CoV-2 (5 MOI) virus in in vitro VeroE6 cells. Vero E6 cells were infected with UV-C

irradiated SARS-CoV-2 virus at a MOI of 5. Culture supernatants were harvested at the indicated times (24, 48 and 72 hours) and

virus titers were measured by absolute copy number quantification (Real-Time PCR, A and B). Viral replication was assessed even

on cell lysate harvested at the end of cell cultures (72 hours) (C). All cell culture conditions were seeded in duplicate. Panel D

reports the plots of the measured virus copies normalized at the untreated sample in the different conditions. For descriptive

purposes, mean values and whiskers representing the observed half-ranges are shown.

94
At the intermediate viral concentration (5 MOI), a significant reduction of copy number starting from the

3.7 mJ/cm^2 dose with a decrease of a factor of 2000 (> 3-log decrease) after 24 hours was observed (Figure

2, panel D). A two-way ANOVA confirmed that this UV-C dose significantly dampened viral replication

(p=0.000796, and P=0.000713 for N1 and N2 copies respectively). Even more important, the copy number

did not increase over time, suggesting an effective inactivation of the virus, which was further confirmed by

cytopathic effect assessment (Figure 3, panel A, B and C).

101

102 **Figure 3. Analyses of virus induced cytopathic effect**. (A) No cytopathic effect was observed in uninfected cultured VeroE6

103 monolayers maintained in 50mJ/cm^2 UV-treated complete medium for 72 hours. (B) In vitro infection of SARS-CoV-2 (5 MOI) UV-C

104 untreated VeroE6 cells resulted in an evident cytopathic effect. (C) SARS-CoV-2 irradiation with 3.7 mJ/cm^2 UV-C rescued the

105 cytopathic effect induced by UV-C untreated virus.

108
Figure 4. Viral replication of UV-irradiated SARS-CoV-2 (1000 MOI) virus in in vitro VeroE6 cells. Vero E6 cells were infected with

UV-C irradiated SARS-CoV-2 virus at a MOI of 1000. Culture supernatants were harvested at the indicated times (24, 48 and 72

hours) and virus titers were measured (Panel A, B) by absolute copy number quantification (Real-Time PCR). Viral replication was

assessed even on cell lysate harvested at the end of cell cultures (72 hours) (Panel C). All cell culture conditions were seeded in

duplicate. Panel D reports the plots of the measured virus copies normalized at the untreated sample in the different conditions.

For descriptive purposes, mean values and whiskers representing the observed half-ranges are shown.

115
Using a high viral input (MOI=1000), the two-way ANOVA confirmed that all the tree UV doses analysed

resulted in a significant suppression of viral replication for both N1 (3.7 mJ/cm^2: p= 0.008455; 16.9 mJ/cm^2:

p=0.004216; and 84.4 mJ/cm^2: p=0.000202) and N2 copies (3.7 mJ/cm^2: p= 6.43E-05; 16.9 mJ/cm^2:

p=1.68E-05; and 84.4 mJ/cm^2: p=1.68E-05)(Figure 4). Notably, a different course of infection was observed,

in which the inhibitory effect was not accompanied by viral suppression for the UV-C dose of 3.7 mJ/cm^2

(Figure 4, panel A, B and D). Indeed, a relevant reduction in N1 e N2 copy numbers was observed in a UV-C

dose-dependent manner as early as 24 hours (by a factor of 10^3 at 3.7 mJ/cm^2 and 10^4 at 16.9 mJ/cm^2,

Figure 4, panel A, B and D), but longer culture times resulted in an increase in N1 and N2 copy numbers for

the UV-C dose of 3.7 mJ/cm^2. This indicates that the residual viral input left by the 3.7 mJ/cm^2 was able to

replicate and suffucient to generate an effective infection. This is not the case in cultures exposed to

higher UV-C doses, as no replication could be detected in these conditions. All the results were further

confirmed by 2-ANOVA statistical analyses performed on viral replication at intracellular level (3.7 mJ/cm^2

vs. Untreated: N1, p=0.008455; N2: p=6.43E-05; 16.9 mJ/cm^2 vs. Untreated: N1, p=0.004216; N2: p=1.68E-

05; 84.4 mJ/cm^2 vs. Untreated: N1, p=0.004216; N2: p=1.68E-05) (Figure 1, panel C; Figure 2, panel C;

Figure 4, panel C).

We compared our results with data available in the literature and observed that our inactivating dose is

much smaller than that reported in Heilingloh et al.[26] (1000 mJ/cm^2 for the complete inactivation). This

discrepancy is likely to be the consequence of the UV-C absorption by the medium used in Heilingloh et al,

which has a 4-fold higher thickness compared to the one used in our experiments. This possibility is

supported by the observation that 200 mJ/cm^2 of UV-A, which is not absorbed by the medium, was

sufficient to reduce viral replication of 1-log. As UV-A light is significantly less efficient (order of

magnitudes) than UV-C, the reported UV-C inactivating dose (100mJ/cm2) seems to be questionable.

Two other papers measured the effect of UV-C light on SARS-CoV-2. In Ruetalo et al.[25], the illumination of

254 nm light was employed on a dried sample of SARS-CoV-2. Complete inactivation was obtained with 20

mJ/cm^2, a value greater than ours, but in the same range. It has to be underlined that in the dried film a

shielding effect by the organic component present in the liquid can occur, reducing the efficiency of the UV-

C light. This was shown by Ratnesar-Shumate et al.[28] , who demonstrated that the dose required to obtain a

similar degree of viral inactivation was twice in dried samples from gMEM compared to the ones

resuspended in simulated saliva. Notably, the two mediums differ for their composition, mainly in terms of

protein and solid percentage, with higher values for the gMEM.

Inagaki et al. used a 285 nm UV LED and showed that a dose of about 38 mJ/cm^2 was sufficient to

completely inactivate SARS-CoV-2. This dose is greater compared to the one we established ; this

discrepancy can be explained by the observation that the 285 nm is less efficient than the 254 nm

wavelength[24]. Finally, in an elegant study Storm et al.[27] compared the virucidal effect of UV-C in wet and

dry systems. Results were based on the use of a very small volume of viral stock in DMEM (5 μl) and

showed that a dose of 3.4 mJ/cm^2 inactivated wet samples, whereas a dose that twice as high was needed

in dried samples. These results are comparable to the ones herein, and the shielding effect in dried samples

is almost evident. Such comparisons show how the experimental conditions adopted significantly impact on

the definition of the dose of UV-C resulting in virus inactivation. It is therefore crucial to accurately describe

all the details of the experiments to perform a reliable comparison.

In conclusion, we report the results of a highly controlled experimental model that allowed us to identify

the UV-C radiation dose sufficient to inactivate SARS-CoV-2. The response depends on both the UV-C dose

and the virus concentration. Indeed, for virus concentrations typical of low-level contaminated closed

environment and sputum of COVID-19 infected patients, a very small dose of less than 4 mJ/cm^2 was

enough to achieve full inactivation of the virus. Even at the highest viral input concentration (1000 MOI),

viral replication was totally inactivated with a dose $\geq$16.9 mJ/cm^2. These results show how the SARS-CoV-2

is extremely sensitive to UV-C light and they are important to allow the proper design and development of

efficient UV based disinfection methods to contain SARS-CoV-2 infection.

Methods

In vitro **SARS-CoV-2 infection assay**

3 x 10^5 VeroE6 cells were cultured in DMEM (ECB7501L, Euroclone, Milan, Italy) with 2 % FBS medium, with

100 U/ml penicillin and 100 μg/ml streptomycin, in a 24-well plate one day before viral infection assay.

SARS-CoV-2 (Virus Human 2019-nCoV strain 2019- nCoV/Italy-INMI1, Rome, Italy) at a multiplicity of

infection (MOI) of 1000, 5 and 0.05 were treated with different doses of UV-C radiation (see the dedicated

section) before inoculum into VeroE6 cells. UV-C-untreated virus served as positive controls. Cell cultures

were incubated with the virus inoculum in duplicate for three hours at 37°C and 5% CO_2. Then, cells were

rinsed three times with warm PBS, replenished with the appropriate growth medium and observed daily for

cytopathic effect. Viral replication in culture supernatants was assessed by an Integrated Culture-

polymerase chain reaction (C-RT-PCR) method[30] at 24, 48, and 72 hours post-infection (hpi) while infected

cells were harvested for RNA collection at 72 hpi. Cell cultures from SARS-CoV-2 at 0.05 MOI were

harvested 6 days post infection. RNA was extracted from VeroE6 cell culture supernatant and cell lysate by

the Maxwell® RSC Instrument with Maxwell® RSC Viral Total Nucleic Acid Purification Kit (Promega,

Fitchburg, WI, USA), quantified by the Nanodrop 2000 Instrument (Thermo Scientific) and purified from

genomic DNA with RNase-free DNase (RQ1 DNase; Promega). One microgram of RNA was reverse

transcribed into first-strand cDNA in a 20-μl final volume as previously described[31,32].

Real-time PCR was performed on a CFX96 (Bio-Rad, CA, USA) using the 2019-nCoV CDC qPCR Probe Assay

emergency kit (IDT, Iowa, USA), which targets two regions (N1 and N2) of the nucleocapsid gene of SARS-

CoV-2. Reactions were performed according to the following thermal profile: initial denaturation (95°C, 10

min) followed by 45 cycles of 15 s at 95°C (denaturation) and 1 min at 60°C (annealing-extension).

Viral copy quantification was assessed by creating a standard curve from the quantified 2019-nCoV_N

positive Plasmid Control (IDT, Iowa, USA).

UV illumination test

The illumination of the virus solution was conducted using a low-pressure mercury lamp mounted in a

custom designed holder, which consist in a box with a circular aperture 50 mm in diameter placed at

approximately 220 mm from the source. The aperture works as a spatial filter to make the illumination of

the area behind more uniform. A mechanical shutter is also present to start the illumination process. The

plate is placed 30 mm below the circular aperture and a single dwell (34.7 mm in diameter), centered in

respect to the 50 mm aperture, has been irradiated from the top. The dwell was filled with 0.976 ml of the

virus suspended in Dulbecco's Modified Eagle's Medium (DMEM) in order to have a 1 mm thick liquid layer.

After the irradiation, the sample was treated as described in the previous section.

The intensity of the lamp and its spectral properties have been measured using an Ocean Optics HR2000+

spectrometer (Ocean Optics Inc., Dunedin, USA). The HR2000+ spectrometer was calibrated against a

reference deuterium–halogen source (Ocean Optics Inc. Winter Park, Winter Park, Florida) and in

compliance with National Institute of Standards and Technology (NIST) practices recommended in NIST

Handbook 150-2E, Technical guide for Optical Radiation Measurements. The last calibration was performed

in March 2019. The detector of our spectrometer is a high-sensitivity 2048-element Charge-Coupled Device

(CCD) array from Sony. The spectral range is 200–1100 nm with a 25 µm wide entrance slit and an optical

resolution of 1.4 nm (FWHM). The cosine-corrected irradiance probe, model CC-3-UV-T, is attached to the

tip of a 1 m long optical fibre and couples to the spectrometer. The intensity of the lamp has been

measured by positioning the spectrometer in five positions: in the center and at the ends of a 20 mm cross

arm after a warming up time of 30 s. The spectra in the five positions are reported in Figure 5 (Panel A)

together with a scheme of the dwell and illuminated area.

Figure 5. Mercury lamp spectrum measured in the five positions (Panel A). Inset: scheme of the illuminated dwell and the measuring

position. UV-vis transmission spectrum of the Dulbecco's Modified Eagle's Medium (DMEM) in a 1 mm quartz cuvette (Panel B).

As expected, the emission is dominated by the UV-C line (Figure 5, Panel A) and its intensity was uniform in

the area with an average value of 1.082 mW/cm^2. The stability of the lamp was evaluated in +/- 11E-3

mW/cm^2 during a 130 s measurement. According to this value, three exposure times were set: 5, 23 and

114 s (with an accuracy of 0.2s), which correspond to following doses: 5.4, 25.0, 123.4 mJ/cm^2. This is the

nominal UV doses provided to the dwell, but we were interested in the effective doses (D_e) reaching the

virus. It was necessary to calculate the effective irradiance (I_e). This step was performed considering both

the reflection losses at the air/water interface (R_w) and the Transmittance (T_s) of the DMEM solution at 254

nm (from the spectrum in figure 5, Panel B, considering the cuvette losses, T_s = 0 .70). It is important to

notice that the spectrum was measured in a quartz cuvette (1 mm thick) by means of a Jasco V770

spectrophotometer and this thickness was the same of the solution in the dwell during the UV irradiation

step.

The reflection loss was computed as follow:

$$R_w = \frac{(n_w-1)^2}{(n_w+1)^2} \tag{1}$$

Where n_w = 1.375 is the refractive index of water at 254 nm. Then, I_e was calculated:

$$I_e = I_n(1 - R_w) \times T_s \tag{2}$$

The final transmission of the DMEM solution was equal to 0.68 and the corresponding effective doses were

derived simply multiplying I_e by the exposure time.

According to this value, the effective doses provided to the viruses were: 3.7±0.15, 16.9±0.2 and 84.4±0.9

mJ/cm^2. We have to notice that we are neglecting here the absorption of the virus at this wavelength and

the possible scattering. Such approximations are valid considering the relative low concentration of the

virus and small thickness of the layer (the solution appeared fully transparent).

Statistical analyses

To assess the effect of the different UV-C doses on N1 and N2 copy numbers, two-way ANOVAs were

performed. For the analysis of intracellular N1 and N2 doses in the supernatant, UV-C dose and MOI

represented the dependent variables, while for the analysis of N1 and N2 in the supernatant, different

analyses were performed for individual MOI, using UV-C dose and time as dependent variables.

References

1. Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N. Engl. J. Med.* **382**, 727–733 (2020).

2. Cobey, S. Modeling infectious disease dynamics. *Science* **368**, 713–714 (2020).

3. Jarvis, M. C. Aerosol Transmission of SARS-CoV-2: Physical Principles and Implications. *Front. Public Heal.* **8**, 590041 (2020).

4. Tang, S. *et al.* Aerosol transmission of SARS-CoV-2? Evidence, prevention and control. *Environ. Int.* **144**, 106039 (2020).

5. van Doremalen, N. *et al.* Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *N. Engl. J. Med.* **382**, 1564–1567 (2020).

6. Chin, A. W. H. *et al.* Stability of SARS-CoV-2 in different environmental conditions. *The Lancet Microbe* **1**, e10 (2020).

7. Aboubakr, H. A., Sharafeldin, T. A. & Goyal, S. M. Stability of SARS-CoV-2 and other coronaviruses in the environment and on common touch surfaces and the influence of climatic conditions: A review. *Transbound. Emerg. Dis.* 10.1111/tbed.13707 (2020) doi:10.1111/tbed.13707.

8. Pirnie, M., Linden, K. G. & Malley, J. P. J. Ultraviolet disinfection guidance manual for the final long term 2 enhanced surface water treatment rule. *Environ. Prot.* **2**, 1–436 (2006).

9. Reed, N. G. The history of ultraviolet germicidal irradiation for air disinfection. *Public Health Rep.* **125**, 15–27 (2010).

10. Kovalski, W. *Ultraviolet germicidal irradiation handbook: UVGI for air and surface disinfection*. (Springer Science & Business Media, 2010).

11. Darnell, M. E. R., Subbarao, K., Feinstone, S. M. & Taylor, D. R. Inactivation of the coronavirus that induces severe acute respiratory syndrome, SARS-CoV. *J. Virol. Methods* **121**, 85–91 (2004).

12. Raeiszadeh, M. & Adeli, B. A Critical Review on Ultraviolet Disinfection Systems against COVID-19 Outbreak: Applicability, Validation, and Safety Considerations. *ACS Photonics* **7**, 2941–2951 (2020).

13. Bosshard, F., Armand, F., Hamelin, R. & Kohn, T. Mechanisms of Human Adenovirus Inactivation by Sunlight and UVC Light as Examined by Quantitative PCR and Quantitative Proteomics. *Appl. Environ. Microbiol.* **79**, 1325 – 1332 (2013).

14. Nishisaka-Nonaka, R. *et al.* Irradiation by ultraviolet light-emitting diodes inactivates influenza a viruses by inhibiting replication and transcription of viral RNA in host cells. *J. Photochem. Photobiol. B Biol.* **189**, 193–200 (2018).

15. Araud, E., Fuzawa, M., Shisler, J. L., Li, J. & Nguyen, T. H. UV Inactivation of Rotavirus and Tulane Virus Targets Different Components of the Virions. *Appl. Environ. Microbiol.* **86**, e02436-19 (2020).

16. Qiao, Z. & Wigginton, K. R. Direct and Indirect Photochemical Reactions in Viral RNA Measured with RT-qPCR and Mass Spectrometry. *Environ. Sci. Technol.* **50**, 13371–13379 (2016).

17. Wigginton, K. R. & Kohn, T. Virus disinfection mechanisms: the role of virus composition, structure, and function. *Curr. Opin. Virol.* **2**, 84–89 (2012).

18. Lytle, C. D. & Sagripanti, J.-L. Predicted Inactivation of Viruses of Relevance to Biodefense by Solar Radiation. *J. Virol.* **79**, 14244 LP – 14252 (2005).

19. Walker, C. M. & Ko, G. Effect of Ultraviolet Germicidal Irradiation on Viral Aerosols. *Environ. Sci.*
Technol. **41**, 5460–5465 (2007).

20. McDevitt, J. J., Rudnick, S. N. & Radonovich, L. J. Aerosol susceptibility of influenza virus to UV-C
light. *Appl. Environ. Microbiol.* **78**, 1666–1669 (2012).

21. Calgua, B. *et al.* UVC Inactivation of dsDNA and ssRNA Viruses in Water: UV Fluences and a qPCR-
Based Approach to Evaluate Decay on Viral Infectivity. *Food Environ. Virol.* **6**, 260–268 (2014).

22. Eickmann, M. *et al.* Inactivation of three emerging viruses – severe acute respiratory syndrome
coronavirus, Crimean–Congo haemorrhagic fever virus and Nipah virus – in platelet concentrates by
ultraviolet C light and in plasma by methylene blue plus visible light. *Vox Sang.* **115**, 146–151 (2020).

23. Duan, S.-M. *et al.* Stability of SARS Coronavirus in Human Specimens and Environment and Its
Sensitivity to Heating and UV Irradiation. *Biomed. Environ. Sci.* **16**, 246–255 (2003).

24. Inagaki, H., Saito, A., Sugiyama, H., Okabayashi, T. & Fujimoto, S. Rapid inactivation of SARS-CoV-2
with Deep-UV LED irradiation. *Emerg. Microbes Infect.* **9**, 1744–1747 (2020).

25. Ruetalo, N., Businger, R. & Schindler, M. Rapid and efficient inactivation of surface dried SARS-CoV-2
by UV-C irradiation. *bioRxiv* 2020.09.22.308098 (2020).

26. Heilingloh, C. S. *et al.* Susceptibility of SARS-CoV-2 to UV irradiation. *Am. J. Infect. Control* **48**, 1273–
1275 (2020).

27. Storm, N. *et al.* Rapid and complete inactivation of SARS-CoV-2 by ultraviolet-C irradiation. *Sci. Rep.*
10, 22421 (2020).

28. Ratnesar-Shumate, S. *et al.* Simulated Sunlight Rapidly Inactivates SARS-CoV-2 on Surfaces. *J. Infect.*
Dis. **222**, 214–222 (2020).

29. Wölfel, R. *et al.* Virological assessment of hospitalized patients with COVID-2019. *Nature* **581**, 465–
469 (2020).

30. Reynolds, K. A. Integrated Cell Culture/PCR for Detection of Enteric Viruses in Environmental
Samples BT - Public Health Microbiology: Methods and Protocols. in (eds. Spencer, J. F. T. & Ragout
de Spencer, A. L.) 69–78 (Humana Press, 2004). doi:10.1385/1-59259-766-1:069.

31. Saulle, I. *et al.* Endoplasmic Reticulum Associated Aminopeptidase 2 (ERAP2) Is Released in the
Secretome of Activated MDMs and Reduces in vitro HIV-1 Infection. *Front. Immunol.* **10**, 1648
(2019).

32. Ibba, S. V *et al.* Analysing the role of stat3 in HIV-1 infection. *Journal of Biological Regulators and*
Homeostatic Agents **33** 1635–1640 (2019).

Acknowledgements

This research was partially supported by a grant from Falk Renewables and it has been carried out in the

context of the activities promoted by the Italian Government and in particular, by the Ministries of Health

and of University and Research, against the COVID-19 pandemic. Authors are grateful to INAF's President,

Prof. N. D'Amico, for the support and for a critical reading of the manuscript.

Author Contribution

P.G., L.L. E.R. and A.Z. designed and produced the illumination system; A.C., C.C. and M.L. performed the lamp calibration; A.B. performed the lamp setup and lamp dosimetry, wrote the main manuscript; M.B. performed biological experiments, analyzed the data, wrote the main manuscript; C.F., I.S. designed and performed some biological tests; E.T, A.A. performed the statistical analysis; D.T. discussed the results; G.P. and M.C. supervised the study and review the manuscript.