Estimating individual risks of COVID-19-associated hospitalization and death using publicly available data

Rajiv Bhatia [1] [¶] *, Jeffrey Klausner [2] [¶]

[1]Assistant Professor of Medicine (affiliated), Stanford University, Stanford, California, USA

[2]Professor of Medicine and Public Health, University of California Los Angeles, Los Angeles, California, USA

* Corresponding Author

Email: drajiv@stanford.edu (RB)

[¶] These authors contributed equally to this work

Abstract

We describe a method to estimate individual risks of hospitalization and death

attributable to non-household and household transmission of SARS-CoV-2 using

available public data on confirmed-case incidence data along with estimates of the

clinical fraction, timing of transmission, isolation adherence, secondary infection risks,

contact rates, and case-hospitalization and case-fatality ratios. Using the method, we

estimate that risks for a 90-day period at the median daily summertime U.S. county

confirmed COVID-19 case incidence of 10.8 per 100,000 and pre-pandemic contact

rates range from 0.4 to 8.9 per 100,000 for the four deciles of age between 20 and 60

years. The corresponding 90-day period risk of hospitalization ranges from 13.7 to 69.2

per 100,000. Assuming a non-household secondary infection risk of 4% and pre-

pandemic contact rates, the share of transmissions attributable to household settings

ranges from 73% to 78%. These estimates are sensitive to the parameter assumptions;

nevertheless, they are reasonably comparable to the COVID-19 hospitalization and

fatality rates observed over the time period. We conclude that individual risk of

hospitalization and death from SARS-CoV-2 infection is calculable from publicly

available data sources. Access to publicly reported infection incidence data by setting

and other exposure characteristics along with setting specific estimates of secondary

infection risk would allow for more precise individual risk estimation.

Introduction

Perceptions of personal risk can modify pandemic disease transmission and population health outcomes by modifying behaviors such as mask wearing and by influencing demand for and compliance with government recommendations. A recent rapid review of studies that examined adherence with "quarantine" found adherence varied from 0% up to 92.8% and was influenced by social norms, perceived benefits of quarantine, perceived risk of the disease, and financial and material needs.[1] Public perceptions of risk also may influence how political leaders act. Ideally, public leaders should communicate risk precisely and transparently.

To be useful to individuals, estimates of risk need to be specific to people, times, places, and activities. Numerous reports have described clusters of SARS-CoV-2 infections in diverse settings, yet there has been little attention to estimating and communicating how personal risk varies by place, time, and population. Surveillance measures used to monitor and characterize the COVID-19 pandemic, including laboratory-based "case" counts of infection and mortality rates describe risk in aggregate. In the United States, public data on laboratory-confirmed infections, hospitalizations and deaths have not disaggregated information based on exposure factors even though case-reports include this information. No large-scale U.S. community transmission studies have comparatively evaluated risk by setting.

Recognizing the limitations of available data sources, we estimate and compare the person-level risks of SARS-CoV-2 infection associated hospitalization and death

63   attributable to household and non-household contacts using public confirmed case

64   incidence data and published transmission parameters. We identify knowledge gaps

65   that, if filled, could make individual risk assessment more precise and useful.

67   Materials and Methods

We conceptualize risk as the product of susceptibility, infection incidence, the timing

and efficiency of transmission from an infected individual, the contact setting, the

contact rate, and the severity of infection. Because the true infection incidence rate is

unknown, we assume a stable multiplicative relationship between the incidence rate of

confirmed infections and the incidence of infections ($I \sim I_{Confirmed}$). We consider the

transmission potential of clinical and subclinical infection separately, assuming that all

symptomatic infected individuals will contribute to transmission during a pre-

symptomatic infectious period and a share of symptomatic individuals will voluntarily

self-isolate, effectively limiting transmission to within their own households. We assume

those with subclinical infection will contribute to transmission for the duration of their

infectious period but have a lower likelihood of forward transmission than those with

symptoms.

Equation (1) estimates the age-specific probability of a confirmed infection attributable

to contact within household settings, *P (Infection | HH)$_i$*, for age-strata *i*, as a function of

the proportion susceptible to infection, *S*, the confirmed case incidence rate, *I*, the

household specific secondary infection risk, B_{HH} , an age-specific household contact

rate, $C_{HH\,i}$, the fraction of symptomatic infections, F_{SX} , and the complementary fraction

of asymptomatic infections, *1- F_{SX}* , multiplied by an estimate of their relative

infectiousness, r.

(1) P (Infection | HH)$_i$ = S I C$_{HHi}$ B$_{HH}$ { F$_{SX}$ + r (1- F$_{SX}$) }

Equation (2) estimates the age-specific probability of a confirmed infection attributable

to contact in non-household settings, *P (Infection | HH)$_i$*, for age-strata *i*. The share of

pre-symptomatic transmission of the clinical fraction is denoted by p. The fraction of

symptomatic infected cases that do not self-isolate, *1-q*, also contribute to transmission

to non-household contacts during the period of post-symptomatic transmission.

(2) P (Infection | NH)$_i$ = S I C$_{NHi}$ B$_{NH}$ { p (F$_{SX}$) +(1-q)(1-p) (F$_{SX}$) + r (1- F$_{SX}$) }

We estimate the age-specific probabilities of hospitalization and death in equations (3)

and (4) from the age specific probabilities of confirmed infection, multiplying these

quantities by the age-specific confirmed case hospitalization and fatality ratios (CFRs),

respectively.

(3) P (Hospitalization)$_i$ = CHR$_i$ × { P (I|HH)$_i$ + P (I|NH)$_i$ }

(4) P (Fatality)$_i$ = CFR$_i$ × { P (I|HH)$_i$ + P (I|NH)$_i$ }

We do not estimate risks for persons under 20 years due to the infrequent incidence of

severe disease events in this age group. We also omit risk estimates for persons over

60 years as case hospital and fatality ratios cannot be reliably calculated for non-

congregate settings using publicly available data. We list our parameters and their

sources in Table 1 and discuss them further below.

Table 1. Parameters used for risk estimation

Symbol	Parameter	Unit	Value	Source
I	Average daily incidence rate of confirmed (reported) infections	Infections per day per person	Computed from data	Complied by the NYT from various sources
S	Proportion of the population susceptible	Unitless proportion	91%	Anand
F_{SX}	Proportion of infections with clinical illness	Unitless proportion	69% (95% CI, 63% - 74%)	Buitrago-Garcia
p	Percentage of transmission occurring prior to symptom onset	Unitless proportion	50%	He, Casey, Ferrriti, US CDC
q	Compliance with household self-isolation	Unitless proportion	75%	Unsourced
r	Relative Risk of infection transmitted from an asymptomatic individual relative to a symptomatic one	Unitless proportion	RR = 0.35 (95% CI, 0.10–1.27)	Buitrago-Garcia
B_{HH}	Household Secondary infection risk	Infections observed per contact[a] during a monitoring period	31.1% (95% CI 19.4% -42.7%)	Madewell
B_{NH}	Non-household Secondary infection risk	Infections observed per contact during a monitoring period	4.0% (95% CI: 2.8%, 5.2%)	Koh
$C_{i\ HH}$	Household contact rate for age group i	Contacts[b] / day	20-29 years: 2.77 30-39: 3.20 40-49: 3.18	Prem

			50-59 3.33	
$C_i N_H$	Non-household contact rate for age group i	Contacts / day	20-29 y 11.88 30-39 11.10 40-49 10.49 50-59 10.53	Prem
CHR_i	Case Hospitalization ratio for age group i	Unitless proportion	20 – 29 years: 1.65 % 30 – 39 years: 3.13 % 40-49 years: 4.75 % 50-59 years: 7.47 %	US CDC surveillance case reports
CFR_i	Age-specific Case Fatality ratio for age group i	Unitless proportion	20 – 29 years: 0.05% 30 – 39 years: 0.15% 40-49 years: 0.37% 50-59 years: 0.97%	US CDC surveillance case reports

[a] Most studies estimating the SIR defined contact as but most commonly either as face to face unprotected contact for greater than 15 minutes, prolonged contact, or household membership

[b] The basis of these estimates, the POLYMOD study, defined contact as "…either skin-to-skin contact such as a kiss or handshake (a physical contact), or a two-way conversation with three or more words in the physical presence of another person but no skin-to-skin contact (a nonphysical contact).

We acquired data on confirmed SARS-CoV-2 infection incidence rates from publicly reported statistics compiled by The New York Times,[2] computing average daily county-level confirmed infection incidence rates for the period of the "summer wave" in the U.S. (June 16th to Sept 15th 2020), then finding the median incidence rate for each quintile of incidence.

Confirmed infection rates underestimate the true incidence of infection as not all people with symptomatic infections obtain tests, test produce false negative results and subclinical infections occur. The US CDC currently estimates that an average of 11

infections may occur for every confirmed case, but the source of this estimate indicates

substantial regional variation.[3] We assume that confirmed infections and total

infections will be proportional and related multiplicatively and that unconfirmed

infections will have similar characteristics to confirmed ones. This assumption removes

the need for application of this parameter in our methodology.

The true susceptible proportion of the population is also unknown. Pandemic risk

assessments at the onset of the pandemic assumed a 100% susceptible population.

Seroprevalence surveys conducted in US populations using convenience samples

demonstrated low (<10%) antibody prevalence after the spring wave but substantial

variability among regions.[3] A systematic study of the U.S. dialysis population

estimated that seroprevalence of SARS-CoV-2 was 9·3% in US adult population,

ranging from 3·5% in the west to 27·2% in the northeast.[4] Understanding the

limitations of antibody-based determinations of susceptibility, we assumed the

prevalence of susceptibility to be 91% based on the seroprevalence of Anand et al. but

did not apply region-specific parameters in our methodology.

The parameters for secondary infection risk (SIR) (alternatively, secondary attack rates)

come from reviews of community transmission studies. Studies included in reviews

identify cases through active and passive surveillance and follow secondary contacts for

a specified duration (typically 14 days), monitoring contacts for symptoms and, in most

cases, testing contacts to confirm infection. Studies estimating the SIR define "contact"

variously but commonly either as face to face unprotected (i.e., mask-less) contact for

greater than 15 minutes, prolonged contact, or household membership. Studies

compute the SIR as the number of infections that occur among the reported contacts of

an ascertained case during the monitoring period. Studies typically do not ascertain the

frequency of contact.

Published reviews of the SIR for SARS-CoV-2 have focused on household

transmission. Koh et al reviewed 20 studies published by May 15, 2020, estimating a

pooled household SIR at 15.4% (95% CI: 12.2%, 18.7%).[5] The authors reported a

higher SIR for adults relative to children (RR 1.40, 95% CI: 1.00, 1.96). This review also

reported a summary non-household SIR of 4.0% (95% CI, 2.8%, 5.2%) which included

several cluster investigations with high SIR estimates.

Lei et al. summarized 24 studies with data on household transmission risk published by

July 1, 2020 including case reports with > 10 households, estimating a pooled SIR of

27% (95% CI, 21-32%).[6] The review found a 3-fold higher risks for household

transmission to adults. Data from ten studies with data on non-household contact,

specifically, provided an SIR of 1.65% (95% CI, 1.43 – 1.87) based on 218 secondary

cases among 13194 non-household contacts.

Madewell et al. estimated a summary SIR for 40 studies reporting SIRs for household

and family contacts published through 31 July 2020, estimating a mean SIR for

household contacts of 19·0% (95% CI, 14·9%–23·1%) and for family contacts of 18·1%

(95% CI: 12·9%–34·8%).[7] The estimated mean SIR for contact to adult household

members was 31.1% (95% CI, 19.4% -42.7%). The authors also estimated a mean SIR

for "close-contacts," which combined household and non-household contacts, as 4·3%

(95% CI: 2·9%–5·6%).

For our risk estimates, we used the 31% summary estimate of the adult household SIR

from Madewell et al. and the 4% summary estimate of the non-household SIR from Koh

et al. We used the 1.65% estimate of the non-household SIR from Lei et al in a

sensitivity analysis.

In the U.S., government agencies have not reported confirmed infection incidence by

symptom status. Estimates of the clinical fraction from published studies have

significant heterogeneity. A non-quantitative review of studies published from 19 April

through 26 May 2020, reported asymptomatic fractions of 43 - 65% in community

populations, 46-8% on cruise ships, 48 - 58% among personnel on aircraft carriers, 63-

88% in the context of homeless shelter outbreaks, and 96% among inmates.[8] A meta-

analysis of studies reported through 10 June estimated the asymptomatic fraction to be

20% (95% confidence interval [CI] 17–25) based on 79 studies with 6616 cases and

31% (95% CI 26%–37%) based on 7 studies with prospective follow-up.[9]

Alternatively, a single large case-ascertained follow up study of 5,484 contacts which

included both RT-PCR and serological examination, which limited under ascertainment

from false negative RT-PCR tests, found that only 32% of test positive contacts

developed symptoms of cough, shortness of breath with the clinical fraction varying

from 18% for those under 20 to 65% for those over 80 years old.[10] Another meta-

analysis of international studies also reported that clinical symptoms vary from 21% to

69% from the youngest to oldest subpopulations.[11] We used the 69% summary

estimate of the clinical fraction reported by Buitrago-Garcia et al. from the subset of

prospective studies and the lower 32% all-age-group clinical fraction reported by Polletti

et al in a sensitivity analysis.

We assume that subclinical infections will contribute to non-household exposure for the

duration of their infectious period while a fraction of those with clinical infections will self-

isolate in their households following symptoms. We found no empirical data to estimate

actual compliance with self-isolation requirements for the U.S. during the COVID-19

period. A recent survey of public attitudes in Israel found that willingness to comply with

self-quarantine rose from 57% without governmental financial compensation to 94%

with compensation.[12] We assumed that 75% of symptomatic individuals would self-

isolate after developing symptoms and tested the sensitivity of our estimate to an

alternative assumption of 50%.

Several transmission studies have estimated that roughly half of infection transmission

occurs pre-symptomatically.[13-15] He et al. estimated that 44% (95% CI, 30–57%) of

secondary cases were infected during the index cases' pre-symptomatic stage.[13]

Based on a review of 17 studies reporting serial intervals or generation times, Casey et

al. estimated that 56.1% of transmission occurred in the pre-symptomatic period.[14]

Ferretti et al. found that the peak of transmission occurred at the time of symptom onset

with 41% of transmission events occurring before symptoms onset and another 35% on

the day of and day after symptom onset.[15] We estimate that 50% of infectiousness

occurs before symptom onset.

Subclinical infections may be less infectious than clinical infections, because of

differences in viral burden or the result of symptoms, such as coughing or sneezing.

The review of Buitrago-Garcia reported that the secondary infection risk was lower

(relative risk 0.35, 95% CI 0.10–1.27) among contacts of asymptomatic cases relative to

symptomatic cases.[9] We considered infections in the subclinical fraction as being 35%

as infectious as the clinical fraction. In a sensitivity analysis, we considered subclinical

and clinical infections to be equally infectious.

We used age specific pre-pandemic estimates of contact rates from the work of Prem et

al. who modeled age and location specific contact rates in 140 countries re-applying

data from the European POLYMOD study.[16] The POLYMOD study defined contact as

"…either skin-to-skin contact such as a kiss or handshake (a physical contact), or a two-

way conversation with three or more words in the physical presence of another person

but no skin-to-skin contact (a nonphysical contact)."[17] This definition differs from the

one applied most commonly for studies estimating the SIR for SARS-CoV-2.   Prem et

al. reported contact rates for home settings as well as for work, school and other

settings. We summed the rates for non-home settings as a non-household contact rate.

(S1 Table 1) As we were unable to obtain age-stratified case incidence data, we did not

account for the age-structure of contacts.

We estimated the period case-hospitalization and case-fatality ratios by decile of age

directly from US CDC surveillance case reports using the total counts of cases and

those flagged for hospitalization or death.[18] We treated missing value in fields for

hospitalization and death as non-hospitalized, non-fatal cases. (S1: Table 2)

We computed risks for hospitalization and death for 90-day period using the average

period confirmed case incidence and the parameters listed in table 1. We utilized a

Monte Carlo simulation to incorporate each equation parameter with an error ranges as

a normally distributed variable, reporting risk as the mean of the resulting distribution.

To validate our estimates, we summed household and non-household risk and

compared it to cumulative national age-specific mortality and hospitalization incidence

rates for the same time period. [19-20] (S1: Tables 3 and Table 4)

Results

Among US Counties, in the 90-day period from June 16th, 2020 to September 15th,

2020, the median daily confirmed case incidence was 10.8 per 100,000, varying by

county from 3.1 per 100,000 in the first quartile to 28.5 per 100,000 in the fifth quartile.

Fig 1 illustrates the 90-day period risks for hospitalization and death attributable to

household and non-household contacts across the range of U.S. county case incidence

values applying the primary assumptions in Table 1.

268
269

Fig 1. Estimated risk of hospitalization and death over a 90-day period at pre-

pandemic U.S. contact rates within the range of recent U.S. county case

incidence under the following assumptions: Household SIR 31.1.%; Non-

household SIR, 4.0%; Clinical fraction, 69%; Relative infectiousness of

asymptomatic infection, 35%; Isolation adherence, 75%

Table 2 combines the household and non-household components of risk of

hospitalization and death for each quintile of confirmed case incidence. The estimated

90-day period risk of death at the period overall median county confirmed case

incidence of 10.8 per 100,000 ranges between 0.4 and 8.9 per 100,000 for each decile

of age between 20 and 60 years. The corresponding 90-day period risk of

hospitalization ranges from 13.7 to 69.2 per 100,000.

Table 2. Estimated risk of death and hospitalization at pre-pandemic U.S. contact

rates over a 90-day time period at the median county incidence of confirmed

infections during the "summer wave' of the COVID-19 pandemic (June 16 –

September 15, 2020).

	Cumulative Period Risk of Death (X 100000)					Cumulative Period Risk of Hospitalization (X 100000)				
Quintile	1	2	3	4	5	1	2	3	4	5
Average Daily Case Incidence	3.1	6.9	10.8	17.1	28.5	3.1	6.9	10.8	17.1	28.5
20 - 29 years	0.1	0.3	0.4	0.7	1.1	4	8.8	13.7	21.6	36.2
30 - 39 years	0.4	0.9	1.3	2.1	3.6	8.2	18.1	28.4	44.9	75
40 – 49 years	0.9	2.1	3.3	5.2	8.8	12.1	27.1	42.4	66.9	111.9
50 - 59 years	2.6	5.7	8.9	14.2	23.7	19.9	44.2	69.2	109.2	182.6

Though computed for different age-strata, our estimates are comparable to observed

U.S. average age-specific mortality rates during the same time period of 0.48, 1.46,

4.39, 10.81, and 24.67 per 100,000, for 10-year age groups from 15-24 years to 55-64

years. Our estimates are also comparable to average age-specific hospitalization rates

among the 98 counties in the USCDC COVID-Net hospital surveillance program (18-29

years, 44.4 per 100,000; 30-39 years, 59.2 per 100,000; 40-49 years 78.5 per 100,000;

50-64 years 109.1 per 100,000).

Table 3 provides the 90-day period risk estimates disaggregated by setting of contact

(household or non-household) at the national median county confirmed case incidence

of 10.8 per 100,000 and illustrates the sensitivity of risk estimates to the alternative

assumptions. Using the primary assumptions in table 1 (scenario 1), for a person aged

40 to 49 years, the 90-day period risk of death attributable to household and non-

household contacts is 2.56 and 0.74 events in 100,000 respectively. For

hospitalizations, the risk is 32.9 and 9.5 in 100,000 for household and non-household

contacts, respectively. Assuming a non-household SIR of 4.0% and pre-pandemic

contact rates, the share of transmissions attributable to household settings ranges from

73% to 78% depending on age.

Table 3. Estimated risk of death and hospitalization at pre-pandemic U.S. contact

rates over a 90-day time period at the median county incidence of confirmed

infections during the "summer wave' of the COVID-19 pandemic (June 16 –

September 15, 2020).

Age Group	Risk of Death (x 100,000)		Risk of Hospitalization (x 100,000)		Household Share
	Household Contacts	Non-Household Contacts	Household Contacts	Non-Household Contacts	
Scenario 1 (Base): Household SIR 31.1.%; Non-household SIR 4.0%, Clinical fraction 69%; Relative infectiousness of asymptomatic infection, 0.35; Isolation compliance 75%					
20 - 29 Years	0.3	0.11	9.96	3.74	73%
30 - 39 Years	1.04	0.32	21.78	6.62	77%
40 - 49 Years	2.56	0.74	32.89	9.5	78%
50 - 59 Years	7.04	1.95	54.18	15	78%

Scenario 2: Household SIR 31.1.%; Non-household SIR 1.65%, Clinical fraction 69%; Relative infectiousness of asymptomatic infection, 0.35; Isolation compliance 75%					
20 - 29 Years	0.3	0.05	9.89	1.55	86%
30 - 39 Years	1.04	0.13	21.62	2.75	89%
40 - 49 Years	2.54	0.31	32.65	3.95	89%
50 - 59 Years	6.98	0.81	53.77	6.23	90%

Scenario 3: Household SIR 31.1.%; Non-household SIR 4.0%%, Clinical fraction 31%; Relative infectiousness of asymptomatic infection, 0.35; Isolation compliance 75%					
20 - 29 Years	0.21	0.09	6.84	3.04	69%
30 - 39 Years	0.72	0.26	14.97	5.39	74%
40 - 49 Years	1.76	0.6	22.6	7.74	74%
50 - 59 Years	4.83	1.59	37.22	12.21	75%

Scenario 4: Household SIR, 31.1.%; Non-household SIR, 4.0%%, Clinical fraction, 69%; Relative infectiousness of asymptomatic infection, 1; Isolation compliance 75%					
20 - 29 Years	0.38	0.16	12.6	5.13	73%
30 - 39 Years	1.32	0.44	27.56	9.08	77%
40 - 49 Years	3.24	1.01	41.62	13.03	78%
50 - 59 Years	8.9	2.67	68.55	20.56	79%

Scenario 5: Household SIR, 31.1.%; Non-household SIR, 4.0%%, Clinical fraction, 69%; Relative infectiousness of asymptomatic infection, 0.35; Isolation compliance 50%					
20 - 29 Years	0.31	0.13	10.15	4.38	72%
30 - 39 Years	1.06	0.37	22.21	7.77	76%
40 - 49 Years	2.61	0.87	33.54	11.15	77%
50 - 59 Years	7.17	2.28	55.23	17.59	78%

Non-household risk falls proportionally using the lower estimate of the SIR of 1.65%

(scenario 2). Decreasing the estimated clinical fraction to 31% (scenario 3) decreases

both household and non-household risk estimates given we assume subclinical

infections to be less infectious. In the second scenario, the relative size of the non-

household share of risk also decreases slightly fewer infectious individuals are isolated.

Assuming sub-clinical infections are equally infectious as clinical infections, (scenario 4)

risks rise for both household and non-household contacts. Assuming adherence with

isolation drops to 50% (scenario 5), risk increases for non-household contacts.

Discussion

We demonstrate a straightforward method to estimate individual risks of COVID-19

associated hospitalization and deaths, using publicly available data on case incidence,

the clinical fraction, transmission timing, secondary infection risk, contact rates, and

case hospitalization and fatality ratios. These estimates of risk reflect the average risk

across a wide range of exposure settings and do not account for individual risk factors

for vulnerability to severe illness other than age. While many parameters have

significant uncertainties, the comparability of estimated risks to observed hospitalization

and fatality incidence rates validates the approach. Aggregated case incidence data,

parameter uncertainty, and the lack of setting specific transmission risk estimates are

limitations of the approach and suggest opportunities to improve individual risk

estimation.

We assumed the susceptible fraction of the population remains high based on

serological findings. The prevalence of detected antibodies varies with region and will

change with time. Furthermore, protective immunity may not be well estimated by

antibody detection alone. Durable lymphocyte responses may persist following

exposure.[21] Observed cellular immune response to COVID-19 among unexposed

individuals suggest prior exposure to related coronaviruses may further contribute to

immunity.[22]

Our estimates assume that prevalent infections are dispersed homogeneously within a

county's geography. This does not account for clustering within chains of transmission

among related or socially connected individuals. We might also expect higher infection

incidence among those living in congregate living facilities, in neighborhoods with more

density or larger households, or among service workers and those working together in

close quarters. Disaggregated public reporting of confirmed infection incidence by age,

symptom status, and neighborhood and congregate living status would allow for risk

estimates to be setting and population specific.

Reviews of the secondary infection risk find significant heterogeneity among estimates.

Most published estimates of the SIR come from observations outside the US and at an

earlier time period in the pandemic, prior to normalization of behaviors intended to

reduce the risk of infection transmission, such as increased hand washing, mask use

and observing physical distance. SIR estimates are dependent on both contextual and

study characteristics. Setting specific characteristics include the characteristics of

human contact. Most studies defined contacts similarly, yet studies varied in the number

of secondary contacts per primary case, suggesting variable interpretations of contact

definitions or variable ability or wiliness to recall and disclose contacts.

Most studies used a combination of RT-PCR and symptom monitoring to identify

infected contacts. RT-PCR methods will produce false negative results.[23] Studies that

ascertained infection with RT-PCR only at the beginning of contact observation could

have missed pre-clinical or subclinical infections.

The large difference between estimates of household and non-household SIR is not

unexpected given that frequent, prolonged and intimate contact is the norm in

households. Consistent with the relative size of the SIR estimates, we found that the

majority of SARS-CoV-2 transmission can be attributed to households. The WHO

mission to China also concluded that most infection transmission occurred in household

settings. [24] An analysis of 1,038 cases identified in Hong Kong up to 28 April

observed transmission to occur most frequently in households although non-household

social settings could involve a larger number of secondary infections.[25]

Relatively fewer studies have examined risk from non-household contact in detail; most

of these occurred in East Asian countries and used data from governmental surveillance

systems. We found insufficient estimates to apply SIRs for particular settings including

workplaces, schools, and public transportation.

There has been little systematic characterization of contacts of confirmed cases of

COVID-19 in the U.S. Reporting of the proportion confirmed cases with known contacts

and the setting of contact or relationships among cases of contacts varies by

jurisdiction; overall, few jurisdictions report this information. Among confirmed cases

identified before 'stay-at-home' orders in nine Colorado counties participating in a

retrospective survey, only 27% reported a contact with an infected individual.[26]

Another survey found only 46% adults testing positive for SARS-CoV-2 infection could

recall a contact with a known COVID-19 patient; of those who could recall a contact,

most recalled contacts were family members (45%) or work colleagues (34%).[27]

Our estimates do not account for individual variation in the SIR. With respiratory viruses,

the number of secondary cases generated by each index case can vary significantly.

[28] Studies suggest that a large share COVID-19 infection might be due to a small

fraction of particularly infectious individuals. [25,29] Characteristics of COVID-19

individual "super-spreaders" are not described but these individuals may make a

disproportionately larger contribution to the spread of infection in non-household, social

settings. [25] Clusters of COVID-19 infections have occurred in diverse settings. [30]

We applied the summary estimate of the non-household SIR by Koh et al which

included estimates of the SIR from cluster investigations thus representing the potential

contribution of "super-spreader" events.

We applied age-specific contact rates without regard to the age structure of contacts

and based on pre-pandemic estimates. While changes in mobility and consumption

suggest that many individuals may have lower contact rates, the nature and distribution

of current social contacts is unknown. Furthermore, limits on social and workplace

contacts outside the home may have increased household contact rates. Overall, we

did not have a satisfactory way to adjust contact rates for post-pandemic conditions.

The definition of contact used for our contact rates, which includes physical touch or

face to face speech, differs from contact definitions used in studies of secondary risk

from SARS-CoV-2, which was usually defined as face to face "unprotected" contact for

greater than 15 minutes. Neither definition may optimally characterize the mode of

COVID-19 transmission. For example, neither definition explicitly consider transmission

from contact via shared contaminated surfaces. Studies of the SIR that define contacts

alternatively, for example, as physical work in close quarters, might produce useful

findings.

We estimated the case hospitalization and fatality ratio by decile of age directly from

line-level CDC surveillance case report data during the same period used to assess

case-incidence. We note that these ratios have varied significantly both with time and by

location over the course of the pandemic; however, our examination of state-level ratios

over time demonstrates that recent period estimates have more homogeneity. (S1: Fig

1)

Our estimates do not account for variation in the risk of hospitalization and death due to

individual co-morbid conditions. [31] Adjusting for these factors would lower estimated

risks for most adults without common chronic disease conditions.

It is unclear how these risk estimates compare to of perceived individual risk among

adults in the U.S as there are few published estimates of perceived risk. In one online

survey conducted in March 2020, the median perceived risk was 10.0% for infection and

5.0% for infection fatality.[32] Another study conducted in March and April found that

adults found that in March, 14% of U.S. adults perceived the fatality risk from SARS-

CoV-2 infection to be greater than a 1% benchmark while 67% reported a lower risk;

however, the perception of risk increased over time. [33] Daily media reports of counts

of confirmed infections and the perceived lack of control over exposure all may be

influencing risk perception.

Public risk perceptions may not be reflecting the relatively larger risk from household

contacts. U.S. government action on restricting non-household social activity and the

publicity surrounding disease clusters in social settings may have amplified perceived

risks from non-household settings. In contrast, other countries took steps to quickly

physically isolate infectious family member to limit transmissions from household

contact. [34,35]

Return to community workplace and social life will require individuals to be have a better

understanding of the personal risks of COVID-19 infection. Accurate setting and

population specific estimates on the individual probabilities hospitalization and death

may contribute to a more accurate risk perception. Systematically collected and publicly

reported data on infection incidence by, for example, the geographic setting of

exposure, residence type, whether a case had a known exposure, would allow more

precise estimation than those possible with currently available public data. Calculation

of secondary infection risks by setting and a more precise knowledge of susceptibility

would improve individual risk estimates.

References

1.   Webster RK, Brooks SK, Smith LE, Woodland L, Wessely S, Rubin GJ. How to improve adherence with quarantine: rapid review of the evidence. Public Health. 2020 May;182:163-169. doi: 10.1016/j.puhe.2020.03.007.

2.   On ongoing repository of data on coronavirus cases and deaths in the U.S. The New York Times Company. [Cited: 2020 Oct 18] Available at: https://github.com/nytimes/covid-19-data

3.   Havers FP, Reed C, Lim T, Montgomery JM, Klena JD, Hall AJ, et al. Seroprevalence of Antibodies to SARS-CoV-2 in 10 Sites in the United States, March 23-May 12, 2020. JAMA Intern Med. 2020 Jul 21. doi: 10.1001/jamainternmed.2020.4130.

4.   Anand S, Montez-Rath M, Han J, Bozeman J, Kerschmann R, Beyer P, Parsonnet J, Chertow GM. Prevalence of SARS-CoV-2 antibodies in a large nationwide sample of patients on dialysis in the USA: a cross-sectional study. Lancet. 2020 Sep 25. doi: 10.1016/S0140-6736(20)32009-2.

5.   Koh WC, Naing L, Chaw L, Rosledzana MA, Alikhan MF, Jamaludin SA, Amin F, Omar A, Shazli A, Griffith M, Pastore R, Wong J. What do we know about SARS-CoV-2 transmission? A systematic review and meta-analysis of the secondary attack rate and associated risk factors. PLoS One. 2020 Oct 8;15(10):e0240205. doi: 10.1371/journal.pone.0240205.

6.   Lei H, Xu X, Xiao S, Wu X, Shu Y. Household transmission of COVID-19-a systematic review and meta-analysis. J Infect. 2020 Aug 25:S0163-4453(20)30571-5. doi: 10.1016/j.jinf.2020.08.033.

478 7. Madewell ZJ, Yang Y, Longini IM, Halloran ME, Dean NE. Household

479 transmission of SARS-CoV-2: a systematic review and meta-analysis of secondary

480 attack rate. medRxiv [Preprint]. 2020 Jul 31:2020.07.29.20164590. doi:

481 10.1101/2020.07.29.20164590.

482 8. Oran DP, Topol EJ. Prevalence of Asymptomatic SARS-CoV-2 Infection : A

483 Narrative Review. Ann Intern Med. 2020 Sep 1;173(5):362-367. doi: 10.7326/M20-3012.

484 9. Buitrago-Garcia D, Egli-Gany D, Counotte MJ, Hossmann S, Imeri H, Ipekci AM,

485 Salanti G, Low N. Occurrence and transmission potential of asymptomatic and

486 presymptomatic SARS-CoV-2 infections: A living systematic review and meta-analysis.

487 PLoS Med. 2020 Sep 22;17(9):e1003346. doi: 10.1371/journal.pmed.1003346.

488 10. Poletti P, Tirani M, Cereda D, et al. Probability of symptoms and critical disease

489 after SARS-CoV-2 infection. [Preprint]. 2020 Jun 22. arXiv:2006.08471v2

490 11. Davies NG, Klepac P, Liu Y, Prem K, Jit M. Age-dependent effects in the

491 transmission and control of COVID-19 epidemics. Nat Med. 2020Jun 16;26(8).

492 doi:10.1038/s41591-020-0962-9

493 12. Bodas M, Peleg K. Self-Isolation Compliance In The COVID-19 Era Influenced

494 By Compensation: Findings From A Recent Survey In Israel. Health Aff (Millwood).

495 2020 Apr 9; 39(6). doi:10.1377/hlthaff.2020.00382

496 13. He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al.. Temporal dynamics in

497 viral shedding and transmissibility of COVID-19. Nat Med. 2020 May;26(5):672-675. doi:

498 10.1038/s41591-020-0869-5. Erratum in: Nat Med. 2020 Sep;26(9):1491-1493.

499 14. Casey, M., Griffin, J., McAloon, C.G., Byrne, A.W., Madden, J.M., McEvoy, D.,

500 Collins, A.B., Hunt, K., Barber, A., Butler, F. and Lane, E.A. (2020). Estimating pre-

symptomatic transmission of COVID-19: a secondary analysis using published data.

[Preprint] medRxiv. 2020 Jun 11. https://doi.org/10.1101/2020.05.08.20094870

15.   Ferretti L, Ledda A, Wymant C, Zhao L, Ledda V, Abeler-Dorner L, et al. The

Timing of COVID-19 Transmission. [Preprint] MedRxiv. 2020 Sep 16. doi:

https://doi.org/10.1101/2020.09.04.20188516.

16.   Prem K, Cook AR, Jit M. Projecting social contact matrices in 152 countries using

contact surveys and demographic data. PLoS Comput Biol. 2017 Sep

12;13(9):e1005697. doi: 10.1371/journal.pcbi.1005697. PMID:

17.   Mossong J, Hens N, Jit M, Beutels P, Auranen K, Mikolajczyk R, et al. Social

contacts and mixing patterns relevant to the spread of infectious diseases. PLoS Med.

2008 Mar 25;5(3):e74. doi: 10.1371/journal.pmed.0050074.

18.   COVID-19 Case Surveillance Public Data Access, Summary, and Limitations.

U.S. Centers for Disease Control and Prevention. [Cited: 2020 Oct 18] Available at:

https://data.cdc.gov/Case-Surveillance/COVID-19-Case-Surveillance-Public-Use-

Data/vbim-akqf

19.   Provisional COVID-19 Death Counts by Sex, Age, and Week. U.S. Centers for

Disease Control and Prevention. [Cited: 2020 Oct 18] Available at:

https://data.cdc.gov/NCHS/Provisional-COVID-19-Death-Counts-by-Sex-Age-and-

W/vsak-wrfu

20.   Laboratory-confirmed Covid-19- Associated Hospitalizations. U.S. Centers for

Disease Control and Prevention. [Cited: 2020 Oct 18] Available at:

https://gis.cdc.gov/grasp/COVIDNet/COVID19_3.html

21.   Robbiani DF, Gaebler C, Muecksch F, Lorenzi JCC, Wang Z, Cho A, et al.

Convergent antibody responses to SARS-CoV-2 in convalescent individuals. Nature.

2020 Aug;584(7821):437-442. doi: 10.1038/s41586-020-2456-9.

22.   Grifoni A, Weiskopf D, Ramirez SI, Mateus J, Dan JM, Moderbacher CR, et al.

Targets of T Cell Responses to SARS-CoV-2 Coronavirus in Humans with COVID-19

Disease and Unexposed Individuals. Cell. 2020 Jun 25;181(7):1489-1501.e15. doi:

10.1016/j.cell.2020.05.015.

23.   Kucirka LM, Lauer SA, Laeyendecker O, Boon D, Lessler J. Variation in False-

Negative Rate of Reverse Transcriptase Polymerase Chain Reaction-Based SARS-

CoV-2 Tests by Time Since Exposure. Ann Intern Med. 2020 Aug 18;173(4):262-267.

doi: 10.7326/M20-1495.

24.   World Health Organization. Report of the WHO-China Joint Mission of

Coronavirus Disease 2019 (COVID-19). 2020 Feb 16-24.

25.   Adam DC, Wu P, Wong JY, Lau EHY, Tsang TK, Cauchemez S, Let al.

Clustering and superspreading potential of SARS-CoV-2 infections in Hong Kong. Nat

Med. 2020 Sep 17. doi: 10.1038/s41591-020-1092-0.

26.   Marshall K, Vahey GM, McDonald E, Tate JE, Herlihy R, Midgley CM, et al.

Exposures Before Issuance of Stay-at-Home Orders Among Persons with Laboratory-

Confirmed COVID-19 - Colorado, March 2020. MMWR Morb Mortal Wkly Rep. 2020 Jul

3;69(26):847-849. doi: 10.15585/mmwr.mm6926e4.

27.   Tenforde MW, Billig Rose E, Lindsell CJ, Shapiro NI, Files DC, et al.

Characteristics of Adult Outpatients and Inpatients with COVID-19 - 11 Academic

Medical Centers, United States, March-May 2020. MMWR Morb Mortal Wkly Rep. 2020

Jul 3;69(26):841-846. doi: 10.15585/mmwr.mm6926e3

28. Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the

effect of individual variation on disease emergence. Nature. 2005 Nov

17;438(7066):355-9. doi: 10.1038/nature04153.

29. Endo A; Centre for the Mathematical Modelling of Infectious Diseases COVID-19

Working Group, Abbott S, Kucharski AJ, Funk S. Estimating the overdispersion in

COVID-19 transmission using outbreak sizes outside China. Wellcome Open Res. 2020

Jul 10;5:67. doi: 10.12688/wellcomeopenres.15842.3.

30. Leclerc QJ, Fuller NM, Knight LE; CMMID COVID-19 Working Group, Funk S,

Knight GM. What settings have been linked to SARS-CoV-2 transmission clusters?

Wellcome Open Res. 2020 Jun 5;5:83. doi: 10.12688/wellcomeopenres.15889.2.

31. Kim L, Garg S, O'Halloran A, Whitaker M, Pham H, Anderson EJ, et al. Risk

Factors for Intensive Care Unit Admission and In-hospital Mortality among Hospitalized

Adults Identified through the U.S. Coronavirus Disease 2019 (COVID-19)-Associated

Hospitalization Surveillance Network (COVID-NET). Clin Infect Dis. 2020 Jul

16:ciaa1012. doi: 10.1093/cid/ciaa1012.

32. Bruine de Bruin W, Bennett D. Relationships Between Initial COVID-19 Risk

Perceptions and Protective Health Behaviors: A National Survey. Am J Prev Med. 2020

Aug;59(2):157-167. doi: 10.1016/j.amepre.2020.05.001.

Niepel C, Kranz D, Borgonovi F, Emslander V, Greiff S. The coronavirus (COVID-19)

fatality risk perception of US adult residents in March and April 2020. Br J Health

Psychol. 2020 Jun 10:10.1111/bjhp.12438. doi: 10.1111/bjhp.12438.

33.   Pan A, Liu L, Wang C, Guo H, Hao X, Wang Q, et al. Association of Public

Health Interventions With the Epidemiology of the COVID-19 Outbreak in Wuhan,

China. JAMA. 2020 May 19;323(19):1915-1923. doi: 10.1001/jama.2020.6130. PMID:

32275295; PMCID: PMC7149375.

34.   Thompson D. What's Behind South Korea's COVID-19 Exceptionalism? The

Atlantic. 2020 May 6.

Supreme Information

Supporting Information

Table 1. Daily contact rates by age and setting.

Table 2. Estimates of the case hospitalization and case fatality ratios.

Table 3. Cumulative COVID-19 associated hospital admissions rates per 100,000 people during the

period June 16 to September 15, 2020

Table 4. Cumulative COVID-19 associated mortality rates per 100,000 people during the period

June 16 to September 15, 2020.

Fig 1. Estimates of the state-specific case fatality ratio by month.