Estimation of Individual Probabilities of COVID-19 Infection, Hospitalization, and Death From A County-level Contact of Unknown infection Status

Rajiv Bhatia MD, MPH *

Assistant Professor of Medicine (affiliated), Stanford University

Physician, VA Palo Alto Health Care System

drajiv@stanford.edu

Jeffrey Klausner, MD, MPH

Professor of Medicine and Public Health, UCLA

JDKlausner@mednet.ucla.edu

Abstract: - 230 words

Manuscript: 2747 words

References: 822 words

Tables: 3

Figures: 2

* Corresponding Author

Version Date: June 23, 2020

Abstract

Objective: Our objective is to demonstrate a method to estimate the probability of a laboratory confirmed COVID19 infection, hospitalization, and death arising from a contact with an individual of unknown infection status.

Methods: We calculate the probability of a confirmed infection, hospitalization, and death resulting from a county-level person-contact using available data on current case incidence, secondary attack rates, infectious periods, asymptomatic infections, and ratios of confirmed infections to hospitalizations and fatalities.

Results: Among US counties with populations greater than 500,000 people, during the week ending June 13,2020, the median estimate of the county level probability of a confirmed infection is 1 infection in 40,500 person contacts (Range: 10,100 to 586,000). For a 50 to 64 year-old individual, the median estimate of the county level probability of a hospitalization is 1 in 709,000 person contacts (Range: 177,000 to 10,200,000) and the median estimate of the county level probability of a fatality is 1 in 6,670,000 person contacts (Range 1,680,000 to 97,600.000).

Conclusions and Relevance: Estimates of the individual probabilities of COVID19 infection, hospitalization and death vary widely but may not align with public risk perceptions. Systematically collected and publicly reported data on infection incidence by, for example, the setting of exposure, type of residence and occupation would allow more precise estimates of probabilities than possible with currently available public data. Calculation of secondary attack rates by setting and better measures of the prevalence of seropositivity would further improve those estimates.

Introduction

During infectious disease epidemics, human perception of risk modifies disease
transmission, and motivates, or not, protective behaviors such as hand hygiene,
wearing masks and social distancing at the individual level and quarantine, travel
restrictions, and restrictions on gatherings at the societal level. Novel infectious agents
such as the COVID-19 virus, with immature understanding of susceptibility,
transmission, and lethality, challenge accurate risk estimation.

Predictive models, government surveillance, and epidemiological studies have
characterized the risk of COVID-19, in terms of aggregate, and not individual,
outcomes, primarily including County-level counts of reported infections,
hospitalizations, and deaths. (1) Research has attempted to estimate case fatality rates
and identified risk factors for adverse outcomes from COVID-19 disease, such age and
chronic disease status. (2, 3) We are not aware of any published research that
estimates individual level probabilities of infection, hospitalization, and death from
exposure in the general community.

In the United States, publicly accessible data does not yet permit estimating the
individual risks of COVID-19 transmission in specific exposure settings, including
workplaces, prisons, nursing homes, hospitals and group residential housing settings.
However, a starting point for estimation can be the average individual-level probability of
acquiring infection across all settings at the level of a county. One can modify those

estimates as data on setting specific infection incidence rates, susceptibility and

secondary attack rates permit.

Here, we contribute to COVID-19 risk assessment by demonstrating a method to

estimate the individual probabilities of acquiring infection, being hospitalized, and dying

in U.S. Counties. We identify areas of available and future knowledge that could make

risk assessment more precise and context specific.

Materials and Methods

Our objective is to estimate the probability of acquiring COVID-19 infection from a

"contact" with a random individual of unknown infection status. We conceptualize this

probability under steady state conditions (e.g. no epidemic growth or decline) as a

function of individual susceptibility, the current reported case incidence, accounting for

undetected infection, the share of infection transmission occurring without a known

contact, the chance of transmission per contact (e.g., the secondary attack rate), and

the duration of infectiousness, accounting for pre-symptomatic transmission.

We use the formulae below to compute probabilities of infection, confirmed infection,

hospitalization, and death.

$(1)\ P_{infection\,|\,contact} = S\ I_{confirmed}\ C\ \beta\ [1+ \alpha/(1-\alpha)\ ]\ [\ \sigma\ D_{Infectious} + \eta\ (1-\sigma)\ D_{Infectious}\ ]$

$(2)\ P_{confirmed\text{-}infection\,|\,contact} = P_{infection|\,contact}/ (1-\alpha)$

$(3)\ P_{hospitalization\,|\,contact} = CHR \times P_{confirmed\text{-}infection\,|\,contact}$

(4) $P_{death \,|\, contact} = CFR \times P_{confirmed\text{-}infection\,|\, contact}$

We describe the parameters used in Table 1 and explain their sources below.

Table 1. Parameters Used in Formulae

Symbol	Parameter Description	Value
S	Proportion of the population susceptible	95%
$I_{confirmed}$	Incidence of confirmed (reported) infections per day	County Data
C	Proportion of infections resulting from unknown contacts	100%
α	Proportion of total infections unconfirmed (undetected and unreported)	75%
B	Probability of infection per exposure	10%
$D_{infectious}$	Average days infectious per infection	8
σ	Proportion of infectious period pre-symptomatic [a]	40%
η	Proportion non-compliant with isolation	25%
CHR	Case Hospitalization Ratio, 0 to 49 years [a]	0.026
CHR	Case Hospitalization Ratio, 50 to 64 years [a]	0.057
CHR	Case Hospitalization Ratio, Over 65 years [a]	0.1
CFR	Case Fatality Ratio, 0 to 49 years [a]	0.001
CFR	Case Fatality Ratio, 50 to 64 years [a]	0.006
CFR	Case Fatality Ratio, Over 65 years [a]	0.032

[a] US CDC COVID-19 Pandemic Planning Scenarios. Scenario 4 Estimates

The prevalence of susceptibility to COVID-19 is unknown. Pre-existing immunity due to

previous COVID-19-related coronaviral infections is plausible but speculative. Reliable

estimates for the proportion of the population who have acquired immunity is unknown

but non-zero. We have conservatively estimated the prevalence of susceptibility to be

95%.

We acquired COVID-19 confirmed infection incidence data from publicly reported

statistics compiled by The New York Times. (4) Confirmed infection rates underestimate

the true incidence of infection in the community because of undetected infections that

can be both symptomatic and asymptomatic. Several studies have estimated the

asymptomatic fraction. (5-7) In one meta-analytic review, the proportion of

asymptomatic infections ranged from 6% to 41%. (8) A weighted mean estimate from

those studies suggests that about 1 out of 6 persons, or 16%, may be asymptomatic.

On the other hand, seroprevalence studies aim to capture both the confirmed and

unconfirmed fractions together without regard to symptoms. Some seroprevalence

studies suggest that up to 90% of infections may be unconfirmed. (9) The US CDC

gives 50% as their most conservative estimate of the proportion of asymptomatic

infections. (10) We estimate 75% of all infections are unconfirmed which includes both

symptomatic and asymptomatic unconfirmed infections.

Limited data is available on the share of reported infections arising without a known

contact. As of this date, the US CDC has not included any statistics on this attribute of

confirmed infections. The State of Oregon currently publicly reports infections without a

known contact a varying between 30 and 50%%. Because of limited national data, we

use 100% for this parameter; we leave the parameter in the equation for the purpose of

future applications.

The attack rate for an exposure varies by exposure intensity, context, proximity, and

duration. For this analysis, contact means any substantive exposures that happen in a

community, household, workplaces, or group living situations. Examples of substantive

contacts outside households might include dining with a friend or business contact,

working in a shared office space or having close or physical contact without the types of

precautions now recommended for prevention of infection transmission (e.g. avoiding

handshaking, embraces, wearing a mask or indoor ventilation). We do not consider

contacts to be short-term events, such as passing by a person on the street. Contact

within households involves habitual and typically unprotected close physical. We

understand that attack rates will vary across such diverse exposure settings.

Overall, secondary attack rates from contact tracing studies on COVID-19 range from

0.7% to 16.3%. One study in Taiwan estimated a mean attack rate of 0.7% with an

attack rate of ~5% among household and non-household family contacts. (11) A Hong

Kong study of the quarantined contacts of visitors from China estimated a secondary

attack rate of 11.7%. (12) Two published study within China found a household attack

rate of 16.3% and 11.2% respectively. (13-14)

We assume an average plausible attack rate across all settings of exposure based on

these range of estimates to be 10% in the absence of more setting and activity specific

data. We acknowledge that this estimate may overestimate the attack rate for a non-

household contact and underestimate it for a household contact.

We estimate the total duration of infectiousness as 8 days. Research suggests that

individuals who develop symptoms may be infectious two to three days before the onset

of symptoms. (15) We apply the US CDC's estimates that the proportion of

infectiousness before symptom onset is 40% of the total duration. (10) Conservatively,

we treat infections in the unreported fraction as being infectious for the same duration

as those with reported infections.

Compliance with self-isolation affects the number of infectious people circulating with

infection after symptoms develop. Current research within the context of the COVID-19

pandemic finds that compliance with isolation ranges from 57% without financial

compensation to 94% with compensation. (16) Given the current US context and the

availability of sick leave compensation, we assume that 75% of individuals with

confirmed infection will voluntarily self-isolate after symptoms develop. We do not alter

the duration of infectiousness for the unconfirmed fraction of infections.

We estimate the probabilities per contact of reported infections, hospitalizations and

deaths as fixed ratios of the estimated but unobserved probability of infection. The

relationship between total infections and confirmed infections is fixed and defined by the

parameter alpha above.

We estimate the probability of hospitalizations and deaths per contact using US CDC

estimates of case hospitalization ratios and case fatality ratios from their pandemic

planning scenario four, which is their most conservative current estimate of disease

severity and transmissibility. (10) (Table 1) Based on these fixed ratios, we expect 18

reported infections and 70 total infections for every hospitalization in an individual in the

50-64 year old age group. And we expect 167 reported infections, and 667 total

infections for every death in the 50-64 year old age group.

These estimated case fatality ratios are within the range of published values in the

United States. One study of deaths through the early part of the epidemic estimated the

case fatality ratio for symptomatic cases to be 1.3% (95% CI: 0.6% to 2.1%). (2)

We compared estimated weekly hospitalization incidence rates produced using the

estimated probabilities per contact against observed incidence rates of laboratory

confirmed COVID-19 hospitalization in several US multi-county regions where the US

CDC conducts active hospital case surveillance. (17) To estimate weekly

hospitalizations, we multiplied our estimates of age specific probabilities of

hospitalization per contact and a modest number of daily contacts equal to the number

of other household members plus one.

Results

Among US Counties with populations greater than 500,000 people (N= 1224), for the week ending June 13, 2020, the median observed county-level daily case incidence is 6.78 per 100,000 (Range, 0.41- 24). In those counties, the median estimate of the county-level probability of a confirmed COVID-19 infection is 1 infection in 40,500 person-contacts (Range: 10,100 - 586,000). These estimates reflect the probability per contact at a single point in time averaged across all types of contacts and settings, within and outside households.

In the same counties, for a 50 to 64 year old individual, , the median estimate of the county-level probability of a hospitalization is 1 in 709,000 person-contacts (Range: 177,000 – 10,200,000) and the median estimate of the county-level probability of a fatality is 1 in 6,670,000 person-contacts (Range 1,680,000 – 97,600.000). Table 2 lists probabilities for other age groups.

203   Table 2. Estimated County-level Probabilities for the Week Ending June 13, 2020 of

204   COVID-19 Events in Counties with Populations Greater Than 500,000 (N = 1224)

Event	Age Group	Average	Median	Minimum	Maximum
Confirmed Case	All	0.002833%	0.002470%	0.000171%	0.009892%
Hospitalization	Under 49 yr	0.000074%	0.000064%	0.000004%	0.000257%
	50- 64 yr	0.000162%	0.000141%	0.000010%	0.000564%
	Over 65 yr	0.000283%	0.000247%	0.000017%	0.000989%
Fatality	Under 49 yr	0.000003%	0.000002%	0.000000%	0.000010%
	50- 64 yr	0.000017%	0.000015%	0.000001%	0.000059%
	Over 65 yr	0.000091%	0.000079%	0.000005%	0.000317%

Figure 1 illustrates the average estimated probabilities of confirmed infection,

hospitalization and fatality per contact as a function of daily case incidence. Figure 2

Illustrates the estimated number of hospitalizations and fatalities per 1 million contacts

in a subset of analyzed US counties with populations greater than 1.5 million.

We found good concordance between the estimated weekly hospitalization rates and

rates from US CDC hospitalization surveillance data in most of our comparison regions

under the assumption that average daily contacts equaled the average number of other

household members plus one. (Table 3) CDC acknowledges that more recent data

values are subject to revision.

Table 3. Estimated weekly hospitalization rates per 100,000 people for the week ending

June 13, 2020 compared to weekly laboratory confirmed hospitalization rates observed

by US CDC regional surveillance.

County	Alameda, CA	Denver CO	Fulton GA	Montgomery, MD	Wayne, MI
Assumption of Contacts Per Day	2.8	2.3	2.47	2.79	2.52
Estimated Weekly New Cases per 100,000	32.94	37.55	28.2	127.4	53.8
Estimated Weekly Hospitalization Rate (per 100,0000 people)					
18-49 years	0.86	0.98	0.7	3.3	1.4
50-64 years	1.88	2.14	1.6	7.3	3.1
Over 65 years	3.29	3.76	2.8	12.7	5.4
COVID-NET Observed Regional Weekly Hospitalization Rates					
18-49 years	1	1.1	1	5.3	0.4
50-64 years	1	3.2	2	6.6	0.8
Over 65 years	3.3	2.2	1.7	13.5	0.5

Discussion

We demonstrate a method to estimate the average county-level probabilities of COVID-

19 confirmed infections, hospitalizations, and deaths in the U.S resulting from a contact

with a random person in the population. Those estimates reflect current reported

COVID-19 confirmed infection incidence in US counties with more than 500,000 people

for the week ending June 6, 2020. Probabilities vary across a wide range reflecting the

varying case incidences in different counties.

Our method is limited by the availability of publicly available data on infection

transmission on COVID-19, including setting and occupational specific case incidence

rates, data on antibody seroprevalence, the share of infection with known contacts, and

activity-specific attack rates. The estimates therefore reflect the average probability

across a wide range of exposure contexts. Nevertheless, observed rates of

hospitalization for laboratory confirmed COVID-19 disease in several US CDC active

surveillance areas generally corroborate our estimates under the assumption of a

modest level of social contact.

Our estimates of the average probabilities per contact do not accurately estimate risks

for specific subsets of people. Infections occur within geographically and socially

constrained chains of transmission, for example, within clusters of related or socially

connected individuals or among those living in congregate living facilities such as

nursing homes. Clusters of COVID-19 infections have been reported associated with

prisons, workers dormitories, religious services, nightclubs, schools, cruise ships,

sporting events, and professional conferences. (18) Disaggregating confirmed infection

incidence rates into fractions representing those with and without known contacts and

those within and without congregate living settings would allow for more setting and

population specific risk estimates. This requires the systematic public reporting of

confirmed infection exposure factors.

As mentioned above, systematic data on the share of confirmed infections arising

without a known contact is not currently available. Appling this fraction to the model

would reduce our estimates based on the assumption of a 100% share.

Available estimates of the secondary attack rate come from observations outside the

US and at an earlier time period in the pandemic, prior to normalization of behaviors

that reduce the risk of infection transmission, such as increased hand washing,

observing physical distance, forgoing physical greetings, and wearing masks.

The limitations of published secondary attack rates and the lack of disaggregated data

on confirmed infection incidence also do not allow our estimates to differentiate between

community and household transmission. Most studies indicate that household attack

rates are higher than for contacts in the community. Systematic public reporting of

anonymized contact tracing data would provide information to assess context specific

attack rates.

We also cannot account for intra-individual variation in the secondary attack rate within

a setting. With respiratory viruses, the number of secondary cases generated by each

index case can vary significantly. (19-20) One recent estimate suggests that 80% of

COVID-19 infections are due to a small fraction (10%) of particularly infectious

individuals. (21)

We have assumed a high fraction of the population remains susceptible as the

prevalence and significance of antibodies to COVID-19 is not known. However, there

may well be factors conferring protection to infection or responsible for differences in

susceptibility. For example, cross immunity due to infection with other coronaviruses

may be occurring. (22) Researchers have also observed cellular immune system

responses to COVID-19 among unexposed individuals likely due to prior exposure to

related coronaviruses. (23) More data will be required before adjusting risk assessment

for population susceptibility or immunity.

We estimated probabilities of hospitalization and death based on CDC estimates of

disease severity. We chose conservative scenarios but acknowledge that estimates of

confirmed infection fatality ratios may further evolve over the course of the pandemic.

Furthermore, individual states may have different criteria for recording a death from

COVID-19. The risk of hospitalization and death varies with other risk factors including

race/ethnicity, level of deprivation, and chronic diseases such as lung disease and

diabetes, that are not reflected in our analysis.

Better data to address the above limitations would likely reduce estimated probabilities

of adverse events for a large fraction of the population; still, the estimated probabilities

reported here may appear considerably lower than those reflected in public opinion

surveys. Scientific uncertainties, media attention, dramatic governmental action and a

subjective perceived lack of control over exposure all may be influencing risk

perception. Exploring methods to communicate risk and the concordance of perceived

risk and risk probabilities would be an appropriate subject for further work.

Avoiding human contact in the setting of an uncertain and lethal epidemic threat is an

expected and self-protective human behavior. Prevalent beliefs today about the sources

of COVID-19 infection include 'contact with infected persons', 'people coming from

abroad' or 'foreign nationals.' (24) Notably, many people assume that contact with a

family member has a lower risk of infection transmission than contact with a stranger.

This may not be true for all subgroups in the population.

In the US, policy makers have taken dramatic and unprecedented steps to control

COVID-19, applying universal contact reductions through home confinement, limits on

travel, closures of schools and businesses and limits on gatherings. While heightened

perception of risk (e.g., fear) motivated these restrictions at the outset of the epidemic,

ongoing restrictions on community activity may be mediating risk perceptions. (25)

Notably, in Wuhan, the government limited exponential growth of the COVID-19

epidemic using isolation and quarantine, mandatory mask wearing, canceled New

Year's celebrations, the curtailment of intra-city and intercity transportation and

extension of the New Year's holiday period. (26) South Korea achieved epidemic control

with scaled up testing, strong contact notification practices, case isolation and strict

quarantine of those exposed and public awareness. (27) The lack of understanding of

how various countries have brought their epidemics under control maybe another

important factor influencing the perception of risk to COVID-19 infection in community

settings in the US.

Return to community workplace and social life will require individuals to be comfortable

with their personal risk of acquiring COVID-19 infection. Estimates on the individual

probabilities of infection, hospitalization and death may contribute to a more accurate

risk perception. Systematically collected and publicly reported data on infection

incidence by, for example, the geographic setting of exposure, residence type, whether

a case had a known exposure, and would allow more precise estimation than those

possible with currently available public data. Calculation of secondary attack rates by

setting and prevalence of seropositivity would further improve these estimates.

Contributions: Rajiv Bhatia and Jeffrey Klausner equally contributed to the design of

the study and writing of the manuscript. Rajiv Bhatia conducted the analysis.

Conflicts of interest: None

Funding: None

References

1.   Centers for Disease Control and Prevention. Covid-19 Forecasts. URL:

337 https://www.cdc.gov/coronavirus/2019-ncov/covid-data/forecasting-us.html Accessed

338 May 24, 2020.

2.   Basu A. Estimating The Infection Fatality Rate Among Symptomatic COVID-19

340 Cases In The United States [published online ahead of print, 2020 May 7]. *Health Aff*

341 (*Millwood). 2020.* doi:10.1377/hlthaff.2020.00455

3.   Zheng Z, Peng F, Xu B, et al. Risk factors of critical & mortal COVID-19 cases: A

343 systematic literature review and meta-analysis [published online ahead of print, 2020

344 Apr 23]. J Infect. 2020; S0163-4453(20)30234-6. doi:10.1016/j.jinf.2020.04.021

4.   On ongoing repository of data on coronavirus cases and deaths in the U.S. New

346 York Times. https://github.com/nytimes/covid-19-data Accessed: June 1, 2020.

5.   Furukawa NW, Brooks JT, Sobel J. Evidence supporting transmission of severe

348 acute respiratory syndrome coronavirus 2 while pre-symptomatic or asymptomatic.

349 Emerg Infect Dis. 2020 Jul. https://doi.org/10.3201/eid2607.201595

6.   He, X., Lau, E.H.Y., Wu, P. et al. Temporal dynamics in viral shedding and

351 transmissibility of COVID-19. *Nat Med* 26, 672–675 (2020). doi.org/10.1038/s41591-

352 020-0869-

7.   Wei WE, Li Z, Chiew CJ, Yong SE, Toh MP, Lee VJ. Presymptomatic

Transmission of SARS-CoV-2 - Singapore, January 23-March 16, 2020. MMWR Morb

Mortal Wkly Rep. 2020;69(14):411-415. Published 2020 Apr 10.

doi:10.15585/mmwr.mm6914e1

8.   Byambasuren O, Cardona M, Bell K, Clark J, McLaws ML, Glasziou P.

Estimating the extent of true asymptomatic COVID-19 and its potential for community

transmission: systematic review and meta-analysis. medRxiv.

doi.org/10.1101/2020.05.10.20097543

9.   Sood N, Simon P, Ebner P, et al. Seroprevalence of SARS-CoV-2-Specific

Antibodies Among Adults in Los Angeles County, California, on April 10-11, 2020

[published online ahead of print, 2020 May 18].   . 2020;e208279.

doi:10.1001/jama.2020.8279US

10.   Centers for Disease Control. Covid-19 Pandemic Planning Scenarios. URL:

https://www.cdc.gov/coronavirus/2019-ncov/hcp/planning-scenarios.html Accessed

May 29, 2020.

11.   Cheng HY, Jian SW, Liu DP, et al. Contact Tracing Assessment of COVID-19

Transmission Dynamics in Taiwan and Risk at Different Exposure Periods Before and

After Symptom Onset [published online ahead of print, 2020 May 1]. *JAMA Intern Med*.

2020;e202020. doi:10.1001/jamainternmed.2020.2020

12.   Kwok KO, Wong VWY, Wei WI, Wong SYS, Tang JW. Epidemiological

characteristics of the first 53 laboratory-confirmed cases of COVID-19 epidemic in Hong

Kong, 13 February 2020. *Euro Surveill*. 2020;25(16). doi:10.2807/1560-

7917.ES.2020.25.16.2000155

13.   Wei Li, Bo Zhang, Jianhua Lu, Shihua Liu, Zhiqiang Chang, Peng Cao, Xinhua

Liu, Peng Zhang, Yan Ling, Kaixiong Tao, Jianying Chen, The characteristics of

household transmission of COVID-19, Clinical Infectious Diseases.

doi.org/10.1093/cid/ciaa450

14.   Bi Q, Wu Y, Mei S, et al. Epidemiology and transmission of COVID-19 in 391

cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study

[published online ahead of print, 2020 Apr 27]. *Lancet Infect Dis*. 2020;S1473-

3099(20)30287-5. doi:10.1016/S1473-3099(20)30287-5

15.   Lauer SA, Grantz KH, Bi Q, et al. The Incubation Period of Coronavirus Disease

2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and

Application. *Ann Intern Med*. 2020;172(9):577-582. doi:10.7326/M20-0504

16.   Bodas M, Peleg K. Self-Isolation Compliance In The COVID-19 Era Influenced

By Compensation: Findings From A Recent Survey In Israel [published online ahead of

print, 2020 Apr 9]. *Health Aff* (Millwood). 2020;101377hlthaff202000382.

doi:10.1377/hlthaff.2020.00382

17.   US CDC. Laboratory-confirmed Covid-19- Associated Hospitalizations. URL:

https://gis.cdc.gov/grasp/COVIDNet/COVID19_3.html Accessed June 2, 2020.

18.   Leclerc QJ, Fuller NM, Knight LE et al. What settings have been linked to SARS-

CoV-2 transmission clusters? Wellcome Open Res 2020, 5:83

doi.org/10.12688/wellcomeopenres.15889.1

19.   Lipsitch M, Cohen T, Cooper B, et al. Transmission dynamics and control of

severe acute respiratory syndrome. *Science*. 2003;300(5627):1966-1970.

doi:10.1126/science.1086616

20.   Lloyd-Smith, J., Schreiber, S., Kopp, P. et al. Superspreading and the effect of

individual variation on disease emergence. *Nature* 438, 355–359 (2005).

https://doi.org/10.1038/nature04153

21.   Endo A, Centre for the Mathematical Modelling of Infectious Diseases COVID-19

Working Group, Abbott S et al. Estimating the overdispersion in COVID-19 transmission

using outbreak sizes outside China [version 1; peer review: 1 approved]. Wellcome

Open Res 2020, 5:67 doi.org/10.12688/wellcomeopenres.15842.1

22.   Kissler SM, Tedijanto C, Goldstein E, Grad YH, Lipsitch M. Projecting the

transmission dynamics of SARS-CoV-2 through the postpandemic period. Science.

2020;368(6493):860-868. doi:10.1126/science.abb5793

23.   Grifoni, A., Weiskopf, D., Ramirez, S.I., Mateus, J., Dan, J.M., Moderbacher,

C.R., Rawlings, S.A., Sutherland, A., Premkumar, L., Jadi, R.S., Marrama, D., de Silva,

A.M., Frazier, A., Carlin, A., Greenbaum, J.A., Peters, B., Krammer, F., Smith, D.M.,

Crotty, S., Sette, A., Targets of T cell responses to SARS-CoV-2 coronavirus in humans

with COVID-19 disease and unexposed individuals, *Cell* (2020).

doi.org/10.1016/j.cell.2020.05.015.

24.   Lohiniva AL, Sane J, Sibenberg K, Puumalainen T, Salminen M. Understanding

coronavirus disease (COVID-19) risk perceptions among the public to enhance risk

communication efforts: a practical approach for outbreaks, Finland, February

2020. *Euro Surveill.* 2020;25(13):2000317. doi:10.2807/1560-

7917.ES.2020.25.13.2000317

25.   Economic Attitudes as the Country Starts to Re-open. The Associated Press-

NORC Center for Public Affairs Research. URL:

http://www.apnorc.org/projects/Pages/Economic-Attitudes-as-the-Country-Starts-to-

Reopen.aspx Accessed: May 25, 2020.

26.   Pan A, Liu L, Wang C, et al. Association of Public Health Interventions With the

Epidemiology of the COVID-19 Outbreak in Wuhan, China [published online ahead of

print, 2020 Apr 10]. *JAMA*. 2020;323(19):1-9. doi:10.1001/jama.2020.6130

27.   Ryu S, Ali ST, Jang C, Kim B, Cowling BJ. Effect of non-pharmaceutical

interventions on transmission of severe acute respiratory syndrome coronavirus 2,

South Korea, 2020. *Emerg Infect Dis*. 2020 Oct. doi.org/10.3201/eid2610.201886

Figure 1. Estimated county-level probabilities of confirmed infection, hospitalization and death per contact as a function of daily case incidence.

Figure 2. Estimated current number of predicted COVID-19 hospitalizations per 1 million

contacts in counties with populations greater than 1.5 million.