Motivation and Cognitive Abilities as Mediators between Polygenic Scores and Psychopathology in Children

Narun Pornpattananangkul[1], Lucy Riglin[2], Richard Anney[2], Yue Wang[1], Deanna M. Barch[3], Anita Thapar[2] and Argyris Stringaris[4]

[1]Department of Psychology, University of Otago, New Zealand
[2]MRC Centre for Neuropsychiatric Genetics and Genomics, Division of Psychological Medicine and Clinical Neurosciences, School of Medicine, Cardiff University, UK
[3]Department of Psychological & Brain Sciences, Washington University, USA
[4]Mood Brain and Development Unit, National Institute of Mental Health, USA

Corresponding author:
Narun Pornpattananangkul, PhD
Department of Psychology, University of Otago
William James Building
275 Leith Walk
Dunedin 9016
New Zealand
Email: narun.pat@otago.ac.nz

Abstract

A fundamental question in psychology and psychiatry concerns the mechanisms that mediate between genetic liability and psychiatric symptoms. Genetic liability for common psychiatric disorders, such as Major Depressive Disorder (MDD) and Attention Deficit Hyperactivity Disorder (ADHD), often confer transdiagnostic risk to develop a wide variety of psychopathological symptoms through yet unknown pathways. Here we examined the three psychological and cognitive pathways that might mediate the relationship between genetic liability and broad psychopathology (as indexed by the P-Factor and its underlying dimensions): punishment sensitivity, reward sensitivity and cognitive abilities. We applied Structural Equation Modeling (SEM) on the Adolescent Brain Cognitive Development (ABCD) dataset (4439 children, 9-10 years old). The association between MDD polygenic scores (PS) and psychopathology was partially mediated by punishment sensitivity (reflected by Behavioral Inhibition System; BIS) and cognitive abilities (reflected by the G-Factor based on 10 neurocognitive tasks): proportion mediated=13.96%. Particularly, this mediating role of punishment sensitivity was specific to the emotional/internalizing dimension of psychopathology. Conversely, the influence of ADHD PS on psychopathology was partially mediated by reward sensitivity (reflected by Behavioral Activation System; BAS) and cognitive abilities (reflected by the G-Factor): proportion mediated=33.19%. This mediating role of reward sensitivity was focusing on the behavioral/externalizing and neurodevelopmental dimensions of psychopathology. These findings may provide a better understanding of how genetic risks for MDD and ADHD confer risk for psychopathology and suggest potential prevention and intervention targets for children at risk.

Keywords: Polygenic Score, Transdiagnostic Psychopathology, BIS/BAS, Cognitive G-Factor, P-Factor, Adolescent Brain Cognitive Development, MDD, ADHD, RDoC

Introduction

The last decade has seen major advances in psychiatric genetics. However fundamental questions remain about the psychological and cognitive links that associate genes with psychopathology[1,2]. In this paper, we examined whether three psychological and cognitive mechanisms-similar to those proposed by the National Institute of Mental Health's Research Domain Criteria (RDoC)[3] - punishment sensitivity, reward sensitivity and cognitive abilities - mediate the relationship between genetic liability and psychiatric symptoms.

Genetic liability to psychiatric disorders, such as major depressive disorder (MDD), attention deficit hyperactivity disorder (ADHD) and schizophrenia, can be captured by a composite of common gene variants identified from genome-wide association studies (GWAS), known as polygenic scores (PS)[4]. Multiple gene variants associated with different psychiatric disorders have been identified, and these cut across current diagnostic classification, such that genes contributing to one disorder also influence other phenotypes, not necessarily specific to such disorder[4–6]. For instance, PSs associated with case-status for MDD[5], ADHD[4,5] and schizophrenia[5,6] in GWAS discovery studies are associated with covariation among multiple psychiatric symptoms not limited to their respective disorders, assessed through a latent variable known as the Psychopathology-Factor or P-Factor[7]. It is unclear, however, what the mechanisms that explain this link between PSs and the P-Factor might be.

These mechanisms are likely to occur at multiple levels, including molecular[8], cellular[9,10], circuit[11], but also psychological and cognitive[7,12,13,14]. The RDoc has emerged as a framework to investigate how components of these different levels contribute to psychopathology[3]. Surprisingly little work has been done to examine which psychological and cognitive mechanisms mediate links between genetic risks and psychopathology. This is unfortunate as 1) at least some psychological and cognitive phenotypes can be measured with relatively high precision[15–17], and 2) some may also be amenable to interventions[18–20]. Understanding such mediating mechanisms may provide a foundation for effective early prevention and intervention strategies, an especially important goal in childhood given that most psychiatric disorders originate early in life. For this reason, our study focused on uncovering psychological and cognitive pathways between genetic liability (as indexed by PS), and psychopathology (as indexed by P-Factor as well as more specific latent variables, such as internalizing/emotional, externalizing/behavioral and neurodevelopmental) in children.

Here we focus on three psychological and cognitive mechanisms that show promise as potential mediators. The first two are related to motivation: punishment and reward sensitivity, or how easily affected individuals are by aversive and appetitive stimuli, respectively. Decades of work in psychopathology have demonstrated the importance of motivation[21]. Many emotional/internalizing symptoms (such as those that typify MDD and anxiety disorders) are associated with punishment sensitivity[12], as measured by the Behavioral Inhibition System (BIS) subscale[22]. Conversely, neurodevelopmental and behavioral/externalizing symptoms (such as those observed in ADHD and conduct disorders[13,14]) are associated with reward sensitivity, as measured by the Behavioral Activation System (BAS) subscale[22]. Accordingly, it is possible, for instance, that the previously found association between MDD PS and the P-Factor[5] might be mediated through punishment sensitivity (the BIS) while the association between the ADHD PS and the P-Factor[4,5] might be mediated through reward sensitivity (the BAS).

The third mechanism is related to cognitive abilities. Cognitive abilities have many forms, ranging from executive functions[23], verbal aptitude[24], learning and memory[25] to spatial reasoning[26,27]. The shared variance across different forms of cognitive abilities is represented by a latent variable, called the General Intelligence Factor, or G-Factor[28]. The G-factor has been associated with multiple different dimensions of psychopathology[7,16,29]. A recent twin study[29] also demonstrated that the G-factor broadly captures genetic propensity for psychopathology. Accordingly, we hypothesized that the G-Factor may mediate between genetic liability of various PSs and the P-Factor.

Recently the P-Factor has been integrated into a transdiagnostic framework that empirically groups related symptoms together in a hierarchical order of dimensions[30,31]. This framework has advantages over classical diagnostic systems (e.g., DSM or ICD) in terms of predicting clinical outcomes[32], such as new onsets of future diagnoses[33,34], suicide attempts[33,35,36] and psychosocial impairments[37–39]. This hierarchical structure has the P-Factor at its apex to represent broad severity across different types of psychopathology[40]. Moreover, the P-Factor is manifested by lower, specific dimensions. Based on a recent large-scale study in children from the Adolescent Brain Cognitive Development (ABCD) dataset[31], at the lower level of this hierarchical structure are five specific dimensions of behavioral/externalizing, neurodevelopmental, emotional/internalizing, somatoform and detachment items. These lower-level, specific dimensions make it possible to study mediating processes in greater granularity. Researchers can test which of the specific dimensions (in addition to the P-factor) are mediated by each mechanism. For instance, while PS for one disorder and the P-Factor might be jointly mediated by two mechanisms, these two mechanisms might mediate different specific dimensions from the other, suggesting dissociable roles in the pathway from genetic risk to psychopathology.

Our aim here is to provide a missing mechanistic link between genetic risk and clinical symptoms. We do by testing the hypotheses that motivational traits and cognitive abilities mediate between polygenic scores (PSs) for major psychiatric disorders and psychopathology (as indexed by the P-Factor and its underlying dimensions). The first step was to identify which of the common psychiatric PSs were associated with the P-Factor. Here we regressed the P-Factor on PSs for common psychiatric disorders: MDD, ADHD, Anxiety, Bipolar, Schizophrenia and Autism. Once identified, we conducted mediation analyses to formally examine the extent to which the relationship between each identified PS and the P-Factor was accounted for by the proposed mechanisms: punishment sensitivity (the BIS), reward sensitivity (the BAS) and cognitive abilities (the G-Factor). Finally, to further investigate the specific roles of the mediators, we conducted follow-up mediation analyses on the five specific dimensions by which the P-Factor was manifested. This allowed us to demonstrate detailed pathways for each mechanism to mediate specific sets of psychopathology.

Methods

Sample

We used the baseline, cross-sectional data from ABCD Release 2.01, collected at 21 sites across the United States between September 2016 and October 2018 (http://dx.doi.org/10.15154/1504041). While the study recruited over 11,000 children across races and ethnicities[41], we only selected children of European ancestry (assessed by multidimensional scaling analysis of their genotype data and compared with the 1000 Genomes Project phase3) to match with the discovery samples of the PSs. We excluded children whose genetic data had quality control issues (e.g., collected from a problematic plate or having incorrect data input) based on the study's recommendations (see

). After additional quality controls (see below), the final sample included 4439 children (2,081 females; M_{age}= 9.93 (SD=.61) years). The ABCD study was approved by the IRB at multiple sites[42], and obtained informed consent (parents) and assent (child)[43].

Polygenic Scores (PS)

Full details of genotyping has been published elsewhere[44]. Briefly, the study used the Smokescreen™ array, containing 733,293 SNPs[45]. The ABCD applied minimal quality control using the Ricopili pipeline. We further quality controlled and excluded children with high genetic relatedness, gender mismatches, minimal or excessive heterozygosity, disproportionate levels of individual missingness (>2%), insufficient sample replication (IBD <0.8) and cryptic relatedness (IBD ≥0.0422). We excluded SNPs based on minor allele frequency (<5% or >4SD discrepancy from the 1000 Genomes Project Reference panel), call rate (<98%) or evidence for violations of Hardy-Weinberg equilibrium (P < 1E-10). Using PRSice[46], we computed each PS as the Z-scored, weighted mean number of risk alleles in approximate linkage equilibrium, derived from imputed autosomal SNPs. We defined these risk alleles as those associated with case-status in large-scale discovery GWASs of six major psychiatric disorders: MDD[47], ADHD[48], Anxiety[49], Bipolar[50], Schizophrenia[51] and Autism[52]. In the main analysis, we focused on risk alleles that passed the p<.05 threshold in the discovery GWASs (see Supplementary for the analyses at other thresholds from p<.5 to .0001). Overall, the effects shown at the p<.05 threshold were also found at multiple thresholds. In our SEM that involved PSs, we also included control variables: (i) the number of SNP included to generate each PS, (ii) ten principal components (to control for population stratification).

Psychopathology: P-Factor and Five Specific Dimensions

We assessed children's psychopathology using the Child Behavior Checklist (CBCL),[53] reported by parents as detailed previously[54]. The CBCL included 119 items on a scale of 0 (Not True) to 2 (Very True or Often True) that reflected emotional, behavioural and ADHD problems occurring in the past 6 months. Following previous work[31], we removed low-frequency items and created composites for items that were highly correlated with each other. We captured the P-Factor and its lower, specific dimensions as latent variables in two confirmatory factor analysis (CFA) models.

First, the higher-order P-Factor model (Fig.1A) allowed us to model the P-Factor in the mediation analyses. Here we had the P-Factor as the 2nd-order latent variable and the five specific dimensions (behavioral/externalizing, neurodevelopmental, emotional/internalizing, somatoform and detachment; as defined previously[31]) as the 1st-order latent variables. In other words, the P-Factor was manifested by the five specific dimensions.

Second, the first-order model (Fig.1B) allowed us to model the five specific dimensions as correlated latent variables in the follow-up mediation analyses. We had the five specific dimensions as the 1st-order, correlated latent variables without the P-Factor. Using this model, we could test associations between variables (PSs and mediators) and each of the five specific dimensions while controlling for the correlations among the specific dimensions.

Motivation: Punishment (BIS) and Reward (BAS) Sensitivity

We assessed children's motivation through the Behavioral Inhibition System and Behavioral Activation System (BIS/BAS) scale[22] modified from PhenX, reported by children as detailed previously[54]. The scale included 20 items (7 for BIS) on 4-point Likert options (0=Not true; 3=Very true) that reflected punishment (BIS) and reward (BAS) sensitivity. The scale has been developed to

have a 4-factor structure: one BIS and three BAS subscales (Fun, Drive, Reward Responsiveness)[22]. Following a recent factor analysis in children[55], we dropped four problematic items: three from the BIS and one from the BAS-Reward-Responsiveness. This resulted in four items per (sub)scale.   To evaluate the latent structure of the BIS/BAS, we ran a CFA model, similar to the classical[22] and revised[55] 4-factor models. In this model (Fig.1C), BAS (as the higher-order variable) underlined the three BAS subscales (as the 1^{st}-order variables). We then allowed the BIS and BAS to covary. We treated The BIS and BAS as latent mediators for the mediation analyses.

Cognitive Abilities: G-Factor

We assessed children's cognitive abilities through various cognitive tasks as detailed previously[56,16]. Children completed these tasks on an iPad during a 70-min in-session visit. We included 10 tasks in our model (including seven tasks from the NIH Toolbox[17]). First, the Flanker task measured inhibitory control[57]. Second, the Card Sort task measured cognitive flexibility[58]. Third, the Pattern Comparison Processing task measured processing speed[59]. Fourth, the Picture Vocabulary task measured language and vocabulary comprehension[24]. Fifth, the Oral Reading Recognition task measured language decoding and reading[17]. Sixth, the Picture Sequence Memory task measured episodic memory[60]. Seventh, the Rey-Auditory Verbal Learning task measured auditory learning, recall and recognition[25]. Eight, the List Sorting Working Memory task measured working-memory[17]. Ninth, the Little Man task measured visuospatial processing via mental rotation[27]. Tenth, the Matrix Reasoning task measured visuospatial problem solving and inductive reasoning[25].

As a preliminary analysis, we followed previous work[16] by applying principal component analysis (PCA) to evaluate the structure of cognitive abilities. As we assume some similarity among the tasks, we used oblique (oblimin) rotations. The 4-component solution appeared to capture the cognitive tasks well, given minimal cross-loading. From this preliminary analysis, we then used CFA to capture the latent variable, the G-factor, the underlying cognitive abilities. In our higher-order G-Factor model (Fig.1D), we had the G-Factor as the 2^{nd}-order latent variable. We also had four 1^{st}-order latent variables in the model: executive functions (capturing the Flanker, Card Sort and Pattern Comparison Processing tasks), verbal (capturing the Picture Vocabulary and Oral Reading Recognition tasks), memory (capturing the Picture Sequence Memory, Rey-Auditory Verbal Learning and List Sorting Working Memory tasks) and spatial (capturing the Little Man and Matrix Reasoning tasks). We treated the G-Factor as a latent mediator for the mediation analyses.

Latent Variable Modelling Configurations

For each CFA structure, we fixed latent factor variances to one, so that we could estimate all factor loadings. We used robust estimators to deal with non-normality of psychopathological phenotypes in this population-based study. To this end, we first used robust maximum likelihood estimation (MLR) with robust (Huber-White) standard errors and scaled test statistics that also dealt with missing values via the Full Information Maximum Likelihood algorithm. However, if we encountered a non-convergent problem with the MLR, we treated data as ordinal and used WLSMV as an estimator instead. The WLSMV uses diagonally weighted least squares to estimate model parameters. To demonstrate model fit, we used scaled comparative fit index (CFI), Tucker-Lewis Index (TLI) and root mean squared error of approximation (RMSEA) with 90% CI. We reported the robust versions of these indicators for the MLR. For CFA, we also reported reliability of the latent variables: OmegaL2[61] for 2^{nd}-order variables and Omega3[62] for 1^{st}-order variables. These reliability indices reflect the internal consistency of the latent variables of interest. Model fits and reliability indices are shown in the figure captions. We ran the analyses in R4.0 on the standardized data

using lavaan[63] (version=.6-6) and semTools[61] along with semPlot[64] and qgraph[65] for visualization (see Supplementary for the script and detailed outputs).

Mediation Analyses with Structural Equation Modeling Steps

In our mediation analyses, we fit a series of latent variable models in successive steps. First, to identify which of the PSs are associated with the P-Factor, we treated the P-Factor from the higher-order P-Factor model as an outcome variable and the six PSs (MDD, ADHD, Anxiety, Bipolar, Schizophrenia and Autism) as explanatory variables. In this regression SEM model, the association between each PS and P-Factor was already controlled for other PSs. Second, to ensure that the proposed mediators were related to the P-Factor, we treated the P-factor as an outcome variable and the three proposed mechanisms (the BIS, the BAS and the G-Factor) as explanatory variables. Third, to demonstrate which of the three mediators were related to the PSs implicated by the first step, we treated the three proposed mediators as outcome variables and each of the selected PSs as an explanatory variable. Fourth, we examined the extent to which the relationship between each of the selected PS and the P-Factor was accounted for by the mediators implicated by steps two and three. Here we treated each implicated PS (step one) as an independent variable, mechanisms (steps two and three) as mediators and the P-Factor as a dependent variable. We then conducted follow-up analyses to further examine the role of the mediators by exploring associations with the five specific dimensions from the first-order model. For these follow-up mediation analyses, we started by examining the association between the five specific dimensions (as outcome variables) and each of the PS that was significantly associated with the P-factor (as an explanatory variable). Because the first-order model separately estimated correlations among the five specific dimensions, here we captured the unique associations between each specific dimension and PS (i.e., controlling for the correlations among the dimensions). Only the five specific dimensions of psychopathology that were associated with each PS were used in the final follow-up mediation analyses. Finally, we tested the extent to which the relationship between each significant PS and the specific dimensions of psychopathology was accounted for by the mediators.

Results

How Well Do the Proposed Latent Variable Models Fit the Data?
Fig.1 shows the results of Confirmatory Factor Analysis (CFA). All of the four proposed latent variable models had adequate model fit indices. Overall, the proposed latent variables (including the P-Factor, five specific dimensions, BIS, BAS, G-Factor) had good reliability.

Figure 1. Confirmatory Factor Analysis for latent variables of interest. Line thickness reflects the magnitude of standardized parameter estimates. The dotted lines indicate marker variables that were fixed to 1. Numbers in square indicate the item number from each scale. 1A and 1B show the higher-order P-Factor and first-order models for psychopathological phenotypes, respectively. The higher-order P-Factor showed good fit: scaled CFI=.923, TLI=.921 and RMSE=.028 (90%CI=.027-.028). The P-Factor in this model also had good reliability: OmegaL2=.875. The first-order model also showed good fit: scaled CFI=.927, TLI=.925 and RMSE=.027 (90%CI=.027-.028). Overall its 1st-order variables had good reliability (total Omega3=.962) while reliability for specific dimensions varied: externalizing (Omega3=.926), neurodevelopmental (Omega3=.874), internalizing (Omega3=.842), somatoform (Omega3=.730) and detachment (Omega3=.657). 1C shows the BIS/BAS model. This model showed good fit: robust, scaled CFI=.939, TLI=.927 and RMSE=.049 (90%CI=.046-.051). The latent variables had good reliability: 1st-order variables (total Omega3=.87) and the BAS (OmegaL2=.825). 1D shows the higher-order G-Factor model. This model showed good fit: robust, scaled CFI=.968, TLI=.953 and RMSE=.044 (90%CI=.039-.049). The G-Factor had good reliability (OmegaL2=.866). P = P-Factor; Ext = Externalizing; Neuro Dev = Neurodevelopmental; Int = Internalizing; Somatic = Somatoform; Detach = Detachment; Rew Res = Reward Responsiveness; G = the G-Factor; EF = Executive Functions; Car Sort = Card Sort; Pattern = Pattern Comparison Processing; Vocab = Picture Vocabulary; Read = Oral Reading Recognition; Pic = Picture Sequence Memory; Rey = Rey-Auditory Verbal Learning; List = List Sorting Working Memory; Matrix = Matrix Reasoning.

Which PSs are Associated with the P-Factor?

Our first Structural Equation Modelling (SEM) tested the relationship between the six psychiatric PSs and the P-Factor from the higher-order P-Factor model (Fig.2A, Fig.S1). Only the MDD (β=.080, SE=.016, z=4.863, p<.001) and ADHD PSs (β=.074, SE=.017, z=4.375, p<.001) showed unique associations with the P-Factor. Accordingly we treated the MDD and ADHD PSs as independent variables in our subsequent mediation analyses.

Are the Proposed Mechanisms Related to the P-Factor?

Next, we evaluated whether the proposed mediators were related to the main dependent variable, P-Factor (Fig.2B). Here, we examined the P-Factor in relationship to the BIS, BAS and G-Factor simultaneously, again allowing for the assessment of unique relationships. The P-Factor was significantly associated with all proposed mediators: the BIS (β=.061, SE=.026, z=2.413, p=.016), BAS (β=.106, SE=.025, z=4.231, p<.001) and G-Factor (β=-.222, SE=.023, z=-9.757, p<.001).

Are the Proposed Mechanisms Related to MDD and ADHD PSs?

We then separately evaluated whether each of the PSs that were associated with the P-Factor (MDD and ADHD PSs) were also related to each of the proposed mediators. MDD PS (Fig.2C, Fig.S2) was significantly related to the BIS (β=.041, SE=.019, z=2.232, p=.026) and the G-Factor (β=-.046, SE=.019, z=-2.407, p=.016), but not the BAS (β=.01, SE=.028, z=.566, p=.572). The BIS and the G-Factor were therefore included as mediators for MDD PS mediation analyses. ADHD PS (Fig.2D, Fig.S3) was significantly related to the BAS (β=.075, SE=.018, z=4.063, p<.001) and the G-Factor (β=-.09, SE=.018, z=-4.897, p<.001), but not the BIS (β=.014, SE=.02, z=.723, p=.47). The BAS and the G-Factor were therefore included as mediators for the ADHD PS mediation analyses.

Figure 2. The relationships between PSs, the P-Factor and the proposed mediators (BIS, BAS and G-Factor). 2A shows SEM testing the relationship between PSs and the P-Factor. This model showed good fit: robust, scaled CFI=.940, TLI=.938 and RMSE=.032 (90%CI=.031-.032). 2B shows SEM testing the relationship between the proposed mediators and the P-Factor. This model showed good fit: scaled CFI=.920, TLI=.918 and RMSE=.022 (90%CI=.022-.023). 2C shows SEM testing the relationship between MDD PS and the proposed mediators. The model for MDD PS showed good fit: robust, scaled CFI=.988, TLI=.986 and RMSE=.027 (90%CI=.026-.028). 2D shows SEM testing the relationship between the ADHD PS and the proposed mediators. This model showed good fit: robust, scaled CFI=.986, TLI=.984 and RMSE=.027 (90%CI=.026-.028). The numbers indicate standardized parameter estimates. Yellow indicates proposed independent variables. Green indicates dependent variables. Blue indicates proposed mediators. Con Vars = PS control variables (SNP included in each PS, ten principal components). *p<.05; **p<.01; ***p<.001.

<u>Do the Proposed Mechanisms Mediate between Each PS and the P-Factor?</u>
We then conducted the mediation SEM separately for MDD and ADHD PSs. For the MDD PS mediation model, the BIS and G-Factor were included as mediators (Fig.3A, Fig.S4, S5). The association between MDD PS and the P-Factor was partially mediated by both the BIS (proportion mediated=4.33%, Indirect β=.004, SE=.002, z=1.984, p=.047) and the G-Factor (proportion mediated=9.62%, Indirect β=.009, SE=.004, z=2.318, p=.020), together explaining 13.96% of the association.

The ADHD PS mediation model included the BAS and G-Factor as mediators (Fig.3B, Fig.S6, S7). The association between the ADHD PS and P-Factor was partially mediated by both the BAS (proportion mediated=7.716 %, Indirect β=.008, SE=.003, z=3.008 p=.003 and G-Factor (proportion mediated=17.544%, Indirect β=.017, SE=.004 , z=4.182, p<.001). Thus, the two mediators together explained 25.260% of the association between the ADHD PS and P-Factor.

*Figure 3. The mediations between the PSs (MDD and ADHD) and the P-Factor. 3A shows SEM testing the mediation between MDD PS and the P-Factor with the BIS and G-Factor as mediators. This model showed good fit: robust, scaled CFI=.911, TLI=.908 and RMSE=.032 (90%CI=.031-.032). 3B shows SEM testing the mediation between the ADHD PS and the P-Factor with BAS and G-Factor as mediators. This model showed good fit: robust, scaled CFI=.904, TLI=.901 and RMSE=.030 (90%CI=.029-.030). The numbers indicate standardized parameter estimates. The parenthesis indicates the direct effect between each of the PS and the P-Factor after accounted for by the mediators. Yellow indicates independent variables. Blue indicates mediators. Green indicates dependent variables. Con Vars = PS control variables (SNP included in each PS and ten principal components) *p<.05; **p<.01; ***p<.001.*

Which of the Five Specific Dimensions are Associated with Each PS?

We then conducted follow-up mediation analyses to investigate the distinct roles of the mediators at the level of five specific dimensions for both the MDD and ADHD PSs. We first tested the relationship between each of the two PSs and the five dimensions. MDD PS was significantly associated with all five specific dimensions (see βs in Fig.4A, Fig.S8): all dimensions were therefore included in the follow-up mediation analyses for MDD PS.

The ADHD PS was statistically associated with externalizing, neurodevelopmental, internalizing and somatoform but not detachment (see βs in Fig.4B, Fig.S9): these four dimensions were therefore included in the follow-up mediation analyses for the ADHD PS.

Figure 4. The relationships between PSs (MDD and ADHD) and specific dimensions of psychopathology. 4A shows SEM testing the relationship between MDD PS and the five specific dimensions from the first-order model. This model showed good fit: robust, scaled CFI=.917, TLI=.913 and RMSE=.035 (90%CI=.035-.036). 4B shows SEM testing the relationship between the ADHD PS and the five specific dimensions from the first-order model. This model showed good fit: robust, scaled CFI=.908, TLI=.904 and RMSE=.035 (90%CI=.035-.036). The numbers indicate standardized parameter estimates. Yellow indicates independent variables. Green indicates dependent variables. Con Vars = PS control variables (SNP included in each PS and ten principal components). *p<.05; **p<.01; ***p<.001.

Do the Proposed Mechanisms Mediate between Each PS and Specific Dimensions?

For the MDD PS follow-up mediation model (Fig.5A, Fig.S10, S11), the BIS specifically mediated the influence of MDD PS on internalizing (proportion mediated=10.80%, Indirect β=.008, SE=.004, z=2.146, p=.032), whereas the G-Factor specifically mediated the influence of MDD PS on externalizing (proportion mediated=13.30%, Indirect β=.009, SE=.004, z=2.327, p=.02) and neurodevelopmental (proportion mediated=15.38%, Indirect β=.012, SE=.005, z=2.353, p=.019) dimensions.

For the ADHD PS follow-up mediation model (Fig.5B, Fig.S12, S13), the BAS specifically mediated the influence of the ADHD PS on externalizing (proportion mediated=11.41%, Indirect β=.011, SE=0.003, z=3.55, p<0.001) and neurodevelopmental (proportion mediated=8.65%, Indirect β=.009,

SE=0.003, z=3.26, p=0.001), whereas the G-Factor mediated the influence of the ADHD PS on all four dimensions: externalizing (proportion mediated=16.10%, Indirect β=0.015, SE=0.004, z=4.124, p=<0.001), neurodevelopmental (proportion mediated=20.38%, Indirect β=0.021, SE=0.004, z=4.372, p=<0.001), internalizing (proportion mediated=13.10%, Indirect β=0.005, SE=0.002, z=2.257, p=.023) and somatoform (proportion mediated=11.30%, Indirect β=0.005, SE=0.002, z=2.182, p=.029).

*Figure 5. The mediation between Polygenic Scores (MDD and ADHD) and specific dimensions of psychopathology. 5A shows SEM testing the mediation between MDD PS and the five specific dimensions of psychopathology with the BIS and G-Factor as mediators. This model showed good fit: robust, scaled CFI=.913, TLI=.909 and RMSE=.031 (90%CI=.031-.032). 5B shows SEM testing the mediation between ADHD PS and the four specific dimensions of psychopathology with BAS and G-Factor as mediators. This model showed adequate fit: robust, scaled CFI=.899, TLI=.895 and RMSE=.030 (90%CI=.030-.031). The numbers overlaid black lines indicate standardized parameter estimates. The numbers on the right side next to each specific dimension of psychopathology indicate proportion mediated for the mediation paths with significant indirect effects. Dotted lines indicate mediation paths with non-significant (p<.05) indirect effects. Yellow indicates independent variables. Blue indicates mediators. Green indicates dependent variables. Con Vars = PS control variables (SNP included in each PS and ten principal components) *p<.05; **p<.01; ***p<.001.*

Discussion

Here we aimed to uncover the psychological and cognitive mechanisms mediating the relationship between genetics and psychopathology. In particular, we tested whether three RDoC-based psychological and cognitive mechanisms, namely punishment sensitivity (BIS), reward sensitivity (BAS) and cognitive abilities (G-Factor), mediated the relationship between polygenic scores (PSs) for different psychiatric disorders and psychiatric symptoms across disorders (the P-Factor and its specific dimensions). We first identified that, among the six common psychiatric PSs, MDD and ADHD PSs were associated with the P-Factor in children, consistent with the previous reports[4,5]. Moreover, these two PSs were related to our proposed mediation mechanisms. Importantly, the proposed mechanisms partially mediated the relationship of the two PSs to the P-Factor and its specific dimensions.

The relationship of MDD PS to the P-factor was mediated by the BIS and G-Factor, whereas the relationship of ADHD PS to the P-Factor was mediated by the BAS and G-Factor. Thus, the influence of MDD and ADHD PSs on psychopathology may be acting through both shared (G-Factor) and unique (BIS versus BAS) routes. Here we demonstrated two routes for MDD PS (punishment sensitivity and cognitive abilities) and two routes for ADHD PS that are partially dissociable from MDD PS (reward sensitivity and cognitive abilities). To further investigate the specificity of these pathways, we conducted follow-up mediation analyses on the five specific dimensions of psychopathology by which the P-Factor was manifested. As discussed in more detail below, our results showed that the proposed psychological and cognitive mechanisms differentially mediated each of the five specific dimensions. Thus, together these data are consistent with the hypothesis that dissociable but complementary pathways mediate the influence of the MDD and ADHD PSs on the P-Factor.

The mediating role of the BIS from MDD PS to the P-Factor is consistent with studies associating the BIS with emotional/internalizing symptoms[23]. When examining its detailed mediating pathways using specific dimensions of psychopathology, we found a high level of specificity in the mediation: the BIS was only significantly related to one PS (MDD PS) and one specific dimension of psychopathology (internalizing). Conversely, the mediating role of the BAS to ADHD PS and the P-Factor is consistent with associating the BAS with neurodevelopmental and behavioral/externalizing symptoms[13,14]. Similar to the BIS, the BAS also showed a high level of specificity in its mediation: it was only significantly related to one PS (ADHD PS) and two specific dimensions (neurodevelopmental and behavioral/externalizing symptoms). Together, these findings suggest that motivation-related mechanisms, punishment and reward sensitivity, mediated the influences of genetics in a specific manner.

The mediating role of the G-Factor to both MDD and ADHD PSs is in line with previous work showing relationships between cognitive abilities and broad psychopathology[7,16]. Unlike the two motivation-related mediators, the G-Factor showed a broader role. That is, the G-Factor mediated the influences between 1) both MDD and ADHD PSs and 2) various specific dimensions of psychopathology. For both MDD and ADHD PS, the G-Factor mediated the contribution of genetic influences to the externalizing and neurodevelopmental dimensions. For ADHD PS, cognitive abilities additionally mediated the link with internalising and somatic dimensions. Accordingly, we found that having genetic liability for MDD and/or ADHD had negative associations with cognitive abilities, which in turn, may enhance the general risk to develop psychopathology. As such,

cognitive abilities played a key role as a non-specific factor that linked genetic liability with broad psychopathology, consistent with previous findings and theoretical perspectives of the P-Factor[7,40].

We believe understanding the roles of the proposed psychological and cognitive mechanisms have research and clinical implications. As we showed here, the model fit indices and reliability (i.e., internal consistency) indices of the proposed mechanisms were relatively high. This means that we can measure these mechanisms in children with precision using latent variable modeling. Moreover, punishment and reward sensitivity and cognitive abilities are shown to be altered via psychotherapy and other environment-altering interventions[18–20]. Accordingly, they can be targeted for effective early prevention and intervention strategies for children at risk.

This study is not without limitations. First, as highlighted previously[66], while using PSs derived from GWAS are more reliable than using only a few common SNPs from selected genes (the candidate-gene approach), PSs still explain only a small proportion of genetic liability to psychiatric disorders. The relatively small effect sizes of our results confirm this notion. Next, in the original studies of the P-Factor that were done in adults[7,40], psychosis experiences were considered as another separate dimension. Our use of CBCL[31], however, did not allow us to investigate psychosis as another dimension. Thus, our definition of the P-Factor may not totally align with the original definition. Previous research cast doubt on the ability to detect psychosis experiences in 9-10 years old[67], and internalizing and externalizing are usually the focus in children studies[68,69,31]. Nonetheless, the difference in definitions of the P-Factor should be noted. Finally, we measured the mediators and psychopathology at the same time, making it difficult to empirically test the directionality of their relationships. Fortunately, the ABCD is an ongoing longitudinal study that will provide additional data from the same children until they are 20 years old[70]. Thus, we believe that our study will lay a foundation for future research to further empirically test the directionality of the effects found here, for instance, using the cross-lagged panel model.

In summary, in a large sample of children, the influences of genetic predispositions for MDD and ADHD on psychopathology were mediated by three RDoC mechanisms: punishment sensitivity, reward sensitivity and cognitive abilities. Therefore, these findings further our understanding of the structure of psychopathology and the pathways through which it relates to genetic architecture.

Acknowledgment

Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive Development (ABCD) Study (https://abcdstudy.org), held in the NIMH Data Archive (NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children age 9-10 and follow them over 10 years into early adulthood. The ABCD Study is supported by the National Institutes of Health and additional federal partners under award numbers U01DA041022, U01DA041028, U01DA041048, U01DA041089, U01DA041106, U01DA041117, U01DA041120, U01DA041134, U01DA041148, U01DA041156, U01DA041174, U24DA041123, U24DA041147, U01DA041093, and U01DA041025. A full list of supporters is available at https://abcdstudy.org/federal-partners.html. A listing of participating sites and a complete listing of the study investigators can be found at https://abcdstudy.org/scientists/workgroups/. ABCD consortium investigators designed and implemented the study and/or provided data but did not necessarily participate in analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD consortium investigators. NP, YW and AS were supported by Otago Medical Research Foundation Grant though M Begg Charitable. Declaration of Interest: The authors declare no competing interests.

References

1. Nolen-Hoeksema S, Watkins ER. A Heuristic for Developing Transdiagnostic Models of Psychopathology: Explaining Multifinality and Divergent Trajectories. *Perspect Psychol Sci*. 2011;6(6):589-609. doi:10.1177/1745691611419672

2. Gottesman II, Gould TD. The Endophenotype Concept in Psychiatry: Etymology and Strategic Intentions. *Am J Psychiatry*. 2003;160(4):636-645. doi:10.1176/appi.ajp.160.4.636

3. Insel T, Cuthbert B, Garvey M, et al. Research Domain Criteria (RDoC): Toward a New Classification Framework for Research on Mental Disorders. *Am J Psychiatry*. 2010;167(7):748-751. doi:10.1176/appi.ajp.2010.09091379

4. Brikell I, Larsson H, Lu Y, et al. The contribution of common genetic risk variants for ADHD to a general factor of childhood psychopathology. *Mol Psychiatry*. Published online June 22, 2018. doi:10.1038/s41380-018-0109-2

5. Riglin L, Thapar AK, Leppert B, et al. Using Genetics to Examine a General Liability to Childhood Psychopathology. *Behav Genet*. Published online December 11, 2019. doi:10.1007/s10519-019-09985-4

6. Jones HJ, Heron J, Hammerton G, et al. Investigating the genetic architecture of general and specific psychopathology in adolescence. *Transl Psychiatry*. 2018;8(1):145. doi:10.1038/s41398-018-0204-9

7. Caspi A, Houts RM, Belsky DW, et al. The p Factor: One General Psychopathology Factor in the Structure of Psychiatric Disorders? *Clin Psychol Sci*. 2014;2(2):119-137. doi:10.1177/2167702613497473

8. Jansen R, Penninx BWJH, Madar V, et al. Gene expression in major depressive disorder. *Mol Psychiatry*. 2016;21(3):339-347. doi:10.1038/mp.2015.57

9. Lin Y-C, Koleske AJ. Mechanisms of Synapse and Dendrite Maintenance and Their Disruption in Psychiatric and Neurodegenerative Disorders. *Annu Rev Neurosci*. 2010;33(1):349-378. doi:10.1146/annurev-neuro-060909-153204

10. van Spronsen M, Hoogenraad CC. Synapse Pathology in Psychiatric and Neurologic Disease. *Curr Neurol Neurosci Rep*. 2010;10(3):207-214. doi:10.1007/s11910-010-0104-8

11. Ward J, Lyall LM, Bethlehem RAI, et al. Novel genome-wide associations for anhedonia, genetic correlation with psychiatric disorders, and polygenic association with brain structure. *Transl Psychiatry*. 2019;9(1):1-9. doi:10.1038/s41398-019-0635-y

12. Katz BA, Matanky K, Aviram G, Yovel I. Reinforcement sensitivity, depression and anxiety: A meta-analysis and meta-analytic structural equation model. *Clin Psychol Rev*. 2020;77:101842. doi:10.1016/j.cpr.2020.101842

13. Becker SP, Fite PJ, Garner AA, Greening L, Stoppelbein L, Luebbe AM. Reward and punishment sensitivity are differentially associated with ADHD and sluggish cognitive tempo symptoms in children. *J Res Personal*. 2013;47(6):719-727. doi:10.1016/j.jrp.2013.07.001

14. Hundt NE, Kimbrel NA, Mitchell JT, Nelson-Gray RO. High BAS, but not low BIS, predicts externalizing symptoms in adults. *Personal Individ Differ*. 2008;44(3):565-575. doi:10.1016/j.paid.2007.09.018

15. Jorm AF, Christensen H, Henderson AS, Jacomb PA, Korten AE, Rodgers B. Using the BIS/BAS scales to measure behavioural inhibition and behavioural activation: Factor structure, validity and norms in a large community sample. *Personal Individ Differ*. 1998;26(1):49-58. doi:10.1016/S0191-8869(98)00143-3

16. Thompson WK, Barch DM, Bjork JM, et al. The structure of cognition in 9 and 10 year-old children and associations with problem behaviors: Findings from the ABCD study's baseline neurocognitive battery. *Dev Cogn Neurosci*. 2019;36:100606. doi:10.1016/j.dcn.2018.12.004

17. Bleck TP, Nowinski CJ, Gershon R, Koroshetz WJ. What is the NIH Toolbox, and what will it mean to neurology? *Neurology*. 2013;80(10):874-875. doi:10.1212/WNL.0b013e3182872ea0

18. Craske MG, Meuret AE, Ritz T, Treanor M, Dour H, Rosenfield D. Positive affect treatment for depression and anxiety: A randomized clinical trial for a core feature of anhedonia. *J Consult Clin Psychol*. 2019;87(5):457-471. doi:10.1037/ccp0000396

19. Rapee RM. The preventative effects of a brief, early intervention for preschool-aged children at risk for internalising: follow-up into middle adolescence: Adolescent effects of brief, early intervention for internalising. *J Child Psychol Psychiatry*. 2013;54(7):780-788. doi:10.1111/jcpp.12048

20. Diamond A, Lee K. Interventions Shown to Aid Executive Function Development in Children 4 to 12 Years Old. *Science*. 2011;333(6045):959-964. doi:10.1126/science.1204529

21. Tang A, Crawford H, Morales S, Degnan KA, Pine DS, Fox NA. Infant behavioral inhibition predicts personality and social outcomes three decades later. *Proc Natl Acad Sci*. Published online April 20, 2020:201917376. doi:10.1073/pnas.1917376117

22. Carver CS, White TL. Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: The BIS/BAS Scales. *J Pers Soc Psychol*. 1994;67(2):319-333. doi:10.1037/0022-3514.67.2.319

23. Miyake A, Friedman NP. The Nature and Organization of Individual Differences in Executive Functions: Four General Conclusions. *Curr Dir Psychol Sci*. 2012;21(1):8-14. doi:10.1177/0963721411429458

24. Gershon RC, Cook KF, Mungas D, et al. Language measures of the NIH toolbox cognition battery. *J Int Neuropsychol Soc*. 2014;20(6):642-651. doi:10.1017/S1355617714000411

25. Daniel MH, Wahlstrom D. Equivalence of Q-interactive™ and Paper Administrations of Cognitive Tasks: WISC®–V. Published online 2014:13.

26. Wechsler D. *Wechsler Intelligence Scale for Children*. Psychological Corporation; 1949.

27. Acker WL, Acker C. *Bexley Maudsley Automated Psychological Screening and Bexley Maudsley Category Sorting Test Manual*. NFER-Nelson, for the Institute of Psychiatry; 1982.

28. Spearman C. "General Intelligence," Objectively Determined and Measured. *Am J Psychol*. 1904;15(2):201. doi:10.2307/1412107

29. Harden KP, Engelhardt LE, Mann FD, et al. Genetic Associations Between Executive Functions and a General Factor of Psychopathology. *J Am Acad Child Adolesc Psychiatry*. Published online May 2019:S0890856719303272. doi:10.1016/j.jaac.2019.05.006

30. Kotov R, Krueger RF, Watson D, et al. The Hierarchical Taxonomy of Psychopathology (HiTOP): A dimensional alternative to traditional nosologies. *J Abnorm Psychol*. 2017;126(4):454-477. doi:10.1037/abn0000258

31. Michelini G, Barch DM, Tian Y, Watson D, Klein DN, Kotov R. Delineating and validating higher-order dimensions of psychopathology in the Adolescent Brain Cognitive Development (ABCD) study. *Transl Psychiatry*. 2019;9(1):261. doi:10.1038/s41398-019-0593-4

32. Conway CC, Forbes MK, Forbush KT, et al. A Hierarchical Taxonomy of Psychopathology Can Transform Mental Health Research: *Perspect Psychol Sci*. Published online March 7, 2019. doi:10.1177/1745691618810696

33. Eaton NR, Krueger RF, Markon KE, et al. The structure and predictive validity of the internalizing disorders. *J Abnorm Psychol*. 2013;122(1):86-92. doi:10.1037/a0029598

34. Kessler RC, Petukhova M, Zaslavsky AM. The role of latent internalizing and externalizing predispositions in accounting for the development of comorbidity among common mental disorders: *Curr Opin Psychiatry*. 2011;24(4):307-312. doi:10.1097/YCO.0b013e3283477b22

35. Sunderland M, Slade T. The relationship between internalizing psychopathology and suicidality, treatment seeking, and disability in the Australian population. *J Affect Disord*. 2015;171:6-12. doi:10.1016/j.jad.2014.09.012

36. Naragon-Gainey K, Watson D. The anxiety disorders and suicidal ideation: accounting for co-morbidity via underlying personality traits. *Psychol Med*. 2011;41(7):1437-1447. doi:10.1017/S0033291710002096

37. Morey LC, Hopwood CJ, Markowitz JC, et al. Comparison of alternative models for personality disorders, II: 6-, 8- and 10-year follow-up. *Psychol Med*. 2012;42(8):1705-1713. doi:10.1017/S0033291711002601

38. Forbush KT, Hagan KE, Kite BA, Chapa DAN, Bohrer BK, Gould SR. Understanding eating disorders within internalizing psychopathology: A novel transdiagnostic, hierarchical-dimensional model. *Compr Psychiatry*. 2017;79:40-52. doi:10.1016/j.comppsych.2017.06.009

39. Van Os J, Gilvarry C, Bale R, et al. A comparison of the utility of dimensional and categorical representations of psychosis. *Psychol Med*. 1999;29(3):595-606. doi:10.1017/S0033291798008162

40. Caspi A, Moffitt TE. All for One and One for All: Mental Disorders in One Dimension. *Am J Psychiatry*. 2018;175(9):831-844. doi:10.1176/appi.ajp.2018.17121383

41. Garavan H, Bartsch H, Conway K, et al. Recruiting the ABCD sample: Design considerations and procedures. *Dev Cogn Neurosci*. 2018;32:16-22. doi:10.1016/j.dcn.2018.04.004

42. Auchter AM, Hernandez Mejia M, Heyser CJ, et al. A description of the ABCD organizational structure and communication framework. *Dev Cogn Neurosci*. 2018;32:8-15. doi:10.1016/j.dcn.2018.04.003

43. Clark DB, Fisher CB, Bookheimer S, et al. Biomedical ethics and clinical oversight in multisite observational neuroimaging studies with children and adolescents: The ABCD experience. *Dev Cogn Neurosci*. 2018;32:143-154. doi:10.1016/j.dcn.2017.06.005

44. Uban KA, Horton MK, Jacobus J, et al. Biospecimens and the ABCD study: Rationale, methods of collection, measurement and early data. *Dev Cogn Neurosci*. 2018;32:97-106. doi:10.1016/j.dcn.2018.03.005

45. Baurley JW, Edlund CK, Pardamean CI, Conti DV, Bergen AW. Smokescreen: a targeted genotyping array for addiction research. *BMC Genomics*. 2016;17(1):145. doi:10.1186/s12864-016-2495-7

46. Euesden J, Lewis CM, O'Reilly PF. PRSice: Polygenic Risk Score software. *Bioinformatics*. 2015;31(9):1466-1468. doi:10.1093/bioinformatics/btu848

47. Wray NR, Ripke S, Mattheisen M, et al. Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nat Genet*. 2018;50(5):668-681. doi:10.1038/s41588-018-0090-3

48. Demontis D, Walters RK, Martin J, et al. Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. *Nat Genet*. 2019;51(1):63-75. doi:10.1038/s41588-018-0269-7

49. Otowa T, Hek K, Lee M, et al. Meta-analysis of genome-wide association studies of anxiety disorders. *Mol Psychiatry*. 2016;21(10):1391-1399. doi:10.1038/mp.2015.197

50. Ruderfer DM, Ripke S, McQuillin A, et al. Genomic Dissection of Bipolar Disorder and Schizophrenia, Including 28 Subphenotypes. *Cell*. 2018;173(7):1705-1715.e16. doi:10.1016/j.cell.2018.05.046

51. Pardiñas AF, Holmans P, Pocklington AJ, et al. Common schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong background selection. *Nat Genet*. 2018;50(3):381-389. doi:10.1038/s41588-018-0059-2

52. Grove J, Ripke S, Als TD, et al. Identification of common genetic risk variants for autism spectrum disorder. *Nat Genet*. 2019;51(3):431-444. doi:10.1038/s41588-019-0344-8

53. Achenbach TM. Achenbach System of Empirically Based Assessment (ASEBA). In: Cautin RL, Lilienfeld SO, eds. *The Encyclopedia of Clinical Psychology*. John Wiley & Sons, Inc.; 2015:1-8. doi:10.1002/9781118625392.wbecp150

54. Barch DM, Albaugh MD, Avenevoli S, et al. Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale and description. *Dev Cogn Neurosci*. 2018;32:55-66. doi:10.1016/j.dcn.2017.10.010

55. Pagliaccio D, Luking KR, Anokhin AP, et al. Revising the BIS/BAS Scale to study development: Measurement invariance and normative effects of age and sex from childhood through adulthood. *Psychol Assess*. 2016;28(4):429-442. doi:10.1037/pas0000186

56. Luciana M, Bjork JM, Nagel BJ, et al. Adolescent neurocognitive development and impacts of substance use: Overview of the adolescent brain cognitive development (ABCD) baseline neurocognition battery. *Dev Cogn Neurosci*. 2018;32:67-79. doi:10.1016/j.dcn.2018.02.006

57. Eriksen BA, Eriksen CW. Effects of noise letters upon the identification of a target letter in a nonsearch task. *Percept Psychophys*. 1974;16(1):143-149. doi:10.3758/BF03203267

58. Zelazo PD, Anderson JE, Richler J, Wallner-Allen K, Beaumont JL, Weintraub S. II. NIH TOOLBOX COGNITION BATTERY (CB): MEASURING EXECUTIVE FUNCTION AND

ATTENTION: NIH TOOLBOX COGNITION BATTERY (CB). *Monogr Soc Res Child Dev*. 2013;78(4):16-33. doi:10.1111/mono.12032

59. Carlozzi NE, Tulsky DS, Kail RV, Beaumont JL. Vi. Nih Toolbox Cognition Battery (cb): Measuring Processing Speed. *Monogr Soc Res Child Dev*. 2013;78(4):88-102. doi:10.1111/mono.12036

60. Bauer PJ, Dikmen SS, Heaton RK, Mungas D, Slotkin J, Beaumont JL. Iii. Nih Toolbox Cognition Battery (cb): Measuring Episodic Memory. *Monogr Soc Res Child Dev*. 2013;78(4):34-48. doi:10.1111/mono.12033

61. Jorgensen TD, Pornprasertmanit S, Schoemann AM, et al. semTools: Useful tools for structural equation modeling. *R Package Version 05-1*. Published online 2018.

62. McDonald RP. *Test Theory: A Unified Treatment*. Taylor & Francis; 1999.

63. Rosseel Y. lavaan: An R Package for Structural Equation Modeling. *J Stat Softw*. 2012;48(2). doi:10.18637/jss.v048.i02

64. Epskamp S. semPlot: Unified visualizations of structural equation models. *Struct Equ Model*. 2015;22(3):474-483. doi:10.1080/10705511.2014.937847

65. Epskamp S, Cramer AOJ, Waldorp LJ, Schmittmann VD, Borsboom D. qgraph: Network Visualizations of Relationships in Psychometric Data. *J Stat Softw*. 2012;48(1):1-18. doi:10.18637/jss.v048.i04

66. Bogdan R, Baranger DAA, Agrawal A. Polygenic Risk Scores in Clinical Psychology: Bridging Genomic Risk to Individual Differences. *Annu Rev Clin Psychol*. 2018;14(1):119-157. doi:10.1146/annurev-clinpsy-050817-084847

67. Miller PM, Byrne M, Hodges A, Lawrie SM, Johnstone EC. Childhood behaviour, psychotic symptoms and psychosis onset in young people at high risk of schizophrenia: early findings from the Edinburgh High Risk Study. *Psychol Med*. 2002;32(1):173-179. doi:10.1017/S0033291701004779

68. Patalay P, Fonagy P, Deighton J, Belsky J, Vostanis P, Wolpert M. A general psychopathology factor in early adolescence. *Br J Psychiatry*. 2015;207(1):15-22. doi:10.1192/bjp.bp.114.149591

69. Waldman ID, Poore HE, van Hulle C, Rathouz PJ, Lahey BB. External validity of a hierarchical dimensional model of child and adolescent psychopathology: Tests using confirmatory factor analyses and multivariate behavior genetic analyses. *J Abnorm Psychol*. 2016;125(8):1053-1066. doi:10.1037/abn0000183

70. Casey BJ, Cannonier T, Conley MI, et al. The Adolescent Brain Cognitive Development (ABCD) study: Imaging acquisition across 21 sites. *Dev Cogn Neurosci*. 2018;32:43-54. doi:10.1016/j.dcn.2018.03.001

Supplementary Information for Motivation and Cognitive Abilities as Mediators between Polygenic Scores and Psychopathology in Children

Figures below show the results from analyses done using risk alleles at different thresholds from p<.5 to .0001.

Figure S1. Standardized coefficients from the SEM (Figure 2A) testing the relationship between the six polygenic scores (PSs) across PS thresholds and the P-Factor.

Figure S2. Standardized coefficients from the SEM (Figure 2C) testing the relationship between MDD PS across PS thresholds and the proposed mediators.

Figure S3. Standardized coefficients from the SEM (Figure 2D) testing the relationship between ADHD PS across PS thresholds and the proposed mediators.

Figure S4. Standardized coefficients of the indirect effects from the SEM (Figure 3A) testing the mediation between MDD PS across PS thresholds and the P-Factor with the BIS and G-Factor as mediators.

Figure S5. Proportion Mediated (%) from the SEM (Figure 3A) testing the mediation between ADHD PS across PS thresholds and the P-Factor with the BIS and G-Factor as mediators.

Figure S6. Standardized coefficients of the indirect effects from the SEM (Figure 3B) testing the mediation between MDD PS across PS thresholds and the P-Factor with the BIS and G-Factor as mediators.

Figure S7. Proportion Mediated (%) from the SEM (Figure 3B) testing the mediation between ADHD PS across PS thresholds and the P-Factor with the BIS and G-Factor as mediators.

Figure S8. Standardized coefficients from the SEM (Figure 4A) testing the
the relationship between MDD PS across PS thresholds and the five specific dimensions
from the first-order model of psychopathology.

Figure S9. Standardized coefficients from the SEM (Figure 4B) testing the
the relationship between ADHD PS across PS thresholds and the five specific dimensions
from the first-order model of psychopathology.

Figure S10. Standardized coefficients of the indirect effects from the SEM (Figure 5A) testing the mediation between MDD PS across PS thresholds and the specific dimensions with the BIS and G-Factor as mediators.

Figure S11. Proportion Mediated (%) from the SEM (Figure 5A) testing the mediation between MDD PS across PS thresholds and the specific dimensions with the BIS and G-Factor as mediators.

Figure S12. Standardized coefficients of the indirect effects from the SEM (Figure 5B) testing the mediation between ADHD PS across PS thresholds and the specific dimensions with the BAS and G-Factor as mediators.

Figure S13. Proportion Mediated (%) from the SEM (Figure 5B) testing the mediation between ADHD PS across PS thresholds and the specific dimensions with the BAS and G-Factor as mediators.

Markdown File S1. MovCogMedPSPFactor_SEM.html shows R scripts for data pre-processing and latent variable modelling and their detailed outputs.