

Quantifying hospital flows and occupancy due to COVID-19 outbreak in France. Was French lockdown effective?

David Ternant, EA 7501, Université de Tours, Tours, France.

Abstract

Context. The spread of SARS-CoV-2 led to a rapid and deadly pandemic which reached almost all countries in the world in a few months. In most countries, rigorous measures of mitigation, including national or subnational lockdowns were established. The present work aimed at quantifying the effect of national lockdown in France on hospital occupancy.

Methods. A compartmental model describing patient hospital flows was developed, where the effect of lockdown was quantified on the hospitalization rate. Model parameters were estimated using nonlinear mixed-effects (NLME) modelling.

Results. French lockdown led to a hospitalization rate decreased by thrice from 12 days after its beginning. However, lockdown may not to have decreased either hospital occupancy or deaths in hospital, which would have been both decreased by 30% and 85% in average if lockdown was started 20 and 30 days before this date, respectively.

Conclusion. The present work showed an intrinsic effect of lockdown to decrease hospital burden, but its efficacy on hospital burden may have been increased if established sooner.

1 Introduction

After an outbreak in December 2019 in Wuhan (Hubei, China), Coronavirus Disease 2019 (COVID-19) reached the rest of the world in less than two months. The World Health Organization (WHO) declared on February 26 of 2020 that all continents were touched by COVID-19 and declared pandemic situation on March 11. On May 22nd, more than 5 million of cases and 330000 of deaths were recorded: most of deaths were recorded in USA (96363), United Kingdom (36042), Italy (32486), France (28215), Spain (27940) and Brazil (20082) [1].

The report 9 of Ferguson et al. [2] suggested that COVID-19 may trigger not only dramatic number of deaths, but also huge hospital overload and saturation of intensive care units (ICU). Therefore, 87 countries have imposed national or subnational lockdowns [3]. In France, the first COVID-19 cases were confirmed at the end of January, and the first death from COVID-19 was recorded on February 28; a national lockdown, including overseas territories and departments, was begun on March 17th and ended on May 11th in France [4].

Previous works that used compartmental deterministic mathematical models suggested and/or confirmed substantial benefit of lockdown [5–9]. The validity of compartmental deterministic models by themselves is usually agreed. However, since they are most often used as infection and/or death count forecasts very early in epidemic histories, their (numerous) parameters have to be fixed based on prior knowledge, which might lead to over- or under-predictions.

Compartmental models have been used in several fields, including medical research and pharmacology, where compartmental models are usually estimated from data to quantify biologic and/or pharmacologic phenomena, while no (or a few) parameter values are fixed from assumptions. In this field, non-linear mixed-effects (NLME) modelling is used to estimate model parameters with methods based on likelihood maximization or Markov-chain Monte-Carlo (MCMC) methods [11–12].

If most of previous works used sound epidemic and/or mechanistic models to describe COVID-19 outbreak, spread and hospital occupancy, none of them attempted to estimate models parameters to quantify actual patient flows in hospital services and ICU or the influence of lockdown on hospital occupancy. Therefore, the present study proposes to develop a compartmental model describing hospital flows between hospital services, ICU, deaths and return home, to estimate transfer flow rate constants, and to quantify the effect of lockdown in France using NLME modelling.

2 Methods

2.1 Data extraction

The present work was made on French data which were collected from official databases [4]. Data that were considered were hospitalizations in (i) non-ICU units and (ii) in ICU units, (iii) deaths in the hospital, and (iv) returns to home. These data were collected daily from March 1st towards May 22nd of 2020. They were assessed by regions (or sub-regions) selected to keep comparable number of patients within each unit. Indeed, regions were divided in smaller areas if effectives were high, as in Ile-de-France, Grand-Est, Hauts-de-France, Auvergne-Rhône-Alpes and Provence-Alpes-Côtes d'Azur. Therefore, the following 26 regions or sub-regions were assessed : Paris, Hauts-de-Seine, Val-de-Marne, Seine-saint-Denis, Essonne, Yvelines, Val d'Oise, Seine-et-Marne, Haut-Rhin, Bas-Rhin, Moselle, Grand-Est (other), Nord, Hauts-De-France (other), Rhône, Auvergne-Rhône-Alpes (other), Bouches-du-Rhône, Provence-Alpes-Côtes d'Azur (other), Bourgogne-Franche-Comté, Centre-Val-de-Loire, Occitanie, Nouvelle-Aquitaine, Pays-de-la-Loire, Normandie, Bretagne, Corse. Overseas departments and territories were not assessed because of low effectives. No focus was made on COVID-19 cases, because no systematic testing policy was implemented in France up to this date. Since COVID-19 cases were not accounted, the model was made under the assumption of a constant ratio between cases leading to hospitalization, and total cases.

2.2 Hospital flow model




Figure 1. Compartmental model describing patient flows between susceptible (S), non-ICU services (H), in ICU (R), deaths in hospital (D) and return to home (G). Abbreviations are found in the text below.

The hospital flow model is described using a system of 9 ordinary differential equations. (figure 1). This model was obtained after having tested numerous possibilities. First, patient hospitalization was described using a Kermack-McKendrick input function:

$$\frac{ds}{dt} = -k_I \cdot S \cdot \frac{H_1}{N} \quad (\text{E.1})$$

Where S is the number of subjects susceptible to be hospitalized, k_I is hospitalization rate constant, H_1 is the number of hospitalized subjects and N is total number of subjects. Hospitalized patients are assumed to be directed to heal, to die or towards ICU:

$$\frac{dH_1}{dt} = k_I \cdot S \cdot \frac{H_1}{N} - (k_{DH} + k_{H1} + k_{H2}) \cdot H_1 \quad (\text{E.2})$$

Where k_{DH} , k_{H1} and k_{H2} are direction rate constants towards hospital death, hospital heal and ICU, respectively. Because of a delay between hospital admission and death, decease in hospital is described using the following equation:

$$\frac{dH_2}{dt} = k_{DH} \cdot H_1 - k_{DH} \cdot H_2 \quad (\text{E.3})$$

Where H_2 is the number of hospitalized patients in non-ICU services bound to die. Input and output rates were described using the same value because were not differentiable. Because early attempts showed that patients bound to live and to die present different kinetics, they were described using separate equations. Therefore, the kinetics of patients that heal from hospitalization is described as follows:

$$\frac{dH_3}{dt} = k_{H1} \cdot H_2 - k_{G1} \cdot H_3 \quad (\text{E.4})$$

Where H_3 is the number of hospitalized patients (not in ICU) bound to heal. Early attempts showed a delay between admission in hospital and in ICU. In addition, as for hospital patients, patients bound to heal or die in ICU present different kinetics and were therefore described in separate equations. Thus, the following compartment described the transfer of patients toward ICU as follows:

$$\frac{dH_4}{dt} = k_{H1} \cdot H_1 - (k_R + k_{RD}) \cdot H_4 \quad (\text{E.5})$$

Where H_4 is the number of patients in tansit compartment which describes a delay of patient direction towards ICU, and k_R and k_{RD} are direction rate constants toward heal or death paths in ICU, respectively. The kinetics of atients bound to heal for ICU is described as follows:

$$\frac{dR_1}{dt} = k_R \cdot H_4 - k_{G2} \cdot R_1 \quad (\text{E.6})$$

Where R_1 is the number of patients bound to heal in ICU and k_{G2} is ICU heal rate constant. The kinetics of patients bound to die in ICU is described as follows:

$$\frac{dR_2}{dt} = k_{RD} \cdot H_4 - k_{DR} \cdot R_2 \quad (\text{E.7})$$

Where R_2 is the number of patients bound to die in ICU and k_{DR} is ICU death rate constant. Dead (D) and healed (G) patients were described using the following equations:

$$\frac{dD}{dt} = k_{DH} \cdot H_2 + k_{DR} \cdot R_2 \quad (\text{E.8})$$

$$\frac{dG}{dt} = k_{G1} \cdot H_3 + k_{G2} \cdot R_1 \quad (\text{E.9})$$

Initial time ($t=0$) was fixed on the December 31st of 2019, to set January 1st of 2020 as day 1. The system was initialized by estimating the time of the first patient to be hospitalized, i.e. t_{first} such as $H_1(t=t_{\text{first}}) = 1$. All other equations were assumed to be equal to 0 at $t = t_{\text{first}}$. The number of hospitalized patients outside ICU (H), in ICU (R) were respectively assumed to be $H = H_2 + H_3$ and $R = R_1 + R_2$. The influence of French lockdown was assumed to decrease the hospitalization rate constant as follows:

$$k_{IL} = k_I \cdot \alpha \text{ if } t > t_{\text{effect}}$$

$$k_{IL} = k_I \text{ if } t \leq t_{\text{effect}}$$

Where α is a coefficient between 0 and 1 and t_{effect} is the time from which lockdown effect was detected.

2.3 Parameter estimation

The following 13 parameters were estimated: t_{first} , N , k_I , k_{DH} , k_{H1} , k_{H2} , k_{G1} , k_R , k_{RD} , k_{DR} , k_{G2} , α and t_{effect} . Parameter estimation was made using non-linear mixed-effects (NLME) modelling. This approach has been used since the early 70's in the field of quantitative pharmacology [10–11]. The main goals of NLME modelling were to determine the distribution of the values of the parameters of interest in a group of individuals, i.e. regions (and sub-regions) of France; data from all regions or sub-regions were computed simultaneously to estimate (i) the “mean” (referred as “typical”) value of each parameter, (ii) the inter-region variability (referred as “interindividual variance”) and (iii) to estimate parameter values for each region using Bayesian methods.

Software. Hospital data were analysed using MONOLIX-Suite 2019 software. (Lixoft®, Antony, France), which combines the stochastic expectation-maximization (SAEM) algorithm and a Markov Chain Monte-Carlo procedure for likelihood maximization. To ensure the best possible convergence, a large number of iterations (15000 for K1 and 2000 for K2) were used. K1 and K2 refer to the SAEM procedure of MONOLIX, called “iterative kernels”. During K1, the sequence of step sizes is constant, which allows the exploration of the parameter space. During K2, the step sizes decrease to ensure convergence. Three Markov chains were used. Parameter values of regions, sub-regions and whole France were obtained by computing conditional distribution.

Structural model design. Model with and without lockdown effect on k_I were compared using Akaike's information criterion (AIC), defined as: $AIC = OFV + 2.p$, where OFV is the value of the objective function and p is the number of model parameters to estimate. The use of AIC is based on the parsimony principle aiming at a satisfactory fitting of the data with a small number of parameters. The model with the lowest AIC was selected.

Statistical (inter-region) model. The inter-region variability of parameters was described using an exponential model: $\theta_i = \theta_{TV} \cdot \exp(\eta_i)$, where θ_i is the estimated individual parameter, θ_{TV} is the typical value of the parameter and η_i is the random effect for the i^{th} patient. The values of η_i were assumed to be normally distributed with mean 0 and variance ω^2 . The variability of parameter α , assumed between 0 and 1, was described using a probit model.

Error model. Additive, proportional and mixed additive-proportional models were tested. For example, the combined additive-proportional model was implemented as follows: $Y_{O,ij} = Y_{P,ij} \cdot (1 + \varepsilon_{\text{prop},ij}) + \varepsilon_{\text{add},ij}$ where $Y_{O,ij}$ and $Y_{P,ij}$ are observed and predicted j^{th} marker measurements for the i^{th} patient, respectively, and $\varepsilon_{\text{prop},ij}$ and $\varepsilon_{\text{add},ij}$ are proportional and additive errors, which are assumed to follow a Gaussian distribution with mean 0 and variances σ_{prop}^2 and σ_{add}^2 , respectively.

Model evaluation. Models were evaluated graphically using goodness-of-fit diagnostic plots: observed vs. population (PRED) and individual (IPRED) fitted concentrations; population (WRES) and individual (IWRES) weighted residuals vs. PRED and IPRED, respectively.

2.4 Simulations

Typical estimates of model parameters (**table 1**) were used to simulate the effect of lockdown on H, R, D and G data in time. The value of N_0 was converted as its value per 100,000 capita ($N_0* = 170$). The following *scenarii* were simulated: no lockdown, actual lockdown (on the March 17th of 2020), and lockdown started 10, 20 and 30 days before this date.

3 Results

Data of patient hospitalization in non-ICU services (H, figure 2A), in ICU (R, figure 2B), deaths in hospital (D, figure 2C) and return to home (G, figure 2D) were satisfactorily described by the hospital flow model. Best error models were mixed additive-proportional for all measurements. All model parameters were estimated with good accuracy (table 1). No estimation collinearity was observed. The plots of predicted vs. observed measurements showed good model fitting (figure 3). Individual weighted residuals showed no obvious bias or model misspecification (figure 4). Accounting for lockdown model led to a real improve in data description, with a decrease in AIC of 1057.02 (AIC = 81072.13 and 80015.11 without and with lockdown effect, respectively).

If the inter-region variability of parameters related to disease itself was low (< 40% for t_{first} , N_0 and k_I), this variability was much higher for parameters related to hospital flows (between 40% and 95%), and surprisingly high for lockdown effect (130% for α). The lower the α , the higher the lockdown effect on k_I (table 2). For instance, lockdown was very effective in Ile-de-France ($\alpha=0.013$), while less efficient in Grand-Est ($\alpha=0.52$) or Provence-Alpes-Côte d'Azur ($\alpha=0.38$). On the global France data, a poor effect of lockdown was observed ($\alpha=0.89$). Of note, no inter-region correlation was found between parameters related to disease and related to lockdown (α , t_{effect}).

The time for which the first patient was hospitalized in France corresponds to the initiation of the system effect such as $H_1(t=t_{\text{first}}) = 1$, at day 47, which corresponds to the February 17th of 2020. French lockdown started one month later, on March 17th of 2020 (day 76); the effect of lockdown was detectable 12 days after this date in whole France (day 88, table 2), variable among (sub)regions.

Simulations suggested that lockdown did not substantially decrease either hospital occupancy or deaths in hospital. They also suggested that lockdown effect would have been increased if started before its actual date. Indeed, hospital occupancy and deaths in hospital would have been both decreased by 30% and 85% in average if lockdown was started 20 and 30 days before this date, respectively. (figure 5).

Table 1. Parameter estimates

Parameter	Structural model (θ)		Statistical model (ω)		Error model (σ)		
	Estimate	R.S.E. (%)	Estimate	R.S.E. (%)		Parameter	Estimate
t_{first}	47	2.77	0.15	12.6	$\sigma_{\text{add},H}$	17.4	4.4
N_0	7070	7.92	0.40	13.6	$\sigma_{\text{prop},H}$	0.027	3.6
k_I	0.31	2.61	0.14	13.9	$\sigma_{\text{add},R}$	7.93	4.1
k_{DH}	0.014	14.9	0.77	17.1	$\sigma_{\text{prop},R}$	0.054	4.0
k_{H1}	0.026	7.53	0.41	14.4	$\sigma_{\text{add},D}$	9.71	2.4
k_{H2}	0.026	8.66	0.48	13.1	$\sigma_{\text{prop},D}$	0.011	5.2
k_{G1}	0.095	10.8	0.60	14	$\sigma_{\text{add},G}$	31.1	3.2
k_R	0.40	11.5	0.63	14.5	$\sigma_{\text{prop},G}$	0.021	3.9
k_{RD}	0.33	17	0.94	13.6			
k_{G2}	0.11	9.9	0.56	12.7			
k_{DR}	0.24	9.93	0.55	12.9			
alpha	0.17	36.7	1.29	17.5			
t_{effect}	86.7	1.3	0.21	16.3			

Legends. θ , structural parameters; ω , inter-region standard deviation of structural parameters; σ error parameters; R.S.E: relative standard error = standard deviation of estimates / estimate x 100. Other abbreviations are found in the text.




Figure 2A. Observed (clear blue dots) and model-fitted (deep blue line) hospitalizations in non-ICU services in different (sub)-regions




Figure 2B. Observed (clear blue dots) and model-fitted (deep blue line) hospitalizations in ICU in different (sub)-regions




Figure 2C. Observed (clear blue dots) and model-fitted (deep blue line) deaths in hospital in different (sub)-regions




Figure 2D. Observed (clear blue dots) and model-fitted (deep blue line) returns to home in different (sub)-regions.




Figure 3. Model-fitted vs. observed values of hospitalizations in non-ICU services (H), in ICU (R), deaths in hospital (D) and returns at home (G) in log-scale.







Figure 4. Weighted “individual” residuals vs. Gaussian law.

Table 2. Parameter estimates in whole France, and all regions and some sub-regions.

Region	t _{first}	Date first admission	N ₀	N _{0*}	k _I	k _{DH}	k _{H1}	k _{H2}	k _{G1}	k _R	k _{RD}	k _{G2}	k _{DR}	alpha	Lockdown effect delay
France	39	2/9/2020	113897	170	0.33	0.0131	0.025	0.018	0.110	0.34	0.37	0.080	0.21	0.886	12
IDF	46	2/16/2020	46203	376	0.34	0.0120	0.024	0.017	0.086	0.35	0.43	0.073	0.18	0.013	16
Paris	49	2/18/2020	9866	451	0.34	0.0126	0.045	0.028	0.063	0.48	0.69	0.048	0.14	0.415	12
H-D-Seine	53	2/22/2020	7864	490	0.35	0.0122	0.021	0.015	0.094	0.37	0.40	0.073	0.17	0.007	14
V-D-Marne	49	2/19/2020	7371	531	0.30	0.0142	0.029	0.016	0.069	0.42	0.77	0.052	0.22	0.007	19
S-S-Denis	53	2/22/2020	6243	385	0.32	0.0095	0.025	0.013	0.079	0.78	1.55	0.052	0.29	0.003	18
Essonne	54	2/23/2020	3549	274	0.32	0.0113	0.026	0.017	0.048	0.59	0.58	0.076	0.17	0.796	11
Yvelines	53	2/23/2020	4050	282	0.32	0.0139	0.022	0.016	0.098	0.50	0.47	0.090	0.28	0.006	14
Val-Oise	54	2/24/2020	4234	345	0.36	0.0167	0.020	0.019	0.205	0.28	0.28	0.123	0.35	0.644	5
Seine-Marne	58	2/28/2020	3738	266	0.36	0.0044	0.025	0.013	0.113	1.35	2.11	0.099	0.14	0.575	19
Grand-Est	45	2/15/2020	18003	327	0.34	0.0090	0.027	0.020	0.072	0.26	0.32	0.082	0.39	0.518	9
Bas-Rhin	52	2/22/2020	4251	378	0.38	0.0079	0.018	0.021	0.056	0.16	0.10	0.099	0.30	0.035	7
Haut-Rhin	44	2/14/2020	5823	762	0.30	0.0119	0.024	0.014	0.115	0.89	1.84	0.050	1.13	0.001	6
Moselle	53	2/23/2020	4277	410	0.36	0.0030	0.032	0.016	0.053	0.19	0.54	0.051	0.26	0.017	10
GE-other	51	2/21/2020	7782	302	0.34	0.0214	0.035	0.032	0.099	0.46	0.48	0.126	0.26	0.496	9
H-D-France	35	2/5/2020	14359	241	0.23	0.0118	0.022	0.025	0.079	0.72	0.54	0.119	0.27	0.194	15
Nord	51	2/20/2020	5600	215	0.28	0.0137	0.014	0.024	0.094	1.03	0.61	0.110	0.21	0.079	13
HDF-autres	35	2/4/2020	7908	235	0.23	0.0108	0.029	0.026	0.074	0.83	0.78	0.124	0.31	0.405	16
Auvergne-RA	47	2/17/2020	23501	293	0.34	0.0337	0.021	0.038	0.156	0.20	0.09	0.155	0.33	0.296	8
Rhone	50	2/19/2020	8480	1854	0.33	0.0306	0.017	0.034	0.102	0.17	0.07	0.141	0.48	0.266	7
ARA-other	48	2/18/2020	18292	241	0.32	0.0303	0.028	0.038	0.158	0.23	0.12	0.158	0.22	0.257	9
PACA	34	2/4/2020	20170	399	0.28	0.0036	0.058	0.066	0.035	0.31	0.13	0.551	0.07	0.378	12
B-D-Rhone	44	2/13/2020	7209	356	0.26	0.0116	0.034	0.019	0.115	0.21	0.18	0.141	0.09	0.187	12
PACA-autres	35	2/5/2020	6197	204	0.29	0.0076	0.070	0.089	0.036	1.07	0.40	0.409	0.09	0.599	16
Bourgogne-FC	49	2/18/2020	8325	299	0.33	0.0365	0.028	0.035	0.343	0.43	0.34	0.132	0.38	0.534	5
Centre-VDL	56	2/25/2020	5352	209	0.32	0.0276	0.013	0.038	0.040	0.46	0.19	0.205	0.29	0.330	12
Occitanie	44	2/13/2020	13515	228	0.26	0.0022	0.046	0.028	0.073	0.93	0.61	0.096	0.11	0.109	10
Aquitaine	57	2/26/2020	18184	303	0.38	0.0405	0.016	0.045	0.547	0.19	0.06	0.149	0.34	0.189	7
Pays-Loire	53	2/23/2020	7941	209	0.32	0.0313	0.016	0.036	0.149	0.29	0.12	0.214	0.25	0.223	10
Normandie	55	2/24/2020	11530	349	0.31	0.0130	0.013	0.025	0.085	0.19	0.10	0.145	0.23	0.003	9
Bretagne	38	2/8/2020	4560	137	0.24	0.0294	0.025	0.047	0.079	0.50	0.20	0.170	0.36	0.328	15
Corse	44	2/14/2020	1679	487	0.25	0.0393	0.035	0.037	0.188	0.27	0.19	0.138	0.23	0.316	5

Legends: N_{0*} : number of patients bound to be hospitalized per 100,000 capita. Date of first admission is an estimate of the date where the first patient was hospitalized such as H₁(t = t_{first}) = 1, lockdown after delay is the number of days after which lockdown effect is detected since lockdown date on March 17th of 2020. Other abbreviations are found in the text.




Figure 5. Simulations of hospitalizations in non-ICU services (H), in ICU (R), deaths in hospital (D) and returns at home (G) for different lockdown strategies. The value of N_0 was converted as its value per 100,000 capita ($N_0^*=170$). Lockdown effect was detectable at day 88, i.e. 12 days after start of lockdown.

4 Discussion

This work has focused on available data to estimate the actual effect of lockdown on hospitalization burden in regions and sub-regions of France. This model allowed an accurate description of patient flows between non-ICU services, ICU, deaths at hospital and return to home. Adding the effect of lockdown allowed an even better description of hospital data.

A previous work suggested that French lockdown would have prevented almost 600,000 hospital admissions, including almost 150,000 in ICU [9]. However, this work has been criticized, because of a possible over-estimation of hospital burden [12]. The principle of lockdown itself remains controversial. Introduced for the first time in human history in 2020 instead of isolation or quarantine, it may lead to a large social cost while being less efficient than isolation or quarantine measures [13]. To deal with this controversy, the present hospital flow model was attempted to either validate or invalidate the following hypothesis: “lockdown has decreased hospital burden in France”. This necessitated a sound description of patient input/flows/output in hospital.

This study showed a significant effect of lockdown on hospitalization rate (k_l), with a global decrease of k_l by thrice. However, the consequences of lockdown appeared to be minor: simulations made from the hospital flow model suggested that lockdown have had no significant influence on hospital burden and would have saved less than 1 life per 100,000 capita, compared to the 22 per 100,000 who have died until the day May 22nd [14]. There may be at least three explanations: first, the effect of lockdown is delayed in time (detected 12 days after start of lockdown in average), second, hospital occupancy was already high at this date and, last, but not least, the risk of infection may have already started to decline [14]. Nevertheless, the effect of lockdown, as well as patient flow parameters, was found to be very variable among regions and sub-regions of France. This variability may be due, at least in part, to population density in each region, different hospital cares, different population characteristics (inter-region differences in mean age, co-morbidities, etc.), and/or to the respect of mitigation strategies.

If an intrinsic effect of lockdown measures seems indisputable, one could question the date of establishment. Indeed, simulations showed that both hospital occupancy and deaths would not have been significantly different if no lockdown was established, but they might have been dramatically decreased if lockdown had been started 20 or 30 days started before the official date of application. Nevertheless, at these days, the first death from COVID-19, and the first case were respectively recorded.

This work has nevertheless limitations. First, all French sub-regions were not considered, in order to keep N_0 sufficiently high in each region unit and to avoid very long time of computation. In particular, overseas territories were not considered, except for Corse. Second, even if accurate values of model parameters were obtained in each region, notably the time of the first hospitalized patient, no spatial description of disease spread was made. Third, this model does not pretend to be an epidemic model: the hospital admission growth was described using a Kermack-McKendrick input function, which may appear oversimplistic. This also concerns the “on/off” model for lockdown effect, which may not account for the complexity of the real effect of

lockdown. Therefore, this model cannot test lockdown ending strategies and their consequences on the risk and amplitude of epidemic rebounds. To describe or forecast these rebounds, a mechanistic epidemic model as Roux et al. [9] could elegantly replace the present input function. Last but not least, this model is “macroscopic”, i.e. does not account for patient specificities (as age distribution or proportion of comorbidities). Taking into account these factors may improve the description of inter-region variability [14]. Given these limitations, the lockdown effect should be considered with caution, this study providing trends more than a sound quantification.

In conclusion, the present model has shown an intrinsic effect of lockdown to decrease hospital burden. However, the efficiency of lockdown may have been increased if established sooner. This may raise important questions, notably regarding social acceptability of these rigorous measures that would have imposed while no strong signal of danger was present.

References

1. COVID-19 Coronavirus Pandemic. 2020. <https://www.worldometers.info/coronavirus/>
2. Neil M Ferguson et al. Report 9: Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality and healthcare demand, 2020. <https://www.imperial.ac.uk/media/imperial-college/medicine/mrc-gida/2020-03-16-COVID19-Report-9.pdf>
3. COVID-19 Pandemic Lockdowns (Wikipedia). https://en.wikipedia.org/wiki/COVID-19_pandemic_lockdowns.
4. Suivi de l'épidémie de COVID-19 en France. 2020. <https://dashboard.covid19.data.gouv.fr/>
5. Patrick GT Walker et al. The Global Impact of COVID-19 and Strategies for Mitigation and Suppression, 2020 <https://www.imperial.ac.uk/media/imperial-college/medicine/sph/ide/gida-fellowships/Imperial-College-COVID19-Global-Impact-26-03-2020v2.pdf>
6. Fizza Farooq et al. Effect of Lockdown on the spread of COVID-19 in Pakistan. 2020. <https://arxiv.org/ftp/arxiv/papers/2005/2005.09422.pdf>
7. Osmar Pinot Neto et al. Mathematical model of COVID-19 intervention in Sao Paulo – Brazil. 2020. <https://arxiv.org/ftp/arxiv/papers/2005/2005.09426.pdf>.
8. R. Adhikari et al. Inference, prediction and optimization of non-pharmaceutical interventions using compartment models: the PyRoss library. 2020. <https://arxiv.org/pdf/2005.09625.pdf>
9. Jonathan Roux et al. COVID-19: One-month impact of the French lockdown on the epidemic burden. 2020. <https://www.ehesp.fr/wp-content/uploads/2020/04/Impact-Confinement-EHESP-20200322v1-1.pdf>
10. Diane R Mould, et al. Basic concepts in population modeling, simulation, and model-based drug development. CPT Pharmacometrics Syst Pharmacol. 2012;1:e6
11. Diane R Mould et al. Basic Concepts in Population Modeling, Simulation, and Model-Based Drug Development-Part 2: Introduction to Pharmacokinetic Modeling Methods CPT Pharmacometrics Syst Pharmacol. 2013;2:e38
12. Tribune : Etude critique d'une modélisation des effet du confinement. Journal international de médecine, 05/02/2020 https://www.jim.fr/medecin/jimplus/e-docs/etude_critique_dune_modelisation_des_effets_du_confinement_182808/document_edito.phtml
13. M. Zelmat. Facing COVID-19 by the confinement :EBM, "MBM" or "SBM" ? 2020. https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3600511
14. Juanjuan Zhang. Fundamental principles of epidemic spread highlight the immediate need for large-scale serological surveys to assess the stage of the SARS-CoV-2 epidemic. 2020. https://www.reddit.com/r/COVID19/comments/fonnt8/fundamental_principles_of_epidemic_spread/