

**TITLE : PRE EXPOSURE HYDROXYCHLOROQUINE PROPHYLAXIS FOR COVID-19
IN HEALTHCARE WORKERS: A RETROSPECTIVE COHORT.**

AUTHOR INFORMATION :

Dr. Raja Bhattacharya ¹, Dr. Sampurna Chowdhury ², Dr. Rishav Mukherjee ², Dr. Anita Nandi ³,
Dr. Manish Kulshrestha ⁴, Dr. Rohini Ghosh ⁴, Mr. Souvik Saha ⁵

1. Associate Professor, Department of General Medicine, Medical College, Kolkata

2. Post Graduate Trainee, Department of General Medicine, Medical College, Kolkata

3. Assistant Professor, Department of Microbiology, Medical College, Kolkata

4. Intern, Medical College, Kolkata

5. MSc Statistics, Department of Statistics, Presidency University

CORRESPONDING AUTHOR

Dr. Anita Nandi, Assistant Professor, Department of Microbiology

Medical College, Kolkata

88, College Street, Kolkata, India

email: anitanandi20121964@gmail.com

Contact number: +91 9163124733/9831018416

RUNNING TITLE : Pre exposure Hydroxychloroquine Prophylaxis for COVID-19

ORCID ID

Dr. Sampurna Chowdhury- <https://orcid.org/0000-0002-5010-7565>

Dr. Rishav Mukherjee- <https://orcid.org/0000-0001-5883-2908>

WORK ATTRIBUTED TO - Department of General Medicine & Department of Microbiology , Medical College Kolkata, India

AUTHOR CONTRIBUTIONS

RB- Concept,Data interpretation,First draft preparation

SC- Concept, Data Collection & Interpretation of results, Literature Review.

RM-Concept,Data Collection,Literature Review, Coordination.

AN- Data Analysis, Literature Review.

MK- Data Collection,Proof Reading & Draft preparation.

RG- Draft Preparation and Proof reading.

SS- Statistical Analysis.

FUNDING

This was a non-sponsored study with no external funding.

ETHICAL CONSIDERATION

The research protocol was approved by the Institutional Ethics Committee, Medical College, Kolkata and all participants gave written informed consent. Data was redacted and anonymised data was stored in a password protected computer.

DISCLOSURE OF CONFLICT OF INTERESTS

The authors have no conflict of interests in this study and have nothing to disclose.

ACKNOWLEDGEMENT

We acknowledge the dedication, commitment, and sacrifice of the staff and personnel in our institution during the COVID-19 crisis and the suffering and loss of our patients as well as in their families and our community. We would like to extend our gratitude to the Department of Medicine and Department of Microbiology, Medical College, Kolkata for supporting us.

ABSTRACT

BACKGROUND OF STUDY-

While several trials are ongoing for treatment of COVID-19, scientific research on chemoprophylaxis is still lacking even though it has potential to flatten the curve allowing us time to complete research on vaccines.

OBJECTIVES OF THE STUDY

To explore the potential of HCQ as a pre-exposure prophylaxis for COVID 19 in health care workers in a tertiary care hospital.

MATERIALS & METHODS

We have conducted a retrospective cohort study among 106 Health Care Workers (HCW) exposed to COVID-19 patients, at a tertiary care hospital in India where there was an abrupt cluster outbreak within on duty personnel. HCWs who had voluntarily taken hydroxychloroquine (HCQ) prior to exposure were considered one cohort while those who had not were considered to be the Control group. All participants with a verifiable high risk contact history were tested for COVID-19 by RT-PCR.

RESULTS

The two cohorts were comparable in terms of age, gender, co-morbidities and exposure. The primary outcome was incidence rates of RT-PCR positive COVID-19 infection amongst HCQ users and Controls. In this retrospective cohort study, 106 healthcare workers were examined of

whom 54 were HCQ users and rest were not. The comparative analysis of incidence of infection between the two groups demonstrated that voluntary HCQ usage was associated with lesser likelihood of developing SARS-CoV-2 infection (4 out of 54 HCW), compared to those who were not on it (20 out of 52 HCW), $\chi^2 = 14.59$, $p < 0.001$. None of the HCQ users noted any serious adverse effects.

CONCLUSIONS

This study demonstrated that voluntary HCQ consumption as pre-exposure prophylaxis by HCWs is associated with a statistically significant reduction in risk of SARS-CoV-2. These promising findings therefore highlight the need to examine this association in greater detail among a larger sample using Randomised Controlled Trials (RCT).

KEYWORDS : Covid19, hydroxychloroquine, pre-exposure prophylaxis, healthcare workers

INTRODUCTION

SARS-CoV-2 is a positive sense RNA virus of the family *Coronaviridae* and the etiological agent responsible for the novel pneumonia (COVID-19) which triggered an outbreak towards the end of 2019 in Wuhan, China. Since then, there has been an unprecedented spread of the disease and COVID-19 was declared a ‘global pandemic’ by the WHO on March 11, 2020. As of May 24, 2020 it has affected 5.4 million people and caused 344,419 deaths.¹ In a rapidly evolving situation such as this, we have often had to resort to repurposing old drugs to expedite the process of prevention or treatment of an unfamiliar disease. For treating viral illnesses, often there is a “one bug - one drug” approach which however fails in times of emerging and re-emerging infections. This is especially true during pandemics - when drug discovery races against time. An alternative approach is, to use a broad spectrum antiviral as a pandemic tool.²

To this effect, even before this recent crisis, at least 108 Broad Spectrum Antivirals (BSA) had already been identified that had shown activity against 78 viruses.³ These BSAs include 4-aminoquinoline compounds (chloroquine, hydroxychloroquine, amodiaquine) amongst others and have been known to prevent viral entry into host cell.^{3,4} In this study we have chosen to focus on Hydroxychloroquine - particularly it’s role, as an antiviral, in prevention of COVID-19. Hydroxychloroquine (HCQ) is a derivative of Chloroquine (CQ), formulated by introducing a hydroxyl group into CQ and was demonstrated to be much less (~40%) toxic than CQ in animals.⁵ Both share similar chemical structures and mechanisms of action as a weak base and

immunomodulator and have exhibited, in vitro, potent antiviral properties against various viruses.³

They are both concentrated in organelles with low pH like the lysosomes and endosomes hence are also called lysosomotropic agents.⁴ CQ increase lysosomal pH and prevent its fusion with auto-phagosomes in vitro.⁶ It can also inhibit endosomal acidification, thus preventing viral entry into host cells.⁷ A proposal to use CQ as a candidate drug for influenza virus was already in place. In fact in Mouse models, CQ and its derivative HCQ had already been used successfully against Avian Influenza.⁸ However this success was unfortunately not replicated in Influenza A or B and RCTs⁹ failed to demonstrate any preventive role in them. In the case of Coronaviruses, for SARS CoV1 , CQ and its derivatives were found to show strong antiviral properties in vitro. Apart from interfering with lysosomal and endosomal activities, it also inhibits terminal glycosylation of ACE2 receptor¹⁰ which is involved in viral entry.^{11,12} Impaired terminal glycosylation of ACE2 may reduce the binding efficiency between ACE2 on host cells and the SARS-CoV spike protein.¹³ Moreover, Both CQ and HCQ blocked the transport of SARS-CoV-2 from early endosomes to early lysosomes,¹⁴ which appears to be a requirement to release the viral genome as in the case of SARS-CoV.¹⁵

Similar to SARS-CoV-1, SARS-CoV-2 also utilises the surface receptor ACE2¹⁶ for cellular entry. In vitro data have also demonstrated that CQ as well as its derivative HCQ are potent inhibitors of SARS-CoV-2.¹⁴ In spite of its in vitro success in inhibiting SARS-CoV-2, much like in the case of SARS-CoV-1 clinical trials have failed to show any benefit of HCQ as a therapy for SARS-CoV-2 infection.¹⁷

However, data on its Prophylactic role is still incomplete. Noting the in vitro data and theoretical benefits of HCQ usage, ICMR had proposed its prophylactic use for Health Care Workers in India.¹⁸

This retrospective cohort study explores the usage of HCQ in a tertiary health care centre in India amongst healthcare workers and investigates its prophylactic potential in prevention of COVID-19 infection.

METHODS

STUDY DESIGN AND PATIENT SELECTION

This is a retrospective cohort study based on an online survey of 106 health care personnel, who worked at Medical College, Kolkata, a tertiary care teaching hospital in India, dealing with COVID-19 patients and non COVID patients (mixed COVID Facility), in the first two weeks of May, 2020. In the given period, a cluster outbreak of cases amongst HCWs in this hospital had occurred - with about 28 HCW testing positive over a period of two weeks.

Since late March, Indian Council of Medical Research (ICMR)¹⁸, which is the apex body of medical research in India, has proposed consumption of HCQ for prophylaxis against COVID-19. In accordance with that guideline, some of the HCWs were voluntarily on Pre-exposure HCQ prophylaxis whereas few others were not. After the outbreak was identified, all those who fulfilled the high risk contact criteria were quarantined and tested for COVID-19 between Day 7-Day 14 of suspected exposure as per Ministry of Health and Family Welfare (MoHFW), Government of India, guidelines.¹⁹ The end results of these tests were recorded and

COVID-19 positive health care workers were sent for isolation and treatment to appropriate facilities after proper contact tracing. We conducted an online survey amongst exposed healthcare workers after their test reports were available. To be included in the survey the participant needed to be a Health Care Worker, currently working at the tertiary care centre and on duty at the same centre during the last 1 month preceding the survey, and was tested for COVID-19 by RT-PCR in the same hospital.

Two groups of HCWs were identified from the date of publishing of the ICMR guidelines¹⁸ for HCQ prophylaxis (March 23, 2020) till the date on which participants were exposed to their first COVID-19 positive patient. During this period those who started taking HCQ prophylaxis were considered as HCQ users and those who did not take HCQ prophylaxis were considered to be the control group. **(FIGURE 1)**

HCQ users included all the health care workers who were high risk contacts of COVID-19 positive cases & were tested for SARS-CoV-2 Infection in the hospital during the study period, and history of intake of at least the loading dose of hydroxychloroquine prophylaxis and were at any point in their scheduled weekly maintenance dose, as per ICMR guidelines.¹⁸

Control group included all the health care workers who were high risk contacts of COVID-19 positive cases and were tested for SARS-CoV-2 Infection in the hospital during the same study period, and had either no history of hydroxychloroquine prophylaxis or had history of inadequate intake of HCQ (inadequate loading dose, inadequate schedule maintenance dose or missed doses) as per ICMR guidelines.¹⁸

The exclusion criteria were refusal to give consent for the study, not being a contact of COVID-19 positive case as per definition given by the National Centre of Disease Control, India (NCDC).¹⁹ The outcome of Interest was to see whether voluntary HCQ prophylaxis was related to positivity rates by RT- PCR in Health Care Workers.




Figure 1: Participant flow chart showing the study cohort.

TESTING FOR SARS-COV-2 INFECTION

Detection of SARS-CoV-2 in clinical specimens were done by RT-PCR using TaqPath COVID-19 Combo Kit (Applied Biosystems). The real time assay uses the TaqMan fluorogenic probe based chemistry that uses the 5' nuclease activity of Taq DNA polymerase and enables the detection of a specific PCR product as it accumulates during PCR cycles. COVID-19 Real Time PCR Assay Multiplex-Multiplexed assays that contain three primer/probe sets specific to different SARS-CoV-2 genomic regions and primers/probes for phage MS2 (Internal process control for nucleic acid extraction).²⁰ Samples with a result of SARS-CoV-2 Inconclusive were tested again and all were found to be negative and thus, were treated as negative.

DATA COLLECTION

An internet-based retrospective cohort study was designed where participants volunteered to provide data either on an online form or over the telephone. The online survey form was circulated via online messenger services amongst health care workers who were encouraged to recirculate the same amongst their colleagues. After satisfying the inclusion criteria, a total of 106 participants were considered.

The survey included data on demographic profile like age, sex, presence of co-morbidities (defined as Hypertension, Diabetes, Coronary Artery Disease, Chronic Kidney Disease & COPD),^{21,22,23} and use of hydroxychloroquine prophylaxis as per ICMR guidelines.¹⁸ Data on type of exposure, nature of contact, use of PPE was also collected in the survey.

STATISTICAL ANALYSIS

Data was analysed by means of R Software and STATA version 13.1 (StataCorp LP, College Station, TX, USA). Statistical analysis was performed by using test of means of two populations viz. Welch two independent samples T-test, Categorical Data Analysis in the form of contingency tables, partial contingency tables, Chi-squared tests of independence, Chi-squared test of homogeneity, large sample tests of proportion viz. one sample test of proportion and two sample test for proportions. Controlling for possible confounders was done by stratification.

A P value of less than 0.05 was considered statistically significant.

RESULTS

The two cohort groups of those taking hydroxychloroquine (HCQ) and those not taking hydroxychloroquine (control) were comparable in terms of age, gender, degree of exposure and type of exposure and co-morbidities. (**TABLE 1**). The mean number of COVID-19 cases with whom the workers had come in contact was also found to be the same in the two groups by a Welch two sample t-test which was found to be 3.

TABLE 1: Baseline characteristics of participants in the two cohort groups

Parameters	HCQ	Control	p value*
	N=54	N=52	
Age (mean ± SD)	26.46 ± 3.93	27.71 ± 7.24	0.13
Gender			
Male	28 (51.85%)	24 (46.15%)	0.22
Female	26 (48.15%)	28 (53.85%)	
Degree of Exposure			
Face to face contact	51 (94.44%)	41 (78.85%)	0.85
Direct contact	29 (53.70%)	22 (42.31%)	0.70
Environmental contact	23 (42.59%)	19 (36.54%)	0.28
Type of Exposure			
Asymptomatic contact	22 (40.74%)	19 (36.54%)	0.48
Symptomatic contact	32 (59.26%)	33 (63.46%)	
Comorbidities	2 (3.70%)	2 (3.85%)	0.92


Footnote: *Continuous data was compared using t-test and categorical data was compared using Chi-squared test for homogeneity.

In the HCQ group 4 out of 54 participants were tested to be COVID19 positive, whereas, in the control group 20 out of 52 participants were found to be COVID19 positive. Distribution of HCQ takers and controls across outcome of COVID19 test in univariate analysis indicated the association of risk (Relative Risk = 0.193; 95% CI = 0.071-0.526; $p = 0.001$) of SARS-CoV-2 infection with lack of pre-exposure HCQ prophylaxis. In this study, it was seen that taking pre-exposure HCQ prophylaxis was associated with an 80.7% reduction in the risk of acquiring a SARS-CoV-2 infection.

Further analysis of incidence of infection between the two groups demonstrated that the incidence of SARS-CoV-2 in those on HCQ pre-exposure prophylaxis was significantly less when compared to the control group with a χ^2 value of 14.59, $p < 0.001$.

Adverse effects, mostly mild, i.e not requiring any intervention medical or otherwise, not requiring any hospitalisation, were noted in 29.8% of those on HCQ Prophylaxis (**FIGURE 2**), thus corroborating previous treatment trials.²⁴ Among those taking hydroxychloroquine prophylaxis, GI upset (19.1%) , skin rash (6.4%) and headache (4.3%) were reported as adverse effects. GI upset was found to be the most common adverse effect by paired proportion tests. None of the study participants reported discontinuing the drug at the time of survey.

Figure 2: Pie chart showing the adverse effects among those taking HCQ Prophylaxis



Univariate analysis of distribution of COVID-19 test results in the two cohort groups for different exposures was done. (**TABLE 2, TABLE 3**). It was also seen that the healthcare workers who were exposed to symptomatic COVID19 positive patients were more likely to develop SARS-CoV-2 infection compared to those exposed to an asymptomatic COVID19 patient (Odds ratio = 6.046; 95% CI = 1.672 to 21.858; $p = 0.006$).

TABLE 2: Univariate analysis of distribution of COVID-19 test results for different exposures.

Parameters		COVID-19 positive (n=24)	COVID-19 negative (n=82)	OR	95% CI of OR	p value
Symptomatic contacts	Yes	21	44	6.046	1.672 to 21.858	0.006
	No	3	38			
Face to face contact	Yes	20	72	0.694	0.197 to 2.451	0.571
	No	4	10			
Direct contact	Yes	14	37	1.703	0.678 to 4.276	0.257
	No	10	45			
Environment contact	Yes	8	34	0.706	0.271 to 1.836	0.475
	No	16	48			
Recommended PPE*	Yes	11	51	0.514	0.205 to 1.289	0.156
	No	13	31			

Footnote: OR: Odds ratio, CI: Confidence interval, *Recommended as per guidelines on rational use of personal protective equipment (PPE) in a hospital setting by the Ministry of Health and Family Welfare, Government of India.³²

TABLE 3: Two way frequency tables showing distribution of COVID-19 test results in the two cohort groups after restricting observations for different parameters

Parameters		COVID-19 positive (n=24)	COVID-19 negative (n=82)	TOTAL	OR	95% CI of OR	p value
Symptomatic contacts	HCQ+	3 (9.38%)	29 (90.63%)	32 (100%)	2.172	0.211-22.369	0.514
	HCQ-	18 (54.55%)	15 (45.45%)	33 (100%)	10.200	2.024-51.413	0.005
Face to face contact	HCQ+	4 (7.84%)	47 (92.16%)	51 (100%)	0.663	0.029-14.972	0.796
	HCQ-	16 (39.02%)	25 (60.98%)	41 (100%)	1.120	0.282-4.450	0.872
Direct contact	HCQ+	3(10.34%)	26(89.66%)	29(100%)	2.769	0.269-28.468	0.392
	HCQ-	11(50.00%)	11(50.00%)	22(100%)	2.333	0.743-7.324	0.147
Environment contact	HCQ+	0(0.00%)	23(100%)	23(100%)	0.130	0.007-2.543	0.179
	HCQ-	8(42.11%)	11(57.89%)	19(100%)	1.273	0.401-4.037	0.682
Recommended PPE*	HCQ+	3(7.69%)	36(92.31%)	39(100%)	1.167	0.112-12.183	0.898
	HCQ-	8(34.78%)	15(65.22%)	23(100%)	0.756	0.244-2.345	0.628

Footnote: OR: Odds ratio, CI: Confidence interval, *Recommended as per guidelines on rational use of personal protective equipment (PPE) in a hospital setting by the Ministry of Health and Family Welfare, Government of India.³²

DISCUSSION

The COVID-19 pandemic has put healthcare systems across the world in crisis with significant social and economic burden on countries. A number of clinical trials²⁵ are underway to test the efficacy of several repurposed drugs including chloroquine (CQ), hydroxychloroquine (HCQ), Ivermectin, Remdesivir, Ritonavir/Lopinavir for treatment of COVID-19. None have so far shown exceptional results.^{26,24,27} Of these, although HCQ has gathered particular worldwide attention based on in vitro results that demonstrate efficacy against SARS-CoV-2, there is lack of evidence to suggest that HCQ offers any significant additional clinical benefit for the treatment of hospitalised COVID-19 patients.

Although unprecedented progress has been made amidst this pandemic towards development of vaccines, including three candidate vaccines already in clinical trials, most estimates place the timeline for the launch of a safe and effective vaccine at least more than a year away. This highlights the need for possible alternative vaccines for preventing COVID-19. Chemoprophylaxis has been demonstrated to be a successful modality in preventive medicine in a number of other infectious diseases including malaria, HIV and influenza. A mathematical model²⁸ exploring effectiveness of prophylaxis vs treatment in an Influenza pandemic predicted that targeted prophylaxis could flatten the curve by 6-18 months during which effective vaccines against the disease could be developed. However, despite chemoprophylaxis being a promising modality, research on this subject in the context of COVID-19 is currently a missing link. The

first in vitro data on prophylaxis against COVID-19 by HCQ or CQ was put forward by Yao et al²⁹ In this study, HCQ was found to be more potent than CQ in achieving EC50. Several other in vitro studies^{14,30} found a preventive role for CQ/ HCQ. As discussed in the introduction of this article, COVID-19 initiates viral entry through a surface ACE2 receptor, initial viral replication is followed by systemic inflammation. CQ/ HCQ can prevent viral entry has been demonstrated by Wang et al³⁰ through their time-of-addition studies. Based on these encouraging preclinical trials and supported further by preliminary internal observational studies,³¹ ICMR, which is the key government body handling India's COVID response, recommended the use of HCQ by high risk individuals such as HCW for prevention of COVID-19. Given the well-established safety profile in rheumatology practice with HCQ being safe even during pregnancy, ICMR's recommendation is justified, taking into consideration the substantial risks faced by healthcare workers.

The current retrospective cohort study therefore aimed to investigate if hydroxychloroquine could be effective as a pre-exposure prophylaxis for COVID-19 among high risk individuals such as healthcare workers. The results from this study, demonstrate that voluntary consumption of HCQ as prophylaxis among high risk individuals was associated with a significantly reduced risk of testing positive for COVID19. Incidentally, one COVID-19 positive participant who was a long-term user of HCQ for Rheumatoid Arthritis, reported discontinuing the drug 1 month prior to exposure. It was also observed that among HCQ takers 32 were exposed to a symptomatic contact out of which only 3 (9.38%) were positive whereas in the control group 33 were exposed to a symptomatic contact out of which 18 (54.55%) tested to be positive. Similarly, it was seen that proportions of those testing to be positive were less in HCQ takers

among those having a face to face contact, direct contact and environmental contact. Those who used recommended³² personal protective equipment (PPE), even among them, 3 out of 39 (7.69%) HCQ takers were tested to be positive whereas 8 out of 23 (34.78%) controls acquired the infection. (Table 3). Those exposed to a symptomatic patient were more likely to acquire the infection than those exposed to asymptomatic patients (OR = 6.046; 95% C I= 1.672 to 21.858; p = 0.006). These observations need to be further analysed by future studies. The current study also validated the known safety profile for HCQ with no serious adverse events reported by the participants. This study represents a clinical evidence on the potential role of HCQ as prophylaxis.

Our study design also aimed to include those members in the control group who were taking inadequate dosage of HCQ, as per ICMR guidelines. However, all participants were either taking HCQ as per guidelines or not taking at all. Thus, no data on inadequate HCQ dosage could be analysed. Despite a strong statistical association, a cause-effect relationship can not be inferred. The data included young , healthy individuals with no or limited relevant underlying health conditions, reflecting the demographics of health care workers . This warrants a conservative interpretation of our findings and the statistical significance needs to be explored further. Thus, we recommend further validation through large scale RCT to interrogate the significant association between voluntary HCQ consumption and reduced clinical risk of contracting COVID-19 that emerged from this study.

LIMITATIONS

Non random sampling was done based on a voluntary response online survey and thus sampling bias and recall bias might be present in the data set. Being a retrospective cohort study it is limited in its ability to measure all confounding factors. Blinding was not done at any level.

At the time this study was carried out, no prevalence data was available in mixed Covid facilities catering to both covid and non-covid patients. Due to an unmet need for data, a sample size calculation was not appropriate. Moreover, transmission of COVID 19 to healthcare workers in this setting was not only possible from patients alone but also from infected coworkers often sharing common areas of the hospital such as lounges, living quarters, dining spaces etc.

A therapeutic dose monitoring for Hydroxychloroquine was not carried out owing to the questionnaire based nature of the study. Study participants were expected to follow Indian Council of Medical Research's guideline on hydroxychloroquine prophylaxis - however, validation of the dose and its frequency was beyond the scope of the present study.

Among those not taking HCQ a higher number of positives might be because of their risk taking behaviour that prevents them from taking prophylaxis and protective measures as well. On the contrary it might be so that there is a false sense of protection among HCQ takers that predisposes them to more risk-taking behaviour.

The adverse effects which require special assessment like bradycardia and prolonged QT interval could not be recorded and the data was based solely on patient history.

CONCLUSION

In this retrospective cohort study involving healthcare workers who had been exposed to SARS-CoV-2, voluntary pre-exposure Hydroxychloroquine use was associated with a statistically significant lowered risk of testing positive for SARS-CoV-2. No serious adverse effects were found among those taking HCQ Prophylaxis.

ABBREVIATIONS

HCQ- Hydroxychloroquine

CQ- Chloroquine

ICMR- Indian Council Of Medical Research

MOHFW- Ministry of Health & Family Welfare

PPE- Personal Protective Equipment

PCR- Polymerase Chain Reaction

REFERENCES

1. Worldometers.info, Dover, Delaware, U.S.A. *COVID-19 Coronavirus Pandemic*. [Online]

2020 [Cited May 27,2020] . Available from:

<https://www.worldometers.info/coronavirus>.

2. Adalja A, Inglesby T. Broad-Spectrum Antiviral Agents: A Crucial Pandemic Tool. *Expert Rev*

Anti Infect Ther. 2019 Jul;17(7):467-470. <https://doi.org/10.1080/14787210.2019.1635009>

3. Ianevski A, Andersen PI, Merits A, Bjørås M, Kainov D. Expanding the activity spectrum of

antiviral agents. *Drug Discov Today*. 2019 May;24(5):1224-1228. [https://doi.org/10.1016/](https://doi.org/10.1016/j.drudis.2019.04.006)

[j.drudis.2019.04.006](https://doi.org/10.1016/j.drudis.2019.04.006)

4. Al-Bari MAA. Targeting endosomal acidification by chloroquine analogs as a promising

strategy for the treatment of emerging viral diseases. *Pharmacol Res Perspect*. 2017 Jan

23;5(1):e00293. <https://doi.org/10.1002/prp2.293>

5. McChesney EW. Animal toxicity and pharmacokinetics of hydroxychloroquine sulfate. *Am J*

Med. 1983 Jul 18;75(1A):11-8. [https://doi.org/10.1016/0002-9343\(83\)91265-2](https://doi.org/10.1016/0002-9343(83)91265-2)

6. Long L, Yang X, Southwood M, Lu J, Marciniak SJ, Dunmore BJ, et al. Chloroquine prevents

progression of experimental pulmonary hypertension via inhibition of autophagy and lysosomal

bone morphogenetic protein type II receptor degradation. *Circ Res*. 2013 Apr 12;112(8):

1159-70. <https://doi.org/10.1161/circresaha.111.300483>

7. Savarino A, Boelaert JR, Cassone A, Majori G, Cauda R. Effects of chloroquine on viral infections: an old drug against today's diseases? *Lancet Infect Dis*. 2003 Nov;3(11):722-7. [https://doi.org/10.1016/s1473-3099\(03\)00806-5](https://doi.org/10.1016/s1473-3099(03)00806-5)
8. Yan Y, Zou Z, Sun Y, Li X, Xu KF, Wei Y, et al. Anti-malaria drug chloroquine is highly effective in treating avian influenza A H5N1 virus infection in an animal model. *Cell Res*. 2013 Feb;23(2):300-2. <https://doi.org/10.1038/cr.2012.165>
9. Borba EF, Saad CG, Pasoto SG, Calich AL, Aikawa NE, Ribeiro AC, et al. Influenza A/H1N1 vaccination of patients with SLE: can antimalarial drugs restore diminished response under immunosuppressive therapy? *Rheumatology (Oxford)*. 2012 Jun;51(6):1061-9. <https://doi.org/10.1093/rheumatology/ker427>
10. Vincent MJ, Bergeron E, Benjannet S, Erickson BR, Rollin PE, Ksiazek TG, et al. Chloroquine is a potent inhibitor of SARS coronavirus infection and spread. *Virology*. 2005 Aug 22;2:69. <https://doi.org/10.1186/1743-422x-2-69>
11. Berne MA, Somasundaran M, Sullivan JL, Luzuriaga K, Greenough TC, Choe H, et al. Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature*. 2003 Nov 27;426(6965):450-4. <https://doi.org/10.1038/nature02145>
12. Freund NT, Roitburd-Berman A, Sui J, Marasco WA, Gershoni JM. Reconstitution of the receptor-binding motif of the SARS coronavirus. *Protein Eng Des Sel*. 2015 Dec;28(12):567-75. <https://doi.org/10.1093/protein/gzv052>

13. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020 Mar;579(7798):270-273. <https://doi.org/10.1038/s41586-020-2012-7>
14. Liu J, Cao R, Xu M, Wang X, Zhang H, Hu H, et al. Hydroxychloroquine, a less toxic derivative of chloroquine, is effective in inhibiting SARS-CoV-2 infection in vitro. *Cell Discov*. 2020 Mar 18;6:16. <https://doi.org/10.1038/s41421-020-0156-0>
15. Mingo RM, Simmons JA, Shoemaker CJ, Nelson EA, Schornberg KL, D'Souza RS, et al. Ebola virus and severe acute respiratory syndrome coronavirus display late cell entry kinetics: evidence that transport to NPC1+ endolysosomes is a rate-defining step. *J Virol*. 2015 Mar;89(5):2931-43. <https://doi.org/10.1128/jvi.03398-14>
16. Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell*. 2020 Apr 16;181(2):271-280.e8. <https://doi.org/10.1016/j.cell.2020.02.052>
17. Magagnoli J, Narendran S, Pereira F, Cummings TH, Hardin JW, Sutton SS, et al. Outcomes of Hydroxychloroquine Usage in United States Veterans Hospitalized with COVID-19. *Med (N Y)*. 2020 Jun 5. <https://doi.org/10.1016/j.medj.2020.06.001>
18. Ministry of Health and Family Welfare, Government of India. *Advisory on the use of Hydroxychloroquine as prophylaxis for SARS-CoV-2 Infection*. [Online] 2020 [Cited March 23, 2020]. Available from:

<https://www.mohfw.gov.in/pdf/>

[AdvisoryontheuseofHydroxychloroquinasprophylaxisforSARSCoV2infection.pdf.](#)

19. National Centre for Disease Control. *SOP for Contact tracing COVID-19 cases*. [Online] 2020 [Cited April 3, 2020]. Available from:

[https://ncdc.gov.in/showfile.php?lid=538.](https://ncdc.gov.in/showfile.php?lid=538)

20. Indian Council of Medical Research. *Multiplex Real-Time PCR for detection of SARS-CoV-2 using TaqPath COVID-19 Combo Kit (Applied Biosystems)*. [Online] 2020 [Cited April 7, 2020]. Available from:

<https://www.icmr.gov.in/pdf/covid/labs/>

[SARS_CoV2_using_TaqPath_COVID19_ComboKit.pdf.](#)

21. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72 314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. 2020 Apr 7;323(13):1239-1242.

<https://doi.org/10.1001/jama.2020.2648>

22. Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, et al; ACTT-1 Study Group Members. Remdesivir for the Treatment of Covid-19 - Preliminary Report. *N Engl J Med*. 2020 May 22;NEJMoa2007764. <https://doi.org/10.1056/nejmoa2007764>

23. Covid19tracker.health.ny.gov. Workbook: NYS-COVID19-Tracker [Online] 2020 [Cited April 7, 2020]. Available from:

<https://covid19tracker.health.ny.gov/views/NYS-COVID19-Tracker/>

[NYSDOHCOVID-19Tracker-Fatalities?%3Aembed=yes&%3Atoolbar=no&%3Atabs=n](https://covid19tracker.health.ny.gov/views/NYS-COVID19-Tracker-Fatalities?%3Aembed=yes&%3Atoolbar=no&%3Atabs=n).

24. Tang W, Cao Z, Han M, Wang Z, Chen J, Sun W, et al. Hydroxychloroquine in patients with mainly mild to moderate coronavirus disease 2019: open label, randomised controlled trial. *BMJ*. 2020 May 14;369:m1849. <https://doi.org/10.1136/bmj.m1849>

25. Hussain A, Kaler J, Dubey AK. Emerging Pharmaceutical Treatments of Novel COVID-19: A Review. *Cureus*. 2020 May 24;12(5):e8260. <https://doi.org/10.7759/cureus.8260>

26. Wang Y, Zhang D, Du G, Du R, Zhao J, Jin Y, et al. Remdesivir in adults with severe COVID-19: a randomised, double-blind, placebo-controlled, multicentre trial. *Lancet*. 2020 May 16;395(10236):1569-1578. [https://doi.org/10.1016/s0140-6736\(20\)31022-9](https://doi.org/10.1016/s0140-6736(20)31022-9)

27. Cao B, Wang Y, Wen D, Liu W, Wang J, Fan G, et al. A Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe Covid-19. *N Engl J Med*. 2020 May 7;382(19):1787-1799. <https://doi.org/10.1056/nejmoa2001282>

28. McCaw JM, McVernon J. Prophylaxis or treatment? Optimal use of an antiviral stockpile during an influenza pandemic. *Math Biosci*. 2007 Oct;209(2):336-60. <https://doi.org/10.1016/j.mbs.2007.02.003>

29. Yao X, Ye F, Zhang M, Cui C, Huang B, Niu P, et al. In Vitro Antiviral Activity and Projection of Optimized Dosing Design of Hydroxychloroquine for the Treatment of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). *Clin Infect Dis*. 2020 Jul 28;71(15):732-739. <https://doi.org/10.1093/cid/ciaa237>

30. Wang M, Cao R, Zhang L, Yang X, Liu J, Xu M, et al. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res.* 2020 Mar;30(3):269-271. <https://doi.org/10.1038/s41422-020-0282-0>

31. Chatterjee P, Anand T, Singh KJ, Rasaily R, Singh R, Das S, et al. Healthcare workers & SARS-CoV-2 infection in India: A case-control investigation in the time of COVID-19. *Indian J Med Res.* 2020 May;151(5):459-467. https://doi.org/10.4103/ijmr.ijmr_2234_20

32. Ministry of Health and Family Welfare, Government of India. Guidelines on rational use of Personal Protective Equipment. [Online] 2020 [Cited March 24,2020].Available from:

<https://www.mohfw.gov.in/pdf/GuidelinesonrationaluseofPersonalProtectiveEquipment.pdf>.