Identification of SARS-CoV-2 in wastewater in Japan by multiple molecular

assays-implication for wastewater-based epidemiology (WBE).

Akihiko Hata[1], Ryo Honda[2,3,*], Hiroe Hara-Yamamura[2], Yuno Meuchi[1]

[1] Faculty of Engineering, Toyama Prefectural University

[2] Faculty of Geosciences and Civil Engineering, Kanazawa University

[3] Graduate School of Engineering, Kyoto University

*Corresponding author: Ryo Honda

Mailing address: *Faculty of Geosciences and Civil Engineering, Kanazawa University*

Kakuma-machi, Kanazawa 920-1192, Japan

Tel: +81-76-264-6393

E-mail: rhonda@se.kanazawa-u.ac.jp

15   **ABSTRACT**

Presence of SARS-coronavirus-2 (SARS-CoV-2) in wastewater sample has been

documented in several countries. Wastewater-based epidemiology (WBE) is

potentially effective for early warning of COVID-19 outbreak. The purpose of this

study was to verify the detection limit of WBE for COVID-19. In total, 27 influent

wastewater samples were collected from four wastewater treatment plants in

Ishikawa and Toyama prefectures in Japan. During the study period, numbers of

the confirmed COVID-19 cases in these prefectures increased from almost 0 to

around 20 per 100,000 peoples. SARS-CoV-2 RNA in the samples were identified

by several PCR-based assays. Among the 27 samples, 7 were positive for

SARS-CoV-2 by at least one out of the three quantitative RT-PCR assays. These

samples were also positive by RT-nested PCR assays. The detection frequency

became higher when the number of total confirmed SARS-CoV-2 cases in 100,000

peoples became above 10 in each prefecture. However, SARS-CoV-2 could also be

detected with a low frequency when the number was below 1.0. Considering that

the number of the confirmed cases does not necessarily reflect the actual

prevalence of the infection at the time point, data on the relationship between the

number of infection cases and concentration in wastewater needs to be

accumulated further.

1. INTRODUCTION

The recent novel coronavirus pneumonia (COVID-19) pandemic caused by the

severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection has led to

more than 6.8 million of the confirmed cases and nearly 400,000 deaths worldwide, as of

June 7, 2020 (WHO, 2020). Currently, this pandemic is likely to be heading for

convergence due to world-wide stay-at-home order. However, there are still a concern of

re-outbreak after lift of the stay-at-home order for reopening businesses, and also the

2nd pandemic caused by seasonal factors. Wastewater-based epidemiology (WBE) is the

effective approach to provide a snapshot of the outbreak situation in the entire catchment

by monitoring pathogens and viruses in wastewater (Choi et al., 2018; Yang et al., 2015).

The WBE could be an early warning tool for the possible re-outbreaks and seasonal

pandemics in the future.

Recently, detection of SARS-CoV-2 in wastewater is reported in Netherlands,

Australia, US, China, France, Israel, Italy, Spain, and Japan (Ahmed et al., 2020; Bar-or

et al., 2020; Haramoto et al., 2020; La Rosa et al., 2020; Medema et al., 2020; Nemudryi

et al., 2020; Randazzo et al., 2020; Rimoldi et al., 2020; WU et al., 2020; Wurtzer et al.,

2020a). SARS-CoV-2 are present in wastewater because they are shed from not only in

respiratory secretion but also feces of patients (Tang et al., 2020; Wölfel et al., 2020a; J.

Zhang et al., 2020; Y. Zhang et al., 2020). Importantly, pre-symptomatic patients and

asymptomatic patients also had a viral load in feces as well as symptomatic patients

(Tang et al., 2020; Wölfel et al., 2020a; Y. Zhang et al., 2020). Ratio of asymptomatic

infection is estimated as 18% (16-20% as 95%-CI) from the cruise ship passengers in

Yokohama (Mizumoto et al., 2020), and 31% (7.7-54% as 95%-CI) from the passengers

of evacuation flight from Wuhan to Tokyo (Nishiura et al., 2020). Moreover, a much

larger population of asymptomatic and mildly symptomatic infections was implied by

the surveillance of antibodies to SARS-CoV-2 in Santa Clara County, California, in

which the estimated number of infection is 50- to 85-fold more than the number of

confirmed cases (Bendavid et al., 2020). However, these people with no symptoms and

mild symptoms may not be included in clinical surveillance because most of them do not

visit clinics or hospitals. Transmission from such asymptomatic or pre-symptomatic

patients makes it difficult to enable the initial containment of COVID-19. Moreover, the

number of the reported COVID-19 cases is possibly biased by circumstances of medical

services in each country, e.g. access to medical services, capacity of PCR tests,

stay-at-home policy for mildly symptomatic patients, situation of medical collapse, etc.

So far, Japan has unique trend of COVID-19 epidemic. The increasing trend of cases is

much slower than other countries in Europe, US and China (Gloeckner et al., 2020).

Some paper argues that the number of cases is underestimated in Japan because of low

capacity of PCR tests (Omori et al., 2020). WBE is able to cover people with

asymptomatic and pre-symptomatic infections and to predict the overall epidemic status

without such biases of medical situations. Therefore, WBE could be more sensitive and

possibly detect outbreak earlier than clinical surveillance and also effective to verify

coverage of clinical surveillance under various medical situations. For application of

WBE as an early warning tool of COVID-19 outbreak, it is necessary to verify

correlations of SARS-CoV-2 concentration in wastewater and the number of COVID-19

cases in each country.

According the recent studies, SARS-CoV-2 were detected in wastewater when the

number of confirmed cases reached 1-100 cases per 100,000 of population (Ahmed et al.,

2020; Bar-or et al., 2020; Medema et al., 2020; Nemudryi et al., 2020; WU et al., 2020;

Wurtzer et al., 2020a). The sensitivity of WBE depends on viral load in feces of infected

people. Detection of SARS-CoV-2 in wastewater could be less sensitive than norovirus,

because the viral load of SARS-CoV-2 in feces is reportedly 1 to 2 logs lower than

norovirus (Hata and Honda, 2020). SARS-CoV-2 is an enveloped virus, while typical

viral pathogens in water, such as noroviruses, are non-enveloped viruses. Then,

commonly used methods for concentrating viruses in water might not be effective to

SARS-CoV-2 (Kitajima et al., 2020). Besides, several RT-PCR assays for SARS-CoV-2

are now available but their applicability to wastewater sample has not been well

determined. These indicate that detection of SARS-CoV-2 in the wastewater is more

challenging than that of norovirus and other enteric viruses. At this time, verifying

concentration methods and downstream RT-PCR assays is necessary to make WBE

reliable.

The objective of this study is to verify the detection limit of WBE for COVID-19 by

comparing the detected concentration of SARS-CoV-2 in wastewater and the COVID-19

cases reported in clinical surveillance. In this study, wastewater samples were taken in

two prefectures, Ishikawa and Toyama, in Japan which had the most reported

COVID-19 cases per population except Tokyo Metropolitan. To deal with the

methodological issues pointed above, sample processing was validated by assessing

virus detection efficiency with process controls. Besides, multiple RT-PCR assays and

nucleotide sequencing were applied to improve the reliability of the positive results. At

the beginning of the study period, numbers of the confirmed COVID-19 cases in

Ishikawa and Toyama prefectures were only 0.3 and 0 per 100,000 peoples (4 and no

confirmed cases), respectively. This enabled us to discuss sensitivity of WBE for

COVID-19 from the reported COVID-19 cases per population in the target area.

2. MATERIALS AND METHODS

2.1 Sample collection.

Influent wastewater samples were collected at four WWTPs, WWTPs A-C and D, in

Ishikawa and Toyama prefecture, respectively. These WWTPs were built to treat

maximum volumes of 12,800, 53,300, 156,000, and 82,500 m^3/day of wastewater in

total and are serving population equivalents of 34,501, 127,100, 153,291, and 169,400,

respectively (Table 1). At each WWTP, 100 mL of a grab sample was collected in the

morning, during the hour of peak flow, weekly or biweekly from March 5 to April 23,

2020. In total, 27 samples were obtained. The samples were kept cool or frozen until

the analysis, which took place within 3 days after the collection. The sample was split

into 80 mL and 20 mL of subsamples, which were then subjected to virus concentration

process and quantification of F-phages, respectively (Fig. 1). At the beginning of the

study period, there were only four and zero COVID-19 confirmed cases in Ishikawa

and Toyama prefectures, respectively. However, the numbers increased during the

period. At the end of April, Ishikawa and Toyama prefectures were ranked as second

and third, respectively, in Japan in the number of infections per 100,000 peoples (21.9

and 18.1 cases, respectively) (Fig. 2).

2.2 Virus concentration by polyethylene glycol precipitation (PEG precipitation).

Prior to the virus concentration, the 80 mL subsample was centrifuged at 5,000×g

for 5 minutes. The supernatant was recovered and mixed with 8.0 g of PEG8000 and 4.7

g of sodium chloride to be final concentrations of 10% and 1 M, respectively. The

mixture was incubated overnight on a shaker at 4°C. Subsequently, the mixture was

centrifuged at 10,000×g for 30 minutes. The supernatant was discarded and the resultant

pellet was resuspended in 500 µL of phosphate buffer as a virus concentrate.

2.3 RNA extraction.

Murine norovirus was used as a molecular process control to monitor the inhibitory

effects on the RNA extraction-RT-qPCR processes. Briefly, 140 µL of the virus

concentrate was spiked with 1.7×10^4 copies of MNV and subjected to RNA extraction

using a QIAamp viral RNA mini kit (Qiagen) to obtain a 60 µL of RNA extract, in

accordance with the manufacturer's instructions. The obtained RNA extract was

subjected to qRT-PCR assays and RT-semi nested PCR assays, for quantitative detection

of the spiked MNV and indigenous SARS-CoV-2 and for qualitative detection of

indigenous SARS-CoV-2, respectively.

2.4 qRT-PCR assay.

TaqMan-based qRT-PCR assays for quantitative detection of spiked MNV and

indigenous SARS-CoV-2 were performed with a qTOWER^3 (Analytik Jena). For

quantification of MNV RNA, a primer and TaqMan probe set described by Kitajima et

al. (2008) was utilized, while for quantification of SARS-CoV-2, two and one primer

and TaqMan probe sets described by Centers for Disease Control and Prevention (CDC)

in the USA (2020) (CDCN2, and CDCN3 assays, which were originally denoted as

2019-nCoV_N1, 2019-nCoV_N2, and 2019-nCoV_N3, respectively) and Shirato et al.

(2020) (NIID assay, which was originally denoted as NIID_2019-nCoV_N), respectively,

were separately used (Table 2). Shirato et al. (2020) described two reverse primers for

NIID assay. In this study, one of them (NIID_2019-nCoV_N_R2ver3, denoted as

NIID-R in Table 2) was used because it does not have mismatch with reference strains

(Wuhan-Hu-1 strain, Gen Bank acc. No. of MN908947.3, for example). A reaction

mixture (20 μL) was prepared using a One Step PrimeScript™ RT-PCR Kit (Perfect

Real Time) (TaKaRa). Briefly, 20 μL of a reaction mixture was prepared by mixing 2 μL

or 5 μL of RNA extract with, of 10 μL a 2×One Step RT-PCR Buffer III, 0.4 μL of a

TaKaRa Ex Taq HS, 0.4 μL of a PrimeScript RT enzyme Mix II, 400 nM each of

forward- and reverse primers, 100 nM of a TaqMan probe, and nuclease-free water.

qRT-PCR was performed under the following thermal cycling conditions: RT reaction at

42°C for 5 min, initial denaturation and inactivation of the RT enzyme at 95°C for 10 s,

followed by 50 cycles of amplification with denaturation at 95°C for 5 s and annealing

and extension at specific temperatures for each assay (Table 2) for 30 s.

To obtain a calibration curve, a 10-fold serial dilution (concentrations ranged from

1.0×10^0 to 1.0×10^6 copies/reaction) of a standard DNA (plasmid DNA or oligo DNA)

containing the target sequence was amplified. In each qPCR assay, 1.0×10^1

copies/reaction of the standard DNA were positive at all times, indicating that the limit

of the quantification values for each qPCR assay was below 1.0×10^1 copies/reaction. If

the resultant Ct value from a sample was corresponding to >1 copy/reaction, the sample

was determined to be positive for SARS-CoV-2.

Absence of the positive signal in the no template control was confirmed in every

qRT-PCR runs to exclude the potential contamination of the template into the reagents.

2.5 RT-nested PCR assays.

The RT reaction was performed with a high-capacity cDNA reverse transcription kit

(Life Technologies, Tokyo, Japan). The obtained cDNA was subjected to nested PCR

assays for qualitative gene detection of SARS-CoV-2 (Fig. 1). The first round of PCR

amplification was performed in 50 μL of a reaction mixture containing 10.0 μL of cDNA,

2×Premix Taq™ (TaKaRa), 400 nM each of CDCN3-F and NIID-R primers, which are

also employed for CDCN3 and NIID qRT-PCR assays, respectively, and expected to

amplify 600 bp gene fragment (Fig. 3 and Table 2), and nuclease-free water. The

amplification was performed under the following thermal cycling conditions using a

Thermal Cycler Dice (TaKaRa): initial denaturation and enzyme activation at 95°C for

10 min, followed by 40 cycles of amplification with denaturation at 95°C for 10 s,

primer annealing at 55°C for 30 s, and an extension reaction at 72°C for 1 min and then

a final extension at 72°C for 5 min. The first round PCR product was diluted with

nuclease free water by 1,000-fold and then subjected to following PCR assays.

For the second round of PCR, four different assays were conducted, one is a

conventional PCR assay (semi-nested PCR assay) followed by gel electrophoresis and

gene sequencing. Others are real-time PCR based on CDCN2, CDCN3, and NIID assays

(nested-real time PCR assays). The semi-nested PCR assay was performed in a 20 μL

reaction mixture containing 2 μL of the product of the first round of PCR amplification,

2×Premix Taq™ (Takara), and 400 nM each of CDCN3-F and CDCN2-R primers (Fig.

3 and Table 2), which are also employed for CDCN3 and CDCN2 qRT-PCR assays,

respectively, and expected to amplify 550 bp gene fragment, and nuclease-free water.

PCR amplification was performed under the following thermal cycling conditions using

a Thermal Cycler Dice (TaKaRa): initial denaturation at 95°C for 10 min, followed by

40 cycles of amplification with denaturation at 95°C for 10 s, primer annealing at 55°C

for 30 s, and extension reaction at 72°C for 30 sec and then a final extension at 72°C for

5 min. The semi nested PCR product was subjected to electrophoresis on a 1.5% agarose

gel and visualized under a UV lamp after being stained with Gel Red™ (FUJIFILM).

For further confirmation of the successful amplification by the first round PCR, the

PCR product was subjected to CDCN2, CDCN3, and NIID assays (nested-real time PCR

assays) with a qTOWER³ (Analytik Jena) (Fig. 3 and Table 2). The real time PCR was

performed in a 20 μL reaction mixture containing 2 μL of the product of the first round

of PCR amplification, 2×Premix Taq™ (TaKaRa), and 400 nM each of primers, 100 nM

of a TaqMan probe, and nuclease-free water. PCR amplification was performed under

the following thermal cycling conditions: initial denaturation at 95°C for 10 min,

followed by 40 cycles of amplification with denaturation at 95°C for 10 s, annealing and

extension at specific temperatures for each assay for 30 s (Table 2).

2.6 Determining efficiencies of virus concentration and RNA extraction-qRT-PCR.

Indigenous F-phage in the sample before and after the concentration process was

quantified by a plate counting assay employing *Salmonella Typhimurium* WG49 as a

host strain, as described previously (Mooijman et al., 2002). The efficiency of the virus

concentration process was estimated by comparing the F-phage concentration in the

sample before the concentration process with that after the concentration process.

The efficiency of RNA extraction-qRT-PCR was estimated by comparing the

observed gene concentration of the spiked MNV in nuclease-free water with that in the

concentrate.

3. RESULTS

3.1 Efficiency of virus detection.

The recovery efficiency of indigenous F-phage by the concentration process was

57% in geometric mean and was always above 10% (n = 27). RNA extraction-qRT-PCR

efficiency determined with the spiked MNV was 66% in geometric mean and was

always above 10% (n = 27). These indicate that the virus concentration and following

molecular processes were conducted efficiency enough (Haramoto et al., 2018).

Presumptive virus detection efficiency, which can be estimated by multiplying these

efficiencies, was 38% in geometric mean and below 10% for 5 out of 27 samples (3.7%

was the lowest).

3.2 Detection of SARS-CoV-2 by qRT-PCR assays.

In this study, three qRT-PCR assays, CDCN2，CDCN3 and NIID assays, were

separately applied. A sample collected at WTTP D on April 3 was positive for

SARS-CoV-2 using all the assays with an observed concentration of around 3.0×10^4

copies/L ($2.1 \times 10^4 – 4.4 \times 10^4$ copies/L) (Fig. 2 and Table 3). The presumptive virus

detection efficiency in the sample was relatively high (estimated as 130%). In addition to

the sample, CDCN2, CDCN3, and NIID assays showed positive signals in 1 (WWTP B

on April 16), 4 (WWTP A on March 19, WWTP B on March 26 and April 23, and

WWTP C on April 21), and 1 (WWTP D on April 24) samples. In total, 7 samples were

determined to be positive for SARS-CoV-2 by at least one of the assays with observed

concentration of $1.2{\times}10^4 - 4.4{\times}10^4$ copies/L. The detection frequency seems to be

higher when the number of total confirmed SARS-CoV-2 cases in 100,000 peoples

became above 10 in each prefecture (Fig. 2 and Table 3). The SARS-CoV-2 detection

frequency was 15% (3 positives out of 20 samples) before the number became >10,

whereas it reached 57% (4 positives out of 7 samples) after the number became >10.

3.3 Detection of SARS-CoV-2 by RT-nested PCR assays and nucleotide sequencing.

The qRT-PCR assays used in this study target a 600 bp fraction of gene coding

N-protein. Then, nested PCR assays could be performed using the primers used in the

qRT-PCR assays for further confirmation of the positive results obtained by the

qRT-PCR assays. A semi-nested PCR assay to amplify 550 bp of a gene fragment

showed a band of around 600 bp size in four out of seven samples, which were collected

during April (Table 3 and Fig. 4). However, direct Sanger sequencing could not confirm

if the amplicon was obtained from SARS-CoV-2 or not, because of too many

heterozygous peaks. Application of real-time PCR to the first round PCR products

showed ambiguous results. CDCN2 and NIID assay did not show any positive results,

while CDCN3 assay showed positive signal from six out of the seven samples. Among

them, samples collected during March resulted in relatively high (>35) Ct values, while

those collected during April resulted in lower (around 30) Ct values.

4. DISCUSSION

Most of previous studies investigated pathogenic and indicator viruses targeted

non-enveloped viruses (Haramoto et al., 2018). Consequently, retrospective virus

concentration methods are optimized for non-enveloped viruses and may not be effective

to enveloped virus like SARS-CoV-2. For instance, adsorption-elution methods using

electropositive or electronegative microfilter and glass wool are surely effective to

non-enveloped viruses (Cashdollar et al., 2013) but resulted in inefficient recovery

efficiency (consistently less than or around 10%) for at least some of enveloped viruses

(Haramoto et al., 2009; Honjo et al., 2010; Deboosere et al., 2011; Francy et al., 2013;

Blanco et al., 2019). A size excusive method using ultrafiltration is also widely applied

for virus monitoring (Cashdollar et al., 2013). Even though its efficacy in concentrating

enveloped viruses has not been fully elucidated yet, some researchers have been

succeeded in detecting enveloped viruses, influenza-A virus and SARS-CoV-2 (Heijnen

and Medema, 2011; Medema et al., 2020; Ahmed et al., 2020; Nemudryi et al., 2020).

One of the drawbacks of the ultrafiltration-based method is the cost of the ultrafiltration

unit. For example, Centricon plus-70, which can process up to 70 mL of a sample at

once, costs approximately 50 USD per unit, and it might not be suitable for frequent use.

On the contrary, PEG precipitation is one of the most cost-effective method for

concentrating virus in a low volume (around 100 mL) of water sample (Lewis and

Metcalf, 1988). The technique is effective not only for viruses but also for variety of

proteins (Atha and Ingham, 1981). Therefore, it might be effective for SARS-CoV-2,

even though only a limited number of studies have applied the technique to enveloped

viruses, including SARS-CoV-2, in water (Honjo et al., 2010; Deboosere et al., 2011;

Wu et al., 2020). For PEG precipitation, several conditions, i.e., concentrations of PEG

and NaCl, have been applied (Jones and Johns, 2009). Among them, we selected

relatively high concentrations of PEG and NaCl (10% and 1.0 M), in accordance with

Jones and Johns (2009). As expected, the method revealed a sufficiently high and stable

recovery efficiency of indigenous F-phages (57% in geometric mean), indicating that

SARS-CoV-2 was also recovered efficiently. An efficient way for recovering

SARS-CoV-2 in wastewater and an appropriate recovery control to ascertain the

efficient recovery of SARS-CoV-2 should be found in future.

Several molecular assays have been designed for detection of SARS-CoV-2

(CDC2020; Shirato 2020; Corman et al., 2020; Chu et al., 2020). Although some studies

revealed effectiveness of these assays in comparative manner (Jung et al., 2020), a gold

standard method has not been determined yet. Especially, wastewater contains variety of

microbes and free nucleic acids, which might lead to false positive due to non-specific

amplification. Therefore, most studies on SARS-CoV-2 in wastewater samples applied

multiple RT-PCR assays, gel electrophoresis, and amplicon sequencing in attempts to

improve reliability (Ahmed et al., 2020; Medema et al., 2020; Nemudryi et al., 2020; Wu

et al., 2020; Wurtzet et al., 2020). In this study, three qRT-PCR assays and additional

RT-nested PCR assays, to confirm the positive results obtained by the qRT-PCR assays,

were applied. Among seven samples positive by at least one of the qRT-PCR assays, one

collected at WWTP D on April 3 were consistently positive by all three qRT-PCR assays,

while other six samples were positive by one of the assays. Low concentrations of

SARS-CoV-2 gene in the concentrates can be one possible reason for the ambiguous

results. The observed concentration was at most 8 copies/reaction. In this case, the target

gene is occasionally absent in the analyte. It is also possible that some of the positive

signals were generated by non-specific amplicons or that some assays failed to amplify

specific SARS-CoV-2 strains.

Then, RT-nested PCR assays sharing primers with the qRT-PCR assays were

conducted to ascertain the positives obtained by the qRT-PCR assays. Most of the

qRT-PCR positive samples again showed positive results by at least one of the RT-nested

PCR assays, indicating that the positive signals obtained by the qRT-PCR assays are

reliable. Some previous studies revealed that CDCN2 assay seems to be less efficient in

detecting SARS-CoV-2 in wastewater samples than CDCN1 and CDCN3 assays

(Medema et al., 2020; Wu et al., 2020; Nemudryi et al., 2020; Randazzo et al., 2020).

Accordingly, CDCN2 assay, as well as NIID assay, seem to be inefficient in this study,

showed no positive results after the first round PCR. Similarly, the first round PCR

followed by CDCN3 assay seems to be better than that followed by the semi nested PCR

assay, in view of the number of positives. These suggest that primers and TaqMan probe

used in CDCN2 and NIID assays might contain mismatches with SARS-CoV-2 strains in

wastewater. Notably, one sample (at WWTP B on April 16) failed to confirm the

presence of SARS-CoV-2 by the second round CDCN3 assay. This sample was also

negative by qRT-PCR with CDCN3 assay. Considering that the sample was positive by

the semi nested assay, the first round PCR was successfully conducted. This implies that

CDCN3 assay may also not be effective in detecting all the circulating SARS-CoV-2

strains.

Detection frequency of SARS-CoV-2 in our wastewater samples seemed to be in

agreement with the numbers of confirmed cases in both prefectures. The detection

frequency seems to be higher when it becomes above 10 in 100,000 peoples (Fig. 2 and

Table 3) Besides, no positive result was obtained in the samples collected in Toyama

prefecture during March, when no confirmed SARS-CoV-2 infection cases were

reported. Our results also suggest that SARS-CoV-2 in wastewater sample could be

identified even when the number of cases per 100,000 peoples is below 1.0, although the

detection frequency becomes low (Fig. 2 and Table 3). One SARS-CoV-2 positive

sample during March was collected at WWTP A. A relatively low number of populations

served by the WWTP (34,501 peoples) might be attributed to the positive result. Smaller

WWTP would have a limited dilution effect and can be more sensitive to the presence of

the number of infected individuals. Previous studies have suggested that the

concentration of SARS-CoV-2 in wastewater is affected by number of peoples infected

in the catchment area (Medema et al., 2020; Ahmed et al., 2020; Wurtzer et al., 2020b).

However, there seems to be a discrepancy in the observed relationships between the

concentration in wastewater and the number of infected people. Ahmed et al. (2020)

could not identify SARS-CoV-2 in a wastewater sample in Queensland until the number

of the confirmed cases reached up to around 100 cases per 100,000 peoples. In addition,

the observed concentration was below the limit of quantification at that time. In contrast,

Wurtzer et al. (2020b) repeatedly obtained positive results in Paris even when the

number of the confirmed cases was below 1 case per 100,000 peoples, in accordance

with our observation. One of the reasons for this discrepancy might be due to the

difference in methods applied; collecting composite samples or grab samples and ways

of sample concentration, nucleic acid extraction, and qRT-PCR. Besides, the number of

the confirmed cases does not necessarily reflect the actual prevalence of the infection at

the time point, which can be better correlated to the concentration in the wastewater. In

Japan, it takes a few days to confirm the case by the clinical surveillance since the

symptom first appeared or the one become infected. There should be asymptomatic

cases, which were not confirmed. In the context of WBE, data on the relationship

between the number of infection cases and concentration in wastewater needs to be

accumulated further.

ACKNOWLEDGEMENTS

This work was supported by Hiramoto-gumi Inc., the civil construction company in

Japan.

REFERENCES

1.   Ahmed, W., Angel, N., Edson, J., Bibby, K., Bivins, A., O'Brien, J.W., Choi, P.M.,

Kitajima, M., Simpson, S.L., Li, J., Tscharke, B., Verhagen, R., Smith, W.J.M.,

Zaugg, J., Dierens, L., Hugenholtz, P., Thomas, K. V., Mueller, J.F., 2020. First

confirmed detection of SARS-CoV-2 in untreated wastewater in Australia: A proof

of concept for the wastewater surveillance of COVID-19 in the community. Sci.

Total Environ. 138764.

2.   Atha, D.H., Ingham, K.C., 1981. Mechanism of precipitation of proteins by

polyethylene glycols. Analysis in terms of excluded volume. J. Biol. Chem. 256,

12108–12117

3.   Bar-or, I., Yaniv, K., Shagan, M., Ozer, E., Erster, O., Mendelson, E., Shirazi, R.,

Kramarsky-winter, E., Nir, O., Abu-ali, H., Ronen, Z., Lewis, Y.E., Friedler, E.,

Bitkover, E., Paitan, Y., Berchenko, Y., Goldstein-goren, S., Science, N., Sheva, B.,

Saba, K., Boker, S., 2020. Regressing SARS-CoV-2 sewage measurements onto

COVID-19 burden in the population: a proof-of-concept for quantitative

environmental surveillance. medRxiv 020.04.26.20073569.

4.   Bendavid, E., Mulaney, B., Sood, N., Shah, S., Ling, E., Bromley-Dulfano, R., Lai,

C., Weissberg, Z., Saavedra, R., Tedrow, J., Tversky, D., Bogan, A., Kupiec, T.,

Eichner, D., Gupta, R., Ioannidis, J., Bhattacharya, J., 2020. COVID-19 Antibody

Seroprevalence in Santa Clara County, California. medRxiv 2020.04.14.20062463.

5.   Blanco, A., Abid, I., Al-Otaibi, N., Pérez-Rodríguez, F.J., Fuentes, C., Guix, S.,

Pintó, R.M., Bosch, A., 2019. Glass Wool Concentration Optimization for the

Detection of Enveloped and Non-enveloped Waterborne Viruses. Food Environ.

Virol. 11, 184–192.

6.   Cashdollar, J.L., Wymer, L., 2013. Methods for primary concentration of viruses

from water samples: A review and meta-analysis of recent studies. J. Appl.

Microbiol. 115, 1–11.

7.   Centers for Disease Control and Prevention, Respiratory Viruses Branch, Division

of Viral Diseases Instructions. 2020. Real-Time RT-PCR Panel for Detection

2019-Novel Coronavirus Centers for Disease Control and Prevention, Respiratory

Viruses Branch, Division of Viral Diseases, accessed: February 28, 2020.

8.   Choi, P.M., Tscharke, B.J., Donner, E., O'Brien, J.W., Grant, S.C., Kaserzon, S.L.,

Mackie, R., O'Malley, E., Crosbie, N.D., Thomas, K. V., Mueller, J.F., 2018.

Wastewater-based epidemiology biomarkers: Past, present and future. TrAC -

Trends Anal. Chem.

9.   Chu, D.K.W., Pan, Y., Cheng, S.M.S., Hui, K.P.Y., Krishnan, P., Liu, Y., Ng,

D.Y.M., Wan, C.K.C., Yang, P., Wang, Q., Peiris, M., Poon, L.L.M., 2020.

Molecular Diagnosis of a Novel Coronavirus (2019-nCoV) Causing an Outbreak of

Pneumonia. Clin. Chem. 7, 1–7.

10.   Corman, V.M., Landt, O., Kaiser, M., Molenkamp, R., Meijer, A., Chu, D.K.,

Bleicker, T., Brünink, S., Schneider, J., Luisa Schmidt, M., GJC Mulders, D.,

Haagmans, B.L., van der Veer, B., van den Brink, S., Wijsman, L., Goderski, G.,

Romette, J.-L., Ellis, J., Zambon, M., Peiris, M., Goossens, H., Reusken, C.,

Koopmans, M.P., Drosten, C., Victor, C.M., Olfert, L., Marco, K., Richard, M.,

Adam, M., Daniel, C.K., Tobias, B., Sebastian, B., Julia, S., Marie Luisa, S.,

Daphne GJC, M., Bart, H.L., der Veer Bas, V., den Brink Sharon, V., Lisa, W.,

Gabriel, G., Jean-Louis, R., Joanna, E., Maria, Z., Malik, P., Herman, G., Chantal,

R., 2020. Detection of 2019-nCoV by RT-PCR. Euro. Surveill. 25, 1–8.

11. Deboosere, N., Horm, S.V., Pinon, A., Gachet, J., Coldefy, C., Buchy, P., Vialette,

M., 2011. Development and validation of a concentration method for the detection

of influenza a viruses from large volumes of surface water. Appl. Environ.

Microbiol. 77, 3802–3808.

12. Francy, D.S., Stelzer, E.A., Brady, A.M.G., Huitger, C., Bushon, R.N., Ip, H.S.,

Ware, M.W., Villegas, E.N., Gallardo, V., Lindquist, H.D.A., 2013. Comparison of

filters for concentrating microbial indicators and pathogens in lake water samples.

Appl. Environ. Microbiol. 79, 1342–1352.

13. Gloeckner, S., Krause, G., Hoehle, M., 2020. Now-casting the COVID-19 epidemic:

The use case of Japan, March 2020. medRxiv 2020.03.18.20037473.

14. Hata, A., Honda, R., 2020. Potential sensitivity of wastewater monitoring for

SARS-CoV-2: comparison with norovirus cases. Environ. Sci. Technol. (in press).

15. Haramoto, E., Kitajima, M., Katayama, H., Ito, T., Ohgaki, S., 2009. Development

of virus concentration methods for detection of koi herpesvirus in water. J. Fish Dis.

32, 297–300.

16. Haramoto, E., Kitajima, M., Hata, A., Torrey, J.R., Masago, Y., Sano, D., Katayama,

H., 2018. A review on recent progress in the detection methods and prevalence of

human enteric viruses in water. Water Res. 135. 168-186.

17. Haramoto, E., Malla, B., Thakali, O., Kitajima, M. 2020. First environmental

surveillance for the presence of SARS-CoV-2 RNA in wastewater and river water in

Japan. medRxiv.

18. Heijnen, L., Medema, G., 2011. Surveillance of influenza A and the pandemic

influenza A (H1N1) 2009 in sewage and surface water in the Netherlands. J. Water

Health 9, 434–442.

19. Honjo, M.N., Minamoto, T., Matsui, K., Uchii, K., Yamanaka, H., Suzuki, A. a,

Kohmatsu, Y., Iida, T., Kawabata, Z., 2010. Quantification of cyprinid herpesvirus

3 in environmental water by using an external standard virus. Appl. Environ.

Microbiol. 76, 161–8.

20. Jones, T.H., Johns, M.W., 2009. Improved detection of F-specific RNA coliphages

in fecal material by extraction and polyethylene glycol precipitation. Appl. Environ.

Microbiol. 75, 6142–6.

21. Jung, Y.J., Park, G.-S., Moon, J.H., Ku, K., Beak, S.-H., Kim, Seil, Park, E.C., Park,

443 D., Lee, J.-H., Byeon, C.W., Lee, J.J., Maeng, J., Kim, S.J., Kim, Seung Il, Kim,

444 B.-T., Lee, M.J., Kim, H.G., 2020. Comparative analysis of primer-probe sets for

445 the laboratory confirmation of SARS-CoV-2. BioRxiv 2020.02.25.964775.

446 22. Kitajima, M., Tohya, Y., Matsubara, K., Haramoto, E., Utagawa, E., Katayama, H.,

447 Ohgaki, S. (2008). Use of murine norovirus as a novel surrogate to evaluate

448 resistance of human norovirus to free chlorine disinfection in drinking water supply

449 system. Environ. Eng. Res. 45, 361–370 (in Japanese).

450 23. Kitajima, M., Ahmed, W., Bibby, K., Carducci, A., Gerba, C.P., Hamilton, A.,

451 Haramoto, E., Rose, J.B. 2020. SARS-CoV-2 in wastewater: State of the knowledge

452 and research needs. Science of the Total Environment. (in press).

453 24. La Rosa, D., Iaconelli, M., Mancini, P., Ferraro, G.B., Veneri, C., Bonadonna, L.,

454 Lucentini, L., Suffredini, E. 2020. First Detection of SARS-CoV-2 in Untreated

455 Wastewaters in Italy. medRxiv.

456 25. Lewis, G.D., Metcalf, T.G., 1988. Polyethylene glycol precipitation for recovery of

457 pathogenic viruses, including hepatitis A virus and human rotavirus, from oyster,

458 water, and sediment samples. Appl. Environ. Microbiol. 54, 1983–8.

459 26. Medema, G., Heijnen, L., Elsinga, G., Italiaander, R., Brouwer, A., 2020. Presence

460 of SARS-Coronavirus-2 RNA in sewage and correlation with reported COVID-19

461 prevalence in the early stage of the epidemic in the Netherlands. Environ. Sci.

462 Technol. (in press).

27. Mizumoto, K., Kagaya, K., Zarebski, A., Chowell, G., 2020. Estimating the

asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board

the Diamond Princess cruise ship, Yokohama, Japan, 2020. Eurosurveillance 25,

1–5.

28. Mooijman, K. a, Bahar, M., Muniesa, M., Havelaar, A.H., 2002. Optimisation of

ISO 10705-1 on enumeration of F-specific bacteriophages. J. Virol. Methods 103,

129–36.

29. Nemudryi, A., Nemudraia, A., Surya, K., Wiegand, T., Buyukyoruk, M., Wilkinson,

R., Wiedenheft, B., 2020. Temporal detection and phylogenetic assessment of

SARS-CoV-2 in municipal wastewater. medRxiv 2020.03.22.20041079.

30. Nishiura, H., Kobayashi, T., Suzuki, A., Jung, S.-M., Hayashi, K., Kinoshita, R.,

Yang, Y., Yuan, B., Akhmetzhanov, A.R., Linton, N.M., Miyama, T., 2020.

Estimation of the asymptomatic ratio of novel coronavirus infections (COVID-19).

Int. J. Infect. Dis. 94, 154-155.

31. Omori, R., Mizumoto, K., Chowell, G., 2020. Changes in testing rates could mask

the novel coronavirus disease (COVID-19) growth rate. Int. J. Infect. Dis. 94,

116–118.

32. Randazzo, W., Truchado, P., Cuevas-Ferrando, E., Simón, P., Allende, A., Sánchez,

G. 2020. SARS-CoV-2 RNA in Wastewater Anticipated COVID-19 Occurrence in a

Low Prevalence Area. Water Res. (in press)

33. Rimoldi, S.G., Stefani, F., Gigantiello, A., Polesello, S., Comandatore, F., Mileto1,

D., Maresca, M., Longobardi, C., Mancon, A., Romeri, F., Pagani, C., Moja, L.,

Gismondo, M. R., Salerno, F. 2020. Presence and vitality of SARS-CoV-2 virus in

wastewaters and rivers. MedRxiv.

34. Shirato, K., Nao, N., Katano, H., Takayama, I, Saito, S., Kato, F., Katoh, H., Sakata,

M., Nakatsu, Y., Mori, Y., Kageyama, T., Matsuyama, S., Takeda, M., 2020.

Development of Genetic Diagnostic Methods for Novel Coronavirus 2019

(nCoV-2019) in Japan. Jpn. J. Infect. Dis. (in press).

35. World Health Organization, 2020. Coronavirus disease (COVID-2019) situation

reports [WWW Document]. URL

https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports

(accessed 6.8.20).

36. Wu, F., Xiao, A., Zhang, J., GU, X., Lee, W., Kauffman, K., Hanage, W., Matus, M.,

Ghaeli, N., Endo, N., Duvallet, C., Moniz, K., Erickson, T., Pr, C., Thompson, J.,

Alm, E., 2020. SARS-CoV-2 titers in wastewater are higher than expected from

clinically confirmed cases. medRxiv 2020.04.05.20051540.

37. Wurtzer, S., Marechal, V., Mouchel, J.M., Moulin, L. 2020a. Time course

quantitative detection of SARS-CoV-2 in Parisian wastewaters correlates with

COVID-19 confirmed cases. medRxiv. 2020.04.12.20062679.

38. Wurtzer, S., Marechal, V., Mouchel, J.M., Maday, Y., Teyssou, R., Richard, E.,

Almayrac, J.L., Moulin L. 2020b. Evaluation of lockdown impact on SARS-CoV-2

dynamics through viral genome quantification in Paris wastewaters.

39. Yang, Z., Kasprzyk-Hordern, B., Frost, C.G., Estrela, P., Thomas, K. V., 2015.

Community sewage sensors for monitoring public health. Environ. Sci. Technol. 49,

5845–5846.

509   **FIGURES**

Figure 1. Flow diagram of sample processing for confirming the presence of

SARS-CoV-2 in wastewater by RT-PCR assays and for assessing virus detection

efficiency with process controls.

Figure 2. Temporal relationship between number of SARS-CoV-2 infection cases and

presence of SARS-CoV-2 in wastewater samples in Ishikawa and Toyama prefectures.

Number of inpatients was calculated by subtracting numbers of discharged and dead

from number of total confirmed cases, which were derived from a databased

summarized by Ministry of Health, Labor and Welfare in Japan

(https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000121431_00086.html).

Figure 3. Location of primers and TaqMan probes employed in this study on a

referential SARS-CoV-2 strain, Wuhan-Hu-1 (Gen Bank accession no: MN908947).

These are localized on a relatively short fraction (600 nt) of N-gene. Therefore,

CDCN3-F and NIID-R primers, for example, can be used as outer primers for nested

assays.

Figure 4. Detection of SARS-CoV-2 in wastewater samples by the RT-semi-nested PCR

assay, which is expected to result in a 550 bp of amplicon. Lanes 1-3: Samples collected

at WWTPs A, B, and D on March 19, 26, and April 3, which did not show the positive

band. Lanes 4-7: Samples collected at WWTPs A, B, C, B, on April 16, 21, 23, and 24,

which showed the positive band. Lane 8: Non target control. Lane 9: 100-bp ladder.

TABLES

TABLE 1. Characteristics of WWTPs studied in this study.

WWTP	Prefecture	Population served by WWTP	Treatment capacity (m^3/d)
A	Ishikawa	34,501	12,800
B	Ishikawa	127,100	53,300
C	Ishikawa	153,291	156,000
D	Toyama	169,400	82,500

TABLE 2. Primer and TaqMan probe sequences used for detection of SARS-CoV-2 in this study.

Assay	Name	Function	Sequence (5' → 3')[a]	Location	temperature for annealing/extension	Reference
2019-nCoV_N2	CDCN2-F	Forward primer	TTACAAACATTGGCCGCAAA	29164-29183	55°C	CDC (2020)
	CDCN2-R[b]	Reverse primer	GCGCGACATTCCGAAGAA	29213-29230		
	CDCN2-P	TaqMan Probe	ACAATTTGCCCCCAGCGCTTCAG-	29188-29210		
2019-nCoV_N3	CDCN3-F[b]	Forward primer	GGGAGCCTTGAATACACCAAAA	28681-28702	55°C	CDC (2020)
	CDCN3-R	Reverse primer	TGTAGCACGATTGCAGCATTG	28732-28752		
	CDCN3-P	TaqMan Probe	AYCACATTGGCACCCGCAATCCTG-	28704-28727		
NIID_2019-nCoV_N	NIID-F	Forward primer	AAATTTTGGGGACCAGGAAC	29125-29144	60°C	Shirato et al. (2020)
	NIID-R[b]	Reverse primer	TGGCACCTGTGTAGGTCAAC	29263-29282		
	NIID-P	TaqMan Probe	ATGTCGCGCATTGGCATGGA	29222-29241		

[a]: All TaqMan probes were labeled with 5'-FAM and 3'-TAMRA.
[b]: CDCN3-F, CDCN2-R, NIID-R were also used for nested PCR assays.

TABLE 3. Summary of samples positive for SARS-CoV-2 by qRT-PCR assays.

Date	WWTP[a]	Total confirmed cases in 100,000 peoples		Efficiency of detection process			qRT-PCR			RT-nested PCR			
		Ishikawa	Toyama	Concentration	RNA extraction-qRT-PCR	Total	CDCN2	CDCN3	NIID	semi-nested	CDCN2	CDCN3	NIID
				%	%	%		copies/L			Ct	Ct	Ct
March 19	A	0.4	0.0	263	72	190	Neg.[b]	2.8×10^4	Neg.	Neg.	Neg.	35.2	Neg.
March 27	B	0.5	0.0	76	50	38	Neg.	1.7×10^4	Neg.	Neg.	Neg.	36.2	Neg.
April 3	D	1.4	0.8	125	102	130	4.4×10^4	1.2×10^4	2.1×10^4	Neg.	Neg.	29.9	Neg.
April 16	B	12.2	5.1	10	49	4.9	1.5×10^4	Neg.	Neg.	Pos.	Neg.	Neg.	Neg.
April 21	C	15.7	10.9	87	41	36	Neg.	1.5×10^4	Neg.	Pos.	Neg.	32.1	Neg.
April 23	B	17.4	11.0	36	50	18	Neg.	1.2×10^4	Neg.	Pos.	Neg.	31.1	Neg.
April 24	D	18.9	14.3	51	31	16	Neg.	Neg.	1.4×10^4	Pos.	Neg.	30.3	Neg.

[a]: WWTPs A-C and D are located in Ishikawa and Toyama prefecture, respectively.

[b]: Neg.: Negative; Pos.: Positive.