Child wasting and concurrent stunting in low- and middle-income countries

Andrew Mertens,[1]* Jade Benjamin-Chung,[2] John M Colford Jr,[1] Alan E Hubbard,[1] Mark J van der Laan,[1] Jeremy Coyle[1], Oleg Sofrygin,[1] Wilson Cai,[1] Wendy Jilek,[1] Sonali Rosete,[1] Anna Nguyen,[1] Nolan N Pokpongkiat,[1] Stephanie Djajadi,[1] Anmol Seth,[1] Esther Jung,[1] Esther O Chung,[1] Ivana Malenica,[1] Nima Hejazi,[1] Haodong Li,[1] Ryan Hafen,[3] Vishak Subramoney,[4] Jonas Häggström,[5] Thea Norman,[6] Parul Christian,[7] Kenneth H Brown,[8] Benjamin F. Arnold,[9,10]* and members of the *ki* Child Growth Consortium**

[1] Division of Epidemiology & Biostatistics, University of California, Berkeley, 2121 Berkeley Way Rm 5302 Berkeley, CA 94720-7360

[2] Department of Epidemiology & Population Health, Stanford University, Stanford University, 259 Campus Drive, HRP Redwood Building T152A, Stanford, CA 94305

[3] Hafen Consulting, LLC, West Richland WA, 99353

[4] DVPL Tech

[5] Cytel Inc, 1050 Winter St Suite 2700 Waltham, MA 02451, USA

[6] Quantitative Sciences, Bill & Melinda Gates Foundation, 500 5th Ave N, Seattle, WA 98109

[7] Center for Human Nutrition, Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD 21205.

[8] Department of Nutrition, University of California, Davis, 3135 Meyer Hall, Davis, CA 95616-5270

[9] Francis I. Proctor Foundation, University of California, San Francisco, 95 Kirkham St, San Francisco, CA 94143

[10] Department of Ophthalmology, University of California, San Francisco, 10 Koret Way, San Francisco, CA 94143

* Corresponding authors are Andrew Mertens (amertens@berkeley.edu) and Benjamin F. Arnold (ben.arnold@ucsf.edu).

** Members of the ki Child Growth Consortium

1.   Tahmeed Ahmed, International Centre for Diarrhoeal Disease Research, Dhaka, Bangladesh

2.   Asad Ali, Aga Khan University, Pakistan, Karachi, Pakistan

3.   France Begín, UNICEF, New York, USA

4.   Mauricio L. Barreto, Center of Data and Knowledge Integration for Health, Fundação Oswaldo Cruz, Salvador, Brazil.

5.   Pascal Obong Bessong, HIV/AIDS & Global Health Research Programme, University of Venda, Thohoyandou, South Africa

6.   Zulfiqar A. Bhutta, Institute for Global Health & Development & Center of Excellence in Women and Child Health, The Aga Khan University South-Central Asia, East Africa & United Kingdom

7.   Robert E. Black, Johns Hopkins University, Bloomberg School of Public Health, Baltimore, USA

8.   Ladaporn Bodhidatta, Armed Forces Research Institute of Medical Sciences, Bangkok, Thailand

9.   William Checkley, Johns Hopkins University, School of Medicine, Baltimore, USA

10.   Jean E. Crabtree, Leeds Institute for Medical Research, St. James's University Hospital, University of Leeds, Leeds, United Kingdom

11.   Rina Das, International Centre for Diarrhoeal Disease Research, Dhaka, Bangladesh

12.   Subhasish Das, International Centre for Diarrhoeal Disease Research, Dhaka, Bangladesh

13. Christopher P. Duggan, Center for Nutrition, Boston Children's Hospital, USA

14. Abu Syed Golam Faruque, International Centre for Diarrhoeal Disease Research, Dhaka, Bangladesh

15. Wafaie W. Fawzi, Department of Global Health and Population, Harvard TH Chan School of Public Health, Cambridge, USA

16. José Quirino da Silva Filho, Federal University of Ceará, Fortaleza, Brazil

17. Robert H. Gilman, Johns Hopkins University, Bloomberg School of Public Health, Baltimore, USA

18. Richard Guerrant, University of Virginia, Charlottesville, USA

19. Rashidul Haque, International Centre for Diarrhoeal Disease Research, Dhaka, Bangladesh

20. Eric R. Houpt, University of Virginia, Charlottesville, USA

21. Najeeha Talat Iqbal, Department of Pediatrics and Child Health, Aga Khan University, Karachi, Pakistan

22. Jacob John, Christian Medical College, Vellore, India

23. Sushil Matthew John, Professor, Low Cost Effective Care Unit, Christian Medical College, Vellore, India

24. Gagandeep Kang, Translational Health Science and Technology Institute, Pali, India

25. Margaret Kosek, University of Virginia, Charlottesville, USA

26. Aldo Ângelo Moreira Lima, Federal University of Ceará, Fortaleza, Brazil

27. Tjale Cloupas Mahopo, Department of Nutrition, School of Health Sciences, University of Venda, Thohoyandou, South Africa

28. Dharma S. Manandhar, Mother and Infant Research Activities (MIRA), Kathmandu, Nepal

29. Karim P. Manji, Department of Pediatrics and Child Health, Muhimbili University School of Health and Allied Sciences, Dar es Salaam, Tanzania

30. Estomih Mduma, Haydom Lutheran Hospital, Haydom, Tanzania

31. Venkata Raghava Mohan, Professor, Community Medicine, Christian Medical College, Vellore, India

32. Sophie E. Moore, Department of Women and Children's Health, Kings College London, UK & MRC Unit The Gambia at London School of Hygiene and Tropical Medicine, United Kingdom

33. Mzwakhe Emanuel Nyathi, Department of Animal Services, School of Agriculture, University of Venda, Thohoyandou, South Africa

34. Maribel Paredes Olortegui, AB PRISMA, Iquitos, Peru

35. William A. Petri, University of Virginia, Charlottesville, USA

36. Prasanna Samuel Premkumar, Christian Medical College, Vellore, India

37. Andrew M. Prentice, MRC Unit The Gambia at London School of Hygiene and Tropical Medicine

38. Najeeb Rahman, Aga Khan University, Karachi, Pakistan

39. Kamran Sadiq, Aga Khan University, Karachi, Pakistan

40. Rajiv Sarkar, Christian Medical College, Vellore, Tamil Nadu, India

41. Naomi M. Saville, Institute for Global Health University of College London

42. Bhim P. Shrestha, Health Research and Development Forum, Kathmandu, Nepal

43. Sanjaya Kumar Shrestha, Walter Reed/AFRIMS Research Unit, University of Bergen Soares, Federal University of Ceará, Fortaleza, Brazil

44.   Alberto Melo Soares, Federal University of Ceará, Fortaleza, Brazil

45.   Bakary Sonko, MRC Unit The Gambia at London School of Hygiene and Tropical Medicine,
London, United Kingdom

46.   Erling Svensen, Haukeland University Hospital, Bergen, Norway

47.   Sana Syed, Department of Pediatrics, Division of Gastroenterology, Hepatology & Nutrition
University of Virginia School of Medicine, Charlottesville, USA, and Aga Khan University

48.   Fayaz Umrani, Aga Khan University, Karachi, Pakistan

49.   Honorine D. Ward, Tufts Medical Center, Tufts University School of Medicine, Boston, USA

50.   Pablo Penataro Yori, University of Virginia, Charlottesville, USA

Abstract limit ~200 words; current count 219
Main text limit ~3000 words; current count 2869

Summary

Sustainable Development Goal 2.2.2, to end malnutrition by 2030, measures progress through
elimination of child wasting, defined as weight-for-length more than 2 standard deviations below
international standards. Prevailing methods to measure wasting rely on cross-sectional surveys that
cannot measure onset, recovery, and persistence — key features of wasting epidemiology that could
inform preventive interventions and disease burden estimates. Here, we show through an analysis of
21 longitudinal cohorts that wasting is a highly dynamic process of onset and recovery, and incidence
peaks between birth and 3 months — far earlier than peak prevalence at 12-15 months. By age 24
months 29.2% of children had experienced at least one wasting episode, more than 5-fold higher than
point prevalence (5.6%), demonstrating that wasting incidence is far higher than cross-sectional
surveys suggest. Children wasted before 6 months were more likely to experience concurrent wasting
and stunting (low height-for-age) later, increasing their risk of mortality. In diverse populations with
seasonal rainfall, population average weight-for-length varied substantially (>0.5 z in some cohorts),
with the lowest mean Z-scores during the rainiest months, creating potential for seasonally targeted
interventions. Our results motivate a new focus on extending preventive interventions for wasting to
pregnant and lactating mothers, and for preventive and therapeutic interventions to include children
below age 6 months in addition to current targets of ages 6-59 months.

Introduction

Wasting is a form of undernutrition that results from a loss of muscle and fat tissue from acute
malnutrition, affecting an estimated 45.4 million children under 5 years (6.7%) worldwide with over half
living in South Asia.[1] Children are considered wasted if their weight-for-length z-score (WLZ) falls 2
standard deviations below the median of international standards.[2] Wasted children have weakened
immune systems,[3] predisposing them to infections and more severe illness once infected.[4] Wasting in
very young children increases their risk of mortality (relative risk: 2.3), with mortality risk further
increasing if they are also stunted and underweight, defined as length-for-age (LAZ) below −2 and
weight-for-age (WAZ) below −2 (relative risk: 12.3).[5] Longitudinal studies have shown that wasting is
also associated with higher risk of stunting[6] and poor neurocognitive development at older ages.[7,8]
Treatment of severely wasted children (WLZ below −3) with high calorie, ready-to-use-therapeutic foods

has proven effective,[9,10] but rates of relapse and mortality remain high in this fragile subgroup.[11,12]
Primary prevention of wasting would be preferable to treatment, but there are scant proven preventive
interventions. Key among them, nutritional interventions have led to only small reductions in wasting,
with breastfeeding support from birth not reducing wasting and small-quantity lipid-based nutrient
supplementation between ages 6-24 months leading to a 14% relative reduction in wasting.[13–15]

Wasting is thought to occur primarily from ages 6-24 months during the critical period for
adequate, appropriate and safe complementary feeding,[16,17] but a more complete understanding of the
epidemiology of wasting and how it varies by age is key to develop and target preventive
interventions.[16,18,19] Unlike the cumulative process of linear growth faltering and stunting,[20] wasting
varies considerably over time, both within-individuals and within-populations.[21,22] The dynamic nature of
wasting means that the number of distinct episodes a child experiences may be poorly captured in
cross-sectional, survey-based estimates.[23] Furthermore, seasonal patterns of wasting onset in relation to
changes in food insecurity or disease can only be measured accurately with longitudinal data.[19,24,25] A
synthesis of longitudinal cohort data across diverse populations provides a unique opportunity to study
the timing, dynamics, and burden of wasting in infants and young children — key knowledge gaps to
inform future prevention efforts.

Pooled longitudinal analyses

Here, we report a pooled analysis of 21 longitudinal cohorts from 10 low- and middle-income
countries (LMICs) in South Asia, Sub-Saharan Africa, and Latin America, that measured length and
weight monthly among children 0-24 months of age. Our primary objectives were to produce the first
large-scale estimates of wasting incidence and recovery and to quantify the temporal and regional
variation. We also assessed the concurrence of wasting and stunting and compared estimates of wasting
prevalence and incidence. We analyzed data from the Bill & Melinda Gates Foundation's Knowledge
Integration (*ki*) database, which has aggregated studies on child growth and development.[26] Inclusion
criteria were defined to select cohorts representative of general populations in LMICs with sufficiently
frequent measurements to investigate the acute, dynamic nature of wasting: 1) conducted in low- or
middle-income countries; 2) enrolled children between birth and age 24 months and measured their
length and weight repeatedly over time; 3) did not restrict enrollment to acutely ill children; 4) had
median birth year of 1990 or later 5) measured anthropometry at least monthly (Extended Data Fig 1).
Twenty-one cohorts met the inclusion criteria, encompassing 11,448 children and 198,154 total
anthropometry measurements from birth to age 24 months (Fig 1).

We calculated WLZ and LAZ-scores using WHO 2006 growth standards.[27] We dropped 385
biologically implausible measurements (0.2%) of WLZ (> 5 or < –5 Z-score) and 212 (0.1%) of LAZ (> 6 or
< –6 Z-score), following WHO recommendations.[28] Most included cohorts did not measure children past
24 months, so analyses focused on birth to 24 months. Cohorts ranged in size from 160 children in the
TDC cohort to 2,931 children from the MRC Keneba cohort (Fig 1). Unless otherwise indicated, we
conducted individual-level analyses within cohorts and then pooled cohort-specific estimates using
random effects models fit with restricted maximum likelihood estimation, a conservative approach
when cohort-specific estimates are heterogeneous. Cohorts were distributed throughout South Asia,
Africa, and Latin America, but did not cover entire regions (Extended Data Fig 2). Most cohorts
measured children every month through age 24 months, but there was some attrition as children aged
and there were four cohorts with few measurements beyond 15 months (Extended Data Fig 3). Pooled
estimates at older ages could be slightly biased if cohorts or children who were not measured were
systematically different from those that were included — for example, if children who were lost were
more likely to be wasted, we might have under-estimated wasting at older ages.

We assessed *ki* cohort representativeness by comparing Z-score measurements to population-based
samples in Demographic and Health Surveys (DHS). Mean z-scores in *ki* cohorts were generally
representative based on country-level DHS estimates, with lower WLZ (overall and by age) in Guatemala,
Pakistan, and South Africa and higher WLZ in Brazil and Guatemala (Extended Data Fig 4). LAZ was
generally lower in *ki* cohorts, so estimates of concurrent wasting and stunting may be higher than
estimates from population-based samples, and rates of wasting recovery in early life may be higher
compared with the general population.
182
Age-specific patterns of wasting

Across all cohorts, mean WLZ was near –0.5 at birth and then increased over the first 6 months
before decreasing until 12 months (Fig 2a). Age-specific patterns in WLZ were similar across geographic
regions, but levels varied substantially by region. WLZ was markedly lower among South Asian cohorts,
reflecting a much higher burden of malnutrition (Fig 2a). Children were wasted for 16,139 (8.6%)
measurements, and severely wasted for 3,391 (1.8%) measurements. Wasting prevalence was highest at
birth (11.9%; 95% CI: 7.0, 19.5) in contrast with prior studies that showed peak wasting prevalence
between 6-24 months old (Fig 2b).[6,29–31] Across regions, wasting prevalence decreased from birth to
three months and then increased until 12 months old, but was far higher in South Asian cohorts. In
South Asian cohorts, where low birthweight is common,[32] at-birth wasting prevalence was 18.9% (95%
CI: 15.0, 23.7), implicating causes of poor fetal growth like maternal malnutrition, maternal morbidities,
and maternal small stature as key regional drivers of wasting.[33,34] Severe wasting followed a similar
pattern but was much rarer (Extended Data Fig 5).

Wasting onset was highest during the first 3 months largely due to its high occurrence at birth (Fig
2c). Overall, 12.9% (95% CI: 7.6, 20.1) of all children experienced wasting by age 3 months, which
accounted for almost half (47.8%) of children who ever experienced wasting in their first two years of
life. A focus on wasting prevalence alone masked how common it was for children to experience wasting
in their first 24 months. After birth, up to 6.5% (95% CI: 4.9, 8.6) of children were wasted at a specific
visit, but 29.2% (95% CI: 17.5, 44.7) of children experienced at least one wasting episode by age 24
months, and in South Asian cohorts the cumulative incidence by age 24 months was 52.2% (95% CI:
43.5, 60.7, Figs 2b, c).

WHO Child Growth Standards overestimate wasting at birth among children born preterm,[35] though
adjustment for gestational age among four cohorts with available data only reduced at-birth stunting
prevalence by 0.8% and increased underweight prevalence by 0.7% (Extended data Fig 6). Even if
preterm birth were to account for some of at-birth wasting, the small birth size, irrespective of cause,
documented in this analysis raises concern because of its consequences for child growth faltering and
mortality during the first 24 months.[34]
210
Seasonality of wasting

We joined monthly rainfall totals to cohorts by location and year, and for each cohort we
estimated a seasonality index based on rainfall (details in Methods). We examined average WLZ over the
calendar year and examined seasonal changes with respect to rainfall, under the hypothesis that
seasonal changes in food availability and infection tied to rainfall could cause seasonal wasting.[36] Mean
WLZ varied dramatically by calendar date in almost all cohorts, with a consistent minimum mean WLZ
coinciding with peak rainfall (Fig 3a). Mean WLZ was -0.16 (95% CI: -0.19, -0.14) lower during the three-
month period of peak rainfall, compared to the mean WLZ in the opposite three-month period of the
year, when pooled across cohorts (Fig. 3b). Mean seasonal decline in WLZ was -0.27 (95% CI: -0.31, -
0.24) in cohorts with a seasonality index > 0.9, but some cohorts evidenced seasonal WLZ declines of

more than –0.5 (Fig. 3b).

South Asian cohorts had temporally synchronous rainfall, so we estimated WLZ at birth by
calendar month, pooled across the 11 South Asian cohorts. Mean WLZ at birth varied by up to 0.72 Z
(95% CI: 0.43, 1.02) depending on the month the child was born (range: -0.5 Z to -1.3 Z; Fig 3c), which
suggests that seasonally-influenced maternal nutrition,[37] likely mediated through intrauterine growth
restriction or preterm birth,[38] was a major determinant of child WLZ at birth. Birth month also
influenced the effect of season on WLZ trajectories that persisted through a child's second year of life
(Extended Data Fig 7).

Wasting incidence and recovery

There was high variability in WLZ across longitudinal measurements, with an average within-
child standard deviation of WLZ measurements of 0.76, compared to 0.64 for LAZ and 0.52 for WAZ. We
thus defined unique wasting episodes by imposing a 60-day recovery period, covering two consequent
monthly measurements when a child's WLZ measurements needed to remain above –2 to be considered
recovered and at risk for a future episode (Fig 4a). Children who were born wasted were considered to
have an incident episode of wasting at birth. The mean number of wasting episodes experienced and
wasting incidence at all ages was higher in South Asia, with highest incidence in the first 3 months with-
or without- including episodes at birth (Fig 4b).

Ultimately, most children recovered from moderate or severe wasting episodes, reinforcing its
status as an acute condition. Children met our definition of "recovered" in 91.5% of episodes of
moderate and 82.5% of episodes of severe wasting. Loss-to-followup may have affected recovery
estimates, recovery can only occur in children surviving the episode, and we did not have data on
wasting treatment, so recovery may be higher in these highly monitored cohorts than in the general
population due to treatment referrals. Pooling across cohorts, children recovered from 39.0% (95% CI:
34.0, 44.3) of wasting episodes within 30 days, 64.7% (95% CI: 59.3, 69.4)) within 60 days, and 71.1%
(95% CI: 66.1, 75.6) within 90 days. Median episode length was 42 (95% CI: 39, 44) days. We examined
the distribution of WLZ in the three-month period after a child was considered recovered from wasting,
stratified by the age at which the episode occurred. We found that children recovered to a higher WLZ
from at-birth wasting or wasting episodes in the 0-6 month age window compared with episodes that
occurred at older ages (Fig 4c). Regression to the mean (RTM) could explain some wasting recovery,
especially the rapid WLZ gain among children born wasted (Figure 4e), but there was catch-up growth
beyond the calculated RTM effect (Extended Data Fig 8).[39] Additionally, we required a 60-day recovery
period to better capture true recovery, and mean WLZ of recovered children did not regress to cohort
means (Figure 4e), especially among older children (Figure 4c). Consistent with larger WLZ increases
during recovery from wasting at younger ages, a larger proportion of children recovered within 30, 60,
and 90 days if the wasting episode occurred before age 6 months (Fig 4d). South Asian cohorts had
lower rates of recovery in the 6-18 month age period compared with other regions (Fig 4d). On average,
however, the WLZ of children born wasted did not catch up to the WLZ of children not born wasted, and
children who were born wasted but recovered had a higher cumulative incidence of wasting after 6
months of age (38.1% cumulative incidence in children born wasted [95% CI: 28.2, 49.2] vs. 24.2%
cumulative incidence in children not born wasted [95% CI: 17.6, 32.4], Fig 4e-f).

Persistent and concurrent growth faltering

We examined more severe forms of growth faltering, including persistent wasting and
concurrent wasting and stunting, because these conditions are associated with higher mortality risk.[5]
We first identified a subset of children who experienced persistent wasting during their first 24 months.

We used a pragmatic definition[40] that classified children as persistently wasted if ≥ 50% of their WLZ
measurements from birth to 24 months fell below –2, capturing both frequent short wasting episodes or
less frequent, longer episodes. Among 10,374 children with at least four measurements, 3.4% (95% CI:
2.0, 5.6) were persistently wasted. Persistent wasting over the first 24 months of life was highest in
South Asia (7.2%, 95% CI: 5.1, 10.4, Extended Data Fig 5a). Among children wasted at birth, 10.9% were
persistently wasted after 6 months (95% CI: 8.7, 13.5) in contrast to 6.3% of children not born wasted
who were persistently wasted after 6 months (95% CI: 4.3, 9.0).

Next, we examined the cumulative incidence of concurrent wasting and stunting and the timing
of their overlap. Overall, 10.6% of children experienced concurrent wasting and stunting before 2 years
of age (Extended data figure 5d), and a further 1.5% experienced concurrent severe wasting and
stunting — far higher than point prevalence estimates in the present study (4.1% at 24 months, 95% CI:
2.6, 6.4; Fig 5a) or the 4.3% prevalence estimated in children under 5 in LMIC from cross-sectional
surveys.[41,42] Concurrent wasting and stunting was most common in South Asia, with highest prevalence
at age 21 months (Fig 5a), driven primarily by increases in stunting prevalence as children aged.[20]
Children wasted and stunting are also underweight, as their maximum possible weight-for-age z-score is
below -2.35.[43] Almost half of children who had prevalent growth faltering met two or more of these
three growth faltering conditions, with 17-22% of all children experiencing multiple conditions after age
12 months (Fig 5b). Longitudinal analyses showed early growth faltering predisposed children to
experience concurrent growth faltering at older ages: children who were wasted by age 6 months were
1.8 (95% CI: 1.6, 2.1) times and children stunted before 6 months were 2.9 (95% CI: 2.5, 3.4) times more
likely to experience concurrent wasting and stunting between ages 18-24 months. A companion article
reports an in depth investigation of key determinants of persistent wasting and concurrent wasting and
stunting, along with their consequences for severe growth faltering and mortality.[34]

Discussion

Nearly all large-scale studies of child wasting including DHS report point prevalence of wasting,
which has enabled broad comparisons between populations but failed to capture the number of
episodes of wasting during the course of childhood.[44,45] By combining information across several
longitudinal cohorts, this study demonstrated that children experience far higher cumulative incidence
of wasting between birth and 24 months (29.2%) than previously known. Children in all included cohorts
experienced higher incidence compared with prevalence due to the episodic nature of acute
malnutrition, but the pattern was most stark in South Asian cohorts where, from birth to 24 months and
pooled across cohorts, 29.2% of children experienced at least one wasting episode, 3.4% were
persistently wasted, and 10.6% had experienced concurrent wasting and stunting. This evidence shows
that the cumulative, child-level burden of wasting is higher than cross-sectional prevalence measures
would suggest, and assessments of wasting and stunting in isolation fail to account for their joint
burden, which can be substantial. This is of particular relevance in South Asia where the largest number
of stunted and wasted children live, and while stunting prevalence is decreasing, wasting prevalence is
not.[1,46]

Our results show consistent patterns in the timing of wasting onset by season and by age, with
important implications for preventive interventions. WLZ varied dramatically by season in most cohorts,
and in South Asia WLZ varied by up to 0.7 z-scores at-birth with consequences that persisted throughout
the first 24 months (Fig 3c, Extended Data Figure 7). The present study has not elucidated the
mechanism that links seasonal rainfall with wasting, indeed mechanisms could differ by geographic and
cultural context. Nevertheless, the high degree of consistency of the rainfall-wasting association
provides strong support for the development of seasonally targeted interventions to prevent wasting in
food-insecure populations, akin to seasonal malaria chemoprevention programs in the Sahel.[47] Wasting

incidence was highest from birth to 3 months, and among children who experienced any wasting during
their first 24 months, 48% experienced their first episode by 3 months. Children born wasted were far
more likely to experience more severe forms of growth faltering at older ages including persistent
wasting from 6-24 months and concurrent wasting and stunting at 18 months. Wasting treatment
programs have traditionally focused on children from 6-59 months.

Sustainable Development Goal 2.2.2 calls to eliminate malnutrition by 2030, with the
elimination of child wasting as its primary indicator.[48] To help achieve this goal, preventive interventions
could complement a historic focus on treatment for acutely malnourished children. Our results should
motivate preventive and therapeutic programs to consider extending efforts to maternal support and
education during pregnancy, and community-based breastfeeding support through infancy.[49,50] If
preventive or therapeutic interventions focus on ages earlier than 6 months, then they must integrate
carefully with current recommendations for exclusive breastfeeding.

326
327
328

References
1. WHO | UNICEF-WHO-The World Bank: Joint child malnutrition estimates - Levels and trends. *WHO*
http://www.who.int/nutgrowthdb/estimates/en/.
2. Onis, M. de. WHO Child Growth Standards based on length/height, weight and age. *Acta Paediatr.*
95, 76–85 (2006).
3. Rytter, M. J. H., Kolte, L., Briend, A., Friis, H. & Christensen, V. B. The Immune System in Children
with Malnutrition—A Systematic Review. *PLOS ONE* **9**, e105017 (2014).
4. Chang, C. Y. *et al.* Children successfully treated for moderate acute malnutrition remain at risk for
malnutrition and death in the subsequent year after recovery. *J. Nutr.* **143**, 215–220 (2013).
5. McDonald, C. M. *et al.* The effect of multiple anthropometric deficits on child mortality: meta-analysis
of individual data in 10 prospective studies from developing countries. *Am. J. Clin. Nutr.* **97**, 896–901
(2013).
6. Schoenbuchner, S. M. *et al.* The relationship between wasting and stunting: a retrospective cohort
analysis of longitudinal data in Gambian children from 1976 to 2016. *Am. J. Clin. Nutr.*
doi:10.1093/ajcn/nqy326.
7. Khandelwal, N. *et al.* Determinants of motor, language, cognitive, and global developmental delay in
children with complicated severe acute malnutrition at the time of discharge: An observational study
from Central India. *PLOS ONE* **15**, e0233949 (2020).
8. Sudfeld, C. R. *et al.* Malnutrition and Its Determinants Are Associated with Suboptimal Cognitive,
Communication, and Motor Development in Tanzanian Children. *J. Nutr.* **145**, 2705–2714 (2015).
9. Bhutta, Z. A. *et al.* Evidence-based interventions for improvement of maternal and child nutrition:
what can be done and at what cost? *The Lancet* **382**, 452–477 (2013).
10. World Health Organization & Nutrition for Health and Development. *Guideline.* (2013).
11. Stobaugh, H. C. *et al.* Relapse after severe acute malnutrition: A systematic literature review and
secondary data analysis. *Matern. Child. Nutr.* **15**, e12702 (2019).
12. O'Sullivan, N. P., Lelijveld, N., Rutishauser-Perera, A., Kerac, M. & James, P. Follow-up between 6
and 24 months after discharge from treatment for severe acute malnutrition in children aged 6-59
months: A systematic review. *PLOS ONE* **13**, e0202053 (2018).
13. Dewey, K. G. *et al.* Characteristics that modify the effect of small-quantity lipid-based nutrient
supplementation on child growth: an individual participant data meta-analysis of randomized
controlled trials. *medRxiv* 2021.02.05.21251105 (2021) doi:10.1101/2021.02.05.21251105.
14. Rana, R., McGrath, M., Sharma, E., Gupta, P. & Kerac, M. Effectiveness of Breastfeeding Support
Packages in Low- and Middle-Income Countries for Infants under Six Months: A Systematic Review.

Nutrients **13**, 681 (2021).

15. Keats, E. C. *et al.* Effective interventions to address maternal and child malnutrition: an update of the
evidence. *Lancet Child Adolesc. Health* **5**, 367–384 (2021).

16. Lenters, L., Wazny, K. & Bhutta, Z. A. Management of Severe and Moderate Acute Malnutrition in
Children. in *Reproductive, Maternal, Newborn, and Child Health: Disease Control Priorities, Third*
Edition (Volume 2) (eds. Black, R. E., Laxminarayan, R., Temmerman, M. & Walker, N.) (The
International Bank for Reconstruction and Development / The World Bank, 2016).

17. Das, J. K., Salam, R. A., Imdad, A. & Bhutta, Z. A. Infant and Young Child Growth. in *Reproductive,*
Maternal, Newborn, and Child Health: Disease Control Priorities, Third Edition (Volume 2) (eds.
Black, R. E., Laxminarayan, R., Temmerman, M. & Walker, N.) (The International Bank for
Reconstruction and Development / The World Bank, 2016).

18. Bhutta, Z. A. *et al.* What works? Interventions for maternal and child undernutrition and survival.
Lancet **371**, 417–440 (2008).

19. Nabwera, H. M., Fulford, A. J., Moore, S. E. & Prentice, A. M. Growth faltering in rural Gambian
children after four decades of interventions: a retrospective cohort study. *Lancet Glob. Health* **5**,
e208–e216 (2017).

20. Benjamin-Chung, J. et. al. (submitted). Early childhood linear growth faltering in low-and middle-
income countries. (2020).

21. Stobaugh, H. C. *et al.* Children with Poor Linear Growth Are at Risk for Repeated Relapse to Wasting
after Recovery from Moderate Acute Malnutrition. *J. Nutr.* **148**, 974–979 (2018).

22. Osgood-Zimmerman, A. *et al.* Mapping child growth failure in Africa between 2000 and 2015. *Nature*
555, 41–47 (2018).

23. Bulti, A. *et al.* Improving estimates of the burden of severe acute malnutrition and predictions of
caseload for programs treating severe acute malnutrition: experiences from Nigeria. *Arch. Public*
Health **75**, (2017).

24. Egata, G., Berhane, Y. & Worku, A. Seasonal variation in the prevalence of acute undernutrition
among children under five years of age in east rural Ethiopia: a longitudinal study. *BMC Public Health*
13, (2013).

25. Kinyoki, D. K. *et al.* Space-time mapping of wasting among children under the age of five years in
Somalia from 2007 to 2010. *Spat. Spatio-Temporal Epidemiol.* **16**, 77–87 (2016).

26. Peppard, T. *et al.* (submitted). Combined longitudinal growth cohorts from infants born in South Asia,
Sub-Saharan Africa and Latin America. (2020).

27. WHO Multicentre Growth Reference Study Group. WHO child growth standards: length/height-for-
age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and
development. *Geneva World Health Organ.* 312 pages (2006).

28. Organization, W. H. & Fund (UNICEF), U. N. C. *Recommendations for data collection, analysis and*
reporting on anthropometric indicators in children under 5 years old. (World Health Organization,
2019).

29. de Onis, M. *et al.* Comparison of the World Health Organization (WHO) Child Growth Standards and
the National Center for Health Statistics/WHO international growth reference: implications for child
health programmes. *Public Health Nutr.* **9**, 942–947 (2006).

30. Garenne, M., Myatt, M., Khara, T., Dolan, C. & Briend, A. Concurrent wasting and stunting among
under-five children in Niakhar, Senegal. *Matern. Child. Nutr.* **15**, (2019).

31. Payandeh, A., Saki, A., Safarian, M., Tabesh, H. & Siadat, Z. Prevalence of Malnutrition among
Preschool Children in Northeast of Iran, A Result of a Population Based Study. *Glob. J. Health Sci.* **5**,
208–212 (2013).

32. Blencowe, H. *et al.* National, regional, and worldwide estimates of low birthweight in 2015, with trends
from 2000: a systematic analysis. *Lancet Glob. Health* **0**, (2019).

33. Ashorn, P. *et al.* Co-causation of reduced newborn size by maternal undernutrition, infections, and
inflammation. *Matern. Child. Nutr.* **14**, e12585 (2018).

34. Mertens, A. *et al.* (submitted). Causes and consequences of child growth faltering in low- and middle-

income countries. (2020).

35. Perumal, N. *et al.* Effect of correcting for gestational age at birth on population prevalence of early childhood undernutrition. *Emerg. Themes Epidemiol.* **15**, 3 (2018).

36. Levy, K., Woster, A. P., Goldstein, R. S. & Carlton, E. J. Untangling the Impacts of Climate Change on Waterborne Diseases: a Systematic Review of Relationships between Diarrheal Diseases and Temperature, Rainfall, Flooding, and Drought. *Environ. Sci. Technol.* **50**, 4905–4922 (2016).

37. Saville, N. M. *et al.* Comprehensive analysis of the association of seasonal variability with maternal and neonatal nutrition in lowland Nepal. *Public Health Nutr.* 1–16 (2021) doi:10.1017/S1368980021003633.

38. Christian, P. *et al.* Risk of childhood undernutrition related to small-for-gestational age and preterm birth in low- and middle-income countries. *Int J Epidemiol* **42**, 1340–1355 (2013).

39. Barnett, A. G., van der Pols, J. C. & Dobson, A. J. Regression to the mean: what it is and how to deal with it. *Int. J. Epidemiol.* **34**, 215–220 (2005).

40. Richard, S. A. *et al.* Wasting Is Associated with Stunting in Early Childhood123. *J. Nutr.* **142**, 1291–1296 (2012).

41. Khara, T., Mwangome, M., Ngari, M. & Dolan, C. Children concurrently wasted and stunted: A meta-analysis of prevalence data of children 6–59 months from 84 countries. *Matern. Child. Nutr.* **14**, e12516 (2018).

42. Victora, C. G. *et al.* Revisiting maternal and child undernutrition in low-income and middle-income countries: variable progress towards an unfinished agenda. *The Lancet* **397**, 1388–1399 (2021).

43. Myatt, M. *et al.* Children who are both wasted and stunted are also underweight and have a high risk of death: a descriptive epidemiology of multiple anthropometric deficits using data from 51 countries. *Arch. Public Health* **76**, (2018).

44. Akombi, B. J. *et al.* Stunting, Wasting and Underweight in Sub-Saharan Africa: A Systematic Review. *Int. J. Environ. Res. Public. Health* **14**, (2017).

45. Harding, K. L., Aguayo, V. M. & Webb, P. Factors associated with wasting among children under five years old in South Asia: Implications for action. *PloS One* **13**, e0198749 (2018).

46. Mutunga, M., Frison, S., Rava, M. & Bahwere, P. The Forgotten Agenda of Wasting in Southeast Asia: Burden, Determinants and Overlap with Stunting: A Review of Nationally Representative Cross-Sectional Demographic and Health Surveys in Six Countries. *Nutrients* **12**, 559 (2020).

47. Baba, E. *et al.* Effectiveness of seasonal malaria chemoprevention at scale in west and central Africa: an observational study. *The Lancet* **396**, 1829–1840 (2020).

48. United Nations. Goal 2. Sustainable Development Knowledge Platform. https://sustainabledevelopment.un.org/sdg2.

49. Kerac, M., Mwangome, M., McGrath, M., Haider, R. & Berkley, J. A. Management of Acute Malnutrition in Infants Aged under 6 Months (MAMI): Current Issues and Future Directions in Policy and Research. *Food Nutr. Bull.* **36**, S30–S34 (2015).

50. von Salmuth, V. *et al.* Maternal-focused interventions to improve infant growth and nutritional status in low-middle income countries: A systematic review of reviews. *PloS One* **16**, e0256188 (2021).

Figure 1 | Summaries of included cohorts.

(**a**) Number of observations (1000s) across cohorts by age in months. (**b**) Mean weight-for-length Z-scores (WLZ) by age in months for each included cohort. Cohorts are sorted by geographic region and overall mean weight-for-length Z-score. The country of each cohort and the start year is printed by each cohort name. (**c**) The number of observations included in each cohort. (**d**) Overall wasting prevalence by cohort, defined as proportion of measurements with WLZ < –2.

Figure 2 | WLZ, prevalence and incidence of wasting by age and region.

(**a**) Mean weight-for-length Z-score (WLZ) by age in 21 longitudinal cohorts, overall (N=21 studies; N=4,165-10,886 observations per month) and stratified by region (Africa: N=4 studies, N=1,067-5,428 observations; Latin America: N=6 studies, N= 569-1718 observations, South Asia 11 studies, N=2,382-4,286 observations).

(**b**) Age-specific wasting prevalence, defined as WLZ < –2, overall (N=3,985-9,906children) and stratified by region (Africa: 1,701-5,017 children, Latin America: N= 290-1397 children, South Asia: N=1,994-3,751 children).

(**c**) Age-specific wasting incidence overall (N=6,199 -10,377 children) and stratified by region (Africa: N=2,249-5,259 children, Latin America: N=763-1,437 children, South Asia: N=3,076-3,966 children). Cumulative incidence measures the proportion of children who have ever experienced wasting since birth, while the new incident cases represent the proportion of children at risk who had an episode of wasting begin during the age period.

Error bars in panels **b** and **c** are 95% confidence intervals for pooled estimates. In each panel, grey curves or points show cohort-specific estimates.

Figure 3 | Mean weight-for-length Z-scores by age and season.

(a) Cubic splines of mean WLZ over day of the year, superimposed over histograms of monthly mean rainfall over study periods, with the seasonality index printed in parentheses beside the cohort name. Panels are sorted by country and splines colored by high (≥ 0.9), medium (< 0.9 and ≥ 0.7), and low seasonality categories (< 0.7). Sample sizes range from 160 children (2,545 measurements) in TDC to 2,545 (40,115 measurements) in the Keneba cohort. **(b)** Mean differences in child WLZ between quarters of the year defined around the adjacent 3-month periods with the highest mean rainfall, pooled across cohorts in panel **a,** overall (N= 21 cohorts, 2,545-40,115 observations) and by seasonality index (high: N=7 cohorts, 3,164-40,115 observations, medium: N=8 cohorts, 2,545-9,202 observations, low: N=6 cohorts, 2,741-29,518observations).

(c) Cubic spline of mean WLZ at birth across birth months among 1,821 children with WLZ measured at birth in ten South Asian cohorts, with mean differences in birth WLZ by month of birth in the inset plot.

(**a**) Example child WLZ trajectory and wasting episode classifications. The age of wasting onset was
assumed to occur halfway between a measurement of WLZ < −2 and the previous measurement of WLZ
≥ −2. Recovery from an episode of wasting or severe wasting occurred when a child had measurements
of WLZ ≥ −2 for at least 60 days, with the age of recovery assumed to be halfway between the last
measurement of WLZ < −2 and the first measurement of WLZ ≥ −2.

(**b**) Wasting incidence rate per 1,000 days at risk, stratified by age and region (Overall: N= 510,070-
822,802 person-days per estimate, Africa: N=245,046-407,940 person-days, Latin America: N= 65,058-
142,318 person-days, South Asia: N=182,694-315,057 person-days). Insets are histograms of the
number of wasting episodes per child by region, with the distribution mean (SD) printed within the plot.

(**c**) The distribution of children's mean WLZ in the three months following recovery from wasting, with
inter-quartile range marked. The median WLZ is annotated, and gray points mark cohort-specific
medians. Children wasted before age 6 months experienced larger improvements in WLZ compared with
children wasted at older ages (p<0.001). The analysis uses 3,686 observations of 2,301 children who

recovered from wasting episodes, with　1,264 observations at birth,　628 observations from 0-6 months,
824 observations from 6-12 months, and　970 observations from 12-18 months.

(**d**) Percentage of children who recovered from wasting within 30, 60, and 90 days of episode onset
(N=21 cohorts,　5,549 wasting episodes).

(**e**) Mean WLZ by age, stratified by wasting status at birth (which includes the first measure of a child
within 7 days of birth), shows that children born wasted (N= 814 children,　14,351 observations) did not
catch up to children not born wasted (N= 3,355 children,　62,568 observations).

(**f**) Higher measures of wasting after 6 months of age among children born wasted (N=814) compared to
those who were not (N=3,355). In panels b, d, and f, vertical lines mark 95% confidence intervals for
pooled means of study specific estimates (light points).

Figure 5| Co-occurrence of wasting, stunting, and underweight.

(**a**) Age-specific prevalence of co-occurrent wasting and stunting overall (N=3,984 to 9,899 children) and stratified by region (Africa: N=1,799-5,014, Latin America: N=290-1,397 children, South Asia: N=1,994-3,747 children). Vertical lines mark 95% confidence intervals for pooled means across study-specific estimates (gray points).

(**b**) Percentage of children classified by different measures of growth faltering, alone or combined. Children classified as "never faltered" had not previously been wasted, stunted, or underweight. Children in the "recovered" category were not wasted, stunted, or underweight but had experienced at least one of these conditions previously. All children who were wasted and stunted were also underweight. Proportions in each category were calculated within cohorts, pooled using random effects, and scaled so percentages added to 100%. The number of children contributing to each age ranges from 3,920 to 9,077 children, with 11,409 total children.

Extended Data Figure 1| *ki* cohort selection. Analyses focused on longitudinal cohorts to enable the
estimation of prospective incidence rates and growth velocity. On July 15, 2018, there were 86
longitudinal studies on GHAP. From this set, we applied five inclusion criteria to select cohorts for
analysis. Our rationale for each criterion follows. (1) *Studies were conducted in lower income or middle-*
income countries. Our target of inference for analyses was children in LMICs, which remains a key target
population for preventive interventions. (2) *Studies measured length and weight between birth and age 24*
months. We were principally interested in growth faltering during the first two years of life including at
birth, thought to be the key window for linear growth faltering [7]. (3) *Studies did not restrict enrollment to*
acutely ill children. Our focus on descriptive analyses led us to target, to the extent possible, the general
population. We thus excluded some studies that exclusively enrolled acutely ill children, such as children
who presented to hospital with acute diarrhea or who were severely malnourished. (4) *Studies enrolled at*
least 200 children. Age-stratified incident episodes of stunting and wasting were sufficiently rare that we
wanted to ensure each cohort would have enough information to estimate rates before contributing to
pooled estimates. (5) *Studies collected anthropometry measurements at least every 3 months*. We limited
studies to those with higher temporal resolution to ensure that we adequately captured incident episodes
and recovery. We further restricted analyses of wasting incidence and recovery to cohorts with monthly
measurements because of high temporal variation in WHZ within individuals.

550
551
552
553

556 **Extended Data Figure 2 | Geographic location of *ki* cohorts.** Locations are
557 approximate and jittered slightly for display.

560 Extended Data Figure 3 | Percentage of enrolled children measured in each ki
561 cohort with monthly measurements.

a

563
564

b

Extended Data Figure 4 | Comparison of cohort anthropometry to population-based samples

(**a**) Kernel density distributions of length-for-age (LAZ), weight-for-age (WAZ), and weight-for-length Z-scores (WLZ) from measurements among children under 24 months old in 21 *ki* longitudinal cohorts (colored line) and among children measured in the most recent population-based, Demographic and Health Survey for each country (black). Sub-Saharan African countries are colored blue, Latin American countries are colored green, and south Asian countries are colored orange. Median Z-scores are denoted with points under the density curves, with open circles for *ki* cohorts and solid points for Demographic and Health Surveys.

(**b**) Mean LAZ, WAZ, and WLZ by age and country among 21 *ki* longitudinal cohorts (dashed lines) and in Demographic and Health Surveys (solid). Means estimated with cubic splines and shaded regions show approximate, simultaneous 95% confidence intervals.

Extended Data Figure 5 | Prevalence of persistent wasting, severe wasting, and underweight by region.

(**a**) Proportion of children persistently wasted (≥50% of measurements from birth to 24 months of age, overall and stratified by region.

(**b**) Prevalence of severe wasting (WLZ < –3) by age and region.

(**c**) Prevalence of underweight (weight-for-age Z-score < –2) by age and region.

(**d**) Incidence proportion of concurrent wasting and stunting by age and region. Across all ages before 24 months, 10.6% of children experienced at least one concurrently wasted and stunted measurement.

Extended Data Figure 6 | Comparison of underweight and stunting prevalence at birth with and without gestational age correction

This figure includes the results from correcting at-birth Z-scores in the *ki* cohorts that measured gestational age (GA). The corrections are using the Intergrowth standards and are implemented using the R *growthstandards* package (https://ki-tools.github.io/growthstandards/). Overall, the prevalence at birth decreased slightly after correcting for gestational age, but the cohort-specific results are inconsistent. Observations with GA outside of the Intergrowth standards range (<168 or > 300 days) were dropped for both the corrected and uncorrected data. Prevalence increased after GA correction in some cohorts due to high rates of late-term births based on reported GA. There were no length measurements at birth in the Tanzania Child 2 cohort, so they do not have stunting estimates. There were 4,449 measurements used in the underweight analysis and 1,931 measurements used in the stunting analysis. Gestational age was estimates based on mother's recall of the last menstrual period in the IRC, CMC-V-BCS-2002, and Tanzania Child cohorts, and was based on the Dubowitz method (newborn exam) in the MRC Keneba cohort.

a

b

Extended Data Figure 7 | Month of birth affects seasonal patterns in WLZ

(**a**) Mean WLZ by calendar month among South Asian cohorts, with children stratified by birth month. The X-axis begins at January 1st of the year of the first January birthday in each cohort. Grey backgrounds indicate the approximate timing of seasonal monsoons in South Asia (June-September) Shaded regions around spline fits indicate 95% simultaneous confidence intervals. Eleven cohorts, 4,040 children, and 78,573 measurements were used to estimate the splines. South Asian children born in July-September had the lowest mean WLZ overall and children born April-September had larger seasonal declines in WLZ during their second year of life than children born October-March.

(**b**) Mean WLZ from birth to age 24 months among children from South Asian cohorts stratified by birth month. Shaded regions around spline fits indicate 95% simultaneous confidence intervals. Data used is the same as in panel (a).

Extended Data Figure 8 | Regression to the mean effects in wasted children

(a) Expected mean Z-scores based on the regression to the mean effect (orange) and observed mean Z-scores (blue) 3 months after wasted children are measured. The lines connecting cohort-specific observed and expected WLZ at each age are colored orange if the expected estimate under RTM was higher than the observed mean (indicating lower than expected change in WLZ under RTM alone), and blue if the observed mean was higher than the expected estimate under RTM (indicating higher than

expected change in WLZ under RTM alone). For examples, most cohorts experienced larger increases in
WLZ than expected in the three-month period beginning in their first month of life (blue lines) and most
cohorts experienced smaller increases in WLZ than expected in the three month periods beginning at
ages 6-9 months (orange lines).
(b) Difference between observed means and expected means under a pure RTM effect by cohort, with
the median differences by age indicated with horizontal lines.
Details on estimation of the RTM effects are in the methods

 Extended Data Table 1| Summary of *ki* cohorts

Region, Study ID	Country	Study Years	Design	Children Enrolled*	Anthropometry measurement ages (months)	Total WLZ measurements*	Primary References
South Asia							
Biomarkers for EE	Pakistan	2013-2015	Prospective cohort	380	Birth, 1, 2, ..., 18	8428	Iqbal et al 2018 Nature Scientific Reports[1]
Resp. Pathogens	Pakistan	2011 - 2014	Prospective cohort	284	Birth, 1, 2, ..., 17	3164	Ali et al 2016 Journal of Medical Virology[2]
Growth Monitoring Study	Nepal	2012 - Ongoing	Prospective cohort	686	Birth, 1, 2, …, 24	13340	Not yet published
MAL-ED	Nepal	2010 - 2014	Prospective cohort	240	Birth, 1, 2, …, 24	5695	Shrestha et al 2014 Clin Infect Dis[3]
CMIN93	Bangladesh	1993 - 1996	Prospective Cohort	272	Birth, 3, 6, ..., 24	5372	Pathela et al. 2006 Acta Paediatrica
TDC	India	2008-2011	Quasi-experimental	160	Birth, 1, 2, ..., 24	3591	Sarkar et al. 2013 BMC Public Health
CMC Birth Cohort, Vellore	India	2002 - 2006	Prospective cohort	373	Birth, 0.5, 1, 1.5, ..., 24	8697	Gladstone et al. 2011 NEJM[4]
MAL-ED	India	2010 - 2012	Prospective cohort	251	Birth, 1, 2, ..., 24	5697	John et al 2014 Clin Infect Dis[5]
Vellore Crypto Study	India	2008 - 2011	Prospective cohort	410	Birth, 1, 2, ..., 24	9729	Kattula et al. 2014 BMJ Open[6]
MAL-ED	Bangladesh	2010 -2014	Prospective cohort	263	Birth, 1, 2, ..., 24	5592	Ahmed et al 2014 Clin Infect Dis[7]
PROVIDE RCT	Bangladesh	2011 -2014	Individual RCT	700	Birth, 6, 10, 12, 14. 17, 18, 24, 39, 40, 52, 53 (weeks)	9202	Kirkpatrick et al 2015 Am J Trop Med Hyg[8]
Africa							
MAL-ED	Tanzania	2009 - 2014	Prospective cohort	261	Birth, 1, 2, ..., 24	5698	Mduma et al 2014 Clin Infect Dis[9]
Tanzania Child 2	Tanzania	2007 - 2011	Individual RCT	2396	1, 2, ..., 20	29518	Locks et al Am J Clin Nutr 2016[10]
MAL-ED	South Africa	2009 - 2014	Prospective cohort	312	Birth, 1, 2, ..., 24	6151	Bessong et al 2014 Clin Infect Dis[11]
MRC Keneba	Gambia	1987 - 1997	Cohort	2920	Birth, 1, 2, ..., 24	40117	Schoenbuchner et al. 2019, AJCN[12]
Latin America							
CMIN Peru95	Peru	1995 - 1998	Prospective Cohort	224	Birth, 1, 2, ..., 24	3978	Checkley et al. 2003 Am J Epidemiol.
CMIN Peru89	Peru	1989 - 1991	Prospective Cohort	210	Birth, 1, 2, ..., 24	2741	Checkley et al. 1998 Am J Epidemiol.
MAL-ED	Peru	2009 - 2014	Prospective cohort	302	Birth, 1, 2, ..., 24	6127	Yori et al 2014 Clin Infect Dis[13]
CONTENT	Peru	2007 - 2011	Prospective cohort	215	Birth, 1, 2, ..., 24	8339	Jaganath et al 2014 Helicobacter[14]

Bovine Serum RCT	Guatemala	1997 - 1998	Individual RCT	315	Baseline, 1, 2, ...,8	2545	Begin et al. 2008, EJCN[15]
MAL-ED	Brazil	2010 - 2014	Prospective cohort	233	Birth, 1, 2, ..., 24	4838	Lima et al 2014 Clin Infect Dis[16]

*Children enrolled is for children with measurements under 2 years of age. Total measurements are number of measurements of anthropometry on children under 2 years of age.

Extended Data Table 2| Summary of baseline characteristics in *ki* cohorts

	CMC-V-BCS-2002, India	CMIN Bangladesh93	CMIN Peru89	CMIN Peru95	CONTENT, Peru	EE, Pakistan	GMS-Nepal	Guatemala BSC	IRC, India	Keneba, The Gambia
	(N=373)	(N=280)	(N=210)	(N=224)	(N=215)	(N=380)	(N=698)	(N=315)	(N=410)	(N=2954)
Sex										
Female	187 (50.1%)	122 (43.6%)	100 (47.6%)	96 (42.9%)	109 (50.7%)	185 (48.7%)	328 (47.0%)	162 (51.4%)	185 (45.1%)	1426 (48.3%)
Male	186 (49.9%)	158 (56.4%)	110 (52.4%)	128 (57.1%)	106 (49.3%)	195 (51.3%)	370 (53.0%)	153 (48.6%)	225 (54.9%)	1528 (51.7%)
Birthweight										
Mean (SD)	2910 (434)	2560 (967)	3550 (492)	3690 (367)	3070 (323)	2640 (506)	2660 (423)		2890 (448)	2970 (421)
Maternal age										
Mean (SD)	24.1 (4.11)					30.0 (3.99)	24.0 (5.09)	25.2 (6.21)	23.7 (3.68)	27.4 (7.16)
Maternal weight										
Mean (SD)										
Maternal education (years)										
Mean (SD)	5.43 (4.10)					0.873 (2.51)	2.48 (4.01)	4.07 (2.95)	5.06 (4.61)	
Number of rooms										
4+	14 (3.75%)				78 (36.3%)		323 (46.3%)		17 (4.17%)	
1	202 (54.2%)				44 (20.5%)		49 (7.02%)		185 (45.3%)	
2	106 (28.4%)				54 (25.1%)		145 (20.8%)		170 (41.7%)	
3	51 (13.7%)				39 (18.1%)		181 (25.9%)		36 (8.82%)	
Number of children <5yrs										
1									89 (21.7%)	

	CMC-V-BCS-2002, India	CMIN Bangladesh93	CMIN Peru89	CMIN Peru95	CONTENT, Peru	EE, Pakistan	GMS-Nepal	Guatemala BSC	IRC, India	Keneba, The Gambia
2+									321 (78.3%)	
Improved sanitation										
1					201 (93.5%)					
0					14 (6.51%)					
Food security level										
Food Secure							479 (71.1%)			
Mildly Food Insecure							106 (15.7%)			
Food Insecure							89 (13.2%)			

	MAL-ED, Bangladesh	MAL-ED, Brazil	MAL-ED, India	MAL-ED, Nepal	MAL-ED, Peru	MAL-ED, South Africa	MAL-ED, Tanzania	PROVIDE, Bangladesh	ResPak, Pakistan	Tanzania Child2	TDC, India
	(N=265)	(N=233)	(N=251)	(N=240)	(N=303)	(N=314)	(N=262)	(N=700)	(N=284)	(N=2400)	(N=160)
Sex											
Female	136 (51.3%)	113 (48.5%)	138 (55.0%)	110 (45.8%)	143 (47.2%)	159 (50.6%)	133 (50.8%)	332 (47.4%)	136 (47.9%)	1184 (49.3%)	75 (46.9%)
Male	129 (48.7%)	120 (51.5%)	113 (45.0%)	130 (54.2%)	160 (52.8%)	155 (49.4%)	129 (49.2%)	368 (52.6%)	148 (52.1%)	1216 (50.7%)	85 (53.1%)
Birthweight											
Mean (SD)	2800 (412)	3340 (483)	2890 (442)	2980 (390)	3130 (430)	3130 (464)	3180 (452)	2780 (371)	2930 (525)	3230 (474)	2910 (450)
Maternal age											
Mean (SD)	24.8 (4.99)	24.8 (5.53)	23.9 (4.10)	26.4 (3.76)	24.2 (6.06)	26.4 (6.86)	28.4 (6.67)	24.7 (4.65)		26.4 (5.04)	
Maternal weight											
Mean (SD)	49.6 (8.54)	61.9 (11.8)	50.4 (9.37)	56.3 (8.27)	55.5 (8.95)	67.5 (14.9)	55.7 (9.11)	49.3 (9.41)		62.2 (11.7)	
Maternal education (years)											
Mean (SD)	5.63 (2.58)	9.18 (2.81)	7.88 (3.17)	8.77 (3.44)	7.81 (2.79)	10.3 (1.94)	6.17 (1.79)	4.33 (3.62)		7.90 (2.27)	
Number of rooms											
4+	12 (4.96%)	127 (60.5%)	25 (10.6%)	131 (55.5%)	139 (51.1%)	196 (76.3%)	108 (43.2%)	23 (3.29%)			5 (3.13%)
1	152 (62.8%)	4 (1.90%)	84 (35.7%)	52 (22.0%)	19 (6.99%)	14 (5.45%)	13 (5.20%)	507 (72.4%)			91 (56.9%)
2	50 (20.7%)	20 (9.52%)	78 (33.2%)	31 (13.1%)	52 (19.1%)	22 (8.56%)	63 (25.2%)	108 (15.4%)			49 (30.6%)
3	28 (11.6%)	59 (28.1%)	48 (20.4%)	22 (9.32%)	62 (22.8%)	25 (9.73%)	66 (26.4%)	62 (8.86%)			15 (9.38%)
Number of children <5yrs											
1								512 (73.1%)		1640 (68.6%)	

	MAL-ED, Bangladesh	MAL-ED, Brazil	MAL-ED, India	MAL-ED, Nepal	MAL-ED, Peru	MAL-ED, South Africa	MAL-ED, Tanzania	PROVIDE, Bangladesh	ResPak, Pakistan	Tanzania Child2	TDC, India
2+								188 (26.9%)		749 (31.4%)	
Improved sanitation											
1	204 (84.3%)	206 (98.1%)	108 (46.4%)	235 (99.6%)	65 (24.7%)	4 (1.60%)		58 (96.7%)			
0	38 (15.7%)	4 (1.90%)	125 (53.6%)	1 (0.424%)	198 (75.3%)	246 (98.4%)		2 (3.33%)			
Food security level											
Food Secure	161 (83.0%)	3 (2.33%)	190 (89.6%)	94 (73.4%)	27 (23.9%)	132 (56.7%)					
Mildly Food Insecure	4 (2.06%)	11 (8.53%)	5 (2.36%)	15 (11.7%)	29 (25.7%)	19 (8.15%)					
Food Insecure	29 (14.9%)	115 (89.1%)	17 (8.02%)	19 (14.8%)	57 (50.4%)	82 (35.2%)					

Supplementary References

1. Iqbal, N. T. *et al.* Promising Biomarkers of Environmental Enteric Dysfunction: A Prospective Cohort study in Pakistani Children. *Sci. Rep.* **8**, 2966 (2018).

2. Ali, A. *et al.* Respiratory viruses associated with severe pneumonia in children under 2 years old in a rural community in Pakistan. *J. Med. Virol.* **88**, 1882–1890 (2016).

3. Shrestha, P. S. *et al.* Bhaktapur, Nepal: The MAL-ED Birth Cohort Study in Nepal. *Clin. Infect. Dis.* **59**, S300–S303 (2014).

4. Gladstone, B. P. *et al.* Protective Effect of Natural Rotavirus Infection in an Indian Birth Cohort. *N. Engl. J. Med.* **365**, 337–346 (2011).

5. John, S. M. *et al.* Establishment of the MAL-ED Birth Cohort Study Site in Vellore, Southern India. *Clin. Infect. Dis.* **59**, S295–S299 (2014).

6. Kattula, D. *et al.* The first 1000 days of life: prenatal and postnatal risk factors for morbidity and growth in a birth cohort in southern India. *BMJ Open* **4**, e005404 (2014).

7. Ahmed, T. *et al.* The MAL-ED Cohort Study in Mirpur, Bangladesh. *Clin. Infect. Dis.* **59**, S280–S286 (2014).

8. Kirkpatrick, B. D. *et al.* The "Performance of Rotavirus and Oral Polio Vaccines in Developing Countries" (PROVIDE) Study: Description of Methods of an Interventional Study Designed to Explore Complex Biologic Problems. *Am. J. Trop. Med. Hyg.* **92**, 744–751 (2015).

9. Mduma, E. R. *et al.* The Etiology, Risk Factors, and Interactions of Enteric Infections and Malnutrition and the Consequences for Child Health and Development Study (MAL-ED): Description of the Tanzanian Site. *Clin. Infect. Dis.* **59**, S325–S330 (2014).

10. Locks, L. M. *et al.* Effect of zinc and multivitamin supplementation on the growth of Tanzanian children aged 6–84 wk: a randomized, placebo-controlled, double-blind trial12. *Am. J. Clin. Nutr.* **103**, 910–918 (2016).

11. Bessong, P. O., Nyathi, E., Mahopo, T. C. & Netshandama, V. Development of the Dzimauli Community in Vhembe District, Limpopo Province of South Africa, for the MAL-ED Cohort Study. *Clin. Infect. Dis.* **59**, S317–S324 (2014).

12. Schoenbuchner, S. M. *et al.* The relationship between wasting and stunting: a retrospective cohort analysis of longitudinal data in Gambian children from 1976 to 2016. *Am. J. Clin. Nutr.* doi:10.1093/ajcn/nqy326.

13. Yori, P. P. *et al.* Santa Clara de Nanay: The MAL-ED Cohort in Peru. *Clin. Infect. Dis.* **59**, S310–S316 (2014).

14. Jaganath, D. *et al.* First Detected Helicobacter pylori Infection in Infancy Modifies the Association Between Diarrheal Disease and Childhood Growth in Peru. *Helicobacter* **19**, 272–279 (2014).

15. Bégin, F., Santizo, M.-C., Peerson, J. M., Torún, B. & Brown, K. H. Effects of bovine serum concentrate, with or without supplemental micronutrients, on the growth, morbidity, and micronutrient status of young children in a low-income, peri-urban Guatemalan community. *Eur. J. Clin. Nutr.* **62**, 39–50 (2008).

745 16. Lima, A. A. M. *et al.* Geography, Population, Demography, Socioeconomic,
746   Anthropometry, and Environmental Status in the MAL-ED Cohort and Case-Control
747   Study Sites in Fortaleza, Ceará, Brazil. *Clin. Infect. Dis.* **59**, S287–S294 (2014).

748
749

Materials and Methods

Study designs and inclusion criteria

We included all longitudinal observational studies and randomized trials available through the *ki* project
on April 2018 that met five inclusion criteria (Extended Data Figure 1): 1) conducted in low- or middle-
income countries; 2) enrolled children between birth and age 24 months and measured their length and
weight repeatedly over time; 3) did not restrict enrollment to acutely ill children; 4) enrolled at least 200
children; 5) collected anthropometry measurements at least monthly. The frequency of measurements
was assessed by calculating the median days between measurements. Our pre-specified analysis
protocol stipulated that if randomized trials found effects of interventions on growth, the analysis would
only include the control arm only; yet, all intervention trials that met the inclusion criteria had null
effects on growth, so all arms were included. We included all children under 24 months of age, assuming
months were 30.4167 days. We excluded extreme measurements of WLZ > 5 or < −5 and of WAZ < -6 or
> 5, consistent with 2006 WHO Growth Standards recommendations.[1] We checked for cohort-wide
anthropometry measurement quality by plotting Z-score densities and calculating the proportion of
length measurements where length decreased beyond the expected technical error of measurement
compared to the last measurement on a child. One cohort, MAL-ED Pakistan, was excluded because
measurements exhibited a multimodal WLZ distribution with scores binned at −2, −1.5, and −1 instead
of continuously distributed.

Outcome definitions

We used the following outcome measures in the analysis:

Weight-for-Length Z-scores were calculated using the 2006 WHO growth standards,[2] and mean WLZ was
calculated within strata of interest. We used the medians of triplicate measurements of lengths and
weights of children from pre-2006 cohorts to re-calculate Z scores to the 2006 standard.

Prevalent wasting was defined as the proportion of measurements within a specific stratum (e.g., age)
below the 2006 WHO standard −2 WLZ, and analogously below the 2006 WHO standard −3 WLZ for
severe wasting. For each age, we included children with WLZ measurements within one month before
and after that age in the point prevalence estimate (e.g., for point prevalence at 6 months, we include
children aged 5-7 months). WLZ is not calculable for children with lengths <45cm (10.2% of children at
birth), so using WLZ underestimates wasting at birth.

Incident wasting episodes were defined as a change in WLZ from above −2 Z in the prior measurement
to below −2 Z in the current measurement. Similarly, we defined severe wasting episodes using −3 Z
cutoff. We assumed a 60-day washout period between episodes of wasting before a new episode of
wasting could occur (Fig 4a). Children were considered at risk for wasting at birth, so children born
wasted were considered to have an incident episode of wasting at birth. Children were also assumed to
be at risk of wasting at the first measurement in studies that enrolled children after birth, so children
wasted at the first measurement in a non-birth cohort were assumed to have incident wasting occurring
at the age halfway between birth and the first measurement.

Incidence proportion of wasting was calculated during a defined age range (e.g. 6-12 months) as the
proportion of children not wasted at the start of the period who became wasted during the age period
(the proportion of children who had the onset of new episodes during the period). This differs from
period prevalence, which would include any children who had any measurement of WLZ < -2 during the
period (including children who began the period wasted). Period prevalence would thus additionally
include in the numerator and denominator children who were wasted at the start of the period,
whereas incidence proportion excludes them.

Recovery from wasting was defined as a WLZ change from below to above –2 Z among children who are
currently wasted or severely wasted. We required a child to maintain WLZ above –2 for 60 days to be
considered "recovered".   Children were only considered "at risk" for recovery if their prior
measurement was below –2 WLZ. We measured the proportion of children who recover from moderate
wasting (WLZ < –2) within 30, 60, and 90 days of the onset of the episode. We assumed recovery from
wasting was spontaneous because we did not have consistent information on referral guidelines across
cohorts in the analysis, but some children with moderate wasting may have been referred to clinical
facilities for outpatient treatment. Wasting recovery may thus be higher in these highly monitored
cohorts than in the general population due to treatment referrals.

Wasting duration was estimated by counting the days between the onset of wasting and recovery within
an individual child. We assumed that the episode started or ended at the midpoint between
measurements. For example, if a child was not wasted at age 40 days, wasted at age 70 days, and not
wasted at age 100 days, the duration of the wasting episode was: (70-40)/2 + (100-70)/2 = 30 days. We
calculated the median duration of wasting episodes among cohorts where children recovered during the
study period from >50% of observed wasting episodes, as the duration of episodes that extended
beyond study period is unknown. Therefore, the median duration could only be calculated when less
than half of episodes were censored. Confidence intervals were calculated using the quantile method.

Incidence rate of wasting was calculated during defined age ranges as the number of incident episodes
of wasting per 1,000 child-days at risk during the age range. Children were considered "at risk" for
incident wasting episodes if they were not currently wasted and were classified as recovered from any
prior wasting episode (beyond the 60-day washout period). Therefore, wasted children, or children
within the washout period, did not contribute to the person-time at risk for wasting used to calculate
wasting incidence. To calculate person-time at risk of wasting, we assumed that the onset of a wasting
episode occurred when the child's age was at the midpoint between the measurement of WLZ ≥ –2 and
the measurement of WLZ < –2, and conversely the time of recovery occurred when the child's age was
at the midpoint between the last measurement of WLZ < –2 and the first measurement of WLZ ≥ –2
within the 60-day washout period (Fig 4a).

Persistent wasting was defined as a >50% longitudinal prevalence of below –2 WLZ (wasting), and
analogously >50% longitudinal prevalence of below –3 WLZ for persistent severe wasting.[3]

Concurrent prevalence of wasting and stunting was defined as the proportion of measurements at a
specific age when a child was both wasted and stunted at the same measurement. For each age, we
included children with WLZ and LAZ measurements within one month before and after that age in the
point prevalence estimate (e.g., for point prevalence at 6 months, we include children aged 5-7 months).

Prevalent underweight was defined as the proportion of measurements at a specific age below the 2006
WHO standard –2 WAZ. For each age, we included children with WAZ measurements within one month
before and after that age in the point prevalence estimate (e.g., for point prevalence at 6 months, we
include children aged 5-7 months).

Subgroups of interest

We stratified the above outcomes of interest within the following subgroups: Child age, grouped into
one, three, or six month intervals, (depending on the outcome); the region of the world (Asia, sub-
Saharan Africa, Latin America); the month of the year, and the combinations of the above categories.
838
839

Statistical analysis

All analyses were conducted in R version 4.0.5. Supplementary pooled, regional, and cohort-specific
results, and sensitivity analyses are available in online supplements at ([https://child-](https://child-growth.github.io/wasting)
[growth.github.io/wasting](https://child-growth.github.io/wasting)).

<u>Fixed and random effects models</u>

We conducted a 2-stage individual participant meta-analysis, estimating wasting outcomes within
specific cohorts and pooling estimates within each age strata using random effects models. For example,
we estimated each age-specific mean using a two-step process. We first estimated the mean in each
cohort, and then pooled age-specific means across cohorts allowing for a cohort-level random effect.
We estimated overall pooled effects, and pooled estimates specific to South Asian, African, or Latin
American cohorts, depending on the analysis. We repeated the pooling of all statistics presented in the
figures using fixed effects models as a sensitivity analysis (https://child-growth.github.io/wasting/fixed-
effects.html). The pooling methods are described in greater detail in Benjamin-Chung (2020).[4] All
pooling was completed using the rma() function from the "metafor" package in the R language (version
3.4.0).[5]

<u>Fitted spline curves</u>

Fitted smoothers used in manuscript figures were fit using cubic splines and generalized cross-
validation.[6] We estimated approximate 95% simultaneous confidence intervals around the cubic splines
using a parametric bootstrap that resampled from the posterior the generalized additive model
parameter variance-covariance matrix.[7]

<u>Seasonality analysis</u>

We compared mean WLZ over day of the year, over child birth day, and over seasons defined by rainfall.
We estimated mean WLZ by cohort over day of the year and child birth day using cubic splines (Fig. 3a,
c).[8] Splines of WLZ over day of the year were plotted over monthly mean rainfall averaged over the
years a study measured child anthropometry below ages 24 months (Fig. 3a). We pulled monthly
precipitation values from Terrraclimate, a dataset that combines readings from WorldClim data, CRU
Ts4.0, and the Japanese 55-year Reanalysis Project.[9] For each study region, we averaged all readings
within a 50 km radius from the study coordinates. If GPS locations were not in the data for a cohort, we
used the approximate location of the cohort based on the published descriptions of the cohort. Monthly
measurements were matched to study data based on the calendar month and year in which
measurements were taken.

The season of peak rainfall was defined as the three-month period with the highest mean
rainfall. Mean differences in WLZ between three-month quarters was estimated using linear regression
models. We compared the consecutive three months of the maximum average rainfall over the study
period, as well as the three months prior and the three months after the maximum-rainfall period, to a
reference level of the three months opposite the calendar year of the maximum-rainfall period. We
used all WLZ measurements of children under two years of age (e.g., if June-August was the period of
maximum rainfall, the reference level is child mean WLZ during January-March).

Estimates were unadjusted for other covariates because we assumed that seasonal effects on
WLZ were exogenous and could not be confounded. Mean differences in WLZ were pooled across
cohorts using random-effects models, with cohorts grouped by the Walsh and Lawler seasonality
index.[10] Cohorts from years with a seasonal index ≥ 0.9 were classified as occurring in locations with high
seasonality, cohorts with a seasonal index < 0.9 and ≥ 0.7 were classified as occurring in locations with
medium seasonality, and cohorts with a seasonal index < 0.7 were classified as occurring in locations
with low seasonality.

888
$$Seasonal\ Index = \frac{1}{R} \sum_{n=1}^{n=12} |X_n - \frac{R}{12}|$$

Where R = total annual precipitation and X_n = monthly precipitation.

Estimation of mean LAZ, WAZ, and WLZ by age in Demographic and Health Surveys and *ki*
cohorts

We downloaded standard DHS individual recode files for each country from the DHS program
website (https://dhsprogram.com/). We used the most recent standard DHS datasets for the individual
women's, household, and length and weight datasets from each country, and we estimated age-
stratified mean LAZ, WAZ, and WLZ from ages 0 to 24 months within each DHS survey, accounting for
the complex survey design and sampling weights. See Benjamin-Chung et. al (2020) for additional details
on the DHS data cleaning and analysis.[4] We compared DHS estimates with mean LAZ, WAZ, and WLZ by
age in the *ki* study cohorts with penalized cubic-splines with bandwidths chosen using generalized cross-
validation.[8] We did not seasonally adjust DHS measurements.

903   .

Sensitivity analyses

We estimated incidence rates and of wasting after excluding children born or enrolled wasted (Fig. 4b).
The rationale for this sensitivity analysis is that incident cases at birth imply a different type of
intervention (i.e., prenatal) compared with postnatal onset of wasting. We estimated the overall and
region stratified prevalence of persistent wasting and of underweight and severe wasting by age
(Extended data fig. 5). Within cohorts that measured child gestational age at birth, we estimated the
prevalence of stunting and underweight at birth both uncorrected and corrected for gestational age at
birth using the Intergrowth standards (Extended data figure 6).[11] We also assessed how much change in
Z-scores after children became wasted could be explained by regression to the mean (RTM) and how
much is catch-up growth beyond that expected by RTM.[12] We calculated the cohort-specific RTM effect
and plotted the mean Z-scores expected from RTM 3 months later among children who were wasted at
each age in months from birth to 21 months. We used the cohort means as the population mean Z-
scores, and compared the expected mean WLZ with the observed mean WLZ (Extended Data Fig. 8).[13,14]
We also compared wasting prevalence defined using middle-upper-arm circumference (MUAC), an
alternative measurement for classifying wasting, with wasting prevalence estimated using WLZ within
the cohort that measured MUAC (https://child-growth.github.io/wasting/muac.html). We also re-
estimate primary results dropping observations of children at birth within the MRC Keneba cohort,
which used a different team to measure child anthropometry at birth from the trained anthropometrists
used in follow-up measurements (https://child-growth.github.io/wasting/no-kenaba.html). We also

examined the effect of shorter (30 day) and longer (90) day washout period when determining if a child
was again at risk when estimating wasting incidence and wasting recovery rates (https://child-
growth.github.io/wasting/ir_sensitivity.html). Lastly, we compared estimates pooled using random
effects models, which are more conservative in the presence of study heterogeneity, with estimates
pooled using fixed effects (inverse variance weighted) models (https://child-
growth.github.io/wasting/fixed-effects.html).

Data and code availability

The data that support the findings of this study are available from the Bill and Melinda Gates Foundation
Knowledge Integration project upon reasonable request. Replication scripts for this analysis are
available here: https://github.com/child-growth/ki-longitudinal-growth.

Methods References

1. Organization, W. H. & Fund (UNICEF), U. N. C. *Recommendations for data collection, analysis and reporting on*
anthropometric indicators in children under 5 years old. (World Health Organization, 2019).

2. WHO Multicentre Growth Reference Study Group. WHO child growth standards: length/height-for-age,
weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and
development. *Geneva World Health Organ.* 312 pages (2006).

3. Richard, S. A. *et al.* Wasting Is Associated with Stunting in Early Childhood123. *J. Nutr.* **142**, 1291–1296 (2012).

4. Benjamin-Chung, J. et. al. (submitted). Early childhood linear growth failure in low-and middle-income
countries. (2020).

5. Viechtbauer, W. Conducting Meta-Analyses in R with the metafor Package. *J. Stat. Softw.* **36**, 1–48 (2010).

6. Cleveland, W. S. & Loader, C. Smoothing by Local Regression: Principles and Methods. in *Statistical Theory and*
Computational Aspects of Smoothing (eds. Härdle, W. & Schimek, M. G.) 10–49 (Physica-Verlag HD, 1996).
doi:10.1007/978-3-642-48425-4_2.

7. Ruppert, D., Wand, M. P. & Carroll, R. J. *Semiparametric Regression*. (Cambridge University Press, 2003).
doi:10.1017/CBO9780511755453.

8. Wood, S. N., Pya, N. & Säfken, B. Smoothing Parameter and Model Selection for General Smooth Models. *J.*
Am. Stat. Assoc. **111**, 1548–1563 (2016).

9. Abatzoglou, J. T., Dobrowski, S. Z., Parks, S. A. & Hegewisch, K. C. TerraClimate, a high-resolution global
dataset of monthly climate and climatic water balance from 1958–2015. *Sci. Data* **5**, 170191 (2018).

10. Walsh, R. P. D. & Lawler, D. M. Rainfall Seasonality: Description, Spatial Patterns and Change Through Time.
Weather **36**, 201–208 (1981).

11. Papageorghiou, A. T. *et al.* The INTERGROWTH-21st fetal growth standards: toward the global integration of
pregnancy and pediatric care. *Am. J. Obstet. Gynecol.* **218**, S630–S640 (2018).

12. Cameron, N., Preece, M. A. & Cole, T. J. Catch-up Growth or Regression to the Mean? Recovery from Stunting
Revisited. *Am. J. Hum. Biol.* **17**, 412–417 (2005).

13. Barnett, A. G., van der Pols, J. C. & Dobson, A. J. Regression to the mean: what it is and how to deal with it. *Int.*
J. Epidemiol. **34**, 215–220 (2005).

14. Linden, A. Assessing regression to the mean effects in health care initiatives. *BMC Med. Res. Methodol.* **13**,
119 (2013).

Acknowledgments

This research was financially supported by a global development grant (OPP1165144)

from the Bill & Melinda Gates Foundation to the University of California, Berkeley, CA, USA. We would also like to thank the following collaborators on the included cohorts and trials for their contributions to study planning, data collection, and analysis: Muhammad Sharif, Sajjad Kerio, Ms. Urosa, Ms. Alveen, Shahneel Hussain, Vikas Paudel (Mother and Infant Research Activities), Anthony Costello (University College London), Benjamin Torun, Lindsey M Locks, Christine M McDonald, Roland Kupka, Ronald J Bosch, Rodrick Kisenge, Said Aboud, Molin Wang, and all other members of the study staffs and field teams. We would also like to thank all study participants and their families for their important contributions.

Author contributions

Conceptualization: A.M., J.B., J.M.C., K.H.B., P.C., B.F.A

Funding Acquisition: J.M.C., A.E.H., M.J.V., B.F.A.

Data curation: A.M., J.B., J.C., O.S., W.C., A.N., N.N.P., W.J., E.J., E.O.C., S.D., N.H., I.M., H.L., R.H., V.S., J.H., T.N.

Formal analyses: A.M., J.B., J.C., O.S., W.C., A.N., N.N.P., W.J., E.J., E.O.C., S.D., N.H., I.M., H.L., V.S., B.F.A

Methodology: A.M., J.B., J.M.C, J.C., O.S., N.H., I.M., A.E.H., M.J.V.,K.H.B., P.C., B.F.A.

Visualization: A.M.,J.B., A.N., N.N.P., S.D., A.S., J.C., R.H., E.J., K.H.B., P.C., B.F.A.

Writing – Original Draft Preparation: A.M., J.B., B.F.A.

Writing – Review & Editing: A.M., J.B., J.M.C., K.H.B., P.C., B.F.A., *ki* Child Growth Consortium members

Competing interest declaration

Thea Norman is an employee of the Bill & Melinda Gates Foundation (BMGF). Kenneth H Brown and Parul Christian are former employees of BMGF. Jeremy Coyle, Vishak Subramoney, Ryan Hafen, and Jonas Häggström work as research contractors funded by the BMGF.

Additional information

Supplementary Information is available for this paper at https://child-growth.github.io/wasting.

Correspondence and requests for materials should be addressed to Andrew Mertens (amertens@berkeley.edu) and Benjamin F. Arnold (ben.arnold@ucsf.edu)