Model-based spatial-temporal mapping of opisthorchiasis in endemic countries of Southeast Asia

Ting-Ting Zhao,[1] Yi-Jing Feng,[1] Pham Ngoc Doanh,[2] Somphou Sayasone,[3] Virak Khieu,[4] Choosak Nithikathkul,[5] Men-Bao Qian,[6,7] Yuan-Tao Hao[1,8] Ying-Si Lai,[1,8*]

[1]Department of Medical Statistics, School of Public Health, Sun Yat-sen University, Guangzhou, Guangdong, People's Republic of China.

[2]Department of Parasitology, Institute of Ecology and Biological Resources, Graduate University of Science and Technology, Vietnam Academy of Sciences and Technology, Cau Giay, Hanoi, Vietnam.

[3]Lao Tropical and Public Health Institute, Ministry of Health, Vientiane Capital, Lao People's Democratic Republic.

[4]National Center for Parasitology, Entomology and Malaria Control, Ministry of Health, Phnom Penh, Cambodia.

[5]Tropical and Parasitic Diseases Research Unit, Faculty of Medicine, Mahasarakham University, Mahasarakham, Thailand.

[6]National Institute of Parasitic Diseases, Chinese Center for Disease Control and Prevention, Shanghai, People's Republic of China.

[7]WHO Collaborating Centre for Tropical Diseases, Key Laboratory of Parasite and Vector Biology, Ministry of Health, Shanghai, People's Republic of China.

[8]Sun Yat-sen Global Health Institute, Sun Yat-sen University, Guangzhou, Guangdong, People's Republic of China.

*Correspondence to: Dr Ying-Si Lai, Department of Medical Statistics, School of Public Health, Sun Yat-sen University, Guangzhou, Guangdong 510080, China. laiys3@mail.sysu.edu.cn

Abstract

Opisthorchiasis is an overlooked danger to Southeast Asia. High-resolution disease risk maps are critical but haven't been available for Southeast Asia. Georeferenced disease data and potential influencing factor data were collected through a systematic review of literatures and open-access databases, respectively. Bayesian spatial-temporal joint models were developed to analyze both point- and area-level disease data, within a logit regression in combination of potential influencing factors and spatial-temporal random effects. The model-based risk mapping identified areas of low, moderate and high prevalence across the study region. Even though the overall population-adjusted estimated prevalence presented a trend down, a total of 12.39 million (95% BCI: 10.10-15.06) people were estimated infected with *O. viverrini* in 2018 in four major endemic countries (i.e., Thailand, Laos, Cambodia, and Vietnam), highlighting the public health importance of the disease in the study region. The high-resolution risk maps provide valuable information for spatial targeting of opisthorchiasis control interventions.

Introduction

End the epidemics of Neglected Tropical Diseases (NTDs) by 2030 embodied in the international set of targets for the Sustainable Development Goals (SDGs) endorsed by the United Nations empowers the efforts made by developing countries to combat the NTD epidemics (*UN, 2015*). To date, 20 diseases have been listed as NTDs, and opisthorchiasis is under the umbrella of food-borne trematodiasis (*Ogorodova et al., 2015*). Two species of opisthorchiasis are of public health significance, that is *Opisthorchis felineus* (*O. felineus*), endemic in eastern Europe and Russia, and *Opithorchis viverrini* (*O. viverrini*) endemic in Southeast Asian countries (*Petney et al., 2013*). The later species is of our interest in the current manuscript.

According to WHO's conservative estimation, an overall disease burden due to opisthorchiasis was

188,346 disability-adjusting life years (DALYs) in 2010 (*Havelaar et al., 2015*). Fürst and colleagues

estimated that more than 99% of the burden worldwide attribute to *O. viverrini* infection in Southeast

Asia (*Fürst et al., 2012*). Five countries in Southeast Asia, Cambodia, Lao PDR, Myanmar, Thailand and

Vietnam, are endemic for opisthorchiasis, with an estimated 67.3 million people at risk (*Keiser &*

Utzinger, 2005). It is well documented that chronic and repeated infection with *O. viverrini* leads to the

development of fatal bile duct cancer (cholangiocarcinoma) (*International Agency for Research on*

Cancer, 1994).

The life cycle of *O. viverrini* involves freshwater snails of the genus *Bithynia* as the first intermediate

host, and freshwater cyprinoid fish as the second intermediate host. Humans and other carnivores (e.g.,

cats and dogs), the final hosts, become infected by consuming raw or insufficiently cooked infected fish

(*Andrews et al., 2008, Saijuntha et al., 2014*). Behavioral, environmental and socio-economic factors

affect the transmission of *O. viverrini* (*Grundy-Warr et al., 2012, Phimpraphai et al., 2017,*

Phimpraphai et al., 2018, Prueksapanich et al., 2018). Raw or insufficiently cooked fish consumption

is the cultural root in endemic countries, shows a strong relationship with the occurrence of the disease

(*Andrews et al., 2008, Grundy-Warr et al., 2012*). Poorly hygienic conditions increase the risk of

infection, especially in areas practicing raw-fish-eating habit (*Grundy-Warr et al., 2012*). In addition,

environmental and climatic factors, such as temperature, precipitation and landscape, affecting either

snail/fish population or growth of the parasites inside the intermediate hosts, can potentially influence

the risk of human infection (*Forrer et al., 2012, Suwannatrai et al., 2017*). Important control strategies

of *O. viverrini* infection include preventive chemotherapy, health education, environmental modification,

improving sanitation, as well as comprehensive approaches with combinations of the above (*Saijuntha*

et al., 2014). For purposes of public health control, WHO recommends implementing preventive

chemotherapy once a year or once every two years depending on the levels of prevalence in population,

with complementary interventions such as health education and improvement of sanitation (*WHO, 2009*).

Understanding the geographical distribution of *O. viverrini* infection risk at high spatial resolution is

critical to prevent and control the disease cost-effectively in priority areas. Thailand conducted national

surveys for *O. viverrini* prevalence in 1981, 1991, 2001, 2009, and 2014 (*Echaubard et al., 2016,*

Suwannatrai et al., 2018), but the results of these surveys were presented at the province-level, which is

less informative for precisely targeting control interventions. Suwannatrai and colleagues, based on

climatic and *O. viverrini* presence data, produced climatic suitability maps for *O. viverrini* in Thailand

using the MaxEnt modeling approach (*Suwannatrai et al., 2017*). The maps brought insights for

identifying areas with a high probability of *O. viverrini* occurrence; however, they did not provide direct

information on prevalence of *O. viverrini* in population (*Elith et al., 2011*). A risk map of *O. viverrini*

infection in Champasack province of Lao PDR were presented by Forrer and colleagues (*Forrer et al.,*

2012). To our knowledge, high-resolution, model-based risk estimates of *O. viverrini* infection are

unavailable in the whole endemic region of Southeast Asia.

Bayesian geostatistical modeling is one of the most rigorous inferential approaches for high-resolution

maps depicting the distribution of the disease risk (*Karagiannis-Voules et al., 2015*). Geostatistical

modeling relates geo-referenced disease data with potential influencing factors (e.g., socioeconomic and

environmental factors) and estimates the infection risk in areas without observed data (*Gelfand &*

Banerjee, 2017). Common geostatistical models are usually based on point-referenced survey data

(*Banerjee et al., 2014*). In practice, disease data collected from various sources often consists of point-

referenced and area-aggregated data. Bayesian geostatistical joint modeling approaches provide a

flexible framework for combining analysis of both kinds of data (*Moraga et al., 2017, Smith et al., 2008*).

In this study, we aimed (1) to collect all available survey data on the prevalence of *O. viverrini* infection

at point- or area-level in Southeast Asia through systematic review; and (2) to estimate the spatial-

temporal distribution of the disease risk at a high spatial resolution, with the application of advanced

Bayesian geostatistical joint modeling approach.

Results

A total of 2,690 references were identified through systematically reviewing peer-review literatures, and

13 additional references were gathered from other sources. According to the inclusion and exclusion

criteria, 168 records were included, resulted in a total of 580 ADM1-level surveys in 174 areas, 210

ADM2-level surveys in 142 areas, 53 ADM3-level surveys in 51 areas, and 251 point-level surveys at

207 locations in five endemic countries (i.e., Cambodia, Lao PDR, Myanmar, Thailand and Vietnam) of

Southeast Asia (Figure 1). Around 70% and 15% of surveys were conducted in Thailand and Lao PDR,

respectively. Only two relevant records were obtained from Myanmar. To avoid large estimated errors,

we did not include this data in the final geostatistical analysis. All surveys were conducted after 1970,

with around 75% done after 1998. Most surveys (95%) are community-based. Around 40% of surveys

used the Kato-Katz technique for diagnosis, while another 42% did not specify diagnostic approaches.

Mean prevalence calculated directly from survey data was 16.74% across the study region. A summary

of survey data is listed in Table 1, and survey locations and observed prevalence in each period are shown

in Figure 2. Area-level data cover all regions in Thailand and Lao PDR, and most regions in Cambodia

and Vietnam, whilst point-referenced data are absent in most areas of Vietnam, the western part of

Cambodia and southern part of Thailand. Around 70% of eligible literatures got a score equal or more

than 7, indicating an overall good quality of eligible literatures in our study (Figure 2-figure supplement 1).

Seven variables were selected for the final model through the Bayesian variable selection process (Table 2). The infection risk was 2.61 (95%BCI: 2.10-3.42) times in the community as much as that in school-aged children. Surveys using FECT as the diagnostic method showed a lower prevalence (*OR* 0.76, 95%BCI: 0.61-0.93) compared to that using Kato-Katz method, while no significant difference was found between Kato-Katz and the other diagnostic methods. Human influence index and elevation were negatively correlated with the infection risk. Each unit increase of the HII index was associated with 0.01 (95%BCI: 0.003-0.02) decrease in the logit of the prevalence. And increase 1 meter in elevation was associated with the 0.003 (95%BCI: 0.001-0.005) decrease in the logit of the prevalence. The spatial range was estimated 83.55 km (95% BCI: 81.34-86.61), the spatial variance σ_ϕ^2 was 12.59 (95% BCI: 11.96-13.56), the variance of beta-likelihood σ_β^2 was 0.15 (95% BCI: 0.14-0.15), and the temporal correlation coefficient ρ was 0.66 (95% BCI: 0.65-0.67). Model validation showed that our model was able to correctly estimate 79.61% of locations within the 95% BCI, indicating the model had a reasonable capacity of prediction accuracy. The ME, MAE, and MSE were 0.24%, 9.06%, and 2.38%, respectively, in the final model, while they were -7.14%, 16.67%, and 5.09%, respectively, in the model only based on point-referenced data, suggesting that the performance of the final model was better than the model only based on point-referenced data. On the other hand, Monte Carlo test for preferential sampling suggested that preferential sampling may exist for survey locations in one third (6/18) of the survey years (Figure 2-source data 2).

The estimated risk maps of *O. viverrini* infection in different selected years (i.e., 1978, 1983, 1988, 1993, 1998, 2003, 2008, 2013, and 2018) are presented in Figure 3. In 2018, the high infection risk (with

prevalence >25%) was mainly estimated in regions of the southern, the central and the north-central parts

of Lao PDR, some areas in the east-central parts of Cambodia, and some areas of the northeastern and

the northern parts Thailand. The southern part of Thailand, the northern part of Lao PDR, and the western

part of Cambodia showed low risk estimates (with prevalence <5%) of *O. viverrini* infection. The central

and several southern parts of Vietnam showed low to moderate risk of *O. viverrini* infection, while there

was no evidence of *O. viverrini* in other parts of Vietnam. High estimation uncertainty was mainly present

in the central part of Lao PDR, the northern and the eastern parts of Thailand, and the central part of

Cambodia and Vietnam (Figure 4).

In addition, the infection risk varies over time across the study region (Figure 5). Areas of northern

Thailand showed an increasing trend in periods 1978-1988 and 1993-2003, while most areas of the

country presented a considerable decrease of infection risk after 2008. The infection risk firstly increased

and then decreased in areas of the north, the central and the southern parts of Lao PDR and the central

parts of Vietnam. The east-central and western part of Cambodia showed an increasing trend in recent

years.

The population-adjusted estimated prevalence over the study region presents a trend down after 1995

(Figure 6 and Figure 6-figure supplement 1-9). At the country-level, the estimated prevalence in Thailand

showed a fast decline after 1995, and took on a gradually decreasing change in Cambodia. In Lao PDR,

the overall prevalence maintained quite stable before 1990, and decreased slightly between 1990 to 1997,

increased significantly after 1997, then decreased from 2006, and became stable after 2011. The

prevalence stable in Vietnam during the whole study period. We estimated that in 2018, the overall

population-adjusted estimated prevalence of *O. viverrini* infection in the whole study region was 6.57%

(95% BCI: 5.35%-7.99%), corresponding to 12.39 million (95% BCI: 10.10-15.06) infected individuals

(Table 3). Lao PDR showed the highest prevalence (35.21%, 95% BCI: 28.50%-40.70%), followed by

Thailand (9.71%, 95% BCI: 7.98%-12.17%), Cambodia (6.15%, 95% BCI: 2.41%-11.73%), and Vietnam

(2.15%, 95% BCI: 0.73%-4.40%). Thailand had the largest numbers of individuals estimated to be

infected with *O. viverrini* (6.71million, 95% BCI: 5.51-8.41), followed by Lao PDR (2.45million, 95%

BCI: 1.98-2.83), Vietnam (2.07million, 95% BCI: 0.70-4.24), and Cambodia (1.00million, 95% BCI:

0.39-1.90).

Discussion

In this study, we produced model-based, high-resolution risk estimates of opisthorchiasis across endemic

countries of Southeast Asia through systematic reviewing and the advanced Bayesian geostatistical joint

modeling approach. The disease is the most important foodborne trematodiasis in the study region (***Sripa***

et al., 2010), taking into account most of the disease burden of opisthorchiasis in the world (***Fürst et al.,***

2012). The estimates were obtained by systematically reviewing all possible geo-referenced survey data

and applying a Bayesian geostatistical modeling approach that jointly analyzes point-referenced and

area-aggregated disease data, as well as environmental and socioeconomic predictors. Our findings will

be important for guiding control and intervention cost-effectively and serve as a baseline for future

progress assessment.

Our estimates suggested that there was an overall decrease of *O. viverrini* infection in Southeast Asia

from 1995 onwards, which may be largely attributed to the decline of infection prevalence in Thailand.

This decline was probably on account of the national opisthorchiasis control program launched by the

Ministry of Public Health of Thailand from 1987 (***Jongsuksuntigul & Imsomboon, 2003,***

Jongsuksuntigul et al., 2003). Our high-resolution risk estimates in Thailand in 2018 showed similar

pattern as the climatic suitability map provided by Suwannatral and colleagues (***Suwannatrai et al.,***

2017). In this case, we estimated the prevalence of the population instead of the occurrence probability

of the parasite, which arms decision makers with more direct epidemiological information for guiding

control and intervention. The national surveys in Thailand reported a prevalence of 8.7% and 5.2% in

2009 and 2014, respectively (***MOPH, 2014, Wongsaroj et al., 2014***). However, we estimated higher

prevalence of 12.44% (95% BCI: 10.79%-14.26%) and 9.34% (95% BCI: 7.88%-11.02%) in 2009 and

2014, respectively. Even though the national surveys covered most provinces in Thailand, estimates were

based on simply calculating the percentage of positive cases among all the participants (***Wongsaroj et al.,***

2014), and the remote areas might not be included (***Maipanich et al., 2004***). Instead, our estimates were

based on rigorous Bayesian geostatistical modeling of available survey data with environmental and

socioeconomic predictors, accounting for heterogeneous distribution of infection risk and population

density when aggregating country-level prevalence.

Our findings suggested that the overall prevalence of *O. viverrini* remained high (>20%) in Lao PDR

during the study periods, consistent with conclusions drawn by Suwannatrai and colleagues

(***Suwannatrai et al., 2018***). We estimated that a total number of 2.45 (95% BCI: 1.98-2.83) million

people living in Lao PDR was infected with *O. viverrini*, equivalent to that estimated by WHO in 2004

(***WHO, 2002***). Besides, our risk mapping for Champasack province shares similarly risk map pattern

produced by Forrer and colleagues (***Forrer et al., 2012***). A national-scaled survey in Cambodia during

the period 2006-2011 reported infection rate of 5.7% (***Yong et al., 2014***), lower than our estimation of

8.34% (95% BCI: 5.25%-14.95%) in 2011. The former may underestimate the prevalence because more

than 77% of participants were schoolchildren (***Yong et al., 2014***). Another large survey in five provinces

of Cambodia suggested a large intra-district variation, which makes the identification of endemic areas

difficult (*Miyamoto et al., 2014*). Our high-resolution estimates for Cambodia help to differentiate the

intra-district risk. However, the estimates should be taken cautious due to large district-wide variances

and a relatively small number of surveys. Indeed, *O. viverrini* infection was underreported in Cambodia

(*Khieu et al., 2019*), and further point-referenced survey data are recommended for more confirmative

results.

Although an overall low prevalence was estimated in Vietnam (2.15%, 95% BCI: 0.73%-4.40%) in 2018,

it corresponds to 2.07 million (95% BCI: 0.70 million-4.24 million) people infected, comparable to the

number in Lao PDR, mainly due to a larger population in Vietnam. The risk mapping suggested moderate

to high risk areas presented in central Vietnam, with a high risk in Phu Yen province for many years,

particularly. This agreed with previous studies considering the province a "hotspot" (*Doanh & Nawa,*

2016). Of note, even though there was no evidence of *O. viverrini* infection in the northern part of the

country, *Clonorchis sinensis*, another important liver fluke species, is endemic in the region

(*Sithithaworn et al., 2012*). We didn't provide estimates for Myanmar in case of large estimated errors.

Indeed, only two relevant papers were identified by our systematic review, where one shows low to

moderate prevalence in three regions of Lower Myanmar (*Aung et al., 2017*), and the other found low

endemic of *O. viverrini* infection in three districts of the capital city Yangon (*Sohn et al., 2019*). Nation-

wide epidemiological studies are urgent for a more comprehensive understanding of the disease in

Myanmar.

We identified several important factors associated with *O. viverrini* infection in Southeast Asia, which

may provide insights for the prevention and control of the disease. The infection risk was higher in the

entire community than that in schoolchildren, consistent with multiple studies (*Aung et al., 2017, Forrer*

et al., 2012, Miyamoto et al., 2014, Van De, 2004, Wongsaroj et al., 2014). A negative association was

found between *O. viverrini* infection and elevation, suggesting the disease was more likely to occur in

low altitude areas, which was consistent with a previous study (***Wang et al., 2013***). HII, a measure of

human direct influence on ecosystems (***Sanderson et al., 2002***), showed a negative relationship with

O. viverrini infection risk, indicating the disease was more likely to occur in areas with low levels of

human activities, which were often remote and economically underdeveloped. The habit of eating raw or

insufficiently cooked fish was more common in rural areas than that in economically developed ones,

which could partially explain our findings (***Grundy-Warr et al., 2012, Jennifer, 2019***). Indeed, this

culturally rooted habit is one of the determinants for human opisthorchiasis (***Kaewpitoon et al., 2008,***

Ziegler et al., 2011). However, the precise geographical distribution of such information is unavailable

and thus we could not use it as a covariate in this study.

Our estimate of the number of people infected with *O. viverrini* is higher than that of the previous study

(12.39 million vs 8.6 million (***Qian & Zhou, 2019***)) emphasizing the public health importance of this

neglected disease in Southeast Asia, and suggesting that more effective control interventions should be

conducted, particularly in the high risk areas. The successful experience in the intervention of Thailand

may be useful for reference by other endemic countries of the region. The national opisthorchiasis control

program, supported by the government of Thailand, applied interrelated approaches, including stool

examination and treatment of positive cases, health education aiming at the promotion of cooked fish

consumption, and environmental sanitation to improve hygienic defecation (***Jongsuksuntigul &***

Imsomboon, 2003). In addition, for areas with difficulties to reduce infection risk, a new strategy was

developed by Sripa and colleagues, using the EcoHealth approach with anthelminthic treatment, novel

intensive health education on both communities and schools, ecosystem monitoring, and active

participation of the community (***Sripa et al., 2015***). This "Lawa model" shows good effectiveness in

Lawa Lake area, where the liver fluke was highly endemic (***Sripa et al., 2015***). Furthermore, common

integrated control interventions (eg, combination of preventive chemotherapy with praziquantel,

improvement of sanitation and water sources, and health education) are applicable not only for

opisthorchiasis but also for other NTDs, such as soil-transmitted helminth infection and schistosomiasis,

which are also prevalent in the study region (***Dunn et al., 2016, Gordon et al., 2019***). Implementation of

such interventions in co-endemic areas could be cost-effective (***Baker et al., 2011, WHO, 2012***).

Frankly, several limitations exist in our study. We collected data from different sources, locations of

which might not be random and preferential sampling may exist. We performed a risk-preferential

sampling test and the results showed that preferential sampling might exist for survey locations in one

third (6/18) of the survey years (Figure 2-source data 2). The corresponding impacts might include

improper variogram estimator, biased parameter estimation and unreliable exposure surface estimates

(***Diggle et al., 2010, Pati et al., 2011, Gelfand et al., 2012***). Nevertheless, a previous study showed that

the INLA method, which we adopted for model fitting, performed well in parameter estimation for both

preferentially and non-preferentially sampled simulated data sets. And in particular, it resulted in better

estimation than that from other methods, such as the Monte Carlo method and the standard MLE

estimation, when fitting the data with preferential sampling (***Dinsdale & Salibian-Barrera, 2019***). To

avoid a more complex model, we didn't take into account the preferential sampling issue for our final

model, as the model validation showed a reasonable capacity of prediction accuracy. However, the

disadvantage of this issue should be well aware.

We set clear criteria for selection of all possible qualified surveys and didn't exclude surveys that reported

prevalence in intervals without exact observed values. Sensitivity analysis showed that the using the

midpoint values of the intervals had little effects on the final results (Figure 3-figure supplement 1). For

surveys across a large area, complex designs, such as randomly sampling from subgroups of the

population under a well-designed scheme, are likely adopted, as it is impractical to draw simple random

samples from the whole area. In such case, respondents may have unequal probabilities to be selected,

thus weighting should be used to generalize results for the entire area. The observed disease data we

collected were from surveys either at point-level (i.e., community or school) or aggregated over areas.

For point-level data, as study areas were quite small, simple sampling design were mostly used in the

corresponding surveys. And for areal-level data, particularly those aggregated across ADM1, complex

designs were likely applied. However, most of the corresponding surveys were only reported raw

prevalence or prevalence without clarifying whether weighting was applied. Thus, we did not have

enough information to address the design effect for each single survey included. On the other hand, as

population density across the study region were different, we calculated the estimated country- and

provincial level prevalence by averaging the estimated pixel-level prevalence weighted by population

density. In this way, we took into account the diversity of population density across areas for regional

summaries of the estimates.

We assumed similar proportions of age and gender in different surveys, as most of which only reported

prevalence aggregated by age and gender. Nevertheless, considering the possible differences in infection

risk between the whole population and schoolchildren, we categorized survey types to the community-

and school-based. Furthermore, our analysis was based on survey data under different diagnostic methods.

The sensitivity and specificity of the same diagnostic method may differ across studies (*Charoensuk et*

al., 2019, Laoprom et al., 2016, Sayasone et al., 2015), while different diagnostic methods may result in

different results in the same survey. To partially taking into account the diversity of diagnostic methods,

we assumed the same diagnostic method has similar sensitivity and specificity, and we considered the

types of diagnostic methods as covariates in the model. Results showed that the odds of infection with

FECT methods was significantly lower than that with Kato-Katz, which was consistent with results found

by Lovis and colleagues (***Lovis et al., 2009***). In addition, most of the diagnostic methods in the surveys

were based on fecal microscopic technique on eggs, which could not effectively distinguish between

O. viverrini and minute intestinal flukes of the family Heterophyidae (eg heterophyid and lecithodendriid)

(***Charoensuk et al., 2019, Sato et al., 2010***). Thus, our results may overestimate the *O. viverrini* infection

risk in areas where heterophyid and lecithodendriid are endemic, such as Phongsaly, Saravane and

Champasak provinces in Lao PDR (***Sato et al., 2010, Chai et al., 2010, Chai et al., 2013***), Nan and

Lampang provinces in Thailand (***Wijit et al., 2013***), and Takeo province in Cambodia (***Sohn et al., 2011***).

There is an urgent need for the application of more powerful diagnostic practices with higher sensitivity

and specificity to better detect the true *O. viverrini* prevalence, such as PCR (***Lovis et al., 2009, Lu et al.,***

2017, Sato et al., 2010). Nevertheless, because of the similar treatment and the prevention strategies of

O. viverrini and minute intestinal flukes (***Keiser & Utzinger, 2010***), our risk mapping is valuable also for

areas co-endemic with the above flukes.

In conclusion, this study contributes to better understand the spatial-temporal characteristics of

O. viverrini infection in major endemic countries of Southeast Asia, providing valuable information

guiding control and intervention, and serving as a baseline for future progress assessment. Estimates were

based on a rigorous geostatistical framework jointly analyzing point- and areal level survey data with

potential predictors. The higher number of infected people we estimated highlights the public health

importance of this neglected disease in the study region. More comprehensive epidemiological studies

are urgently needed for endemic areas with scant survey data.

Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Software, algorithm	R Project for Statistical Computing	R Project for Statistical Computing	RRID:SCR_001905	
Software, algorithm	ArcGIS for Desktop Basic	ArcGIS for Desktop Basic	RRID:SCR_011081	
Software, algorithm	R-INLA Project	R-INLA Project		https://www.r-inla.org/
Software, algorithm	"PStestR" R Package	"PStestR" R Package		https://github.com/joenomiddlename/PStestR

Search strategy, selection criteria and data extraction

We collected relevant publications reporting prevalence data of opisthorchiasis in Southeast Asia through a systematic review (registered in the International Prospective Register of Systematic Reviews, PROSPERO, No.CRD42019136281), and reported our systematic review according to the PRISMA guidelines (Supplementary File 1A) (*Moher et al., 2009*). We searched PubMed and ISI Web of Science from inception to February 9, 2020, with search terms: (liver fluke* OR Opisthorchi*) AND (Southeast Asia OR Indonesia OR (Myanmar OR Burma) OR Thailand OR Vietnam OR Malaysia OR Philippines OR Lao PDR OR Cambodia OR Timor OR Brunei OR Singapore). We set no limitations on language, date of survey, or study design in our search strategy. For literatures not found by the above methods, we also reviewed reports from governments or Ministry of Health, theses, relevant books and documents.

We followed a protocol (Figure 1-figure supplement 1) for inclusion, exclusion and extraction of survey data. Firstly, we screened titles and abstracts to identify potentially relevant articles. Publications on in vitro studies, or absence of human studies or absence of disease studies were excluded. Quality control was undertaken by re-checking 20% of randomly selected irrelevant papers. Second, the full-text review

was applied to potentially relevant articles. We excluded publications with following conditions: absence

of prevalence data; studies done in specific patient groups (eg, prevalence on patients with specific

diseases), in specific population groups (eg, travelers, military personnel, expatriates, nomads, displaced

or migrating population), under specific study designs (eg, case report studies, case-control studies,

clinical trials, autopsy studies); drug efficacy or intervention studies (except for baseline data or control

groups), population deworming within one year, the survey time interval more than ten years, data only

based on the direct smear method (due to low sensitivity) or serum diagnostics (due to unable to differ

the past and the active infection). During the full-text review, the potential relevant cited references of

the articles were also screened. Studies were included if they reported survey data at provincial level and

below, such as administrative divisions of level one (ADM1: province, state, etc.), two (ADM2: city,

etc.), and three (ADM3: county, etc.), and at point-level (village, town, school, etc.). Duplicates were

checked and removed. The quality assessment of each individual record included in the final

geostatistical analysis was performed by two independent reviewers, based on a nine-point quality

evaluation checklist (Figure 2-figure supplement 1-source data 1).

We followed the GATHER checklist (Supplementary File 1B) (*Stevens et al., 2016*) for the data

extraction. Detailed information of records was extracted into a database, which includes literature

information (eg, journal, authors, publication date, title, volume, and issue), survey information (eg,

survey type: community- or school-based, year of survey), location information (eg, location name,

location type, coordinates), and disease related data (eg, species of parasites, diagnostic method,

population age, number of examined, number of positive, and percentage of positive). The coordinates

of the survey locations were obtained from Google Maps (https://www.google.com/maps/). For surveys

reported prevalence in intervals without exact observed values, the midpoints of the intervals were

assigned.

Environmental, socioeconomic, and demographic data

The environmental data (i.e., annual precipitation, distance to the nearest open water bodies, elevation,

land cover, land surface temperature in the daytime and at night and normalized difference vegetation

index), socioeconomic data (i.e., human influence index, survey type, and travel time to the nearest big

city), and demographic data of Southeast Asia were downloaded from open data sources (Figure 7-source

data 1). Land cover data was summarized by the most frequent category within each pixel over the period

of 2001-2018. We combined similar land cover classes and re-grouped them into five categories: (i)

Croplands; (ii) Forests; (iii) Shrub and grass; (iv) Urban; (vi) Others. Land surface temperature (LST) in

the daytime and at night, as well as normalized difference vegetation index (NDVI) were averaged over

the period of 2000-2018. All data were aligned over a 5×5 km grid across the study region (Figure 7).

Data at point-referenced survey locations were extracted. We linked the data to the divisions (i.e., ADM1,

ADM2 or ADM3) reported aggregated outcome of interest (i.e., infection prevalence) by averaging them

within the corresponding divisions. The above data processing was done using the package "ratser"

(https://cran.r-project.org/web/packages/raster) through R (version 3.5.0).

Model fitting and variable selection

As our outcome of interest derived from both point-referenced and area-aggregated surveys, a bivariate

Bayesian geostatistical joint modeling approach was applied to analyze the area-level and point-level

survey data together (*Moraga et al., 2017, Utazi et al., 2019*), and account for both disease data reporting

numbers of examined and positive, and those reporting only prevalence.

We defined p_{it} the probability of infection at location i and time period t, where i is the index either

for the location of point-referenced data or of the area for area-level data. Based on the probability theory,

for data reported with numbers of examined and positive, we assumed that the number of examined Y_{it}

followed a binomial distribution $Y_{it} \sim Bin(p_{it}, N_{it})$, where N_{it} denoted the number of examined; and

for data only reported with the observed prevalence, we assumed that the observed prevalence ob_{it}

followed a beta distribution $ob_{it} \sim Be(p_{it}, \sigma_\beta^2)$. The period of this study were from 1978 to 2018. We

modelled predictors on a logit scale of p_{it}.

We referred to the method proposed by Cameletti and colleagues (***Krainski, 2019, Cameletti et al., 2013***)

to build a spatial-temporal model combined with covariates, which was defined as a SPDE model for the

spatial domain and an AR1 model for the time dimension. A standard grid of 5×5 km^2 was overlaid to

each survey area resulting in a certain number of pixels representing the area. We assumed that survey

locations and pixels within survey areas shared the same spatial-temporal process. In addition, we

assumed the infection risk the same within 1-year period for the same areas. Different observations from

the same year in the same areas can be treated as realizations of the randomized spatial-temporal process.

Let $i = 1, \dots, n_A, n_A + 1, \dots, n_A + n_p$, here n_A was the total number of areas for area-level surveys and

n_p the total number of locations for point-referenced surveys. Regarding to area-level data, $\text{logit}(p_{it}) =$

$\beta_0 + \widetilde{x_t'}\beta + |A_i|^{-1} \int_{A_i} \omega(s,t)dsdt$, where $i = 1, \dots, n_A$, $\widetilde{x_t}$ the vectors of covariate values for i^{th} area

in time period t with $\widetilde{x_{tt}} = |A_i|^{-1} \int_{A_i} x(s,t)dsdt$ and β_0 and β are the intercept and the

corresponding regression coefficients. $|A_i| = \int_{A_i} 1ds$ was the size of the ith area and $\omega(s,t)$ the

spatial-temporal random effects of pixels within the area. For point-referenced data, $\text{logit}(p_{it}) = \beta_0 +$

$x_{it}'\beta + \omega(s_i, t)$, where $i = n_A + 1, \dots, n_A + n_p$, x_{it}' the vectors of covariate values and $\omega(s_i, t)$ the

spatial-temporal random effect for i^{th} location in time period t. To decrease the computational burden,

under the SPDE framework, we built the GMRF on regular temporal knots, that is $\omega =$

$(\omega_{t=1978}, \omega_{t=1983}, \omega_{t=1988}, \omega_{t=1993}, \omega_{t=1998}, \omega_{t=2003}, \omega_{t=2008}, \omega_{t=2013}, \omega_{t=2018})'$ (**Cameletti et al.,**

2013, Krainski, 2019). We assumed the spatio-temporal random effect $\omega(s,t)$ follow a zero-mean

Gaussian distribution, that is $\omega \sim GP(0, \boldsymbol{K}_{space} \otimes \boldsymbol{K}_{time})$, where the spatial covariance matrix $\boldsymbol{K}_{space}$

was defined as a stationary Matérn covariance function $\sigma_\phi^2 (\kappa \boldsymbol{D})^v K_v(\kappa \boldsymbol{D})/(\Gamma(v)2^{v-1})$ and the temporal

covariance matrix as $\boldsymbol{K}_{time} = \rho^{|t_u - t_o|}$ with $|\rho| < 1$, corresponding to the autoregressive stochastic

process with first order (AR1). And the spatio-temporal random effect $\omega(s,t)$ was assumed

independent of each other in different times and locations, that is $\mathrm{Cov}(\omega_{it}, \omega_{jt'}) = \begin{cases} 0, & if \ t \neq t' \\ \sigma_\phi^2, & if \ t = t' \end{cases}$. Here

$\boldsymbol{D}$ donates the Euclidean distance matrix, κ is a scaling parameter and the range $r = \sqrt{8v}/\kappa$,

representing the distance at which spatial correlation becomes negligible (<0.1), and K_v is the modified

Bessel function of the second kind, with the smoothness parameter v fixed at 1. The latent fields

corresponding to other years are approximated by projection of ω using the B-spline basis function of

degree two, that is $B_{i,1}(t) = \begin{cases} 1, & t_i \le t < t_{i+1} \\ 0, & \text{otherwise} \end{cases}$ and $B_{i,m}(t) = \frac{t-t_i}{t_{i+m-1}-t_i} B_{i,m-1}(t) +$

$\frac{t_{i+m}-t}{t_{i+m}-t_{i+1}} B_{i+1,m-1}(t)$, where m is the degree of two (**Cameletti et al., 2013, Krainski, 2019**).

We formulated the model in a Bayesian framework. Minimally informative priors were specified for

parameters and hyper parameters as following: $\boldsymbol{\beta} \sim N(0, 10^5 \boldsymbol{I})$, $\log(1/\sigma_\beta^2) \sim \log\mathrm{Gamma}(1, 0.1)$,

$\log(1/\sigma_\phi^2) \sim \log\mathrm{Gamma}(1, 0.01)$, $\log((1+\rho)/(1-\rho)) \sim N(0, 0.15)$, and $\log(\kappa) \sim N(\log(\sqrt{8}/d), 1)$,

where d is the median distance between the predicted grids.

Additionally, we applied variable selection procedure to identify the best set of predictors for a

parsimonious model. Firstly, the best functional form (continuous or categorical) of continuous variables

was selected, by fitting univariate Bayesian spatial-temporal models with either form as the independent

variable and selecting the form with the lowest log score (**Pettit, 1990**). Secondly, the best subset method

was used to identify the best combination of predictors for the final model. According to previous studies

(**Aung et al., 2017, Forrer et al., 2012, Miyamoto et al., 2014, Wongsaroj et al., 2014**), the infection risk

in community and school may be different, and using different diagnostic methods may differ the

observed prevalence. Thus, the survey type (i.e., community- or school-based) and the diagnostic

methods (i.e., Kato-Katz, FECT, or other methods) were kept in all potential models, while the other ten

environmental and socioeconomic variables were put forth into the Bayesian variable selection process.

The model with the minimum log score was chosen as the final model.

Model fitting and variable selection process were conducted through INLA-SPDE approach (**Lindgren**

et al., 2011, Rue et al., 2009), using integrated nested Laplace approximation (INLA) package in R

(version 3.5.0). Estimation of risk for *O. viverrini* infection in each year of the study period was done

over a grid with cell size of 5×5 km^2. And the relative changes of the prevalence were also calculated

using a formula as $(pp_{st_i} - pp_{st_j})/pp_{st_i}$ for pixel s between the former year t_i and the later year t_j,

where pp indicates the median of the posterior estimated distribution of infection risk. The

corresponding risk maps and the prevalence changing maps were produced using ArcGIS (version 10.2).

In addition, as population density across the study region were different, the population-adjusted

estimated prevalence and number of infected individuals in 2018 were calculated at the country- and

provincial levels averaging the estimated pixel-level prevalence weighted by population density, that is

$\widehat{pp}_A = \sum_{i \in A} \widehat{pp}_i w_i / \sum_{i \in A} w_i$. Here $\widehat{pp}_A$, $\widehat{pp}_i$ and w_{it} are the estimated prevalence in area A, estimated

prevalence at pixel i and population density at pixel i, respectively, where i belongs to area A. Based

on previous studies, for the provinces in Vietnam where there was no evidence of *O. viverrini* infection,

we multiplied the estimated results by zero as the final estimated prevalence (**Doanh & Nawa, 2016**).

The R code used for model fitting is publicly available in GitHub (https://github.com/SYSU-

Opisthorchiasis).

Model validation, sensitivity analysis and test of preferential sampling

Model validation was conducted using the five-fold out-of-sample cross-validation approach. Mean error

(ME $= \frac{1}{N}\sum(ob_{it} - pp_{it})$), mean absolute error (MAE $= \frac{1}{N}\sum|ob_{it} - pp_{it}|$), mean square error (MSE $=$

$\frac{1}{N}\sum(ob_{it} - pp_{it})^2$), and the coverage rate of observations within 95% Bayesian credible intervals (BCI)

were calculated to evaluate the performance of the model. Furthermore, a Bayesian geostatistical model

only based on point-referenced data was fitted and validated, to compared its performance with our joint

modeling approach. In addition, a sensitivity analysis was conducted to evaluate the effects of using the

midpoint values of the intervals as the observed prevalence in one literature from Suwannatrai and

colleagues (***Suwannatrai et al., 2018***), reporting observed prevalence of *O. viverrini* infection in intervals.

Sensitivity analysis was done by using the lower and the upper limits of the intervals in the modeling

analysis.

Considering that the data in this study were sourced from different studies, preferential sampling may

exist. We performed a test for preferential sampling of the data. To our knowledge, there hasn't been

method developed for preferential sampling test on observations combined at point- and areal levels. To

compromise, we took centers of the areas with survey data as their locations for the test of preferential

sampling. A fast and intuitive Monte Carlo test developed by Watson was adopted for its advantage of

fast speed and feasibility of data arising from various distributions. We assumed S_t (i.e., the collection

of sampled points at time t) a realization from an inhomogeneous Poisson processes (IPP) under the

condition of $\omega(s,t)$ (i.e., the spatial-temporal Gaussian random field), that is $[S_t|\omega(s,t)] =$

IPP$(\lambda(s,t))$, and $\log(\lambda(s,t)) = \alpha_0 + h(\omega(s,t))$, where h is a monotonic function of $\omega(s,t)$. When

$h \equiv 0$, the sampling process is independent from $\omega(s,t)$, thus the preferential sampling is not significant.

In this way, the problem of detecting preferential sampling can be transformed into the hypothesis testing

of $h \equiv 0$. If $h \equiv 0$ is false, for example, in case that h is a monotonic increasing function of $\omega(s,t)$,

then the point patterns S_t are expected to exhibit an excess of clustering in areas with higher $\omega(s,t)$,

thus positive association can be detected between the localized amount of clustering and estimated

$\omega(s,t)$. Firstly, we used the mean of the distances to the K nearest points (D_K) to measure the clustering

of locations, and calculated the rank correlation $r_{t(K)}$ between D_K and the estimated $\omega(s,t)$ for survey

year t. Here the estimated $\omega(s,t)$ was obtained from fitting the Bayesian spatial-temporal joint model.

Next, the Monte Carlo method was used to sample realizations from the IPP under the null hypothesis

(i.e., $h \equiv 0$), following which, a set of rank correlations $r_{t(K)}^M$ were calculated, approximating the

distribution of the rank correlations $\rho_{t(K)}$ under $h \equiv 0$. In this way, the nonstandard sampling

distribution of the test statistic can be approximated. Finally, we computed the desired empirical p-value

by evaluating the proportion of the Monte Carlo-sampled $r_{t(K)}^M$ which are more extreme than $r_{t(K)}$. We

set a sample size of 1000 for each Monte Carlo sampling. We also considered K from 1 to 8 to measure

the clustering of locations and resulted in eight p-values respective to different K for each survey year. If

one of the p-values is smaller or equal to 0.05, we considered preferential sampling existing in the

corresponding survey year. Since our model could estimate the disease risk each year of the study period,

this test was done for each survey year with number of locations more than or equal to 10 (i.e., 1978,

1981, 1991, 1995, 1998, 2000, 2001, 2004, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, and

2016). The test was conducted using the package "PStestR" in R (version 3.6.3) (***Watson, 2020***).

Additional information

Funding

Funder	Grant reference number	Author
The Sun Yat-Sen University One Hundred Talent Grant	-	Ying-Si Lai
The Natural Science Foundation of Guangdong Province	2017A030313704	Ying-Si Lai
The National Natural Science Foundation of China	81703320	Ying-Si Lai
The CMB Open Competition Program	17-274	Ying-Si Lai

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Ethics statement

This work was based on survey data pertaining to the prevalence of opisthorchiasis extracted from open

published peer-reviewed literatures. All data were aggregated and did not contain any information at the

individual or household levels. Therefore, there were no specific ethical issues warranted special

attention.

Acknowledgments

We are grateful to Dr Roy Burstein from Institute for Disease Modeling, Bellevue, Washington, USA for

providing very good suggestions for the manuscript.

Competing interests

The authors declare that no competing interests exist.

References

Andrews RH, Sithithaworn P, Petney TN. 2008. *Opisthorchis viverrini*: an underestimated parasite in world health. *Trends in Parasitology* **24**:497-501. doi:10.1016/j.pt.2008.08.011

Aung WPP, Htoon TT, Tin HH, Thinn KK, Sanpool O, Jongthawin J, Sadaow L, Phosuk I, Rodpai R, Intapan PM, Maleewong W. 2017. First report and molecular identification of *Opisthorchis viverrini* infection in human communities from Lower Myanmar. *PLOS ONE* **12**:e177130. doi:10.1371/journal.pone.0177130

Baker M, Hanson C, Ottesen EA, Linehan M, Torres S, Sankara D, Zoerhoff KL, Weaver A, Kabore A. 2011. Integrated Implementation of Programs Targeting Neglected Tropical Diseases through Preventive Chemotherapy: Proving the Feasibility at National Scale. *The American Journal of Tropical Medicine and Hygiene* **84**:5-14. doi:10.4269/ajtmh.2011.10-0411

Banerjee S, Carlin BP, Gelfand AE. 2014. Hierarchical Modeling and Analysis for Spatial Data, Second Edition. Boca Raton, FL: Chapman & Hall/CRC Press.

Cameletti M, Lindgren F, Simpson D, Rue H. 2013. Spatio-temporal modeling of particulate matter concentration through the SPDE approach. *Asta-Advances in Statistical Analysis* **97**:109-31. doi:10.1007/s10182-012-0196-3

Chai JY, Yong TS, Eom KS, Min DY, Jeon HK, Kim TY, Jung BK, Sisabath L, Insisiengmay B, Phommasack B, Rim HJ. 2013. Hyperendemicity of *Haplorchis taichui* Infection among Riparian People in Saravane and Champasak Province, Lao PDR. *The Korean Journal of Parasitology* **51**:305-311. doi:10.3347/kjp.2013.51.3.305

Chai JY, Yong TS, Eom KS, Min DY, Jeon HK, Kim TY, Jung BK, Sisabath L, Insisiengmay B, Phommasack B, Rim HJ. 2010. Prevalence of the Intestinal Flukes *Haplorchis taichui* and *H. yokogawai* in a Mountainous Area of Phongsaly Province, Lao PDR. *The Korean Journal of Parasitology* **48**:339-342. doi:10.3347/kjp.2010.48.4.339

Charoensuk L, Subrungruang I, Mungthin M, Pinlaor S, Suwannahitatorn P. 2019. Comparison of stool examination techniques to detect *Opisthorchis viverrini* in low intensity infection. *Acta Tropica* **191**:13-16. doi:10.1016/j.actatropica.2018.12.018

Diggle PJ, Menezes R, Su TL. 2010. Geostatistical inference under preferential sampling. *Journal of the Royal Statistical Society Series C-Applied Statistics* **59**:191-232. doi:10.1111/j.1467-9876.2009.00701.x

Dinsdale D, Salibian-Barrera M. 2019. Methods for preferential sampling in geostatistics. *Journal of the Royal Statistical Society Series C-Applied Statistics* **68**:181-98. doi:10.1111/rssc.12286

Doanh PN, Nawa Y. 2016. *Clonorchis sinensis* and *Opisthorchis spp.* in Vietnam: current status and prospects. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **110**:13-20. doi:10.1093/trstmh/trv103

Dunn JC, Turner HC, Tun A, Anderson RM. 2016. Epidemiological surveys of, and research on, soil-transmitted helminths in Southeast Asia: a systematic review. *Parasites & Vectors* **9**:31. doi:10.1186/s13071-016-1310-2

Echaubard P, Sripa B, Mallory FF, Wilcox BA. 2016. The role of evolutionary biology in research and
control of liver flukes in Southeast Asia. *Infection, Genetics and Evolution* **43**:381-397.
doi:10.1016/j.meegid.2016.05.019

Elith J, Phillips SJ, Hastie T, Dudk M, Chee YE, Yates CJ. 2011. A statistical explanation of MaxEnt
for ecologists. *Diversity and Distributions* **17**:43-57. doi:10.1111/j.1472-4642.2010.00725.x

Forrer A, Sayasone S, Vounatsou P, Vonghachack Y, Bouakhasith D, Vogt S, Glaser R, Utzinger J,
Akkhavong K, Odermatt P. 2012. Spatial distribution of, and risk factors for, *Opisthorchis viverrini*
infection in southern Lao PDR. *PLOS Neglected Tropical Diseases* **6**:e1481.
doi:10.1371/journal.pntd.0001481

Fürst T, Keiser J, Utzinger J. 2012. Global burden of human food-borne trematodiasis: a systematic
review and meta-analysis. *The Lancet Infectious Diseases* **12**:210-221. doi:10.1016/S1473-
3099(11)70294-8

Gelfand AE, Banerjee S. 2017. Bayesian Modeling and Analysis of Geostatistical Data. *Annual Review*
of Statistics and Its Application **4**:245-266. doi:10.1146/annurev-statistics-060116-054155

Gelfand AE, Sahu SK, Holland DM. 2012. On the effect of preferential sampling in spatial prediction.
Environmetrics **23**:565-78. doi:10.1002/env.2169

Gordon C, Kurscheid J, Williams G, Clements A, Li Y, Zhou X, Utzinger J, McManus D, Gray D. 2019.
Asian Schistosomiasis: Current Status and Prospects for Control Leading to Elimination. *Tropical*
Medicine and Infectious Disease **4**:40. doi:10.3390/tropicalmed4010040

Grundy-Warr C, Andrews RH, Sithithaworn P, Petney TN, Sripa B, Laithavewat L, Ziegler AD. 2012.
Raw attitudes, wetland cultures, life-cycles: Socio-cultural dynamics relating to *Opisthorchis viverrini*
in the Mekong Basin. *Parasitology International* **61**:65-70. doi:10.1016/j.parint.2011.06.015

Havelaar AH, Kirk MD, Torgerson PR, Gibb HJ, Hald T, Lake RJ, Praet N, Bellinger DC, de Silva NR,
Gargouri N, Speybroeck N, Cawthorne A, Mathers C, Stein C, Angulo FJ, Devleesschauwer B. 2015.
World Health Organization Global Estimates and Regional Comparisons of the Burden of Foodborne
Disease in 2010. *PLOS Medicine* **12**:e1001923. doi:10.1371/journal.pmed.1001923

International Agency for Research on Cancer. 1994. Schistosomes, liver flukes and Helicobacter pylori.
IARC Monographs on the Evaluation of Carcinogenic Risks to Humans **61**:1-241.
https://www.ncbi.nlm.nih.gov/books/NBK487782/

Jongsuksuntigul P, Imsomboon T. 2003. Opisthorchiasis control in Thailand. *Acta Tropica* **88**:229-232.
doi:10.1016/j.actatropica.2003.01.002

Jongsuksuntigul P, Manatrakul D, T TW, Krisanamara P, Sawatdimongkol S, Wongsaroj S. 2003.
Evaluation of the Helminthiasis Control Program in Thailand at the End of the 8th Health Development
Plan. *The Journal of Tropical Medicine and Parasitology* **26**:18-45. https://li01.tci-
thaijo.org/index.php/JTMP/article/view/14757

Kaewpitoon N, Kaewpitoon SJ, Pengsaa P. 2008. Opisthorchiasis in Thailand: review and current status.
World Journal of Gastroenterology **14**:2297-2302. doi:10.3748/wjg.14.2297

Karagiannis-Voules D, Biedermann P, Ekpo UF, Garba A, Langer E, Mathieu E, Midzi N, Mwinzi P,

Polderman AM, Raso G, Sacko M, Talla I, Tchuenté LT, Touré S, Winkler MS, Utzinger J, Vounatsou P. 2015. Spatial and temporal distribution of soil-transmitted helminth infection in sub-Saharan Africa: a systematic review and geostatistical meta-analysis. *The Lancet Infectious Diseases* **15**:74-84. doi:10.1016/S1473-3099(14)71004-7

Keiser J, Utzinger J. 2010. The Drugs We Have and the Drugs We Need Against Major Helminth Infections. *Advances in Parasitology* **73**:197-230. doi:10.1016/S0065-308X(10)73008-6

Keiser J, Utzinger J. 2005. Emerging Foodborne Trematodiasis. *Emerging Infectious Diseases* **11**:1507-1514. doi:10.3201/eid1110.050614

Keiser, Jennifer. 2019. Highlighting Operational and Implementation Research for Control of Helminthiasis. GB: Academic Press.

Khieu V, Furst T, Miyamoto K, Yong TS, Chai JY, Huy R, Muth S, Odermatt P. 2019. Is *Opisthorchis viverrini* Emerging in Cambodia? *Advances in Parasitology* **103**:31-73. doi:10.1016/bs.apar.2019.02.002

Krainski ET. 2019. Advanced spatial modeling with stochastic partial differential equations using R and INLA. Boca Raton: CRC Press, Taylor & Francis Group.

Laoprom N, Laithavewat L, Kopolrat K, Kiatsopit N, Kaewkes S, Chantalux S, Mongkolsin C, Chanmaha B, Andrews RH, Petney TN, Sithithaworn P. 2016. Evaluation of a commercial stool concentrator kit compared to direct smear and formalin-ethyl acetate concentration methods for diagnosis of parasitic infection with special reference to *Opisthorchis viverrini* sensu lato in Thailand. *Southeast Asian Journal of Tropical Medicine and Public Health* **47**:890-900. PMID: 29620342

Lindgren F, Rue H, Lindström J, Kent JT, Diggle PJ, Illian JB, Simpson DP, Gneiting T, Scheuerer M, Furrer R, Furrer E, Nychka D, Fearnhead P, Challenor P, Andrianakis Y, Stephenson G, Møller J, Hu X, Simpson D, Bolin D, Brown PE, Cameletti M, Martino S, Cooley D, Hoeting JA, Crujeiras RM, Prieto A, Marco ARF, Fuglstad G, Gelman A, Guttorp P, Das B, Haaland B, Nott DJ, Haslett J, Joshi C, Garreta V, Höhle M, Ippoliti L, Martin RJ, Bhansali RJ, Jandhyala VK, Fotopoulos SB, Jun M, Kongsgård HW, Lasinio GJ, Pollice A, Lee C, Lee WT, Kaufman CG, Li B, Genton MG, Lindgren G, Mardia KV, Mateu J, Mondal D, Müller WG, Waldl H, Ottavi A, Papaspiliopoulos O, Porcu E, Saez M, Schmidt AM, Stein A, Switzer P, Turkman K, Wikle CK, Hooten MB. 2011. An explicit link between Gaussian fields and Gaussian Markov random fields: the stochastic partial differential equation approach. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **73**:423-498. doi:10.1111/j.1467-9868.2011.00777.x

Lovis L, Mak TK, Phongluxa K, Soukhathammavong P, Sayasone S, Akkhavong K, Odermatt P, Keiser J, Felger I. 2009. PCR Diagnosis of *Opisthorchis viverrini* and *Haplorchis taichui* Infections in a Lao Community in an Area of Endemicity and Comparison of Diagnostic Methods for Parasitological Field Surveys. *Journal of Clinical Microbiology* **47**:1517-1523. doi:10.1128/JCM.02011-08

Lu Y, Gonzales G, Chen S, Li H, Cai Y, Chu Y, Ai L, Chen M, Chen H, Chen J. 2017. Urgent needs in fostering neglected tropical diseases (NTDs) laboratory capacity in WHO Western Pacific Region: results from the external quality assessment on NTDs diagnosis in 2012-2015. *Infectious Diseases of Poverty* **6**:106-108. doi:10.1186/s40249-017-0319-x

Maipanich W, Waikagul J, Watthanakulpanich D, Muennoo C, Sanguankiat S. 2004. Intestinal

Parasitic Infections among Inhabitants of the North, West-central and Eastern Border Areas of Thailand. *The Journal of Tropical Medicine and Parasitology* **27**:51-58. https://scholar.google.com/scholar_lookup?journal=J+Trop+Med+Parasitol&title=Intestinal+parasitic+infections+among+inhabitants+of+the+North,+West-central+and+Eastern+border+areas+of+Thailand&author=W+Maipanich&author=J+Waikagul&author=D+Watthanakulpanich&author=C+Muennoo&author=S+Sanguankiat&volume=27&publication_year=2004&pages=51-58&

Miyamoto K, Kirinoki M, Matsuda H, Hayashi N, Chigusa Y, Sinuon M, Chuor CM, Kitikoon V. 2014. Field survey focused on *Opisthorchis viverrini* infection in five provinces of Cambodia. *Parasitology International* **63**:366-373. doi:10.1016/j.parint.2013.12.003

Moher D, Liberati A, Tetzlaff J, Altman DG. 2010. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *International Journal of Surgery* **8**:336-341. doi:10.1016/j.ijsu.2010.02.007

MOPH. 2014. National Survey on Helminthiasis in Thailand. Department of Disease Control, Ministry of Public Health, Bangkok, Thailand

Moraga P, Cramb SM, Mengersen KL, Pagano M. 2017. A geostatistical model for combined analysis of point-level and area-level data using INLA and SPDE. *Spatial Statistics* **21**:27-41. doi:10.1016/j.spasta.2017.04.006

Ogorodova LM, Fedorova OS, Sripa B, Mordvinov VA, Katokhin AV, Keiser J, Odermatt P, Brindley PJ, Mayboroda OA, Velavan TP, Freidin MB, Sazonov AE, Saltykova IV, Pakharukova MY, Kovshirina YV, Kaloulis K, Krylova OY, Yazdanbakhsh M. 2015. Opisthorchiasis: An Overlooked Danger. *PLOS Neglected Tropical Diseases* **9**:e3563. doi:10.1371/journal.pntd.0003563

Pati D, Reich BJ, Dunson DB. 2011. Bayesian geostatistical modelling with informative sampling locations. *Biometrika* **98**:35-48. doi:10.1093/biomet/asq067

Petney TN, Andrews RH, Saijuntha W, Wenz-Mücke A, Sithithaworn P. 2013. The zoonotic, fish-borne liver flukes *Clonorchis sinensis*, *Opisthorchis felineus* and *Opisthorchis viverrini*. *International Journal for Parasitology* **43**:1031-1046. doi:10.1016/j.ijpara.2013.07.007

Pettit LI. 1990. The Conditional Predictive Ordinate for the Normal Distribution. Journal of the Royal Statistical Society. Series B (Methodological) **52**:175-184. doi:10.1111/j.2517-6161.1990.tb01780.x

Phimpraphai W, Tangkawattana S, Kasemsuwan S, Sripa B. 2018. Social Influence in Liver Fluke Transmission: Application of Social Network Analysis of Food Sharing in Thai Isaan Culture. *Advances in Parasitology* **101**:97-124. doi:10.1016/bs.apar.2018.05.004

Phimpraphai W, Tangkawattana S, Sereerak P, Kasemsuwan S, Sripa B. 2017. Social network analysis of food sharing among households in opisthorchiasis endemic villages of Lawa Lake, Thailand. *Acta Tropica* **169**:150-156. doi:10.1016/j.actatropica.2017.02.001

Prueksapanich P, Piyachaturawat P, Aumpansub P, Ridtitid W, Chaiteerakij R, Rerknimitr R. 2018. Liver Fluke-Associated Biliary Tract Cancer. *Gut and Liver* **12**:236-245. doi:10.5009/gnl17102

Qian M, Zhou X. 2019. Human liver flukes in China and ASEAN: Time to fight together. *PLOS Neglected Tropical Diseases* **13**:e7214. doi:10.1371/journal.pntd.0007214

Rue H, Martino S, Chopin N. 2009. Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **71**:319-392. doi:10.1111/j.1467-9868.2008.00700.x

Saijuntha W, Sithithaworn P, Kaitsopit N, Andrews RH, Petney TN. 2014. Liver flukes: Clonorchis and Opisthorchis. *Advances in Experimental Medicine and Biology* **766**:153-199. doi:10.1007/978-1-4939-0915-5_6

Sanderson EW, Jaiteh M, Levy MA, Redford KH, Woolmer G. 2002. The Human Footprint and the Last of the Wild: The human footprint is a global map of human influence on the land surface, which suggests that human beings are stewards of nature, whether we like it or not. *BioScience* **52**:891-904. doi:10.1641/0006-3568(2002)052[0891:THFATL]2.0.CO;2

Sato M, Pongvongsa T, Sanguankiat S, Yoonuan T, Dekumyoy P, Kalambaheti T, Keomoungkhoun M, Phimmayoi I, Boupha B, Moji K, Waikagul J. 2010. Copro-DNA diagnosis of *Opisthorchis viverrini* and *Haplorchis taichui* infection in an endemic area of Lao PDR. *The Southeast Asian Journal of Tropical Medicine and Public Health* **41**:28. PMID: 20578479

Sayasone S, Utzinger J, Akkhavong K, Odermatt P. 2015. Repeated stool sampling and use of multiple techniques enhance the sensitivity of helminth diagnosis: A cross-sectional survey in southern Lao People's Democratic Republic. *Acta Tropica* **141**:315-321. doi:10.1016/j.actatropica.2014.09.004

Sithithaworn P, Andrews RH, Van De N, Wongsaroj T, Sinuon M, Odermatt P, Nawa Y, Liang S, Brindley PJ, Sripa B. 2012. The current status of opisthorchiasis and clonorchiasis in the Mekong Basin. *Parasitology International* **61**:10-16. doi:10.1016/j.parint.2011.08.014

Smith BJ, Yan J, Cowles MK. 2008. Unified Geostatistical Modeling for Data Fusion and Spatial Heteroskedasticity with R Package ramps. *Journal of Statistical Software* **25**. doi:10.18637/jss.v025.i10

Sohn WM, Jung BK, Hong SJ, Lee KH, Park JB, Kim HS, Cho S, Htoon TT, Tin HH, Chai JY. 2019. Low-Grade Endemicity of Opisthorchiasis, Yangon, Myanmar. *Emerging Infectious Diseases* **25**:1435-1437. doi:10.3201/eid2507.190495

Sohn WM, Shin EH, Yong TS, Eom KS, Jeong HG, Sinuon M, Socheat D, Chai JY. 2011. Adult *Opisthorchis viverrini* Flukes in Humans, Takeo, Cambodia. *Emerging Infectious Diseases* **17**:1302-1304. doi:10.3201/eid1707.102071

Sripa B, Kaewkes S, Intapan PM, Maleewong W, Brindley PJ. 2010. Food-Borne Trematodiases in Southeast Asia: Epidemiology, Pathology, Clinical Manifestation and Control. *Advances in Parasitology* **72**:305-350. doi:10.1016/S0065-308X(10)72011-X

Sripa B, Tangkawattana S, Laha T, Kaewkes S, Mallory FF, Smith JF, Wilcox BA. 2015. Toward integrated opisthorchiasis control in northeast Thailand: The Lawa project. *Acta Tropica* **141**:361-367. doi:10.1016/j.actatropica.2014.07.017

Stevens GA, Alkema L, Black RE, Boerma JT, Collins GS, Ezzati M, Grove JT, Hogan DR, Hogan MC, Horton R, Lawn JE, Marušić A, Mathers CD, Murray CJL, Rudan I, Salomon JA, Simpson PJ, Vos T, Welch V. 2016. Guidelines for Accurate and Transparent Health Estimates Reporting: the GATHER statement. *PLOS Medicine* **13**:e1002116. doi:10.1371/journal.pmed.1002056

Suwannatrai A, Pratumchart K, Suwannatrai K, Thinkhamrop K, Chaiyos J, Kim CS, Suwanweerakamtorn R, Boonmars T, Wongsaroj T, Sripa B. 2017. Modeling impacts of climate change on the potential distribution of the carcinogenic liver fluke, *Opisthorchis viverrini*, in Thailand. *Parasitology Research* **116**:243-250. doi:10.1007/s00436-016-5285-x

Suwannatrai A, Saichua P, Haswell M. 2018. Epidemiology of *Opisthorchis viverrini* Infection. *Advances in Parasitology* **101**:41-67. doi:10.1016/bs.apar.2018.05.002

UN. 2015. Transforming our world: the 2030 Agenda for Sustainable Development. United Nations, New York, NY, USA

Utazi CE, Thorley J, Alegana VA, Ferrari MJ, Nilsen K, Takahashi S, Metcalf C, Lessler J, Tatem AJ. 2019. A spatial regression model for the disaggregation of areal unit based data to high-resolution grids with application to vaccination coverage mapping. *Statistical Methods in Medical Research* **28**:3226-3241. doi:10.1177/0962280218797362

Van De N. 2004. Fish-borne trematodes in Vietnam. *Southeast Asian Journal of Tropical Medicine and Public Health* **35**:299-301. https://pdfs.semanticscholar.org/4952/fe365526c6cf9bb2e8157 1d151450d18ef0f.pdf

Wang YC, Feng CC, Sithithaworn P. 2013. Environmental determinants of Opisthorchis viverrini prevalence in northeast Thailand. *Geospat Health* **8**:111-23. doi:10.4081/gh.2013.59

Watson J. 2020. A fast Monte Carlo test for preferential sampling. *arXiv*. https://arxiv.org/abs/2003.01319v2 [Preprint]

WHO. 2002. Joint WHO/FAO Workshop on Foodborne Trematode Infections in Asia. World Health Organization, Geneva

WHO. 2009. Report of the WHO Expert Consultation on Foodborne Trematode Infections and Taeniasis/Cysticercosis. World Health Organization, Geneva

WHO. 2012. Accelerating work to overcome the global impact of neglected tropical diseases. World Health Organization, Geneva

Wijit A, Morakote N, Klinchid J. 2013. High Prevalence of Haplorchiasis in Nan and Lampang Provinces, Thailand, Proven by Adult Worm Recovery from Suspected Opisthorchiasis Cases. *The Korean Journal of Parasitology* **51**:767-769. doi:10.3347/kjp.2013.51.6.767

Wongsaroj T, Nithikathkul C, Rojkitikul W, Nakai W, Royal L, Rammasut P. 2014. National survey of helminthiasis in Thailand. *Asian Biomedicine* **8**:779-783. doi:10.5372/1905-7415.0806.357

Yong T, Chai J, Sohn W, Eom KS, Jeoung H, Hoang E, Yoon C, Jung B, Lee S, Sinuon M, Socheat D. 2014. Prevalence of Intestinal Helminths among Inhabitants of Cambodia (2006-2011). *The Korean Journal of Parasitology* **52**:661-666. doi:10.3347/kjp.2014.52.6.661

Ziegler AD, Andrews RH, Grundy-Warr C, Sithithaworn P, Petney TN. 2011. Fighting Liverflukes with Food Safety Education. *Science* **331**:282-283. doi:10.1126/science.331.6015.282-b

711 **Tables**

712 **Table 1. Overview of opisthorchiasis survey data in Southeast Asia**

	Cambodia	Lao PDR	Myanmar	Thailand	Vietnam	Total
Relevant papers	14	43	2	97	15	168
Total surveys/locations	91/73	156/99	6/6	770/335	71/64	1094/574
Survey type (surveys/locations)						
School	33/31	4/4	0/0	13/13	0/0	50/48
Community	58/46	152/94	6/6	757/325	71/64	1044/535
Location type (surveys/locations)						
Point-level	55/43	63/51	3/3	125/105	5/5	251/207
ADM3-level	0/0	0/0	0/0	53/51	0/0	53/51
ADM2-level	14/11	35/27	0/0	159/102	2/2	210/142
ADM1-level	22/19	58/18	3/3	433/77	64/57	580/174
Period	1998-2016	1989-2016	2015-2016	1978-2018	1991-2015	1978-2018
Year of Survey (surveys/locations)						
1978-1982	0/0	0/0	0/0	123/115	0/0	123/115
1983-1987	0/0	0/0	0/0	7/6	0/0	7/6
1988-1992	0/0	2/2	0/0	97/89	1/1	100/92
1993-1997	0/0	9/5	0/0	18/18	6/2	33/25
1998-2002	25/22	28/22	0/0	103/103	2/2	158/149
2003-2007	3/2	26/24	0/0	15/15	1/1	45/42
2008-2012	62/48	75/54	0/0	166/153	9/8	312/263
2013-2018	1/1	16/16	6/6	241/201	52/52	316/276
Diagnostic methods (surveys/locations)						
Kato-Katz	86/70	128/83	3/3	212/166	7/7	436/329
FECT	2/2	8/7	3/3	109/99	0/0	122/111
Stoll's	0/0	0/0	0/0	38/28	0/0	38/28
PCR	0/0	5/4	0/0	1/1	0/0	6/5
Combined	3/3	14/13	0/0	14/12	0/0	31/28
Others	0/0	1/1	0/0	6/6	0/0	7/7
NS[*]	0/0	5/5	0/0	391/111	64/57	460/173
Mean prevalence	10.56%	39.50%	4.93%	14.25%	2.65%	16.74%

713 [*]NS: not stated or missing

715   **Table 2. Posterior summaries of model parameters**

	Estimated median (95% BCI)	OR	Prob (%)[†]
Intercept	-4.51 (-5.08, -3.94)		
Survey type			
School-based survey	Ref	Ref	-
Community-based survey	0.96 (0.70, 1.23)	2.61 (2.10, 3.42)	> 99.99%
Diagnostic methods			
Kato-Katz	Ref	Ref	Ref
FECT	-0.28 (-0.49, -0.07)	0.76 (0.61, 0.93)	0.80
Other methods	0.01 (-0.07, 0.10)	1.01 (0.93, 1.12)	64.20
Land surface temperature (LST) in the daytime (°C)			
< 30.65	Ref	Ref	Ref
30.65 - 32.07	0.25 (-0.001, 0.50)	1.28 (0.999, 1.65)	97.40
> 32.07	0.07 (-0.18, 0.33)	1.07 (0.84, 1.39)	73.40
Human influence index	-0.01 (-0.02, -0.003)	0.99 (0.98, 1.00)	0.80
Distance to the nearest open water bodies (km)	0.24 (-1.45, 1.94)	1.27 (0.23, 6.96)	60.20
Elevation (m)	-0.003 (-0.005, -0.001)	0.997 (0.995, 0.999)	< 0.10
Travel time to the nearest big city (min)	0.0001 (-0.002, 0.002)	1.00 (0.998, 1.002)	56.60

716   [†]Posterior probability of *OR*>1.

718   **Table 3. Population-adjusted estimated prevalence and number of individuals infected with**

719   ***O. viverrini* in endemic countries of Southeast Asia in 2018[*]**

	Population (×10³)	Prevalence (%)	No. infected (×10³)
Cambodia	16227.39	6.15% (2.41%, 11.73%)	997.95 (390.46, 1903.46)
Lao PDR	6960.28	35.21% (28.50%, 40.70%)	2450.54 (1983.38, 2832.96)
Thailand	69112.64	9.71% (7.98%, 12.17%)	6708.68 (5514.87, 8411.98)
Vietnam	96421.69	2.15% (0.73%, 4.40%)	2073.72 (703.46, 4244.85)
Total	188722.01	6.57% (5.35%, 7.98%)	12389.69 (10099.29, 15060.18)

[*]Estimates were based on gridded population of 2018 and the median and 95% BCI of the posterior

estimated distribution of the infection risk in 2018.

List of Figures

Figure 1. Data search and selection flow chart.

Figure 2. Survey locations and observed prevalence of *O. viverrini* infection in endemic countries

of Southeast Asia. (A) 1978-1982, (B) 1983-1987, (C) 1988-1992, (D) 1993-1997, (E) 1998-2002, (F)

2003-2007, (G) 2008-2012, and (H) 2013-2018. (Source Data: Figure 2-source data 1, and Figure 2-

source data 2)

Figure 3. Model-based estimated risk maps of *O. viverrini* infection in endemic countries of

Southeast Asia in different years. Estimated prevalence based on the median of the posterior estimated

distribution of infection risk in (A) 1978, (B) 1983, (C) 1988, (D) 1993, (E) 1998, (F) 2003, (G) 2008,

(H) 2013, and (I) 2018. (Source Data: Figure 3-source data 1)

Figure 4. The estimation uncertainty in endemic countries of Southeast Asia in different years. (A)

1978, (B) 1983, (C) 1988, (D) 1993, (E) 1998, (F) 2003, (G) 2008, (H) 2013, and (I) 2018. (Source Data:

Figure 4-source data 1)

Figure 5. Changes of *O. viverrini* infection risk across time periods. Changes were calculated by the

median of the posterior estimated distribution of infection risk for the former time period minus that for

the latter time period divided by that for the former time period. The risk changes (A) between 1978 and

2018; (B) between 1978 and 1983; (C) between 1983 and 1988; (D) between 1988 and 1993; (E) between

1993 and 1998; (F) between 1998 and 2003; (G) between 2003 and 2008; (H) between 2008 and 2013;

and (I) between 2013 and 2018. (Source Data: Figure 5-source data 1)

Figure 6. Trends in estimated prevalence of *O. viverrini* infection in Southeast Asia. (Source Data:

Figure 6-source data 1)

Figure 7. Images of spatial covariates used in the present study. (Source Data: Figure 7-source data

1)

List of figure supplements for primary figures

Figure 1-figure supplement 1. Study protocol.

Figure 2-figure supplement 1. Result of quality assessment of eligible studies. (Source Data: Figure

2-figure supplement 1-source data 1)

Figure 3-figure supplement 1. Model-based estimated risk maps of *O. viverrini* infection in 2018

under different values assigned to prevalence for surveys only reported prevalence in intervals. (A)

the midpoint values of the intervals, (B) the lower limits and (C) the upper limits of the intervals were

assigned to prevalence. (Source Data: Figure 3-figure supplement 1-source data 1)

Figure 6-figure supplement 1. The population-adjusted estimated prevalence (median±95% BCI) in 2018 in four countries at administrative division of level one. (A) Cambodia, (B) Laos, (C) Thailand, (D) Vietnam. (Source Data: Figure 6-figure supplement 1-source data 1)

Figure 6-figure supplement 2. The population-adjusted estimated prevalence (median±95% BCI)

in 2013 in four countries at administrative division of level one. (A) Cambodia, (B) Laos, (C) Thailand,

(D) Vietnam. (Source Data: Figure 6-figure supplement 2-source data 1)

Figure 6-figure supplement 3. The population-adjusted estimated prevalence (median±95% BCI)

in 2008 in four countries at administrative division of level one. (A) Cambodia, (B) Laos, (C) Thailand,

(D) Vietnam. (Source Data: Figure 6-figure supplement 3-source data 1)

Figure 6-figure supplement 4. The population-adjusted estimated prevalence (median±95% BCI)

in 2003 in four countries at administrative division of level one. (A) Cambodia, (B) Laos, (C) Thailand,

(D) Vietnam. (Source Data: Figure 6-figure supplement 4-source data 1)

Figure 6-figure supplement 5. The population-adjusted estimated prevalence (median±95% BCI)

in 1998 in four countries at administrative division of level one. (A) Cambodia, (B) Laos, (C) Thailand,

(D) Vietnam. (Source Data: Figure 6-figure supplement 5-source data 1)

Figure 6-figure supplement 6. The population-adjusted estimated prevalence (median±95% BCI)

in 1993 in four countries at administrative division of level one. (A) Cambodia, (B) Laos, (C) Thailand,

(D) Vietnam. (Source Data: Figure 6-figure supplement 4-source data 1)

Figure 6-figure supplement 7. The population-adjusted estimated prevalence (median±95% BCI)

in 1988 in four countries at administrative division of level one. (A) Cambodia, (B) Laos, (C) Thailand,

(D) Vietnam. (Source Data: Figure 6-figure supplement 7-source data 1)

Figure 6-figure supplement 8. The population-adjusted estimated prevalence (median±95% BCI)

in 1983 in four countries at administrative division of level one. (A) Cambodia, (B) Laos, (C) Thailand,

(D) Vietnam. (Source Data: Figure 6-figure supplement 8-source data 1)

Figure 6-figure supplement 9. The population-adjusted estimated prevalence (median±95% BCI)

in 1978 in four countries at administrative division of level one. (A) Cambodia, (B) Laos, (C) Thailand,

(D) Vietnam. (Source Data: Figure 6-figure supplement 9-source data 1)

List of Source Data Files, for Figures

Figure 2-source data 1

Figure 2-source data 2

Figure 3-source data 1

Figure 4-source data 1

Figure 5-source data 1

Figure 6-source data 1

Figure 7-source data 1

Figure 2-figure supplement 1-source data 1

Figure 3-figure supplement 1-source data 1

Figure 6-figure supplement 1-source data 1

Figure 6-figure supplement 2-source data 1

Figure 6-figure supplement 3-source data 1

Figure 6-figure supplement 4-source data 1

Figure 6-figure supplement 5-source data 1

Figure 6-figure supplement 6-source data 1

Figure 6-figure supplement 7-source data 1

Figure 6-figure supplement 8-source data 1

Figure 6-figure supplement 9-source data 1

The above source data files for figures can be available from the corresponding author: Dr Ying-Si Lai,

laiys3@mail.sysu.edu.cn.

List of Supplementary Files

840   Supplementary File 1. PRISMA 2009 Checklist and GATHER checklist