Underlying causes of death with changing mortality among adults in the United States, 2013-2017

Xin Hu, MSPH[1,#]; Yong Lin, PhD[2,3]; Lanjing Zhang, MD[3,4,5,6]

Yale School of Public Health, New Haven, Connecticut; [2]Rutgers Cancer Institute of New Jersey, New Brunswick, New Jersey; [3]Department of Biostatistics, School of Public Health, Rutgers University, Piscataway, New Jersey; [4]Department of Pathology, Princeton Medical Center of Princeton, Plainsboro, New Jersey; [5]Department of Biological Sciences, Rutgers University, Newark, New Jersey; [6]Department of Chemical Biology, Ernest Mario School of Pharmacy, Rutgers University, Piscataway, New Jersey.
Correspondence author: Lanjing Zhang, lanjing.zhang@rutgers.edu, 609-853-6833
#Current address: Emory University School of Public Health, Atlanta, GA, USA

No conflict of interest declared by any of the authors.

Abstract

Overall mortality among U.S. adults was stable in the past years, while racial disparity was found in 10 leading causes of death or age-specific mortality in U.S. Blacks or African Americans. However, the trends in sex- and race-adjusted age-standardized cause-specific mortality are poorly understood. This study was aimed at identifying the UCD with sex- and race-adjusted, age-standardized mortality that was changing in recent years. We extracted the data of underlying causes of death (UCD) from the Multiple Cause of Death database of the Centers for Disease Control and Prevention (CDC). Multivariable log-linear regression models were used to estimate trends in sex- and race-adjusted, age-standardized mortality during 2013-2017. A total of 31,029,133 deaths were identified. Among the list of 113 UCD compiled by the CDC, there were 29 UCD with upward trend, 33 UCD with downward trend and 56 UCD with no significant trend. The 2 UCD with largest annual percent change were both nutrition related (annual percent change= 17.73, 95% CI [15.13-20.33] for malnutrition and annual percent change= 17.49, 95% CI [14.94-20.04] for Nutritional deficiencies), followed by Accidental poisoning and exposure to noxious substances. This study thus reported the UCD with changing mortality in recent years, which was sex- and race-adjusted and age-standardized. More efforts and resources should be focused on understanding, prevention and control of the mortality linked to these UCD. Continuous monitoring of mortality trends is recommended.

O verall mortality among U.S. adults was stable in the past years.[1, 2] Recent works documented the racial disparity in 10 leading causes of death or age-specific mortality in U.S. Blacks or African Americans.[3, 4] and disparities of race, age, and sex in the trends of suicide mortality.[5] However, the trends in sex- and race-adjusted age-standardized cause-specific mortality are largely unknown, despite its significance in public health, policy making and disease prevention. We therefore aimed to describe the UCD with sex- and race-adjusted, age-standardized mortality that was changing in recent years.

Methods

Multiple Cause of Death database of the Centers for Disease Control and Prevention (CDC) contains mortality and population counts for all U.S. counties.[6] Its data were extracted from death certificates for all eligible U.S. residents. The age-standardized mortality rates during 2013-2017 were estimated for adults (25+ years) in the U.S., using the U.S. standard population of year 2000 and the CDC Multiple Cause of Death database (1999-2017).[6] Underlying causes of death (UCD) were the single, underlying cause of death reported on each death certificate, and here classified using the list of 113 UCD compiled by the CDC.[1, 2] Age-standardized mortality rates by sex and race (White versus Non-white) and multivariable log-linear regression models were used for computing annual percent change (APC) of sex- and race-adjusted, age-standardized mortality rates, according to the CDC guidelines on using NHCS data.[7] The Stata software (version 15, StataCorp, College Station, TX) was employed for statistical analyses. An Institutional Review Board (IRB) approval is not required due to the use of de-identified and publicly available data on the deceased subjects. All P values were 2-sided and considered significant if <.05. The trends during 2015-2017 were also computed to assess potential influence of

implementation of the edition of International Classification of Diseases (ICD-10) in 2015.

Results
Among 31,027,401 deaths with un-suppressed mortality rate recorded during 2013-2017 (99.9% of the total deaths), there were 14,814,842 women (47.8%) and 26,499,625 Whites (85.4%) (**Table 1**). In the listed 113 UCD, 29 UCD had an upward trend in the sex- and race-adjusted age-standardized morality during 2013-2017 (**Table 2**), including in the ranking order of APC: Malnutrition/nutritional deficiencies, Accidental poisoning and exposure to noxious substances, Other nutritional deficiencies, Complications of medical and surgical care, Hypertensive heart and renal disease, Acute and rapidly progressive nephritic and nephrotic syndrome, Other and unspecified events of undetermined intent and their sequelae, Nontransport accidents, Alzheimer disease, Events of undetermined intent, Accidents (unintentional injuries), Parkinson disease, Alcoholic liver disease and others. In these 29 UCD, the 2 UCD with largest APC were both nutrition related, followed by Accidental poisoning and exposure to noxious substances. A total of 7 accident-related UCD, 4 heart-related UCD, 3 hypertension-related UCD and 2 liver-related UCD had increasing sex- and race-adjusted age-standardized mortality. There were also 33 UCD with downward trend in sex- and race-standardized mortality (**Table 3**), including mostly infection, malignancies and ischemic-heart diseases. There were 56 UCD with no significant trend, and 15 UCD with statistically-suppressed morality rate. The multivariable sensitivity analysis identified 16 UCD with upward mortality trend and 13 UCD with downward mortality trend during 2015-2017 (data not shown), which all had similar trends during 2013-2017.

Discussion
We here report the UCD with growing mortality and those with decreasing mortality among the adults in the U.S. during 2013-2017, which was sex- and race-adjusted and age-standardized. Sex- and race-adjustment reduces the biases associated with these disparity factors in mortality.[4, 5, 8] The 2 UCD with largest APC were both nutrition related (APC= 17.73 for Malnutrition and APC= 17.49 for Nutritional deficiencies), followed by Accidental poisoning and exposure to noxious substances. Future policy making and public health

resources perhaps should be focused on the identified UCD with growing mortality.

Despite the overall downward trends, cardiovascular diseases, heart diseases, and stroke mortality all had a decelerated downward trend substantially during 2011-2014.[9] It is therefore possible, but not certain, that some of these UCD might be linked to increasing mortality in recent years. Moreover, trend analyses on mortality were rightfully focused on leading causes of death.[3, 9, 10] Few studies systematically delineated recent trends of UCD with increasing mortality. It is probably most effective to decrease overall mortality by identifying and then reversing or neutralizing the increasing morality of certain UCD. Finally, age-standardized mortality had significantly different trends from those of crude mortality.[8] Given the variations of morality trends by race, sex and age and the lack of related data, this study is important for reporting understand the sex- and race-adjusted age-standardized mortality, and identifying the UCD linked to changing mortality. It thus fills the knowledge gap of recent trends in sex- and race-adjusted age-standardized mortality, and provides early data for policy making and public health focus area on projecting and reducing future mortality among adults in the U.S.

Malnutrition/nutrition deficiency was one of the top-10 leading UCD among pediatric populations of American Indians/Alaska natives or Asian/Pacific Islanders in 2013,[1] and Hispanic females aged 5-9 years in 2016.[2] Given its ~17.5% of APC, our data suggest malnutrition/nutrition deficiency will likely become a leading UCD in adults in the near future. The increasing mortality linked to UCD of accidents, heart, hypertension and liver are also concerning. Hypertension, for example, was among top-10 leading UCD in U.S. adults aged >85 years in 2016, but not in 2013.[1, 2] Future studies should continue monitoring UCD-specific mortality, and better understand and control these upward trends.

Several limitations of this study are noteworthy. First, the UCD on some death certificates may be misclassified or missed. For example, coders more likely include the UCD which the patient recently presented than the ones remotely presented. Second, there was a transition of ICD-9 to ICD-10 coding in 2015. Some cases may have different ICD-9 and ICD-10 classifications, but CDC has shown that ICD-9 and ICD-19 classifications are overall

consistent. In addition, the list of 113 UCD were unlikely influenced by such a transition because of its relatively broad definition of each UCD (none of them had decimals). Finally, there were missing/suppressed data in some UCD due to the small number of deaths linked to them, which were excluded in the analysis.

In summary, this study reports the UCD that are associated with upward or downward trend in sex- and race-adjusted, age-standardized U.S. mortality rate during 2013-2017. Some of them are reported recently and consistent with our findings.[10] These data will help better prioritize efforts on reducing mortality among U.S. adults.

Rererences

1. Heron MP. Deaths: leading causes for 2013. *National vital statistics reports.* 2016;65.
2. Heron MP. Deaths: Leading causes for 2016. *National vital statistics reports* 2018;67.
3. Cunningham TJ, Croft JB, Liu Y, Lu H, Eke PI, Giles WH. Vital Signs: Racial Disparities in Age-Specific Mortality Among Blacks or African Americans - United States, 1999-2015. *MMWR. Morbidity and mortality weekly report.* 2017;66:444-456.
4. Chang MH, Moonesinghe R, Athar HM, Truman BI. Trends in Disparity by Sex and Race/Ethnicity for the Leading Causes of Death in the United States-1999-2010. *Journal of public health management and practice : JPHMP.* 2016;22 Suppl 1:S13-24.
5. Ivey-Stephenson AZ, Crosby AE, Jack SPD, Haileyesus T, Kresnow-Sedacca MJ. Suicide Trends Among and Within Urbanization Levels by Sex, Race/Ethnicity, Age Group, and Mechanism of Death - United States, 2001-2015. *Morbidity and mortality weekly report. Surveillance summaries (Washington, D.C. : 2002).* 2017;66:1-16.
6. Centers for Disease Control and Prevention (CDC) NCfHSN. Multiple Cause of Death 1999-2017 on CDC WONDER Online Database, released 2018. Data are compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Vol 2019: U.S. HHS; 2018.
7. Ingram DD, Malec DJ, Makuc DM, et al. National Center for Health Statistics Guidelines for Analysis of Trends. *Vital Health Stat 2.* 2018:1-71.
8. Murphy SL, Xu J, Kochanek KD. Deaths: final data for 2010. *Natl Vital Stat Rep.* 2013;61:1-117.
9. Sidney S, Quesenberry CP, Jr., Jaffe MG, et al. Recent Trends in Cardiovascular Mortality in the United States and Public Health Goals. *JAMA cardiology.* 2016;1:594-599.
10. Tapper EB, Parikh ND. Mortality due to cirrhosis and liver cancer in the United States, 1999-2016: observational study. *BMJ (Clinical research ed.).* 2018;362:k2817.

Table 1. Baseline characteristics of deaths among adults in the U.S. by year, 2013-2017

	2013 (n=5,992,801)	2014 (n=6,049,415)	2015 (n=6,240,640)	2016 (n=6,299,895)	2017 (n=6,446,382)	Total (n=31,029,133)	*P*
Sex							<0.001
Female	2,891,245	2,900,958	2,990,172	2,988,697	3,044,725	14,815,797	
(%)	48.25	47.95	47.91	47.44	47.23	47.75	
Male	3,101,556	3,148,457	3,250,468	3,311,198	3,401,657	16,213,336	
(%)	51.75	52.05	52.09	52.56	52.77	52.25	
Race							<0.001
Non-White	849,131	867,071	903,395	938,445	970,678	4,528,720	
(%)	14.17	14.33	14.48	14.9	15.06	14.6	
White	5,143,670	5,182,344	5,337,245	5,361,450	5,475,704	26,500,413	
(%)	85.83	85.67	85.52	85.1	84.94	85.4	

Age, year							<0.001
25-34	45,463	47,177	51,517	57,616	60,215	261,988	
(%)	1.79	1.84	1.94	2.15	2.19	1.99	
35-44	69,573	70,996	73,088	77,792	79,796	371,245	
(%)	2.74	2.77	2.76	2.9	2.9	2.82	
45-54	177,724	175,917	174,494	173,516	170,142	871,793	
(%)	7.01	6.86	6.59	6.48	6.19	6.62	
55-64	338,127	348,808	357,785	366,445	372,006	1,783,171	
(%)	13.34	13.6	13.51	13.68	13.53	13.53	
65-74	454,429	471,541	495,016	512,080	531,610	2,464,676	
(%)	17.92	18.38	18.69	19.12	19.33	18.7	
75-84	625,013	624,504	637,566	636,916	657,759	3,181,758	
(%)	24.65	24.35	24.07	23.78	23.92	24.14	
85+	825,198	826,226	859,701	854,462	878,035	4,243,622	
(%)	32.55	32.21	32.45	31.9	31.93	32.2	

Table 2. The underlying causes of death with increasing sex- and race-adjusted age-standardized mortality among adults in the U.S., 2013-2017

Underlying cause of death[¶]	2013 mortality*	2017 mortality*	Death, n	APC (95% CI)	P[‡]
Malnutrition (E40-E46)	1.37	2.95	25806	17.73 (15.13 to 20.33)	<0.001
#Nutritional deficiencies (E40-E64)	1.43	3.06	26788	17.49 (14.94 to 20.04)	<0.001
Accidental poisoning and exposure to noxious substances (X40-X49)	17.28	28.54	230327	13.89 (10.94 to 16.84)	<0.001
#Complications of medical and surgical care (Y40-Y84,Y88)	1.18	1.75	15203	10.36 (6.36 to 14.35)	<0.001
Hypertensive heart and renal disease (I13)	1.75	2.86	26402	10.01 (6.98 to 13.04)	<0.001
Acute and rapidly progressive nephritic and nephrotic syndrome (N00–N01,N04)	0.16	0.23	2550	8.96 (5.03 to 12.89)	<0.001
Other and unspecified events of undetermined intent and their sequelae (Y10–Y21,Y25–Y34,Y87.2,Y89.9)	1.79	2.25	21213	7.63 (2.31 to 12.95)	0.011
Nontransport accidents (W00-X59,Y86)	39.14	52.04	501465	7.59 (5.44 to 9.74)	<0.001
#Alzheimer disease (G30)	36.38	48.01	526372	7.52 (6.13 to 8.91)	<0.001
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	1.91	2.34	22257	7.44 (2.25 to 12.64)	0.011
#Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53.08	67.47	660823	6.57 (4.54 to 8.59)	<0.001
#Parkinson disease (G20-G21)	11.26	13.01	140968	4.67 (3.92 to 5.41)	<0.001
Alcoholic liver disease (K70)	7.88	9.17	102510	3.97 (2.26 to 5.68)	<0.001

Cause of death (ICD-10)					
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4..	12.91	14.39	147771	3.93 (2.59 to 5.26)	<0.001
Transport accidents (V01-V99,Y85)	13.93	15.44	159358	3.79 (2.56 to 5.02)	<0.001
Heart failure (I50)	27.80	31.56	367490	3.54 (2.95 to 4.13)	<0.001
Hypertensive heart disease (I11)	15.86	18.78	208511	2.96 (1.53 to 4.39)	0.001
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60–X71,X75–X84,Y87.0)	8.16	9.15	92679	2.91 (1.35 to 4.47)	0.002
#Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16.89	18.27	190649	2.68 (1.66 to 3.70)	<0.001
Other and unspecified infectious and parasitic diseases and their sequalae (A00,A05,A20–A36,A42–A44,A48–A49,A54–A79,A81–A82,A85.0–A85.1,A85.8,A86–B04,B06–B09,B25–B49,B55–B99)	2.49	2.90	31630	2.66 (0.70 to 4.63)	0.015
Water, air and space, and other and unspecified transport accidents and their sequelae (V90–V99,Y85)	0.65	0.67	7376	2.55 (0.39 to 4.70)	0.032
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	3.97	4.28	50862	1.91 (1.42 to 2.39)	<0.001
Other complications of pregnancy, childbirth and the puerperium (O10–O99)	0.49	0.54	4881	1.87 (0.41 to 3.33)	0.033
#Pregnancy, childbirth and the puerperium (O00-O99)	0.49	0.54	4976	1.73 (0.24 to 3.23)	0.049
Other diseases of respiratory system (J00-J06,J30- J39,J67,J70-J98)	15.11	16.42	187819	1.71 (0.75 to 2.67)	0.003
Other and unspecified nontransport accidents and their sequelae (W20–W31,W35–W64,W75–W99,X10–X39,X50–X59,Y86)	6.57	6.85	80381	1.64 (0.67 to 2.62)	0.004
#Essential hypertension and hypertensive renal disease (I10,I12,I15)	13.17	13.92	161640	1.41 (0.24 to 2.57)	0.029
#Chronic liver disease and cirrhosis (K70,K73-K74)	15.68	16.80	197021	1.38 (0.37 to 2.38)	0.015
Other heart diseases (I26-I51)	84.05	87.94	1052221	0.91 (0.34 to 1.47)	0.005

Note: ¶ The list of 113 underlying causes of death compiled by the U.S. Centers for Disease Control and Prevention, with the 10th edition of the International classification of Diseases (ICD-10) in parentheses; *Age-standardized mortality per 100,000 in the U.S.; APC, annual percent change; CI, confidence intervals; ‡P for sex- and race-adjusted trend in the mortality which was age-standardized using the U.S. standard population of year 2000; # indicates a "rankable" cause of death, from the National Center for Health Statistics (NCHS) list of rankable causes of death. The rankable causes are a subset of the 113 selected causes of death.

Table 3. The underlying causes of death with decreasing sex- and race-adjusted age-standardized mortality among adults in the U.S., 2013-2017

Underlying cause of death [¶]	2013 mortality*	2017 mortality*	Death, n	APC	(95% CI)	P[‡]
#Viral hepatitis (B15-B19)	3.31	2.13	35696	-11.71	(-13.31 to -10.11)	<0.001
Bronchitis, chronic and unspecified (J40-J42)	0.26	0.20	2672	-8.26	(-13.57 to -2.95)	0.007
Emphysema (J43)	3.59	2.78	37036	-7.11	(-12.14 to -2.07)	0.012
#Human immunodeficiency virus (HIV) disease (B20-B24)	3.17	2.43	31524	-7.10	(-13.48 to -0.72)	0.042
#Atherosclerosis (I70)	2.83	2.17	30369	-6.55	(-7.75 to -5.34)	<0.001
#Enterocolitis due to Clostridium difficile (A04.7)	3.32	2.41	35022	-6.47	(-8.6 to -4.33)	<0.001
Hodgkin disease (C81)	0.48	0.39	5133	-5.69	(-9.28 to -2.09)	0.006
Certain other intestinal infections (A04,A07-A09)	4.49	3.41	47250	-5.55	(-7.38 to -3.72)	<0.001
Respiratory tuberculosis (A16)	0.16	0.14	1808	-5.27	(-10.13 to -0.4)	0.047
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	67.06	56.63	760393	-4.68	(-6.78 to -2.57)	<0.001
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00–R99)	14.68	11.74	149792	-4.13	(-6.39 to -1.88)	0.002
Pneumonia (J12-J18)	22.72	19.31	251383	-4.07	(-5.39 to -2.75)	<0.001
Acute myocardial infarction (I21-I22)	50.11	43.36	566631	-3.94	(-5.29 to -2.58)	<0.001
#Influenza and pneumonia (J09-J18)	24.25	21.81	273633	-3.27	(-4.8 to -1.74)	<0.001
Ischemic heart diseases (I20-I25)	158.67	143.56	1830094	-2.83	(-3.93 to -1.73)	<0.001
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	81.63	73.80	937255	-2.82	(-4.27 to -1.37)	0.001
Malignant neoplasm of stomach (C16)	4.91	4.44	56378	-2.58	(-3.48 to -1.68)	<0.001
Other forms of chronic ischemic heart disease (I20,I25)	106.84	98.48	1242985	-2.39	(-3.4 to -1.38)	<0.001
Leukemia (C91-C95)	10.02	9.14	112661	-2.25	(-3.41 to -1.1)	0.001
Malignant neoplasms of kidney and renal pelvis (C64-C65)	5.94	5.44	69758	-2.22	(-3.14 to -1.31)	<0.001
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81–C96)	24.60	22.79	282027	-2.2	(-3.25 to -1.15)	0.001
Non-Hodgkin lymphoma (C82-C85)	8.77	8.13	100796	-2.2	(-2.99 to -1.41)	<0.001
#Aortic aneurysm and dissection (I71)	4.31	4.00	49207	-2.14	(-3.29 to -0.98)	0.002
#Malignant neoplasms (C00-C97)	250.98	234.46	2955924	-2.01	(-2.4 to -1.63)	<0.001
Malignant neoplasms of colon, rectum and anus (C18-C21)	22.55	21.19	264027	-1.77	(-2.44 to -1.11)	<0.001
Malignant neoplasm of ovary (C56)	6.11	5.57	70677	-1.67	(-2.39 to -0.95)	0.001
Other diseases of circulatory system (I71-I78)	8.22	7.80	96630	-1.59	(-2.48 to -0.7)	0.002
All other diseases (Residual)	136.30	131.52	1599086	-1.30	(-2.07 to -0.54)	0.004

Cause of death	*	*	Deaths	APC	CI	P
Other chronic liver disease and cirrhosis (K73-K74)	7.80	7.68	94511	-1.26	(-2.39 to -0.13)	0.042
#Septicemia (A40-A41)	16.39	16.13	197318	-1.15	(-1.89 to -0.41)	0.007
Atherosclerotic cardiovascular disease, so described (I25.0)	25.26	24.65	305730	-1.12	(-1.83 to -0.42)	0.006
#Diseases of heart (I00-I09,I11,I13,I20-I51)	261.74	254.45	3133923	-1.01	(-1.56 to -0.45)	0.002
Major cardiovascular diseases (I00-I78)	341.71	336.23	4111472	-0.60	(-0.97 to -0.23)	0.005

Note: [¶] The list of 113 underlying causes of death compiled by the U.S. Centers for Disease Control and Prevention, with the 10th edition of the International classification of Diseases (ICD-10) in parentheses; *Age-standardized mortality per 100,000 in the U.S.; APC, annual percent change; CI, confidence intervals; [‡]P for sex- and race-adjusted trend in the mortality which was age-standardized using the U.S. standard population of year 2000; # indicates a "rankable" cause of death, from the National Center for Health Statistics (NCHS) list of rankable causes of death. The rankable causes are a subset of the 113 selected causes of death.