

1 **Title**

2 First SARS-CoV-2 detection in river water: implications in low sanitation countries

3

4 **Author names and affiliations**

5 Laura Guerrero-Latorre^{1,2}, Isabel Ballesteros¹, Irina M. Villacrés³, M. Genoveva

6 Granda³Byron P. Freire³ & Blanca Ríos-Touma^{1,*}

7

8 1. Grupo de Investigación en Biodiversidad, Medio Ambiente y Salud. Facultad de
9 Ingenierías y Ciencias Aplicadas. Universidad de Las Américas, Ecuador.

10 2. Department of Biology, Area of Microbiology, University of Girona, Spain.

11 3. Laboratorios de Investigación. Universidad de Las Américas, Ecuador.

12 *corresponding author: blanca.rios@udla.edu.ec

13

14 **Abstract**

15 Since the beginning of SARS-CoV-2 pandemic studies on viral shedding have reported
16 that this virus is excreted in feces in most patients. High viral loads are found at the
17 sewage pipeline or at the entrance of wastewater treatment plants from cities where
18 the number of COVID-19 cases are significant. In Quito (Ecuador) as in many other
19 cities worldwide, wastewater is directly discharged into natural waters. The aim of this
20 study was to evaluate SARS-CoV-2 presence in urban streams from a low sanitation
21 context. Three river locations along the urban rivers of Quito were sampled on the 5th
22 of June during a peak of COVID-19 cases. River samples were evaluated for water
23 quality parameters and afterwards, concentrated for viral analysis using skimmed milk
24 flocculation method. The viral concentrates were quantified for SARS-CoV-2 (N1 and
25 N2 genes) and Human Adenovirus as a human viral indicator. The results showed that
26 SARS-CoV-2 was detected for both target genes in all samples analyzed in a range of
27 2,91E+05 to 3,19E+06 GC/L for N1 and from 2,07E+05 to 2,22E+06 GC/L for N2. The

28 high values detected in natural waters from a low sanitation region have several
29 implications in health and ecology that should be further assessed.

30

31 **Keywords**

32 COVID-19; Andean Rivers; wastewater; viral dissemination; Ecuador; HadV

33

34 1. Introduction

35

36 Since the first detected case of unknown pneumonia in Wuhan (China), in half a year,
37 SARS-CoV-2 has caused almost 8 million confirmed cases and more than 400.000
38 deaths worldwide (WHO, 2020). Currently, Latin America is considered the epicenter of
39 the pandemic (UN, 2020). Brazil reported Latin America's first case in late February
40 2020, and since then, COVID-19 cases and deaths have increased widely throughout
41 nearly every country in the region. In Ecuador, official reports declare more than 45.000
42 cases and 3.800 deaths in the second week of June (SNGRE, 2020).

43 SARS-CoV-2 transmission in humans is mainly airborne (Zhang *et al.*, 2020) but recent
44 studies suggest that extended shedding of SARS-CoV-2 in feces is a potential health
45 risk (Lodder and Roda Husman, 2020). As most COVID-19 cases fecally excreted the
46 virus, SARS-CoV-2 has been detected in wastewater treatment plants from several
47 cities around the world and some countries have started to monitor the viral load in
48 sewage systems in an attempt to design early warnings for future outbreaks (Bivins *et*
49 *al.*, 2020).

50 However, the fecal shedding in wastewater could be especially relevant in low
51 sanitation countries where insufficient or nonexistent treatments of wastewater are
52 applied.

53 Studies analyzing SARS-CoV-2 in sewage have tested the efficiency after wastewater
54 treatments applied and in most cases effluent waters did not show the presence of
55 SARS-CoV-2 (Randazzo *et al.*, 2020a). Moreover, in Japan, Haramoto *et al.* 2020
56 looked for SARS-CoV-2 in rivers impacted by sewage, and did not detect its presence.

57 In most countries of Latin America an average of 30% of wastewaters are treated
58 before discharging in water bodies (Rodriguez *et al.*, 2020). In particular, Ecuador has
59 less than 20% of treatment coverage for its wastewaters (Rodriguez *et al.*, 2020) and
60 its capital city, Quito with almost 3 million inhabitants only treats 3% of its sewage
61 (EPMAPS, 2020). Therefore, the urban rivers of Quito, whose receive the fecal waters

62 from the whole city, are highly contaminated by human microorganisms and other
63 pollutants, disseminating them along the river basin and crossing from up 2800 masl to
64 the Pacific coast (Guerrero-Latorre et al., 2018; Voloshenko et al 2015). Previous
65 studies have described the viral diversity of human pathogens present in the urban
66 rivers of Quito and highlighting the health risk of populations that can use downstream
67 waters for irrigation or recreational purposes (Guerrero-Latorre et al 2018). In the
68 context of the COVID-19 pandemic, very little information has emerged from low
69 sanitation countries about the presence of SARS-Cov-2 in sewage and natural waters.
70 Therefore, this is the first study in a low sanitation country that attempted to analyze the
71 presence of SARS-CoV-2 in river waters impacted by urban wastewater.

72

73 **2. Material and methods**

74 **2.1 Sampling**

75

76 The 5th of June 2020 three locations from Quito's river were sampled. Sampling
77 locations are representative from south-center (M1), north-center (M2) and north (M3)
78 accumulated wastewater discharges into the natural streams passing through the city
79 (Figure 1, EPMAPS, 2011). At each location, 2 L samples were collected for viral
80 analysis and 2 L for other biological and chemical measurements: DBO₅, DQO, pH and
81 conductivity. Samples were preserved at 4 °C and taken to the laboratory in less than 3
82 hours.

83

84 **2.2 Water concentration and RNA extraction**

85

86 Collected water was concentrated for viral analysis using an adapted Skimmed Milk
87 Flocculation method using 2 L of water (Fernandez-Cassi *et al.*, 2018). Briefly, sample
88 was preconditioned at pH 3.5 and a conductivity above 1.5 mS/cm². Then, 10 ml of

89 pre-flocculated skimmed milk solution were added. After 8 h of stirring, flocks were
90 centrifuged at 8,000× g for 40 min, and the pellet was suspended in 10 mL of
91 phosphate buffer. Nucleic acids were extracted using AccuPrep® Universal RNA
92 Extraction Kit - Bioneer from 200 ul of the viral concentrates into 40 ul of eluted
93 solution.

94

95 **2.3 qRT-PCR analysis**

96 Nucleic acid extractions from viral concentrates were used to analyze by RT-qPCR
97 SARS-CoV-2 using N1 and N2 primer sets (CDC, 2020) and Human Adenovirus as a
98 microbial indicator of human fecal contamination (Allard *et al.*, 2001).

99 The volume of original water sample analyzed in each PCR reaction after concentration
100 and extraction steps is 3 µL for SARS-CoV-2 using TaqMan Fast Virus 1-Step Master
101 Mix (ThermoFisher) and 10 µL for HAdV using Environmental TaqMan MasterMix
102 (ThermoFisher). Quantitative PCRs were run in Bio-Rad CFX96 qPCR thermocycler.
103 The inhibition was analysed by testing the direct and 1/10 diluted extractions for each
104 qPCR or RT-qPCR.

105

106 **2.4 Cases per sampling point**

107

108 In order to relate SARS-CoV-2 viral load in river water samples with reported COVID-
109 19 cases in Quito, we gathered information from the sectors of Quito that discharge
110 sewage to the sampling points according to the Potable Water and Sanitation
111 Enterprise of Quito (EPMAPS, 2011). Since the city of Quito reports its cases by these
112 sectors we gather current and historical information from 14 days prior to the sampling
113 points (<https://twitter.com/MunicipioQuito/>). This information was validated with the
114 official government reports (SNGRE, 2020).

115

116 **3. Results**

117

118 Water quality parameters of samples analyzed show an important anthropogenic
119 impact in the rivers (Table 1). Sampling points M2 and M3 do not meet the national
120 requirement for aquatic life preservation for DQO and DBO₅ (Ministerio del Ambiente
121 de Ecuador, 2015). Human Adenovirus, a viral indicator used to evaluate the microbial
122 impact in river water present high quantities corresponding to strong impact (Rusiñol *et*
123 *al.* 2014)

124 SARS-CoV-2 genes (N1 and N2) were present in the three locations and show
125 concentrations ranging from 2,84E+05 to 3,19E+06 GC/L for N1 target gene and from
126 2,07E+05 to 2,23E+06 for N2 target gene (Table 2).

127 These values are clearly related to COVID-19 cases reported in the contributing areas
128 for each point, showing higher values where higher active cases were registered (Table
129 2). The collection of samples was during a peak of the outbreak, since the active cases
130 (considered the notified cases in the last 14 days prior to the sampling day) represent
131 25% of the total cases of COVID19 reported in the city since the beginning of the
132 outbreak.

133

134 **4. Discussion**

135

136 This is the first study that reports SARS-CoV2, and in important quantities, in rivers,
137 however other studies have attempted to find it without success (Haramoto *et al.*,
138 2020). The urban river of Quito is impacted by the direct discharge of sewage water
139 from a population of almost 3 million inhabitants. The water quality parameters
140 described in Table 1 indicate the high levels of organic and inorganic contamination
141 from those urban rivers. The water parameters measured in 2 of the 3 locations studied
142 were above the national regulations for aquatic life preservation (Ministerio del

143 Ambiente de Ecuador, 2015). However, all environmental values are higher than those
144 registered for reference sites in this same basin (Ríos-Touma *et al.*, 2014)

145

146 Viral human contamination analyzed by HAdV indicator shows the great impact of
147 human excreta in the urban rivers of Quito. Values are very similar from the samples
148 collected in the same locations in 2017 (Guerrero-Latorre *et al.*, 2018). In that study, a
149 metagenomic analysis was carried out in the same locations and sequences from 26
150 human viral pathogens were obtained, revealing the microbial consequences of
151 sewage discharge without previous treatment into natural streams.

152

153 Nowadays, in the peak of COVID-19 pandemic in Ecuador, levels of SARS-CoV-2
154 found in urban rivers of Quito are similar as found in sewage of Valencia (Spain) with
155 more than 5000 active cases (Randazzo *et al.* 2020b) and Paris in the epidemic peak
156 of cases with more than 10000 hospitalized cases (Wurtzel *et al.*, 2020). However, at
157 the sampling date in the urban area of Quito only 750 COVID-19 active cases were
158 notified, suggesting an important underdiagnose as seen in many other regions
159 worldwide.

160

161 The implications from our study can be extended to other cities where the sewage is
162 directly discharged into natural streams. Firstly, SARS-CoV-2 presence as other
163 microbial pathogens discharged in open river water might pose a risk of infection for
164 the population in contact with the water downstream. The risk has been recently
165 approached in a QMRA study of WWTP workers exposed to a different concentration
166 of SARS-CoV-2 in sewage (Zaneti *et al.*, 2020). However, it is important to remark that
167 only genomic material has been detected in waters and the viability of the virus is not
168 known for polluted waters. Secondly, the viral dissemination into the environment has
169 an unknown impact into livestock and wildlife health, as the spillover events of zoonotic
170 links are frequent in the coronaviridae family (Franklin and Bevins, 2020).

171 Finally, the presence of the virus can be also used as a surveillance tool for an early
172 warning system using main sewage discharges along the city helping to control the
173 pandemic where diagnostic tools are limited (Bivins *et al.*, 2020). Also, the method
174 implemented in the present publication can be used in other cities where sewage is not
175 possible to sample and wastewaters are discharged to streams or rivers.

176

177 **5. Conclusions**

178

179 We have detected for the first time important viral loads of SARS-CoV-2 from rivers in
180 urban streams of Quito. The loads found suggest that cases are probably a lot higher
181 than the official data. Also, implications in dissemination of SARS-CoV-2 in low
182 sanitation countries by polluted freshwaters should be further considered. The low
183 degree of wastewater treatment coverage in the region, might be a factor of increased
184 risk for COVID-19 pandemic.

185

186 **Acknowledgements**

187 Special thanks to Xavier Amigo and Nature Experience for its invaluable service during
188 the sampling campaign.

189

190 **Funding**

191 This work was supported by research funds from Universidad de Las Américas,
192 Ecuador.

193

194 **6. References**

195 Allard, A., Albinsson, B., Wadell, G., Albinsson, B.O., 2001. Rapid Typing of
196 Human Adenoviruses by a General PCR Combined with Restriction Endonuclease
197 Analysis Rapid Typing of Human Adenoviruses by a General PCR Combined with

- 198 Restriction Endonuclease Analysis. *J. Clin. Microbiol.* 39, 498–505.
- 199 <https://doi.org/10.1128/JCM.39.2.498>
- 200 Bivins, A., North, D., Ahmad, A., Ahmed, W., Alm, E., Been, F., et al, 2020.
- 201 Wastewater-Based Epidemiology: Global Collaborative to Maximize Contributions
- 202 in the Fight Against COVID-19. *Environ. Sci. Technol.*
- 203 doi:10.1021/acs.est.0c02388
- 204 CDC, 2020. 2019-Novel Coronavirus (2019-nCoV) Real-Time RT-PCR Diagnostic
- 205 Panel. US Centers for Disease Control and Prevention.
- 206 <https://www.cdc.gov/coronavirus/2019-ncov/lab/rt-pcr-panel-primer-probes.html>
- 207 [Accessed 3 Mar 2020].
- 208 Empresa Pública Metropolitana de Agua Potable y Saneamiento. (2011). Estudios
- 209 de Actualización del Plan Maestro Integrado de Agua Potable y Alcantarillado
- 210 para el DMQ. Recuperado el 4 de febrero del 2018 de
- 211 [https://www.aguaquito.gob.ec/sites/default/files/documentos/plan_ma](https://www.aguaquito.gob.ec/sites/default/files/documentos/plan_maestro_agua_potable.pdf)
- 212 [estro_agua_potable.pdf](https://www.aguaquito.gob.ec/sites/default/files/documentos/plan_maestro_agua_potable.pdf)
- 213 Empresa Pública Metropolitana de Agua Potable y Saneamiento. (2020).
- 214 Programa para la descontaminación de los ríos de Quito.
- 215 [https://www.aguaquito.gob.ec/programa-para-la-descontaminacion-de-los-rios-de-](https://www.aguaquito.gob.ec/programa-para-la-descontaminacion-de-los-rios-de-quito/)
- 216 [quito/](https://www.aguaquito.gob.ec/programa-para-la-descontaminacion-de-los-rios-de-quito/). [Accessed June 13th, 2020]
- 217 Fernandez-Cassi , X., Timoneda, N., Martínez-Puchol, S., Rusiñol, M., Rodriguez-
- 218 Manzano, J., Figuerola, N., *et al.* 2018. Metagenomics for the Study of Viruses in
- 219 Urban Sewage as a Tool for Public Health Surveillance. *Sci Total Environ.* Mar
- 220 15;618:870-880. <https://doi.org/10.1016/j.scitotenv.2017.08.249>

- 221 Franklin, A. B., & Bevins, S. N. (2020). Spillover of SARS-CoV-2 into novel wild
222 hosts in North America: A conceptual model for perpetuation of the pathogen. *The*
223 *Science of the total environment*, 733, 139358. Advance online publication.
224 <https://doi.org/10.1016/j.scitotenv.2020.139358>
- 225 Guerrero-Latorre, L., Romero, B., Bonifaz E., Timoneda, N., Rusiñol, M., Girones,
226 R. and Rios-Touma, B. 2018. Quito's virome: Metagenomic analysis of viral
227 diversity in urban streams of Ecuador's capital city. *Sci Total Environ*. 645, pp.
228 1334 - 1343; doi: <https://doi.org/10.1016/j.scitotenv.2018.07.213>
- 229 Haramoto, E., Malla, B., Thakali, O., Kitajima, M., 2020. First environmental
230 surveillance for the presence of SARS-CoV-2 RNA in wastewater and river water
231 in Japan. medRxiv 2020.06.04.20122747; doi:
232 <https://doi.org/10.1101/2020.06.04.20122747>
- 233 Lodder, W., de Roda Husman, A.M., 2020. SARS-CoV-2 in wastewater: potential
234 health risk, but also data source. *Lancet Gastroenterol. Hepatol*. 5, 533–534.
235 [https://doi.org/10.1016/S2468-1253\(20\)30087-X](https://doi.org/10.1016/S2468-1253(20)30087-X)
- 236 Ministerio del Ambiente de Ecuador. 2015. 097-A Refórmese el Texto Unificado
237 de Legislación Secundaria. Registro Oficial. Año III - No 387. Quito, miércoles 4 de
238 noviembre de 2015. P 6 a 78.
- 239 Randazzo, W., Truchado, P., Cuevas-Ferrando, E., Simon, P., Allende, A. and
240 Sanchez, G. 2020a. SARS-CoV-2 RNA in wastewater anticipated COVID-19
241 occurrence in a low prevalence area. *Water Research*. Volume 181, 15 August
242 2020, 115942. <https://doi.org/10.1016/j.watres.2020.115942>
- 243 Randazzo, W., Cuevas-Ferrando, E., Sanjuan, R., Domingo-Calap, P., Sanchez,
244 G., 2020b. Metropolitan Wastewater Analysis for COVID-19 Epidemiological

- 245 Surveillance. medRxiv 2020.04.23.20076679.
- 246 <https://doi.org/10.1101/2020.04.23.20076679>
- 247 Ríos-Touma, B., Acosta, R., Prat, N., 2014. The Andean biotic index (ABI):
248 Revised tolerance to pollution values for macroinvertebrate families and index
249 performance evaluation. *Rev. Biol. Trop.* 62, 249–273.
- 250 Rodriguez Hector, Alexander Delgado, Anna Nolasco, Daniel Saltiel, Gustavo, D.
251 J. S. 2020. From Waste to Resource. Water Papers. World Bank.
252 <https://doi.org/doi:10.1596/33436>
- 253 Rusiñol, M., Fernandez-Cassi, X., Hundesa, A., Vieira, C., Kern, A., Eriksson, I. *et*
254 *al.* (2014). Application of human and animal viral microbial source tracking tools in
255 fresh and marine waters from five different geographical areas. *Water Research.*
256 59, pp. 119-129. <https://doi.org/doi:10.1016/j.watres.2014.04.013>
- 257 SNGRE 2020. Informes de Situación e Infografías – COVID 19 – desde el 29 de
258 Febrero del 2020. [https://www.gestionderiesgos.gob.ec/informes-de-situacion-](https://www.gestionderiesgos.gob.ec/informes-de-situacion-covid-19-desde-el-13-de-marzo-del-2020/)
259 [covid-19-desde-el-13-de-marzo-del-2020/](https://www.gestionderiesgos.gob.ec/informes-de-situacion-covid-19-desde-el-13-de-marzo-del-2020/)
- 260 UN 2020. Central and South America now ‘intense zones’ for COVID-19
261 transmission. 1 June 2020 <https://news.un.org/en/story/2020/06/1065252>
- 262 Voloshenko-Rossin, A., Gasser, G., Cohen, K., Gun, J., Cumbal-Flores, L., Parra-
263 Morales, W., Sarabia, F., Ojeda, F., Lev, O., 2015. Emerging pollutants in the
264 Esmeraldas watershed in Ecuador: discharge and attenuation of emerging organic
265 pollutants along the San Pedro–Guayllabamba–Esmeraldas rivers. *Environ. Sci.*
266 *Process. Impacts* 17, 41–53. <https://doi.org/10.1039/C4EM00394B>
- 267 WHO. WHO Coronavirus Disease (COVID-19) Dashboard. 2020.
268 <https://covid19.who.int/>
- 269

270 Wurtzer, S., Marechal, V., Mouchel, JM., Maday, Y., Teysou, R., Richard, E., et
271 al. 2020. Evaluation of lockdown impact on SARS-CoV-2 dynamics through viral
272 genome quantification in Paris wastewaters. medRxiv 2020.04.12.20062679; doi:
273 <https://doi.org/10.1101/2020.04.12.20062679>

274 Zaneti,R.N., Girardi, V., Spilki, F.R., Mena, K., Campos-Westphalen, A.P., da
275 Costa Colares, W.R. et al. 2020. QMRA of SARS-CoV-2 for workers in
276 wastewater treatment plants. medRxiv 2020.05.28.20116277; doi:
277 <https://doi.org/10.1101/2020.05.28.20116277>

278 Zhang, R., Li, Y., Zhang, A.L., Wang, Y., Molina, M.J., 2020. Identifying airborne
279 transmission as the dominant route for the spread of COVID-19. Proc. Natl. Acad.
280 Sci. 202009637. doi:10.1073/pnas.2009637117


281

282

283

284

285 **Figure 1. Left.** Sampling locations in urban rivers from Quito (Guerrero-Latorre *et al.*
286 2018). M1 (0°14'00.27"S; 78°30'42.43"O), M2 (0°11'52.47"S;78°28'06.46"O), M3
287 (0°00'42.98"S; 78°26'25.86"O). **Right.** Picture of direct discharge of wastewater into
288 Machángara River (sampling point M1).



289

290

291 **Table 1.** Water quality parameters at sampling locations

Sampling points	pH	Conductivity	DBO₅ mg O₂/L	DQO mg O₂/L	HAdV GC/L
M1	7,73	653,07	13	35.706	1,13E+04
M2	8,16	703,2	28*	63.353*	2,18E+04
M3	7,83	696,48	16	43.941*	2,60E+05

292

293 *Parameters that do not comply with the Ecuadorian normative for aquatic life

294 preservation (Ministerio del Ambiente de Ecuador, 2015).

295

296

297 **Table 2.** Viral loads of SARS-CoV-2 per location and cumulative and active COVID-19

298 cases in the influence area.

299

Sampling points	SARS-CoV-2 N1 GC/L	SARS-CoV-2 N2 GC/L	Cumulative cases (total COVID-19 cases until 5th June)	Active cases (COVID-19 cases from the prior 14 days)
M1	3,19E+06	2,23E+06	2077	579
M2	2,84E+05	2,07E+05	580	81
M3	2,91E+06	8,55E+05	358	90

300

301