# Differential impact of antiretroviral therapy initiated before or during pregnancy on placenta pathology in HIV-positive women.

Nadia M IKUMI[1†], Thokozile R MALABA[2†], Komala PILLAY[3,4], Marta C COHEN[5,6], Hlengiwe P MADLALA[2], Mushi MATJILA[7], Dilly ANUMBA[8], Landon MYER[2], Marie-Louise NEWELL[9,10], and Clive M GRAY[1, 4*], for the PIMS Study Group

[1]Division of Immunology, Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Cape Town, South Africa
[2]Division of Epidemiology and Biostatistics, School of Public Health and Family Medicine, University of Cape Town, Cape Town, South Africa
[3]Division of Anatomical Pathology, Department of Pathology, University of Cape Town, Cape Town, South Africa
[4]National Health Laboratory Services, Groote Schuur Hospital, Cape Town, South Africa
[5]Department of Histopathology, Sheffield Children's NHS Foundation Trust , Sheffield, UK
[6]Department of Oncology and Metabolism, University of Sheffield, Sheffield, UK
Department of Obstetrics and Gynaecology, University of Cape Town, Cape Town South Africa
[7]Department of Obstetrics and Gynaecology, University of Cape Town, Cape Town South Africa
[8]Academic Unit of Reproductive and Developmental Medicine, University of Sheffield, Sheffield, United Kingdom
[9]School of Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, United Kingdom
[10]School of Public Health, Faculty of Health Sciences, University of Witwatersrand, Johannesburg, South Africa

**Conflict of interest statement**
The authors report no conflict of interest.

**Funding Support:** The research reported in this publication was supported by the Eunice Kennedy Shriver National Institute of Child Health & Human Development of the National Institutes of Health under Award Number R01 HD080385 (PIMS Prematurity Immunology in HIV-Infected Women and their Infants Study); also in part by the National Institute for Health Research (NIHR) (17/63/26) using UK Government aid to support global health research and by the Academy of Medical Sciences networking grant. The views expressed in this publication are those of the author(s) and not necessarily those of the NICH NIH, NIHR or the UK Department of Health and Social Care.

[†]Co-First Authors

**\*Corresponding Author**
Clive M Gray, Division of Immunology, Institute of Infectious Disease and Molecular Medicine, Falmouth Building, Faculty of Health Sciences, University of Cape Town, Anzio Road, Observatory, Cape Town, South Africa, 7925; Tel: +27 21406 6616; Email: clive.gray@uct.ac.za

**Short Title**
ART and placenta pathology

**Word count:** Body: 3220; Abstract: 239

53    **ABSTRACT**
54    **Objective:**
55    To examine the association between timing of ART initiation in HIV-infected women and
56    placental histopathology.
57
58    **Design:**
59    A nested sub-study in a larger cohort of HIV-infected women which examined the
60    association between ART status and birth outcomes.
61
**Methods:**
Placentas (n=130) were examined for histopathology from two ART groups: *stable* (n=53),
who initiated ART before conception and *initiating* (n=77), who started ART during
pregnancy (median [IQR] 15 weeks gestation [11-18]). Using binomial regression we
quantified associations between ART initiation timing with placental histopathology and
pregnancy outcomes.
68
**Results:**
One-third of all placentas were <10$^{th}$ percentile weight-for-gestation and there was no
significant difference between ART groups. Placental diameter, thickness, cord insertion
position and fetal-placental weight ratio were also similar by group. However, placentas
from the *stable* group showed increased maternal vascular malperfusion (MVM) (39.6% *vs*
19.4%), and decreased weight (392g vs 422g, p=0.09). MVM risk was twice as high (RR 2.03
[95% CI: 1.16 - 3.57]; p=0.01) in the stable group; the increased risk remaining significant
when adjusting for maternal age (RR 2.04 [95% CI: 1.12 - 3.72]; p=0.02). Furthermore, MVM
was significantly associated with pre-term delivery (PTD) and low birth weight (LBW)
(p=0.002 and p<0.0001, respectively).
79
80    **Conclusion:**
81    Preconception initiation of ART was associated with an increased MVM risk, and may
82    contribute to placental dysfunction. The association between MVM with PTD and LBW
83    suggests that a placenta-mediated mechanism likely links the putative association between
84    long-term use of ART and adverse birth outcomes.
85
86    **Keywords:** HIV, placenta pathology, timing of ART initiation, pre-term delivery, small for
87    gestational age
88
89

## Introduction

Antiretroviral treatment (ART) during pregnancy is not only of benefit for both the mother and infant, substantially decreasing MTCT, but is also proven to be most effective in reducing maternal and child mortality.[1,2] Like many other countries, South Africa has now fully adopted universal ART for life for all HIV-positive women upon diagnosis, including pregnant women.[3] However, the extensive rollout and uptake of ART from before or early pregnancy has led to a rapid rise in fetal exposure to ART; where such *in-utero* drug exposure has been associated with increased risk of adverse birth outcomes: including preterm delivery (PTD)[4–6], low birth weight (LBW)[7] and small-for-gestational age (SGA) infants[4,8].

HIV infection in pregnancy, even in the absence of ART, has been associated with an increased risk of adverse birth outcomes which is thought to be through inflammation-mediated disruption of placental development and function[9,10]. How maternal HIV infection impacts placental structure and function is thus important to consider as it can affect both maternal health, as seen in eclampsia[11], and fetal health, as seen with PTD and SGA[9,12–14]. Uncharacteristic placental shapes and sizes may indicate decreased placental efficiency, and maternal supply-line deprivation[15,16]. Histologic chorioamnionitis reflects maternal and fetal inflammatory responses to ascending infection[17,18], and is a major cause of spontaneous PTD[19–21]. Despite recognition of the importance of the placenta in adverse birth outcomes, there is scant data on the impact of *in-utero* ART exposure on placental development and pathology, and subsequent adverse birth outcomes.

Within a large cohort of HIV-positive women, we set out to systematically examine the histopathology of placentas collected at delivery from HIV-positive women initiating ART before or during pregnancy, to inform understanding of the impact of ART initiation timing on placental pathology. We conjectured that initiation of ART during pregnancy would result in greater inflammation-linked placental pathology than ART initiated before pregnancy.

## Methods

We performed a nested sub-study in the Prematurity Immunology in HIV-infected Mothers and their infants Study (PIMS) cohort in Cape Town, South Africa[22]. PIMS was a prospective cohort study investigating the association between ART use and adverse birth outcomes, in particular PTD, SGA and LBW[22]. HIV-positive pregnant women aged ≥18 years seeking antenatal care (ANC) services at the Gugulethu Midwife Obstetric Unit (MOU) were recruited between April 2015 and October 2016, at ≤24 weeks gestation. Whole placentas were collected at delivery and only those graded as good quality (grades 1-3)[23] were processed within 24 hours. Included were women who initiated ART before pregnancy (stable group), or during pregnancy (initiating group). ART regimens consisted predominately of tenofovir (TDF) + lamivudine (3TC)/emtricitabine (FTC) + efavirenz (EFV)[22].

*Ethical considerations*

The PIMS study protocol, informed consent forms and all data collection tools were approved by the University of Cape Town's, Faculty of Health Sciences Human Research

135   Ethics Committee (UCT FHS-HREC) (HREC 739/2014) and the University of Southampton
136   Faculty of Medicine Ethics Committee on (12542 PIMS). Written informed consent was
137   obtained from all study participants, including placenta collection for the nested cohort.
138

*Data collection*
At first ANC visit, gestational age (GA) was determined by ultrasound scan (USS) by a
research sonographer using a standardized assessment protocol. Obstetric outcomes,
including date of delivery and birthweight, were obtained from clinical records. Maternal
CD4 and viral load measurements were obtained from hospital records at the time closest
to the first ANC visit. Body Mass Index (BMI) was calculated from weight and height
measurements; blood pressure were measured at the first ANC visit (baseline) and during
follow-up.
147

*Placenta collection and processing*
Whole placentas were collected at delivery and fixed in 10% buffered formalin.  Placental
histopathological examination were performed in the Department of Pathology, Red Cross
Memorial Children's Hospital, Cape Town, South Africa. Specimens were examined from the
umbilical cord, membranes and placental disk and scored following the Amsterdam
Placental Workshop Group Consensus Statement[24]. Briefly, four blocks were prepared from
each placenta, one block included a roll of the extraplacental membranes and two cross
sections of the umbilical cord, one from the placental insertion, and one close to the fetal
end. In addition, three blocks each contained a full-thickness section of the placenta
parenchyma, one from a visibly abnormal area if present and two sections from
macroscopically normal tissue including one from the cord insertion. The tissue sections
were 3-5µm thick and were cut and stained with hematoxylin and eosin for histological
examination. Macroscopic and microscopic features were combined to report  placenta
pathology. For the sake of pathology scoring rigour, sections were also assesed in the
Department of Histopathology, Sheffield Children Hospital NHS Foundation Trust, Sheffield,
UK.
164

*Variables of interest*
Placenta pathology was based on placenta weight-for-gestational age, fetal-placental weight
ratio and features of placenta pathology. All placenta weights were reported as trimmed
weights, after removal of umbilical cords and membranes. Placenta weight percentiles by
gestational age and fetal-placenta ratios were categorized as small ($<10^{th}$), appropriate (10-
$90^{th}$) and large ($>90^{th}$) for gestational age.[25] Microscopically, the placenta was examined for
features of meconium exposure, based on the presence and amount of meconium-laden
macrophages in the placental membranes, with some cases showing features of prolonged
exposure characterized by meconium deep in the chorionic plate and umbilical cord;
amnion degeneration and chorion vascular muscle necrosis[26]. Histologic chorioamnionitis
was defined as the presence of acute inflammation in the placenta, with or without a fetal
inflammatory response[27,28]. Maternal vascular malperfusion (MVM) was defined as a
constellation of macroscopic and histologic features representing a pattern of injury in the
placenta as a result of maternal vascular abnormalities. The presence of decidual
vasculopathy was indicative of maternal vascular malperfusion. In the absence of decidual

vasculopathy, MVM was indicated by a small placenta for gestational age accompanied by
at least two villous changes (villous infarct, retroplacental haematoma, accelerated villous
maturation or distal villous hypoplasia). In the case of normal placenta weights for
gestational age, at least 3-4 villous features were noted to be placed in the category of
MVM.[24,29,30] Decidual arteriopathy was defined as abnormal or incomplete remodelling of
decidual spiral arterioles resulting in fibrinoid necrotic lesions[24,29,30]. Chronic deciduitis was
defined as abnormal infiltration of lymphocytes and plasma cells in the decidua[24,31].
Placenta disk dimensions were recorded as maximum linear dimension (*length*), greatest
dimension of the axis perpendicular to the *length* (*width*) and the thickness. Cord insertion
referred to site of insertion on the placenta in relation to the margin of the placenta
classified as central, off-centre and marginal based on the distance from the margin.
191
Secondary outcomes included PTD, SGA and LBW. PTD was defined as delivery at <37
weeks' GA[32]. Birthweight was categorized as low (LBW, <2500g), normal (NBW, 2500-
4000g) and high (HBW, >4000g)[33]. Size for gestational age, based on gender-specific
INTERGROWTH-21[st] project standards, was defined as small for gestational age (SGA) for
infants with birthweight <10[th] percentile for gestational age, appropriate for gestational age
(AGA) for those between 10-90[th] percentile and large for gestational age (LGA) for those
>90[th] percentiles[34].
199
*Statistical analysis*
Statistical analyses were performed using STATA version 12.0 (Stata Corporation, College
Station, Texas, USA) and R. Analyses focused on placenta pathology outcomes by ART
initiation before or during pregnancy. The impact of ART timing on features of placental
pathology was summarized using an Alluvial plot generated using the *ggalluvial* package in
R[35]. Secondary analyses focused on birth outcomes (PTD, LBW and SGA) related to features
of placenta pathology. In bivariable analyses, proportions were compared using chi-
squared/Fishers exact test and rank-sum tests. Associations between baseline
characteristics and placenta pathology outcomes (small placental weight for GA, MVM and
chorioamnionitis) were assessed using binomial regression. Results were presented as risk
ratios (RR) with 95% confidence intervals (CI) generated using the forestplot package in R (R
package version 1.9)[36]. For secondary analyses of birth outcomes, given the limited sample
size, only bivariate associations were assessed. As placenta weight and MVM abnormalities
were likely to originate in early gestation, we explored associations between timing of ART
initiation and the placenta pathologies. For MVM, as advanced maternal age has been
shown to increase the risk of placental dysfunction,[37] we applied a multivariable regression
model to adjust for maternal age differences recorded at booking between ART groups. To
assess potential risk factors for PTD, SGA and LBW we compared differences between
groups using the chi$^2$ test or Fisher's exact test. Missing values were excluded from these
analyses.

225 **Results**
226 A total of 130 placentas were examined: 53 (40.8%) from the stable group and 77 (59.2%)
227 from the initiating group of women. Maternal and obstetric characteristics are summarised
228 in Table 1.
229
*Maternal characteristics*
Women in the stable group were older (median 33 years, IQR 28-37) than initiating women
(median 28, IQR 25-32). Median GA at first ANC visit was 15 weeks (IQR 11-18), with no
differences by ART initiation timing group (Table 1). Overall, 119 (91.5%), women were on a
fixed dose regimen of Tenofovir (TDF) + Lamivudine (3TC) + Efavirenz (EFV), 2 (1.5%) were
on Zidovudine (AZT) + Lamividune (3TC) + Nevirapine (NVP), 6 (4.6%) were on AZT +
Emtricitabine or Lamivudine (XTC) + EFV with no difference between the groups; only 3
(2.3%; all in the *stable* group) were on a protease inhibitor (PI)-based regimen. Stable
women had significantly higher median CD4 T cell counts at enrolment (455 cells/ml, IQR
381 - 602) than initiating women (369.5 cells/ml, IQR 251 - 534.5). Most women (78.4%)
were overweight or obese, not significantly different by ART group. Initiating women were
more likely to be pre-hypertensive than stable women (28.6% vs 11.3%). Alcohol was the
most used substance, with 22% of women reporting alcohol use < 30 days prior to first ANC
visit (Supp. Table 1). There was no difference in smoking or recreational drug use between
the two groups.
245
*Infant characteristics*
Of the 130 live singleton births, 58 (44.6%) were female and 72 (55.3%) male (Table 2), with
no sex difference by ART timing group. Overall, median gestation at delivery was 39 weeks
(IQR 38 - 40), and PTD incidence was low (6.9%) with no difference by timing of ART
initiation. While median birthweight was also similar between ART groups, there was a
tendency towards differences in the percentage of LBW (11.3% vs 6.5%) and HBW (1.9% vs
9.1%) deliveries in the stable and initiating group respectively; but not reaching statistical
significance. The tendency towards increased >4000g macrosomia births in the ART
initiation during pregnancy group was not associated with birth complications. Differences
were also observed in size for gestational age, where 12 (9.2%) infants were SGA overall,
with 8 (15.1%) in the stable and 4 (5.2%) in the initiating group.
257
*Placental Pathology.*
Table 3 shows placenta characteristics and pathology stratified by timing of ART initiation,
and Figures 1A-D show representative images of some of the features of placenta pathology
included in our analysis: chorioamnionitis (Figure 1A), neutrophilic infiltrate involving the
cord vein representing a fetal response (Figure 1B) and decidual artery atherosis (Figure 1C).
Median placenta weight was 412g (IQR 350 - 480) overall, 393g in the stable and 422g in the
initiating group (p=0.09). Overall, 43 (33%) placentas were <10$^{th}$ percentile weight-for-
gestation, with no significant differences between ART groups; placental diameter, thickness
or cord insertion were also similar in each group as was the Fetal-Placental weight ratio.
Meconium exposure was identified in 22 (19.2%) cases, similar in both ART groups.
Chorioamnionitis was present in 19 (14.6%) cases, with a higher percentage in initiating
than in stable women, but not reaching statistical significance. MVM was significantly higher

in placentas from stable than initiating women (39.2% vs 19.5%; p=0.01). There were no
cases of chronic basal villitis, chronic chorioamnionitis, eosinophilic/T-cell vasculitis or
chronic histiocytic intervillositis. Chronic decidual perivasculitis was included in cases of
chronic deciduitis.

To discount selection bias due to collection of placentas from a sub-group of women, we
compared maternal characteristics in "no-placenta" versus "placenta" collection for each
ART group (Supp Table 2) in the larger PIMS cohort. Importantly, blood pressure was no
different in the wider cohort (n=421), irrespective of ART group, indicating that the
observed MVM in the smaller placenta stable group was not due to any bias in a pre- or
hypertensive state, which could have impacted on our observations.

Figures 2A-C shows the relative risk of placenta pathology by ART group and maternal
characteristics. In unadjusted models, women who were overweight (RR 0.46 [95% CI: 0.27-
081]; p=0.007) and obese (RR 0.37 [95% CI: 0.21 - 0.64]; p<0.0001) were less likely to have
small-for-gestational age placentas (Figure 2A). None of the variables explored were
associated with significant risk of chorioamionitis (Figure 2B). The risk of MVM (Figure 2C)
was twice as high (RR 2.03 [95% CI: 1.16 - 3.57]; p=0.01) in stable than in initiating women
and remained significant when adjusting for maternal age (RR 2.04 [95% CI: 1.12 - 3.72];
p=0.02). The impact of ART timing on placental pathology is summarized in the Alluvial plot
in Figure 2D, emphasizing the occurrence of MVM in women initiating ART pre-conception.
The association between placenta pathology and PTD, SGA and LBW (Table 4) was
significant for MVM (PTD: p=0.002; LBW: p<0.0001), and borderline for SGA (p=0.09). In
addition, SGA deliveries were more likely to present with meconium exposure, indicating
fetal distress (p=0.01).

**Discussion**
We examined the association between timing of ART initiation in HIV-positive women and
placental pathology, and with subsequent birth outcomes. Placentas from women who had
initiated ART pre-conception, but not those initiating during pregnancy, showed significantly
increased maternal vascular malperfusion (MVM). There was no evidence for overt
maternal or fetal inflammation in either ART group, contrary to our initial supposition. As
MVM was significantly associated with PTD and LBW, our findings suggest that associations
between ART and adverse birth outcomes may be placenta-mediated, especially so for
women initiating ART before pregnancy, where there is antiretroviral drug exposure during
placental implantation and development.

Previous studies of HIV-positive women initiating ART before or during pregnancy have
focused on quantifying risk of PTD[1,4,7,38–41], SGA[4,7,41,42], LBW[7,38,39], stillbirth[43] and/or
neonatal death[42]; virologic failure[44]; MTCT[45] maternal plasma angiogenic factors[46], and
maternal virologic and immunologic responses[47]. This is the first study to investigate
placenta pathology in association with timing of ART initiation. In our study, women who
initiated ART before pregnancy were significantly older, which could have accounted for the
observed MVM. However, when we controlled for maternal age, MVM remained

significantly associated with pre-conception ART initiation. Thus, ART initiation pre-
conception is a substantial risk factor for MVM.

MVM represents a pattern of placental injury which affects the maternal vasculature and
circulation and often associates with significantly increased risk of adverse effects to the
fetus[24,29]. We found that the high incidence of MVM in our study was consistent with
findings from a previous HIV-positive pregnancy cohort[48], albeit MVM in our study
associated with PTD and LBW. We posit that prolonged ART, initiated pre-conception, is
likely resulting in perturbed placental development and poor perfusion and function giving
rise to adverse birth outcomes. Interestingly, there was a non-significantly higher proportion
of macrosomia births in the women initiating ART during pregnancy, perhaps reflecting the
higher proportions of mothers with a high BMI in this group. Although macrosmia has been
associated with shoulder dystocia, neonatal and maternal trauma,[49] hypogycaemia,[50] this
was not observed in our larger PIMS cohort when we stratified the groups by time of ART
initiation (before versus during pregnancy).[51]

Although HIV-positive individuals on ART have been shown to have a high prevalence of
hypertension[52], women initiating ART prior to pregnancy in our study were less likely to be
pre- or hypertensive than those initiating during pregnancy. Additionally, we did not find
any associations between MVM and hypertensive disorders in our cohort, although
hypertensive disorders of pregnancy, including preeclampsia and chronic hypertension, are
the classical maternal clinical features associated with an increased risk of MVM[53–55]. Our
data suggests that placental research should form an integral part when exploring
interventions aimed at mitigating adverse birth outcome associated with ART use in
pregnancy.

The impact of ART exposure on the placenta is under-explored and we speculate that
exposure of ART before or during early pregnancy results in systemic maternal inflammation
that may disrupt vasculogenesis and angiogenesis[9]. In our cohort, the underlying
mechanism leading to this disruption may be dysregulation of pro-angiogenic mediators
including placental growth factor (PIGF), soluble endoglin (sEng) and angiopoietin-1 (Ang-1).
Although data is scarce, previous findings demonstrate that maternal HIV infection and ART
exposure is associated with an altered pro-angiogenic state; including lower PIGF, higher
sEng, decreased Ang-2 and increased Ang-1[10,56–58]. Based on previous evidence linking ART
with dysregulated angiogenesis and compromised placental vascular development in early
pregnancy, we postulate that the higher incidence of MVM in women who initiate ART
before pregnancy was likely due to ART-associated shifts towards an anti-angiogenic state
[59]. This may account for the increased MVM in women intiating ART pre-conception;
suggestive that ART exposure could have detrimental effects on placental development
characterized by altered uterine and intervillous blood flow dynamics.

Of potential importance for the interpretation of our findings was that over three quarters
of the women were either overweight or obese at first antenatal visit; increased BMI at that
time was associated with a decreased likelihood of a small placenta at delivery. We did not
have reliable assessments of BMI pre-pregnancy, but in other studies[53,54] elevated pre-

pregnancy BMI has been associated with large for gestational age placentas and altered
placental metabolic function. We recently showed from our larger PIMS cohort, maternal
obesity can increase the risk of having high birthweight and large for gestational age infants
and that incremental gestational weight gain increased the risk of having a spontaneous
preterm delivery.[55] Beyond the effect of high BMI during pregnancy on the placenta, it is
also associated with childhood obesity and cardiovascular diseases in later life.[56,57]
Although our study was limited in this regard, interventions targeting the risks associated
with the high rate of obesity among women of child-bearing age or during gestation should
not be ignored
368
The main strength of our study is the cohort design allowing us to compare two groups of
pregnant women receiving the same NNRTI-based ART regimen and enrolled early in
gestation, with data collected and samples investigated according to a strict protocol.
Additional strength lies in the ultrasound-based gestational age assessment, leading to
accurate PTD determination. Limitations include the non-random nature of the selection of
the nested cohort, giving rise to potential differences in maternal characteristics between
ART groups or placentas collected. However, we doubt this would affect our findings as: a)
there was no significant difference in maternal characteristics when we compared the
placenta sub-group with the overall cohort and b) the rates of stillbirths and miscarriages
were less than 5% in the larger PIMS cohort, with no differences in pregnancy loss when
groups were stratified by time of ART initiation. All staff in the three clinics where women
delivered were trained in placenta collection (with no exclusion criteria), and placentas were
transported to the laboratory for processing immediately.
382
In conclusion, we show women receiving ART pre-conception were at increased risk of
MVM, suggestive of placental dysfunction. In turn, the association between MVM with PTD
and LBW suggests that a placenta-mediated mechanism links the putative association
between ART and adverse birth outcomes. Our results further intimate that monitoring
placental function in women who initiated ART prior to conception is warranted and that
adjunctive therapies that target placental insufficiency in these women can alleviate the
burden of adverse birth outcomes.
390

391 **ACKNOWLEDGEMENTS**
392 We thank all the study participants in this study; and all the members of the PIMS study. We
393 also wish to thank Sue Ford, Colin Newel for data management and Michael Zulu for
394 specimen processing.
395

**Author contributions**
Clive M Gray, Marie-Louise Newell, Thokozile R Malaba,  Landon Myer: Conceptualization
and design of the study
Thokozile R Malaba, Hlengiwe P Madlala: study conduct, data collection and specimen
collection
Nadia M Ikumi: Laboratory and sample preparation
Komala Pillay, Marta Cohen: Histopathology scoring and interpretation
Nadia M Ikumi, Thokozile R Malaba: Statistical analysis
Nadia M Ikumi, Thokozile R Malaba, Clive Gray, Hlengiwe P Madlala, Marie-Louise Newell,
Mushi Matjila, Dilly Anumba, Landon Myer: Writing the manuscript, review and editing
406
407

**REFERENCES**

1.  Aniji CD, Towobola OA, Hoque ME, Mashamba TJ, Monokoane S. Impact of antiretroviral therapy on pregnancy outcomes. South Afr J HIV Med [Internet]. 2013 Nov 22;14(4):176. Available from: http://www.sajhivmed.org.za/index.php/sajhivmed/article/view/834

2.  Fowler MG, Qin M, Fiscus SA, Currier JS, Flynn PM, Chipato T, et al. Benefits and Risks of Antiretroviral Therapy for Perinatal HIV Prevention. N Engl J Med [Internet]. 2016;375(18):1726–37. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27806243

3.  World Health Organization. Programmatic Update Use of Antiretroviral Drugs for Treating Pregnant Women and Preventing HIV Infection in Infants Executive Summary. Vol. 45, World Health Organization. Geneva; 2012.

4.  Chen JY, Ribaudo HJ, Souda S, Parekh N, Ogwu A, Lockman S, et al. Highly active antiretroviral therapy and adverse birth outcomes among HIV-infected women in Botswana. J Infect Dis [Internet]. 2012 Dec 1;206(11):1695–705. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23066160

5.  Short C-ES, Douglas M, Smith JH, Taylor GP. Preterm delivery risk in women initiating antiretroviral therapy to prevent HIV mother-to-child transmission. HIV Med [Internet]. 2014 Apr;15(4):233–8. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24025074

6.  Thorne C, Patel D, Newell M-L. Increased risk of adverse pregnancy outcomes in HIV-infected women treated with highly active antiretroviral therapy in Europe. AIDS [Internet]. 2004 Nov;18(17):2337–9. Available from: http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00002030-200411190-00019

7.  Li N, Sando MM, Spiegelman D, Hertzmark E, Liu E, Sando D, et al. Antiretroviral Therapy in Relation to Birth Outcomes among HIV-infected Women: A Cohort Study. J Infect Dis [Internet]. 2016 Apr 1;213(7):1057–64. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26265780

8.  Zash R, Souda S, Chen JY, Binda K, Dryden-Peterson S, Lockman S, et al. Reassuring Birth Outcomes With Tenofovir/Emtricitabine/Efavirenz Used for Prevention of Mother-to-Child Transmission of HIV in Botswana. J Acquir Immune Defic Syndr [Internet]. 2016 Apr 1;71(4):428–36. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26379069

9.  Weckman AM, Ngai M, Wright J, McDonald CR, Kain KC. The Impact of Infection in Pregnancy on Placental Vascular Development and Adverse Birth Outcomes. Front Microbiol [Internet]. 2019 Aug 22;10. Available from: https://www.frontiersin.org/article/10.3389/fmicb.2019.01924/full

10. Conroy AL, McDonald CR, Gamble JL, Olwoch P, Natureeba P, Cohan D, et al. Altered angiogenesis as a common mechanism underlying preterm birth, small for gestational age, and stillbirth in women living with HIV. Am J Obstet Gynecol [Internet]. 2017 Dec;217(6):684.e1-684.e17. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0002937817311857

11. Fisher SJ. Why is placentation abnormal in preeclampsia? Am J Obstet Gynecol [Internet]. 2015 Oct;213(4):S115–22. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0002937815009126

12. Romero R, Kusanovic JP, Chaiworapongsa T, Hassan SS. Placental bed disorders in preterm labor, preterm PROM, spontaneous abortion and abruptio placentae. Best Pract Res Clin Obstet Gynaecol [Internet]. 2011 Jun;25(3):313–27. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21388889

13. Doyle RM, Alber DG, Jones HE, Harris K, Fitzgerald F, Peebles D, et al. Term and preterm labour are associated with distinct microbial community structures in

placental membranes which are independent of mode of delivery. Placenta
[Internet]. 2014 Dec;35(12):1099–101. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0143400414008194
14. Morgan T. Role of the Placenta in Preterm Birth: A Review. Am J Perinatol [Internet].
2016 Jan 5;33(03):258–66. Available from: http://www.thieme-
connect.de/DOI/DOI?10.1055/s-0035-1570379
15. Salafia CM, Yampolsky M, Misra DP, Shlakhter O, Haas D, Eucker B, et al. Placental
surface shape, function, and effects of maternal and fetal vascular pathology.
Placenta [Internet]. 2010 Nov;31(11):958–62. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0143400410003395
16. Freedman AA, Hogue CJ, Marsit CJ, Rajakumar A, Smith AK, Goldenberg RL, et al.
Associations Between the Features of Gross Placental Morphology and Birthweight.
Pediatr Dev Pathol [Internet]. 2019 May 16;22(3):194–204. Available from:
http://journals.sagepub.com/doi/10.1177/1093526618789310
17. Redline RW. Inflammatory response in acute chorioamnionitis. Semin Fetal Neonatal
Med [Internet]. 2012 Feb;17(1):20–5. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S1744165X11000916
18. Ategeka J, Wasswa R, Olwoch P, Kakuru A, Natureeba P, Muehlenbachs A, et al. The
prevalence of histologic acute chorioamnionitis among HIV infected pregnant women
in Uganda and its association with adverse birth outcomes. Robboy SJ, editor. PLoS
One [Internet]. 2019 Apr 11;14(4):e0215058. Available from:
http://dx.plos.org/10.1371/journal.pone.0215058
19. Galinsky R, Polglase GR, Hooper SB, Black MJ, Moss TJM. The Consequences of
Chorioamnionitis: Preterm Birth and Effects on Development. J Pregnancy [Internet].
2013;2013:1–11. Available from: http://www.hindawi.com/journals/jp/2013/412831/
20. Prince AL, Ma J, Kannan PS, Alvarez M, Gisslen T, Harris RA, et al. The placental
membrane microbiome is altered among subjects with spontaneous preterm birth
with and without chorioamnionitis. Am J Obstet Gynecol [Internet]. 2016
May;214(5):627.e1-627.e16. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0002937816003057
21. Wortham JM, Hansen NI, Schrag SJ, Hale E, Van Meurs K, Sánchez PJ, et al.
Chorioamnionitis and Culture-Confirmed, Early-Onset Neonatal Infections. Pediatrics
[Internet]. 2016 Jan;137(1):e20152323. Available from:
http://pediatrics.aappublications.org/lookup/doi/10.1542/peds.2015-2323
22. Malaba TR, Gray CM, Myer L, Newell M-L. Cohort Profile: Prematurity Immunology in
HIV-infected Mothers and their infants Study (PIMS). medRixv [Internet].
2020;2020.03.18.20033654. Available from:
http://medrxiv.org/content/early/2020/03/23/2020.03.18.20033654.abstract
23. Malaba TR, Gray CM, Myer L, Newell M-L. Cohort Profile: Prematurity Immunology in
HIV-infected Mothers and their infants Study (PIMS). medRxiv [Internet]. 2020 Jan
1;2020.03.18.20033654. Available from:
http://medrxiv.org/content/early/2020/03/23/2020.03.18.20033654.abstract
24. Khong TY, Mooney EE, Ariel I, Balmus NCM, Boyd TK, Brundler M-A, et al. Sampling
and Definitions of Placental Lesions: Amsterdam Placental Workshop Group
Consensus Statement. Arch Pathol Lab Med [Internet]. 2016 Jul;140(7):698–713.
Available from: http://www.archivesofpathology.org/doi/10.5858/arpa.2015-0225-
CC
25. Kraus FT. Placenta Pathology. American Registry of Pathology, editor. American
Registry of Pathology; 2004.
26. Kim B, Oh S, Kim J-S. Placental Lesions in Meconium Aspiration Syndrome. J Pathol
Transl Med [Internet]. 2017 Sep 15;51(5):488–98. Available from:
http://jpatholtm.org/journal/view.php?doi=10.4132/jptm.2017.07.20

27. Kallapur S, Presicce P, Rueda C, Jobe A, Chougnet C. Fetal Immune Response to Chorioamnionitis. Semin Reprod Med [Internet]. 2014 Jan 3;32(01):056–67. Available from: http://www.thieme-connect.de/DOI/DOI?10.1055/s-0033-1361823

28. Kim CJ, Romero R, Chaemsaithong P, Chaiyasit N, Yoon BH, Kim YM. Acute chorioamnionitis and funisitis: definition, pathologic features, and clinical significance. Am J Obstet Gynecol [Internet]. 2015 Oct;213(4):S29–52. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0002937815009102

29. Ernst LM. Maternal vascular malperfusion of the placental bed. APMIS [Internet]. 2018 Jul;126(7):551–60. Available from: http://doi.wiley.com/10.1111/apm.12833

30. Heider A. Fetal Vascular Malperfusion. Arch Pathol Lab Med [Internet]. 2017 Nov;141(11):1484–9. Available from: http://www.archivesofpathology.org/doi/10.5858/arpa.2017-0212-RA

31. Ziadie M. Chronic deciduitis [Internet]. PathologyOutlines.com website. [cited 2020 Mar 30]. Available from: https://www.pathologyoutlines.com/topic/placentachronicdecid.html

32. World Health Organization. Preterm Birth [Internet]. [cited 2019 Mar 30]. Available from: https://www.who.int/news-room/fact-sheets/detail/preterm-birth

33. Ota E, Haruna M, Suzuki M, Anh DD, Tho LH, Tam NTT, et al. Maternal body mass index and gestational weight gain and their association with perinatal outcomes in Viet Nam. Bull World Health Organ [Internet]. 2011 Feb 1;89(2):127–36. Available from: http://www.who.int/bulletin/volumes/89/2/10-077982.pdf

34. Papageorghiou AT, Kennedy SH, Salomon LJ, Altman DG, Ohuma EO, Stones W, et al. The INTERGROWTH-21 st fetal growth standards: toward the global integration of pregnancy and pediatric care. Am J Obstet Gynecol [Internet]. 2018 Feb;218(2):S630–40. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0002937818300127

35. Brunson JC. ggalluvial: Alluvial Plots in "ggplot2" [Internet]. 2020. Available from: https://cran.r-project.org/package=ggalluvial

36. Gordon M, Lumley T. forestplot: Advanced Forest Plot Using "grid" Graphics. 2019.

37. Lean SC, Heazell AEP, Dilworth MR, Mills TA, Jones RL. Placental Dysfunction Underlies Increased Risk of Fetal Growth Restriction and Stillbirth in Advanced Maternal Age Women. Sci Rep [Internet]. 2017;7(1):9677. Available from: http://www.ncbi.nlm.nih.gov/pubmed/28852057

38. Machado ES, Hofer CB, Costa TT, Nogueira SA, Oliveira RH, Abreu TF, et al. Pregnancy outcome in women infected with HIV-1 receiving combination antiretroviral therapy before versus after conception. Sex Transm Infect [Internet]. 2009 Apr;85(2):82–7. Available from: http://www.ncbi.nlm.nih.gov/pubmed/18987014

39. Chetty T, Thorne C, Coutsoudis A. Preterm delivery and small-for-gestation outcomes in HIV-infected pregnant women on antiretroviral therapy in rural South Africa: Results from a cohort study, 2010-2015. PLoS One [Internet]. 2018;13(2):e0192805. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29470508

40. Stoner MCD, Cole SR, Price J, Winston J, Stringer JSA. Timing of Initiation of Antiretroviral Therapy and Risk of Preterm Birth in Studies of HIV-infected Pregnant Women: The Role of Selection Bias. Epidemiology [Internet]. 2018;29(2):224–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29045283

41. Malaba TR, Phillips T, Le Roux S, Brittain K, Zerbe A, Petro G, et al. Antiretroviral therapy use during pregnancy and adverse birth outcomes in South African women. Int J Epidemiol [Internet]. 2017 Oct 1;46(5):1678–89. Available from: https://academic.oup.com/ije/article/46/5/1678/4055952

42. Santosa WB, Staines-Urias E, Tshivuila-Matala COO, Norris SA, Hemelaar J. Perinatal outcomes associated with maternal HIV and antiretroviral therapy in pregnancies with accurate gestational age in South Africa. AIDS [Internet]. 2019 Aug;33(10):1623–33. Available from: http://insights.ovid.com/crossref?an=00002030-201908010-

00010

43.   Msukwa MT, Keiser O, Jahn A, van Oosterhout JJ, Edmonds A, Phiri N, et al. Timing of
combination antiretroviral therapy (cART) initiation is not associated with stillbirth
among HIV-infected pregnant women in Malawi. Trop Med Int Health [Internet].
2019 Jun;24(6):727–35. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/30891866

44.   Onoya D, Sineke T, Brennan AT, Long L, Fox MP. Timing of pregnancy, postpartum risk
of virologic failure and loss to follow-up among HIV-positive women. AIDS [Internet].
2017;31(11):1593–602. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/28463877

45.   Dinh T-H, Mushavi A, Shiraishi RW, Tippett Barr B, Balachandra S, Shambira G, et al.
Impact of Timing of Antiretroviral Treatment and Birth Weight on Mother-to-Child
Human Immunodeficiency Virus Transmission: Findings From an 18-Month
Prospective Cohort of a Nationally Representative Sample of Mother–Infant Pairs
During the Transition Fro. Clin Infect Dis [Internet]. 2018 Feb 1;66(4):576–85.
Available from: https://academic.oup.com/cid/article/66/4/576/4159392

46.   Djeha A, Girard S, Trottier H, Kakkar F, Soudeyns H, Boucher M, et al. No association
between early antiretroviral therapy during pregnancy and plasma levels of
angiogenic factors: a cohort study. BMC Pregnancy Childbirth [Internet]. 2019 Dec
9;19(1):482. Available from: http://www.ncbi.nlm.nih.gov/pubmed/31815612

47.   Melekhin V V, Shepherd BE, Stinnette SE, Rebeiro PF, Barkanic G, Raffanti SP, et al.
Antiretroviral therapy initiation before, during, or after pregnancy in HIV-1-infected
women: maternal virologic, immunologic, and clinical response. PLoS One [Internet].
2009 Sep 9;4(9):e6961. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/19742315

48.   Kalk E, Schubert P, Bettinger JA, Cotton MF, Esser M, Slogrove A, et al. Placental
pathology in HIV infection at term: a comparison with HIV-uninfected women. Trop
Med Int Health [Internet]. 2017;22(5):604–13. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/28214384

49.   Pillai S, Cheyney M, Everson CL, Bovbjerg ML. Fetal macrosomia in home and birth
center births in the United States: Maternal, fetal, and newborn outcomes. Birth
[Internet]. 2020 Oct 15;birt.12506. Available from:
https://onlinelibrary.wiley.com/doi/10.1111/birt.12506

50.   Bouvier D, Forest J-C, Dion-Buteau E, Bernard N, Bujold E, Pereira B, et al. Association
of Maternal Weight and Gestational Weight Gain with Maternal and Neonate
Outcomes: A Prospective Cohort Study. J Clin Med [Internet]. 2019 Nov
27;8(12):2074. Available from: https://www.mdpi.com/2077-0383/8/12/2074

51.   Madlala HP, Malaba TR, Newell M, Myer L. Elevated body mass index during
pregnancy and gestational weight gain in HIV-infected and HIV-uninfected women in
Cape Town, South Africa: association with adverse birth outcomes. Trop Med Int Heal
[Internet]. 2020 Jun 23;25(6):702–13. Available from:
https://onlinelibrary.wiley.com/doi/10.1111/tmi.13387

52.   Fahme SA, Bloomfield GS, Peck R. Hypertension in HIV-Infected Adults. Hypertension
[Internet]. 2018 Jul;72(1):44–55. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S1473309914708965

53.   Kovo M, Bar J, Schreiber L, Shargorodsky M. The relationship between hypertensive
disorders in pregnancy and placental maternal and fetal vascular circulation. J Am Soc
Hypertens [Internet]. 2017 Nov;11(11):724–9. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S1933171117303054

54.   Bustamante Helfrich B, Chilukuri N, He H, Cerda SR, Hong X, Wang G, et al. Maternal
vascular malperfusion of the placental bed associated with hypertensive disorders in
the Boston Birth Cohort. Placenta [Internet]. 2017 Apr;52:106–13. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0143400417301595

55.  Naderi S, Tsai SA, Khandelwal A. Hypertensive Disorders of Pregnancy. Curr
Atheroscler Rep [Internet]. 2017 Mar 22;19(3):15. Available from:
http://link.springer.com/10.1007/s11883-017-0648-z

56.  Govender N, Naicker T, Moodley J. Maternal imbalance between pro-angiogenic and
anti-angiogenic factors in HIV-infected women with pre-eclampsia. Cardiovasc J Afr
[Internet]. 2013 Jun;24(5):174–9. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/24217170

57.  Graham SM, Rajwans N, Jaoko W, Estambale BBA, McClelland RS, Overbaugh J, et al.
Endothelial activation biomarkers increase after HIV-1 acquisition. AIDS [Internet].
2013 Jul;27(11):1803–13. Available from:
http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00002
030-201307170-00014

58.  Graham SM, Rajwans N, Tapia KA, Jaoko W, Estambale BB, McClelland RS, et al. A
prospective study of endothelial activation biomarkers, including plasma
angiopoietin-1 and angiopoietin-2, in Kenyan women initiating antiretroviral therapy.
BMC Infect Dis [Internet]. 2013 Dec 4;13(1):263. Available from:
https://bmcinfectdis.biomedcentral.com/articles/10.1186/1471-2334-13-263

59.  Mohammadi H, Papp E, Cahill L, Rennie M, Banko N, Pinnaduwage L, et al. HIV
antiretroviral exposure in pregnancy induces detrimental placenta vascular changes
that are rescued by progesterone supplementation. Sci Rep [Internet]. 2018 Dec
26;8(1):6552. Available from: http://www.nature.com/articles/s41598-018-24680-w

**TABLES**

**Table 1: Maternal and obstetric characteristics of the pregnant women at enrolment.**

| | Total N=130 | [1]ART Initiation before pregnancy N=53 | ART Initiation during pregnancy N=77 | p-Value |
|---|---|---|---|---|
| **Age, years** | | | | **<0.0001** |
| ≤24 | 24 (18.5) | 5 (9.4) | 19 (24.7) | |
| 25-29 | 36 (27.7) | 10 (18.9) | 26 (33.8) | |
| 30-34 | 37 (28.5) | 14 (26.4) | 23 (29.9) | |
| ≥35 | 33 (25.4) | 24 (45.3) | 9 (11.7) | |
| Median (IQR) | 31 (26 - 35) | 33 (28 - 37) | 28 (25 - 32) | |
| **Median Gestation at Enrolment** | 15 (11-18) | 14 (11-18) | 15 (11-18) | 0.4 |
| **Gravidity** | | | | 0.1 |
| 1 | 14 (10.8) | 4 (7.5) | 10 (13.0) | |
| 2 | 50 (38.5) | 16 (30.2) | 34 (44.2) | |
| ≥3 | 66 (50.8) | 33 (62.3) | 33 (42.9) | |
| Median (IQR) | 3 (2-3) | 3 (2-4) | 2 (2-3) | |
| **[2]Previous Preterm** | | | | 1.0 |
| Yes | 7 (5.4) | 4 (7.5) | 3 (3.9) | |
| **ART regimen** | | | | 0.2 |
| TDF-3TC-EFV | 119 (91.5) | 47 (88.7) | 72 (93.5) | |
| AZT-3TC-NVP | 2 (1.5) | 1 (1.9) | 1 (1.9) | |
| AZT-XTC-EFV | 6 (4.6) | 2 (3.8) | 4 (5.2) | |
| PI-based regimen | 3 (2.3) | 3 (5.7) | 0 (0) | |
| **[3]CD4 cell count, cells/ml** | | | | **0.04** |
| ≤ 200 | 9 (6.9) | 1 (1.9) | 8 (10.4) | |
| 201-350 | 32 (24.6) | 8 (15.1) | 24 (31.2) | |
| 351-500 | 29 (22.3) | 14 (26.4) | 15 (19.5) | |
| >500 | 41 (31.5) | 20 (37.7) | 21 (27.3) | |
| Median (IQR) | 416 (311 - 589) | 455 (381 - 602) | 369.5 (251 - 534.5) | |
| *Missing* | *19 (14.6)* | *10 (18.9)* | *9 (11.7)* | |
| **[3]Viral load, copies/ml** | | | | 0.1 |
| <400 | 84 (64.6) | 45 (84.9) | 39 (50.6) | |
| 401-1000 | 2 (1.5) | 0 (0) | 2 (2.6) | |
| >1000 | 7 (5.4) | 2 (3.8) | 5 (6.5) | |
| Median (IQR) | 20 (20 - 27) | 20 (20 - 27) | 20 (20 - 34) | |
| *Missing* | *37 (28.5)* | *6 (11.3)* | *31 (40.3)* | |
| **[4]Body Mass Index, kg/m$^2$** | | | | 0.5 |
| Normal (18.5-24.9) | 27 (20.7) | 12 (22.6) | 15 (19.5) | |
| Overweight (25.0-29.9) | 41 (31.5) | 18 (34.0) | 23 (29.9) | |
| Obese (>30.0) | 61 (46.9) | 22 (41.5) | 39 (50.6) | |
| Median (IQR) | 29.45 (25.5 - 33.6) | 29.2 (25.8 - 32.2) | 30 (25.4 -35.2) | |
| *Missing* | *1 (0.8)* | *1 (1.9)* | *0 (0)* | |
| **Blood Pressure, Enrolment (mmHG)** | | | | **0.03** |
| Normal (Sys <120 and Dia <80) | 99 (76.1) | 46 (86.8) | 53 (68.8) | |
| Prehypertensive (Sys 120-139 or Dia 80-89) | 28 (21.5) | 6 (11.3) | 22 (28.6) | |
| Hypertensive (Sys >140 or Dia >90) | 3 (2.3) | 1 (1.9) | 2 (2.6) | |

643    [1]Differences between the two groups were assessed using a two-sample t-test or Wilcoxon rank-sum test
644    depending on the distribution for numerical data and the chi$^2$ test or Fisher's exact test for categorical data.
645    [2]Among women with a previous pregnancy
646    [3]CD4 and viral load results abstracted from routine records and are the nearest in time to the first ANC visit.
647    [4]At enrolment or first ANC
648

649    **Table 2: Infant characteristics**

| | Total N=130 | ART Initiation before pregnancy N=53 | ART Initiation during pregnancy N=77 | p-Value |
|---|---|---|---|---|
| Sex | | | | 0.6 |
|   Female | 58 (44.6) | 25 (47.2) | 33 (42.9) | |
|   Male | 72 (55.4) | 28 (52.8) | 44 (57.1) | |
| Median Delivery Gestation (completed weeks) | 39 (38 - 40) | 39 (38 - 39) | 39 (38 - 40) | 0.1 |
| Birthweight, grams | | | | 0.2 |
|   Low birthweight (<2500g) | 11 (8.5) | 6 (11.3) | 5 (6.5) | |
|   Normal (2500 - 4000g) | 111 (85.4) | 46 (86.8) | 65 (84.4) | |
|   High (>4000g) | 8 (6.1) | 1 (1.9) | 7 (9.1) | |
|   Median (IQR) | 3180 (2880 - 3540) | 3140 (2690 - 3360) | 3240 (2940 - 3580) | |
| Preterm (<37 weeks Gestational Age) | | | | 1.0 |
|   Yes | 9 (6.9) | 4 (7.5) | 5 (6.5) | |
|   No | 121 (93.1) | 49 (92.5) | 72 (93.5) | |
| Size for Gestational Age, <10[th] centile | | | | 0.07 |
|   Yes | 12 (9.2) | 8 (15.1) | 4 (5.2) | |
|   No | 118 (90.8) | 45 (84.9) | 73 (94.8) | |
| Mode of Delivery | | | | 0.2 |
|   Vaginal Delivery | 97 (74.6) | 42 (79.2) | 55 (71.4) | |
|   Elective Caesarean | 24 (18.5) | 4 (7.6) | 4 (5.2) | |
|   Emergency Caesarean | 8 (6.1) | 6 (11.3) | 18 (23.4) | |
|   *Missing* | *1 (0.8)* | *1 (1.9)* | *0 (0)* | |

651    [1]Differences between the two groups were assessed using a two-sample t-test or Wilcoxon rank-sum test

652    depending on the distribution for numerical data and the chi$^2$ test or Fisher's exact test for categorical data.

654 **Table 3: Placental characteristics and pathology at delivery among women initiating ART**
655 **before or during pregnancy.**

| | Total<br>N=130 | [1]ART Initiation<br>before pregnancy<br>N=53 | ART Initiation<br>during pregnancy<br>N=77 | P-value |
|---|---|---|---|---|
| Placental weight (g), median (IQR) | 412 (346 - 480) | 392 (346 − 440) | 422 (344 − 490) | 0.09 |
| Placental weight (g) | | | | 1.0 |
| Small (<10[th] percentile) | 43 (33.1) | 17 (32.1) | 26 (33.8) | |
| Appropriate (10 − 90[th] percentile) | 85 (65.4) | 35 (66.0) | 50 (64.9) | |
| Large (>90[th] percentile) | 1 (0.8) | 0 (0) | 1 (1.3) | |
| *Missing* | *1 (0.8)* | *1(1.9)* | *0(0)* | |
| Fetal-Placenta weight ratio | | | | 0.9 |
| Small (<10[th] percentile) | 3 (2.3) | 1 (1.9) | 2 (2.6) | |
| Appropriate (10 − 90[th] percentile) | 81 (62.3) | 32 (60.4) | 49 (63.6) | |
| Large (>90[th] percentile) | 45 (34.6) | 19 (35.8) | 26 (33.8) | |
| *Missing* | *1 (0.8)* | *1(1.9)* | *0(0)* | |
| Placenta greatest diameter (mm), median (IQR) | 170 (160 − 180) | 175 (161 − 189) | 170 (160 − 180) | 0.1 |
| Placenta thickness (mm), median (IQR) | 17.5 (10 − 25) | 15 (10 − 22) | 20 (10 - 25) | 0.4 |
| Cord insertion | | | | 0.1 |
| Central | 20 (15.4) | 6 (11.3) | 14 (18.2) | |
| Off centre | 88 (67.7) | 34 (64.1) | 54 (70.1) | |
| Marginal | 16 (12.3) | 10 (18.9) | 6 (7.8) | |
| *Missing* | *6 (4.6)* | *3(5.7)* | *3(3.9)* | |
| **Meconium exposure** | | | | 0.9 |
| + | 25 (19.2) | 10 (18.9) | 15 (19.5) | |
| - | 105 (80.8) | 43 (81.1) | 62 (80.5) | |
| **Chorioamnionitis** | | | | |
| *Maternal Inflammatory Response* | | | | 0.8 |
| + | 19 (14.6) | 7 (13.2) | 12 (15.6) | |
| - | 111 (85.4) | 46 (86.3) | 65 (84.4) | |
| *Fetal Inflammatory Response* | | | | 0.1 |
| + | 20 (15.4) | 5 (9.4) | 15 (19.5) | |
| - | 110 (84.6) | 48 (90.6) | 62 (80.5) | |
| **Chronic Deciduitis** | | | | 0.8 |
| + | 14 (10.8) | 5 (9.4) | 9 (11.7) | |
| - | 116 (89.2) | 48 (90.6) | 68 (88.3) | |
| **Maternal Vascular Malperfusion (MVM)** | | | | *0.01* |
| + | 36 (27.7) | 21 (39.6) | 15 (19.5) | |
| - | 94 (72.3) | 32 (60.4) | 62 (80.5) | |

656 [1]Differences between the two groups were assessed using a two-sample t-test or Wilcoxon rank-sum test
657 depending on the distribution for numerical data and the chi$^2$ test or Fisher's exact test for categorical data.

659    **Table 4: Relationship between placenta pathology and preterm delivery, size for gestational age and low birthweight.**

| | [1]Preterm delivery | | | [1]Size for gestational age (<10th centile) | | | [1]Low birthweight (<2500g) | | |
|---|---|---|---|---|---|---|---|---|---|
| | Y (%) | N (%) | p | Y (%) | N (%) | p | Y (%) | N (%) | p |
| **Percentile placental weight** | | | 0.7 | | | 0.1 | | | 1.0 |
| <10th | 2 (23) | 41 (34) | | 7 (58) | 36 (30) | | 4 (36) | 39 (33) | |
| >10th | 6 (67) | 80 (66) | | 5 (42) | 81 (69) | | 7 (64) | 79 (66) | |
| *Missing* | *1 (11)* | *0 (0)* | | *0(0)* | *1(1)* | | *0 (0)* | *1(1)* | |
| **Cord insertion** | | | 0.2 | | | 1.0 | | | 0.1 |
| Central | 3 (33) | 17 (14) | | 2 (17) | 18 (15) | | 4 (36) | 16 (13) | |
| Off centre | 5 (56) | 83 (69) | | 8 (67) | 80 (68) | | 7 (64) | 81 (68) | |
| Marginal | 0 (0) | 16 (13) | | 1 (8) | 15 (13) | | 0 (0) | 16 (14) | |
| *Missing* | *1 (11)* | *5 (4)* | | *1 (8)* | *5 (4)* | | *0 (0)* | *6 (5)* | |
| **Meconium exposure** | | | 1.0 | | | *0.01* | | | 0.7 |
| + | 1 (11) | 24 (19) | | 6 (50) | 19 (16) | | 1 (9) | 24 (20) | |
| - | 8 (89) | 97 (80) | | 6 (50) | 99 (84) | | 10 (91) | 95 (78) | |
| **Chorioamnionitis** | | | | | | | | | |
| *Maternal Inflammatory Response* | | | 0.4 | | | 0.4 | | | 0.4 |
| + | 0 (0) | 19 (16) | | 3 (25) | 16 (14) | | 0 | 19 (16) | |
| - | 9 (100) | 102 (84) | | 9 (75) | 102 (86) | | 11 (100) | 100 (84) | |
| *Fetal Inflammatory Response* | | | 0.4 | | | 1.0 | | | 0.2 |
| + | 0 | 20 17) | | 2 (17) | 18 (15) | | 0 | 20 (17) | |
| - | 9 (100) | 101 (83) | | 10 (83) | 100 (85) | | 11 (100) | 99 (83) | |
| **Chronic Deciduitis** | | | 1.0 | | | 0.4 | | | 0.4 |
| + | 1 (11) | 13 (11) | | 0 | 14 (12) | | 2 (18) | 12 (10) | |
| - | 8 (89) | 108 (89) | | 12 (100) | 104 (88) | | 9 (82) | 107 (90) | |
| **Decidual arteriopathy** | | | 0.07 | | | 0.5 | | | 0.1 |
| + | 2 (22) | 5 (4) | | 1 (8) | 6 (5) | | 2 (18) | 5 (4) | |
| - | 7 (78) | 116 (96) | | 11 (92) | 112 (95) | | 9 (82) | 114 (96) | |
| **Maternal Vascular Malperfusion (MVM)** | | | *0.002* | | | 0.09 | | | *<0.0001* |
| + | 7 (78) | 29 (24) | | 6 (50) | 30 (25) | | 10 (91) | 26 (22) | |
| - | 2 (23) | 92 (76) | | 6 (50) | 88 (75) | | 1 (9) | 93 (78) | |

661    [1]To assess potential risk factors for preterm delivery (<37 weeks gestational age), size for gestational age (<10th percentile) and low birth weight (<2500g), we compared
662    differences between groups using the $chi^2$ test or Fisher's exact test.

663 **Supplementary Table 1: Substance use among study participants.**

| | Total N=130 | ART Initiation before pregnancy N=53 | ART Initiation during pregnancy N=77 | p-Value |
|---|---|---|---|---|
| Substance Use† | | | | |
| Alcohol | | | | 0.2 |
| Yes | 22 (16.9) | 6 (11.3) | 16 (20.8) | |
| No | 105 (80.8) | 45 (84.9) | 60 (77.9) | |
| *Missing* | *3 (2.3)* | *2 (3.8)* | *1 (1.3)* | |
| Smoking | | | | 0.1 |
| Yes | 7 (5.4) | 5 (9.4) | 2 (2.6) | |
| No | 121 (93.1) | 46 (86.8) | 75 (97.4) | |
| *Missing* | *2 (1.5)* | *2 (3.8)* | *0 (0)* | |
| Drugs | | | | 0.5 |
| Yes | 2 (1.5) | 0 (0) | 2 (2.6) | |
| No | 126 (96.9) | 51 (96.2) | 75 (97.4) | |
| *Missing* | *2 (1.5)* | *2 (3.8)* | *0 (0)* | |

664 †Substance use is self-reported over the last 30 days

665 [1]Differences between the two groups were assessed using a two-sample t-test or Wilcoxon rank-sum test

666 depending on the distribution for numerical data and the $chi^2$ test or Fisher's exact test for categorical data.

668 **Supplementary Table 2: Maternal and obstetric characteristics of pregnant women at enrolment: comparing no placenta vs placenta**

669 **collection.**

| | Total | No Placenta | Placenta | p-value | No Placenta N=421 | | p-value | Placenta N=130 | | p-value |
|---|---|---|---|---|---|---|---|---|---|---|
| | N=551 | N=421 | N=130 | | ART Initiation before pregnancy N=208 | ART Initiation during pregnancy N=213 | | ART Initiation before pregnancy N=53 | ART Initiation during pregnancy N=77 | |
| Age, years | | | | 0.6 | | | <0.001 | | | <0.001 |
| ≤24 | 92 (16.7) | 68 (16.2) | 24 (18.5) | | 20 (9.6) | 48 (22.5) | | 5 (9.4) | 19 (24.7) | |
| 25-29 | 156 (28.3) | 120 (28.5) | 36 (27.7) | | 48 (23.1) | 72 (33.8) | | 10 (18.9) | 26 (33.8) | |
| 30-34 | 179 (32.5) | 142 (33.7) | 37 (28.5) | | 74 (35.6) | 68 (31.9) | | 14 (26.4) | 23 (29.9) | |
| ≥35 | 124 (22.5) | 91 (21.6) | 33 (25.4) | | 66 (31.7) | 25 (11.7) | | 24 (45.3) | 9 (11.7) | |
| Median (IQR) | 30 (26-34) | 30 (26-34) | 31 (26-35) | | 32 (28-35) | 29 (25-32) | | 33 (28 - 37) | 28 (25 - 32) | |
| Median Gestation at Enrolment | 14 (10-18) | 13 (10-17) | 14 (11-18) | 0.04 | 13 (9-16) | 14 (10-18) | 0.02 | 13 (10-18) | 15 (11-18) | 0.2 |
| Gravidity | | | | 0.14 | | | 0.001 | | | 0.1 |
| 1 | 89 (16.1) | 75 (17.8) | 14 (10.8) | | 50 (23.5) | 25 (12.0) | | 4 (7.5) | 10 (13.0) | |
| 2 | 189 (34.3) | 139 (33.0) | 50 (38.5) | | 76 (35.7) | 63 (30.3) | | 16 (30.2) | 34 (44.2) | |
| ≥3 | 273 (49.6) | 207 (49.2) | 66 (50.8) | | 87 (40.8) | 120 (57.7) | | 33 (62.3) | 33 (42.9) | |
| ART regimen | | | | 0.1 | | | <0.001 | | | 0.2 |
| TDF-3TC-EFV | 499 (90.6) | 380 (90.3) | 119 (91.5) | | 173 (83.2) | 207 (97.2) | | 47 (88.7) | 72 (93.5) | |
| TDF-3TC-NVP | 4 (0.7) | 4 (0.9) | 0 (0) | | 2 (0.9) | 2 (0.9) | | 0 (0) | 0 (0) | |
| AZT-3TC-NVP | 10 (1.8) | 8 (1.9) | 2 (1.6) | | 8 (3.9) | 0 (0) | | 1 (1.9) | 1 (1.9) | |
| AZT-XTC-EFV | 13 (2.4) | 7 (1.7) | 6 (4.6) | | 5 (2.4) | 2 (0.9) | | 2 (3.8) | 4 (5.2) | |
| PI-based regimen | 25 (4.5) | 22 (5.2) | 3 (2.3) | | 20 (9.6) | 2 (0.9) | | 3 (5.7) | 0 (0) | |
| [1]Body Mass Index, kg/m$^2$ | | | | 0.4 | | | 0.3 | | | 0.6 |
| Normal (18.5-24.9) | 115 (20.9) | 88 (20.9) | 27 (20.8) | | 38 (18.3) | 50 (23.5) | | 12 (22.6) | 15 (19.5) | |
| Overweight (25.0-29.9) | 149 (27.0) | 108 (25.7) | 41 (31.5) | | 58 (27.9) | 50 (23.5) | | 18 (34.0) | 23 (29.9) | |
| Obese (>30.0) | 282 (51.2) | 221 (52.5) | 61 (46.9) | | 111 (53.4) | 110 (51.6) | | 22 (41.5) | 39 (50.6) | |
| Missing | 5 (0.9) | 4 (0.9) | 1 (0.8) | | 1 (0.5) | 3 (1.41) | | 1 (1.9) | 0(0) | |
| Blood Pressure, Enrolment (mmHG) | | | | 0.4 | | | 0.3 | | | 0.03 |
| Normal (Sys <120 and Dia <80) | 405 (73.5) | 306 (72.7) | 99 (76.2) | | 156 (75.0) | 150 (70.4) | | 46 (86.8) | 53 (68.8) | |
| Prehypertensive (Sys 120-139 or Dia 80-89) | 123 (22.3) | 95 (22.6) | 28 (21.5) | | 45 (21.6) | 50 (23.5) | | 6 (11.3) | 22 (28.6) | |
| Hypertensive (Sys >140 or Dia >90) | 23 (4.2) | 20 (4.7) | 3 (2.3) | | 7 (3.4) | 13 (6.1) | | 1 (1.9) | 2 (2.6) | |

[1]At enrolment or first ANC

**FIGURE LEGENDS**

Figure 1A: Digital scan of a representative Haematoxylin and Eosin (H&E) stained section

showing chorioamnionitis of the membranes (maternal response).

Figure 1B: Digital scan of a representative H&E stained section showing a neutrophilic

infiltrate involving the cord vein wall extending into Wharton's jelly of the cord (fetal

response).

Figure 1C: Digital scans of representative H&E stained sections of the placenta showing an

infarct. Insert 1 shows a magnified image of the infarct and Insert 2 shows a magnified foam

cell (denoted by the arrow) within the decidual artery demonstrating atherosis.

Figure 2A: Forest plot depicting the relative risk with the 95% confidence intervals of HIV

infected women presenting with a small placenta (<10th percentile for gestational age).

Figure 2B: Forest plot depicting the relative risk with the 95% confidence intervals of HIV

infected women presenting histologic chorioamnionitis.

Figure 2C: Forest plot depicting the relative risk with the 95% confidence intervals of HIV

infected women presenting with placental maternal vascular malperfusion.

Figure 2D: Alluvial plot depicting the impact of ART timing on the placental pathology

among women initiating treatment before pregnancy (*Initiating)* or during pregnancy

(*stable*). The plot depicts frequencies from a subset of women with abnormal placenta

pathology only. These include CA (Chorioamnionitis depicting FIR Fetal Inflammatory

Response; MIR, Maternal Inflammatory Response), CD (Chronic Deciduitis), MVM (Maternal

Vascular Malperfusion) and Meconium (Meconium Exposure) and the placenta size i.e. <10th

percentile for gestational age (small) and > 10th percentile for gestational age (Normal)

among women initiating treatment before pregnancy (*Initiating)* or during pregnancy

(*stable*).

Differential impact of antiretroviral therapy initiated before or during pregnancy on placenta pathology in HIV-positive women.

Figures

Figure 1A

Figure 1A: Digital scan of a representative Haematoxylin and Eosin (H&E) stained section showing chorioamnionitis of the membranes (maternal response).

Figure 1B

Figure 1B: Digital scan of a representative H&E stained section showing a neutrophilic infiltrate involving the cord vein wall extending into Wharton's jelly of the cord (fetal response).

Figure 1C

Figure 1C: Digital scans of representative H&E stained sections of the placenta showing an infarct. Insert 1 shows a magnified image of the infarct and Insert 2 shows a magnified foam cell (denoted by the arrow) within the decidual artery demonstrating atherosis.

Figure 2A

Figure 2A: Forest plot depicting the relative risk with the 95% confidence intervals of HIV infected women presenting with a small placenta (<10th percentile for gestational age).

Figure 2B

Figure 2B: Forest plot depicting the relative risk with the 95% confidence intervals of HIV infected women presenting histologic chorioamnionitis.

Figure 2C

**Maternal Vascular Malperfusion**

| Outcome: Malperfusion | N (%) | Risk Ratio (95% CI) | p-Value |
|---|---|---|---|
| ART status | | | |
| Initiating | 15(41.7) | | |
| Stable | 21(58.3) | 2.03(1.16 - 3.57) | 0.01 |
| Maternal age (years) | | | |
| <=24 | 8(22.2) | | |
| 25-29 | 8(22.2) | 0.67(0.29 - 1.53) | 0.34 |
| >=30 | 20(55.6) | 0.86(0.43 - 1.68) | 0.65 |
| Maternal BMI, enrolment | | | |
| Normal | 9(25) | | |
| Overweight | 14(38.9) | 1.02(0.52 - 2.03) | 0.94 |
| Obese | 12(33.3) | 0.59(0.28 - 1.23) | 0.16 |
| Booking Blood Pressure | | | |
| Normal | 29(80.6) | | |
| Prehypertensive/Hypertensive | 7(19.4) | 0.77(0.37 - 1.58) | 0.48 |
| Any substance use | | | |
| No substance use | 28(77.8) | | |
| Substance use | 8(22.2) | 1.09(0.56 - 2.11) | 0.8 |

Figure 2C: Forest plot depicting the relative risk with the 95% confidence intervals of HIV infected women presenting with placental maternal vascular malperfusion.

Figure 2D

Figure 2D: Alluvial plot depicting the impact of ART timing on the placental pathology clusters among women initiating treatment before pregnancy (*Initiating)* or during pregnancy (*stable*). The plot depicts frequencies from a subset of women with abnormal placenta pathology only. These include CA (Chorioamnionitis depicting FIR Fetal Inflammatory Response; MIR, Maternal Inflammatory Response), CD (Chronic Deciduitis), MVM (Maternal Vascular Malperfusion) and Meconium (Meconium Exposure) and the placenta size i.e. <10th percentile for gestational age (small) and > 10th percentile for gestational age (Normal) among women initiating treatment before pregnancy (*Initiating)* or during pregnancy (*stable*).