# Evaluation of a Novel Therapeutic Repetitive Transcranial Magnetic Stimulation Technique Optimized for Increased Accessibility in Major Depression

Jean-Philippe Miron [1,2,3,4,5,6,7,*], Helena Voetterl [1,8], Linsay Fox [1], Molly Hyde [1], Farrokh Mansouri [1,3], Sinjin Dees [9], Ryan Zhou [1], Jack Sheen [1,3], Arsalan Mir-Moghtadaei [1,3], Daniel M. Blumberger [3,4,10], Zafiris J. Daskalakis [3,4,10], Fidel Vila-Rodriguez [11], Jonathan Downar [1,2,3,4]

[1] Krembil Research Institute, University Health Network, Toronto, ON, Canada;

[2] Poul Hansen Family Centre for Depression, University Health Network, Toronto, ON, Canada;

[3] Institute of Medical Science, Faculty of Medicine, University of Toronto, Toronto, ON, Canada;

[4] Department of Psychiatry, Faculty of Medicine, University of Toronto, Toronto, ON, Canada;

[5] Unité de Neuromodulation Psychiatrique (UNP), Département de Psychiatrie, Centre Hospitalier de l'Université de Montréal (CHUM), Montréal, QC, Canada;

[6] Centre de Recherche du CHUM (CRCHUM), Université de Montréal, Montréal, QC, Canada;

[7] Département de psychiatrie, Faculté de Médecine, Université de Montréal, Montréal, QC, Canada;

[8] Department of Cognitive Neuroscience, Maastricht University, Maastricht, Limburg, Netherlands;

[9] Faculty of Engineering, McMaster University, Hamilton, ON, Canada;

[10] Temerty Centre for Therapeutic Brain Intervention at the Centre for Addiction and Mental Health, Toronto, ON, Canada;

[11] Non-Invasive Neurostimulation Therapies Laboratory, Department of Psychiatry, University of British Columbia, Vancouver, BC, Canada.

* Corresponding author

ABSTRACT WORD COUNT: 244

MANUSRIPT WORD COUNT: 3824

TABLES AND FIGURES: 9

**ABSTRACT**

**BACKGROUND**

Repetitive transcranial magnetic stimulation (rTMS) is effective in major depressive disorder (MDD). However, technical complexity and operational costs might have been barriers for its wide use and implementation in some jurisdictions, thereby decreasing accessibility.

**OBJECTIVE**

Our main goal was to test the feasibility of a novel rTMS protocol optimized for practicality, scalability and cost-effectiveness. We hypothesized that our novel rTMS protocol would be simple to implement and well-tolerated, but less costly and allow for more treatment capacity.

**METHODS**

Treatment was administered in an open-room setting, allowing a single technician to attend to multiple patients. Large non-focal parabolic coils held by custom-built arms allowed simple yet efficient and accurate placement. We employed a low-frequency (LF) 1 Hz stimulation protocol (360 pulses per session), delivered on the most affordable FDA-approved devices. MDD participants received an initial accelerated rTMS course (arTMS) of 6 sessions/day over 5 days (30 total), followed by a tapering course of daily sessions (up to 25) to decrease the odds of relapse. The self-reported Beck Depression Inventory II (BDI-II) was used to measure severity of depression.

**RESULTS**

Forty-eight (48) patients completed the arTMS course. No serious adverse events occurred, and all patients reported manageable pain levels. Response and remission rates were 35.4% and 27.1% on the BDI-II, respectively, at the end of the tapering course.

**CONCLUSION**

If rTMS could be delivered for lower cost at higher volume, while preserving efficacy, safety and tolerability, it could warrant further investigation of this treatment as a first-line intervention in MDD.

**TRIAL REGISTRATION**

ClinicalTrials.gov Identifier: NCT04376697

**INTRODUCTION**

Major depressive disorder (MDD) is a common and disabling illness. Up to 50% of patients experience a chronic or recurrent course, and 30 to 40% develop treatment-resistant depression [1]. Furthermore, although antidepressants offer high convenience and simplicity of

administration, discontinuation rates approach 50% after 3 months of use due to concern over side effects and non-response [2]. Alternative treatments are therefore needed.

Repetitive transcranial magnetic stimulation (rTMS) is recognized as an effective intervention in MDD, with recent studies and meta-analyses reporting response and remission rates of up to 50-55% and 30-35%, respectively [3]. rTMS has an advantageous side effect profile, with lower discontinuation rates than medication (~5% vs ~25%) [3]. Unfortunately, its widespread adoption is impeded by several obstacles that limit clinical accessibility.

The main issue concerns high acquisition and operation costs, with average cost per remission estimated to be of up to $6,146 in the US [4]. This is partly a result of the equipment needed to deliver high-frequency (HF) protocols, as well as the need for continuous 1:1 technician-patient supervision during treatment. Technical challenges stem from the complexity of the treatment, such as the need for precise positioning of the widely used figure-of-eight (Fo8) focal coils over the target region.

Another area that remains to be refined is treatment course optimization. The current treatment paradigm forces patients to travel to a treatment center every day for 6 weeks in order to receive a full 30 session course. This can be discouraging, especially for working individuals and those who have families. Several studies have explored "accelerated" rTMS (arTMS), whereby multiple stimulation sessions are delivered per day to shorten the overall treatment course duration. Several large trials of arTMS have consistently reported similar or better remission and response rates than conventional once-daily rTMS [5–16]. Furthermore, the total cumulative number of sessions needed to achieve maximal effect is still debated. So far, findings from large rTMS trials have suggested a plateau in clinical response at ~30 sessions, on average [17]. This has recently been challenged by a secondary analysis [18] from the THREE-D trial [19], demonstrating differences in patients' response trajectories to rTMS. Indeed, some participants were still showing signs of improvement at 30 sessions (not reaching a plateau), suggesting that a treatment extension might be needed in some individuals.

To address these issues, we developed a novel rTMS technique optimized for maximum practicality, scalability and effectiveness, while minimizing costs. We employed a safe, well-tolerated low-frequency (LF) 1 Hz stimulation protocol on the lowest cost FDA-approved devices, delivered via large parabolic coils held by low-cost custom arms enabling simple yet accurate placement, in an open-room setting enabling supervision of multiple simultaneous sessions. We also tested the effects of 1 Hz arTMS followed by a tapering course of once-daily treatment. These technical refinements are designed to enable the provider to accelerate, increase and maintain treatment response beyond what 'standard of care' rTMS can achieve, while achieving higher patient volumes at lower cost.

We hypothesized that our novel rTMS technique would be safe, well-tolerated and effective, while allowing cost-saving opportunities and therefore increasing accessibility.

**METHODS**

**Participants**

We conducted a prospective, single-arm, open-label feasibility study. Participants were recruited after referral to the Poul Hansen Family Centre for Depression neurostimulation specialty clinic, located at the Toronto Western Hospital, an academic healthcare centre which is part of the University Health Network (UHN) in Toronto, Canada.

Adult (18-85 years of age) outpatients were included for study participation if they 1) had a Mini International Neuropsychiatric Interview (MINI) confirmed MDD diagnosis (single or recurrent episode) and 2) maintained a stable medication regimen from 4 weeks before treatment start to the end of the study. We did not require participants to have had a minimal number of failed antidepressant treatment in their current depressive episode. Exclusion criteria were: 1) history of substance dependence or abuse within the last 3 months; 2) concomitant major unstable medical illness; 3) cardiac pacemaker or implanted medication pump; 4) active suicidal intent; 5) diagnosis of any personality disorder as assessed by a study investigator to

be primary and causing greater impairment than MDD; 6) diagnosis of any psychotic disorder; 7) any significant neurological disorder or insult (including, but not limited to: any condition likely to be associated with increased intracranial pressure, space occupying brain lesion, any history of seizure confirmed diagnostically by neurological assessment [except those therapeutically induced by ECT], cerebral aneurysm, Parkinson's disease, Huntington's chorea, dementia, stroke, neurologically confirmed diagnosis of traumatic brain injury, or multiple sclerosis); 8) if participating in psychotherapy must have been in stable treatment for at least 3 months prior to entry into the study (with no anticipation of change in the frequency of therapeutic sessions, or the therapeutic focus over the duration of the study); 9) any clinically significant laboratory abnormality in the opinion of the investigator; 10) a dose of more than lorazepam 2 mg daily (or equivalent) currently (or in the last 4 weeks) or any dose of an anticonvulsant due to the potential to limit rTMS efficacy and 11) any non-correctable clinically significant sensory impairment. All participants provided informed consent and this study was approved by the Research Ethics Board of the University Health Network.

**Study design and procedure**

Treatment was delivered in an open room setting, allowing up to 4 participants to receive treatment simultaneously, with the help of one or two technicians (**Fig. 1**). This was facilitated by an easy-to-use coil placement system, with the Anchored Articulating Arm (AAA) concept at its core (**Fig. 2**). AAAs are made out of readily available camera tripod equipment and are anchored at their base to the corner of the treatment chair. Coils are held securely by a clamp locking it to the AAA, maintaining position until treatment has been completed, at which time they can be easily repositioned out of the way.

rTMS was delivered through MagPro R20 stimulators equipped with parabolic MMC-140 coils (MagVenture, Farum, Denmark). We recently published a case series on the safety, tolerability, and effectiveness of those coils in MDD [20]. The resting motor threshold (rMT) was

determined according to standard clinical practice, with the additional need to use the coil's

middle ring for stimulation, where the electromagnetic field strength is at its highest [20,21]. We

used a previously published 1 Hz rTMS protocol (60 s on and 30 s off, 6 trains, 8.5 min total

stimulation time, 360 pulses/sessions, 120% rMT) over the right dorsolateral prefrontal cortex

(DLPFC) with the coil centred on the F4 EEG location (right-flipped adjusted BeamF3) [20,22].

Treatment consisted of an arTMS course of 6 sessions/day (50 min inter-sessions

intervals) over 5 days (on weekdays), thus totalling 30 sessions. After a 1-week gap, arTMS

was followed by a tapering course of once-daily stimulation (minimum of 20, maximum of 25

sessions), 3-5 days per week, to decrease the odds of relapse. *Baseline* visits were conducted

the week prior to arTMS initiation and consisted of a clinical assessment by a trained research

staff, navigation cap fitting and motor threshold calibration. Participants were clinically

reassessed at 3 subsequent timepoints: *post-acute* visits were conducted during the one-week

gap following arTMS; *post-tapering* visits were conducted within 7 days of tapering termination;

*1-month follow-up* visits were conducted 1-month (± 7 days) after the last tapering session.

Participants who missed any sessions of the arTMS course were withdrawn. All participants

were encouraged to complete the tapering course. The study timeline is illustrated in **Fig. 3**.

At each clinical assessment, participants completed the self-rated Beck Depression

Inventory – II (BDI-II), and research staff administered the Hamilton Rating Scale for Depression

17-item (HRSD-17). On each treatment day, participants also completed the BDI-II and were

queried about any adverse events. Self-rated pain intensity of the rTMS procedure was also

recorded on a verbal analogue scale (VRS – from 1 [no pain] to 10 [intolerable pain]). We also

recorded the number of serious adverse events and reasons for treatment discontinuation when

such events occurred. To ensure tolerability, stimulation intensity was adaptively titrated upward

as quickly as possible to the target intensity of 120% RMT, without exceeding maximum

tolerable pain. We recorded the number of sessions required to reach 120% rMT.

**Outcomes**

The primary outcome was the degree of improvement following the tapering course as illustrated by response and remission rates on the BDI-II. Secondary outcomes included score changes and percent improvement. The same outcomes are reported on the HRSD-17. Response was defined as score reductions of $\geq 50\%$ from baseline. Remission was defined as a score of $\leq 12$ [23] on the BDI-II and $\leq 7$ on the HRSD-17 [24]. We also assessed the number of participants maintaining remission and response from at various timepoints, as well as BDI-II scores trajectory across the entire study. Regarding suicidality, we assessed the number of participants showing remission (number of participants starting with a suicidality score greater than zero at baseline and decreasing to zero at reassessment). Suicidality scores are assessed by item #9 on the BDI-II and item #3 on the HRSD-17.

**RESULTS**

From March 18 to September 27, 2019, 57 participants with MDD were assessed for eligibility, 7 of whom were ineligible or declined to participate. 50 participants were thus enrolled and started the study. Of these, 2 discontinued treatment during arTMS because of lack of perceived efficacy and were lost to follow-up. 48 participants moved on to the tapering course and were thus included in the final analysis. Of these, 3 participants discontinued before having completed at least 20 sessions of the tapering course because of the lack of perceived efficacy. 45 participants thus completed the entire study, and 35 followed up at 1 month (**Fig. 4**).

**Table 1** provides the baseline characteristics of the study participants. Mean age was $41.8 \pm 12.2$, with 56.3% female participants. Mean age of depression onset was $23.8 \pm 10.8$ years old, with average length of current episode $51.7 \pm 78.2$ months. 70.8% of participants were receiving psychopharmacotherapy during the trial, with 58.3% taking at least one antidepressant during the study. All participants had tried antidepressant medication in the past,

with a mean of 5.2 ± 3.6 lifetime trials. Average Antidepressant Treatment History Form (ATHF) total score was 4.2 ± 4.3. Average number of trials on the ATHF in the current episode was 1.4 ± 2.6, with 39/48 (81.3%) of participants having had at least one adequate antidepressant trial in their current depressive episode.

Regarding the primary outcome, 35.4% and 27.1% of participants had achieved response and remission on the BDI-II, respectively. Secondary outcomes are presented in **Table 2**. On the BDI-II, 10 out 12 (83.3%) who were responders *post-acute* treatment maintained response *post-tapering*, and 6 out of 7 (85.7%) participants who were in remission *post-acute* treatment maintained remission *post-tapering*. On the HRSD-17, 10 out of 11 (90.9%) maintained response, and 3 out 3 (100.0%) maintained remission. BDI-II scores over the duration of the study are presented in **Fig. 5**. Responders showed rapid improvement during the accelerated course to meet criteria for clinical response, having achieved on average response *post-acutely*, and continued to show slow but steady additional improvement up to the end of the tapering course. Data regarding suicidality are shown in **Table 3**. No participants reported active suicidal intent at any point during the study.

Safety and tolerability outcomes are presented in **Table 4**. No serious adverse events (AE) were reported at any point of the study. One patient was withdrawn at session 51 since he reported visual symptoms suggestive of possible retinal detachment but was assessed and cleared by ophthalmology (subsequent diagnosis of migraine equivalent). Overall, 39.6% of participants reported at least one occurrence of an AE at some point during treatment, the most common one having been headache (22.9%). All AEs were reported exclusively during the arTMS course, with the exception of the previously mentioned participant with visual symptoms at session 51. Pain ratings went from 5.6 ± 2.0 on the first treatment, down to 1.9 ± 1.9 on the last treatment. The average rMT was 34.6 ± 5.7% of maximum stimulator output, resulting in a mean target stimulation intensity (120%) of 41.8 ± 6.8%. All participants were able to reach their

target stimulation intensity, averaging 2.4 ± 3.0 sessions (1.2 ± 0.4 days) to do so, with 33/48 (68.8%) able to achieve it during their first session of treatment. Average total number of sessions was 54.1 ± 2.9, and mean overall treatment length, including the week gap between arTMS and tapering course, was 8.39 ± 0.93 weeks.

Regarding data collection quality, all participants were reassessed within the established time frame *post-acutely*. For *post-tapering* reassessments, the established boundary was within 7 days of the last treatment, and 4/48 (8.3%) participants were reassessed beyond that objective. Finally, for *1-month follow-up* visits, the established boundary was ± 7 days 1-month post- last tapering rTMS sessions, with 1/35 patient having been reassessed below that boundary (1 day too early), and 8/35 having been reassessed beyond that boundary (most delayed by 12 days).

## DISCUSSION

We present preliminary data for a novel rTMS technique optimized for widespread use. Currently, most guidelines recommend rTMS use solely in treatment-resistant depression (TRD). Specifically, the Canadian Network for Mood and Anxiety Treatments (CANMAT) recommends use in MDD participants who have failed to respond to at least 1 antidepressant [3], a decision mirrored in other international guidelines [17,25]. However, access could potentially be broadened if treatments could be delivered safely in more centres, at higher volumes and lower costs, with a faster onset of response. Our study thus attempted to resolve these issues by offering several technical refinements.

First, treatment was delivered in an open-room setting, allowing one or two technicians to treat up to 4 participants in parallel, compared to the usual one-on-one approach. Even though such a setting could bring about some concerns about technician fatigue and negative patient feedback, this was not reported. We also did not observe any negative

interactions between participants that could have jeopardized treatment outcome, and several participants anecdotally gave positive feedback about the more flexible scheduling of treatment sessions enabled by the open setting. Overall, this approach merits qualitative assessment with standardized feedback questionnaires for both technicians and participants. If successful on a larger scale, we believe that this approach may reduce staffing demands, increase clinic capacity and flexibility, and improve cost and accessibility. Given the current COVID-19 pandemic, reduce staffing might be a potential advantage in order to observe physical distancing measures, but it would need to be conjugated with stringent infection control procedures.

One innovation that permitted such a treatment setting was the design and use of our novel AAAs. Standard articulated arms can be more expensive, while offering a limited range of motion. Our AAAs are made of widely available camera tripod equipment and have been both functional and reliable, with no material malfunctions observed. Their durable yet bendable extended branch allowed for the treatment technician to easily and accurately position the coil without tedious joint adjustments. Their anchoring to the treatment chair also eliminated any standalone equipment or cart, creating a more spacious, barrier-free clinic. The mechanical advantage of the anchoring point gives the patient the flexibility to raise or lower the incline of their treatment chair without placement adjustment, as the coil is now anchored to the shifting point of reference, thereby maintaining the coils' targeted area of focus. This custom solution decreased operational complexity, allowing for better patient care, ease of use for the technician, and more efficient use of space and equipment. Similar coil positioning equipment would be relatively straightforward for rTMS manufacturers to adopt in future.

The use of 1 Hz stimulation is another major component allowing for simplified rTMS delivery. Firstly, 1 Hz rTMS is the medically safest stimulation pattern available, having even shown anti-seizure properties in epileptic patients [26]. Further, there exists empirical evidence that 1 Hz rTMS is better tolerated than HF protocols [27], and also shows similar efficacy [28–

30]. With preliminary data potentially showing a higher safety and tolerability profile, LF 1 Hz could be considered in the future as a first line rTMS protocol. Beyond these factors, the greatest advantage of 1 Hz rTMS lies in the low cost of the equipment it requires. HF rTMS require higher-cost stimulators, with expensive cooling systems. 1 Hz rTMS on the other hand can be delivered on more simple stimulators with coils that do not require cooling. This setup is more affordable than the ones required for HF, with the possibility of additional cost reductions in the future through higher volume use.

Parabolic coils also allow simplification of rTMS delivery. We previously published a case report on their use [20], delineating their potential advantage over Fo8 coils. Indeed, large parabolic coils may require less precise placement given their large electromagnetic field compared to the target brain region – a factor also present with larger helmet-shaped coils [31]. Also, their central opening allows direct visualization of underlying landmarks, resulting in easier accurate placement over the marked target. We also hypothesized that, given their possibly weaker stimulation of the DLPFC owed to the aforementioned opening, parabolic coils may function by stimulating adjacent neural substrates, such as the orbitofrontal cortex (OFC). Our group also previously published a case series on 1 Hz OFC, which is theorized to be involved in depression [32–34].

Another novel aspect of our study was the piloting of accelerated 1 Hz rTMS. We are aware of only one small (n = 7) negative trial with limited stimulation course and duration (18 sessions over 10 days) [35]. Our results are encouraging, but overall limited response and remission rates after the arTMS course would suggest that increasing the number of sessions per day and the number of pulses delivered at each session may be necessary [5]. Indeed, a recent arTMS pilot (n = 6) with highly refractory MDD patients using intermittent theta-burst stimulation (iTBS), 1800 pulses/session, 10 sessions per day over 5 days, reported remission rates of 66% [14]. A larger follow-up study (n = 21) in a less refractory MDD population reported remission rates of up to 90% [36]. The rapid improvement in responders *post-acutely* (**Fig. 5**) is

encouraging, showing that it is indeed possible to rapidly accelerate response to 1 Hz stimulation in some individuals. This could justify the further study of this rTMS technique in more severe inpatient cases, where suicidality is of concern and rapid response is at a premium. Overall, final outcomes *post-taper* compared favorably to another large trial [19], as well as a landmark meta-analysis [37], which may be explained by the high number of total sessions. Supporting this observation, a recent study that offered up to 21 weeks of rTMS reported remission rates of 72% [38]. Since there is a lack of data about 1 Hz arTMS alone without the subsequent tapering period, we are unable to determine the impact or the necessity of such an extension. It may well be possible that 1 Hz arTMS has a delayed effect, where participants would have kept on improving without the extension course, as suggested in other arTMS protocols [12]. Conversely, high relapse rates following arTMS are also of concern [14,39].

We also included data regarding suicidality scores for each scale since arTMS has also been studied in the treatment of acute suicidality [6,8]. In the future, suicidal ideation should be assessed with either the Beck Scale of Suicide ideation (BSI) or the Columbia-Suicide Severity Rating Scale (C-SSRS).

This preliminary study has several limitations. Of primary concern is the open-label design and the absence of a sham control arm. It is to be expected that the various estimates of effectiveness might be higher than what would be obtained in a randomized, sham-controlled trial, which would represent the next logical step. In addition, the naturalistic approach of the study does not allow estimates of efficacy, but only effectiveness. The design of the study, with multiple reassessments throughout, presented logistical challenges, mainly with patient adherence. Indeed, several participants did not adhere to the planned schedule, and were thus reassessed outside of the planned boundaries. Additionally, we did not qualitatively assess our open-room setting, as discussed previously. We also had 10 participants lost to follow-up at 1-month, limiting interpretation of results at that timepoint. In future studies, this would need to be

corrected to achieve more standardized measures. Furthermore, frequency of tapering course varied between individuals. Although we recommended daily sessions for participants (5 sessions/week), a minimum of 3 sessions/week was tolerated, given the fact that this approach has proven effective on its own [40]. Moreover, results did vary between the self-rated BDI-II and the clinician-rated HRSD-17. Indeed, remission and response rates were higher *post-acutely* and lower *post-tapering* and at *1-month follow-up* visits on the BDI-II compared to HRSD-17. Discrepancies between self- and clinician-rated scales are not unique to our studies and have often been reported in MDD [41], showing the general limitation of mood scales, and the need for objective biomarkers in psychiatry [42]. As a further matter, the weaker central field in the parabolic coils used may have under-stimulated the DLPFC [20]; this may be correctable in future by moving the central opening to an intermediate mark between F4, Fz and Cz. Due to their less widespread use in clinical settings, results with 1 Hz rTMS on parabolic coils may not generalize to more commonly used Fo8 coils. The 1 Hz protocol used, although highly tolerable for participants, may also have been less effective due to its limited amount of pulses per session. Given that increased number of pulses has been associated with increased response in 1 Hz rTMS in some studies [43], it may be advantageous to increase the number of pulses within the same time frame (e.g., 600 pulses in 10 min). Finally, although we did not require participants to meet the usual requirement of TRD in our trial, the majority (>80%) of participants had failed at least one adequate trial. We also did not have a minimal cut-off regarding depression severity on the mood scales for study inclusion, but average baseline scores on the BDI-II was in the severe range. Average HRSD-17 score was also higher than in a recent large trial from our group [19].

Given its established efficacy in MDD and its lower side-effect profile compared to medication, efforts should be made to make rTMS more accessible to the general public. The current trend depicts increased costs and complexity associated with rTMS administration [14,36]. However, the techniques demonstrated here may be suitable for safe use in a wider

range of settings, and potentially could be adapted in future to enable home use of 1 Hz rTMS. Besides eliminating the need for in-clinic visits, particularly relevant during the current COVID-19 pandemic [44], home-based rTMS could potentially be delivered for as little as $10/day, comparable to some medication regimens. Thus, if the presently described technique does indeed achieve efficacy while preserving superior safety and tolerability, rTMS may serve as a viable first-line treatment technique for treatment-naïve MDD patients or a home-based maintenance technique for rTMS responsive patients.

## CONFLICTS OF INTEREST

The authors declare no financial interests relative to this work. **JPM** reports research grants from the Brain & Behavior Research Foundation NARSAD Young Investigator Award and salary support for his graduate studies from the Branch Out Neurological Foundation. **HV**, **LF**, **MH**, **FM**, **SD**, **RZ, JS** and **AMM** do not report any conflict of interest. **DMB** receives research support from CIHR, NIH, Brain Canada and the Temerty Family through the CAMH Foundation and the Campbell Family Research Institute. He received research support and in-kind equipment support for an investigator-initiated study from Brainsway Ltd. He is the site principal investigator for three sponsor-initiated studies for Brainsway Ltd. He also receives in-kind equipment support from Magventure for investigator-initiated research. He received medication supplies for an investigator-initiated trial from Indivior. **ZJD** has received research and equipment in-kind support for an investigator-initiated study through Brainsway Inc and Magventure Inc. His work was supported by the Ontario Mental Health Foundation (OMHF), the Canadian Institutes of Health Research (CIHR), the National Institutes of Mental Health (NIMH) and the Temerty Family and Grant Family and through the Centre for Addiction and Mental Health (CAMH) Foundation and the Campbell Institute. **FVR** receives research support from CIHR, Brain Canada, Michael Smith Foundation for Health Research, Vancouver Coastal Health Research Institute, and in-kind equipment support for this investigator-initiated trial from

MagVenture. He has received honoraria for participation in advisory board for Janssen. **JD** reports research grants from CIHR, the National Institute of Mental Health, Brain Canada, the Canadian Biomarker Integration Network in Depression, the Ontario Brain Institute, the Weston Foundation, the Klarman Family Foundation, the Arrell Family Foundation, and the Buchan Family Foundation, travel stipends from Lundbeck and ANT Neuro, in-kind equipment support for investigator-initiated trials from MagVenture, and is an advisor for BrainCheck, TMS Neuro Solutions, and Restorative Brain Clinics.

**ACKNOWLEDGEMENT**

**JPM** would like to thank the **Brain & Behavior Research Foundation** and the **Branch Out Neurological Foundation** for their financial support of this project. We would like to thank **Terri Cairo**, **Julian Kwok**, **Meaghan Todd**, **Nuno Ferreira**, **Thomas Russell** and **Eileen Lam** for their involvement and organizational support throughout this project.

## REFERENCES

[1] Lam RW, McIntosh D, Wang J, Enns MW, Kolivakis T, Michalak EE, et al. Canadian Network for Mood and Anxiety Treatments (CANMAT) 2016 Clinical Guidelines for the Management of Adults with Major Depressive Disorder. Can J Psychiatry 2016;61:510–23. doi:10.1177/0706743716659416.

[2] Kennedy SH, Lam RW, McIntyre RS, Tourjman SV, Bhat V, Blier P, et al. Canadian Network for Mood and Anxiety Treatments (CANMAT) 2016 Clinical Guidelines for the Management of Adults with Major Depressive Disorder. Can J Psychiatry 2016;61:540–60. doi:10.1177/0706743716659417.

[3] Milev RV, Giacobbe P, Kennedy SH, Blumberger DM, Daskalakis ZJ, Downar J, et al. Canadian Network for Mood and Anxiety Treatments (CANMAT) 2016 Clinical Guidelines for the Management of Adults with Major Depressive Disorder. Can J Psychiatry 2016;61:561–75. doi:10.1177/0706743716660033.

[4] Mendlowitz AB, Shanbour A, Downar J, Vila-Rodriguez F, Daskalakis ZJ, Isaranuwatchai W, et al. Implementation of intermittent theta burst stimulation compared to conventional repetitive transcranial magnetic stimulation in patients with treatment resistant depression: A cost analysis. Plos One 2019;14:e0222546-13. doi:10.1371/journal.pone.0222546.

[5] Baeken C. Accelerated rTMS: A Potential Treatment to Alleviate Refractory Depression. Front Psychol 2018;9:2017. doi:10.3389/fpsyg.2018.02017.

[6] George MS, Raman R, Benedek DM, Pelic CG, Grammer GG, Stokes KT, et al. A Two-site Pilot Randomized 3 Day Trial of High Dose Left Prefrontal Repetitive Transcranial Magnetic Stimulation (rTMS) for Suicidal Inpatients. Brain Stimulation 2014;7:421–31. doi:10.1016/j.brs.2014.03.006.

[7] Dardenne A, Baeken C, Crunelle CL, Bervoets C, Matthys F, Herremans SC. Accelerated HF-rTMS in the elderly depressed: A feasibility study. Brain Stimul 2018;11:247–8. doi:10.1016/j.brs.2017.10.018.

[8] Desmyter S, Duprat R, Baeken C, Autreve SV, Audenaert K, Heeringen K van. Accelerated Intermittent Theta Burst Stimulation for Suicide Risk in Therapy-Resistant Depressed Patients: A Randomized, Sham-Controlled Trial. Frontiers in Human Neuroscience 2016;10:480. doi:10.3389/fnhum.2016.00480.

[9] Duprat R, Desmyter S, Raedt RD, Heeringen K van, Abbeele DV den, Tandt H, et al. Accelerated intermittent theta burst stimulation treatment in medication-resistant major depression_ A fast road to remission? J Affect Disorders 2016;200:6–14. doi:10.1016/j.jad.2016.04.015.

[10] Holtzheimer PE, McDonald WM, Mufti M, Kelley ME, Quinn S, Corso G, et al. Accelerated repetitive transcranial magnetic stimulation for treatment-resistant depression. Depress Anxiety 2010;27:960–3. doi:10.1002/da.20731.

[11] Fitzgerald PB, Hoy KE, Elliot D, McQueen RNS, Wambeek LE, Daskalakis ZJ. Accelerated repetitive transcranial magnetic stimulation in the treatment of depression. Neuropsychopharmacol 2018;43:1–8. doi:10.1038/s41386-018-0009-9.

[12] Sonmez AI, Camsari DD, Nandakumar AL, Voort JLV, Kung S, Lewis CP, et al. Accelerated TMS for Depression: A systematic review and meta-analysis. Psychiat Res 2019;273:770–81. doi:10.1016/j.psychres.2018.12.041.

[13] McGirr A, Eynde FV den, Tovar-Perdomo S, Fleck MPA, Berlim MT. Effectiveness and acceptability of accelerated repetitive transcranial magnetic stimulation (rTMS) for treatment-resistant major depressive disorder_ An open label trial. J Affect Disorders 2015;173:216–20. doi:10.1016/j.jad.2014.10.068.

[14] Williams NR, Sudheimer KD, Bentzley BS, Pannu J, Stimpson KH, Duvio D, et al. High-dose spaced theta-burst TMS as a rapid-acting antidepressant in highly refractory depression. Brain 2018;141:e18–e18. doi:10.1093/brain/awx379.

[15] Baeken C, Vanderhasselt M-A, Remue J, Herremans S, Vanderbruggen N, Zeeuws D, et al. Intensive HF-rTMS treatment in refractory medication-resistant unipolar depressed patients. Journal of Affective Disorders 2013;151:625–31. doi:10.1016/j.jad.2013.07.008.

[16] Schulze L, Feffer K, Lozano C, Giacobbe P, Daskalakis ZJ, Blumberger DM, et al. Number of pulses or number of sessions? An open-label study of trajectories of improvement for once- vs. twice-daily dorsomedial prefrontal rTMS in major depression. Brain Stimul 2017;11:1–10. doi:10.1016/j.brs.2017.11.002.

[17] Perera T, George MS, Grammer G, Janicak PG, Pascual-Leone A, Wirecki TS. The Clinical TMS Society Consensus Review and Treatment Recommendations for TMS Therapy for Major Depressive Disorder. Brain Stimul 2016;9:336–46. doi:10.1016/j.brs.2016.03.010.

[18] Kaster TS, Downar J, Vila-Rodriguez F, Thorpe KE, Feffer K, Noda Y, et al. Trajectories of Response to Dorsolateral Prefrontal rTMS in Major Depression: A THREE-D Study. Am J Psychiat 2019;176:appi.ajp.2018.1-9. doi:10.1176/appi.ajp.2018.18091096.

[19] Blumberger D, Vila-Rodriguez F, Thorpe K, Feffer K, Noda Y, Giacobbe P, et al. Effectiveness of theta burst versus high-frequency repetitive transcranial magnetic stimulation in patients with depression (THREE-D): a randomised non-inferiority trial. Lancet 2018;391:1683–92. doi:10.1016/s0140-6736(18)30295-2.

[20] Miron J-P, Voetterl H, Mansouri F, Blumberger DM, Daskalakis ZJ, Downar J. A case series of a novel 1 Hz right-sided dorsolateral prefrontal cortex rTMS protocol in major depression. Brain Stimul 2019. doi:10.1016/j.brs.2019.11.006.

[21] McClintock SM, Reti IM, Carpenter LL, McDonald WM, Dubin M, Taylor SF, et al. Consensus Recommendations for the Clinical Application of Repetitive Transcranial Magnetic Stimulation (rTMS) in the Treatment of Depression. J Clin Psychiatry 2017;79:1–16. doi:10.4088/jcp.16cs10905.

[22] Brunelin J, Jalenques I, Trojak B, Attal J, Szekely D, Gay A, et al. The efficacy and safety of low frequency repetitive transcranial magnetic stimulation for treatment-resistant depression: the results from a large multicenter French RCT. Brain Stimul 2014;7:855–63. doi:10.1016/j.brs.2014.07.040.

[23] Riedel M, Möller H-J, Obermeier M, Schennach-Wolff R, Bauer M, Adli M, et al. Response and remission criteria in major depression--a validation of current practice. J Psychiatr Res 2010;44:1063–8. doi:10.1016/j.jpsychires.2010.03.006.

[24] Zimmerman M, Posternak MA, Chelminski I. Derivation of a definition of remission on the Montgomery–Asberg depression rating scale corresponding to the definition of remission on the Hamilton rating scale for depression. J Psychiatr Res 2004;38:577–82. doi:10.1016/j.jpsychires.2004.03.007.

[25] Lefaucheur J-P, Aleman A, Baeken C, Benninger DH, Brunelin J, Lazzaro VD, et al. Evidence-based guidelines on the therapeutic use of repetitive transcranial magnetic stimulation (rTMS): An update (2014–2018). Clinical Neurophysiology 2020. doi:10.1016/j.clinph.2019.11.002.

[26] Sun W, Mao W, Meng X, Wang D, Qiao L, Tao W, et al. Low-frequency repetitive transcranial magnetic stimulation for the treatment of refractory partial epilepsy: a controlled clinical study. Epilepsia 2012;53:1782–9. doi:10.1111/j.1528-1167.2012.03626.x.

[27] Kaur M, Michael JA, Fitzgibbon BM, Hoy KE, Fitzgerald PB. Low-frequency rTMS is better tolerated than high-frequency rTMS in healthy people: Empirical evidence from a single session study. J Psychiatr Res 2019;113:79–82. doi:10.1016/j.jpsychires.2019.03.015.

[28] Cao X, Deng C, Su X, Guo Y. Response and Remission Rates Following High-Frequency vs. Low-Frequency Repetitive Transcranial Magnetic Stimulation (rTMS) Over Right DLPFC for Treating Major Depressive Disorder (MDD): A Meta-Analysis of Randomized, Double-Blind Trials. Frontiers Psychiatry 2018;9:1106–7. doi:10.3389/fpsyt.2018.00413.

[29] Brunoni AR, Chaimani A, Moffa AH, Razza LB, Gattaz WF, Daskalakis ZJ, et al. Repetitive Transcranial Magnetic Stimulation for the Acute Treatment of Major Depressive Episodes. Jama Psychiat 2017;74:143–10. doi:10.1001/jamapsychiatry.2016.3644.

[30] Fitzgerald PB, Hoy KE, Reynolds J, Singh A, Gunewardene R, Slack C, et al. A pragmatic randomized controlled trial exploring the relationship between pulse number and response to repetitive transcranial magnetic stimulation treatment in depression. Brain Stimul 2019. doi:10.1016/j.brs.2019.09.001.

[31] Levkovitz Y, Isserles M, Padberg F, Lisanby SH, Bystritsky A, Xia G, et al. Efficacy and safety of deep transcranial magnetic stimulation for major depression: a prospective multicenter randomized controlled trial. World Psychiatry 2015;14:64–73. doi:10.1002/wps.20199.

[32] Rao VR, Sellers KK, Wallace DL, Lee MB, Bijanzadeh M, Sani OG, et al. Direct Electrical Stimulation of Lateral Orbitofrontal Cortex Acutely Improves Mood in Individuals with Symptoms of Depression. Curr Biol 2018;28:1–25. doi:10.1016/j.cub.2018.10.026.

[33] Downar J. Orbitofrontal Cortex: A `Non-rewarding' New Treatment Target in Depression? Curr Biol 2019;29:R59–62. doi:10.1016/j.cub.2018.11.057.

[34] Rolls ET. A non-reward attractor theory of depression. Neurosci Biobehav Rev 2016;68:47–58. doi:10.1016/j.neubiorev.2016.05.007.

[35] Tor P-C, Galvez V, Goldstein J, George D, Loo CK. Pilot Study of Accelerated Low-Frequency Right-Sided Transcranial Magnetic Stimulation for Treatment-Resistant Depression. J Ect 2016;32:180–2. doi:10.1097/yct.0000000000000306.

[36] Cole EJ, Stimpson KH, Bentzley BS, Gulser M, Cherian K, Tischler C, et al. Stanford Accelerated Intelligent Neuromodulation Therapy for Treatment-Resistant Depression. Am J Psychiatry 2020:appiajp201919070720. doi:10.1176/appi.ajp.2019.19070720.

[37] Berlim MT, Eynde F van den, Daskalakis ZJ. A systematic review and meta-analysis on the efficacy and acceptability of bilateral repetitive transcranial magnetic stimulation (rTMS) for treating major depression. Psychol Med 2012;43:2245–54. doi:10.1017/s0033291712002802.

[38] Stubbeman WF, Zarrabi B, Bastea S, Ragland V, Khairkhah R. Bilateral neuronavigated 20Hz theta burst TMS for treatment refractory depression: An open label study. Brain Stimul 2018;11:953–5. doi:10.1016/j.brs.2018.04.012.

[39] Caulfield KA. Is Accelerated, High-Dose Theta Burst Stimulation a Panacea for Treatment-Resistant Depression? J Neurophysiol 2019;3:185. doi:10.1152/jn.00537.2019.

[40] Galletly C, Gill S, Clarke P, Burton C, Fitzgerald PB. A randomized trial comparing repetitive transcranial magnetic stimulation given 3 days/week and 5 days/week for the treatment of major depression: is efficacy related to the duration of treatment or the number of treatments? Psychol Med 2012;42:981–8. doi:10.1017/s0033291711001760.

[41] Dunlop BW, Li T, Kornstein SG, Friedman ES, Rothschild AJ, Pedersen R, et al. Concordance between clinician and patient ratings as predictors of response, remission, and recurrence in major depressive disorder. J Psychiatr Res 2010;45:96–103. doi:10.1016/j.jpsychires.2010.04.032.

[42] Kennis M, Gerritsen L, Dalen M van, Williams A, Cuijpers P, Bockting C. Prospective biomarkers of major depressive disorder: a systematic review and meta-analysis. Mol Psychiatr 2019;25:321–38. doi:10.1038/s41380-019-0585-z.

[43] Berlim MT, Eynde FV den, Daskalakis ZJ. Clinically Meaningful Efficacy and Acceptability of Low-Frequency Repetitive Transcranial Magnetic Stimulation (rTMS) for Treating Primary

Major Depression: A Meta-Analysis of Randomized, Double-Blind and Sham-Controlled Trials. Neuropsychopharmacol 2012;38:543–51. doi:10.1038/npp.2012.237.

[44] Caulfield KA, George MS. Treating the Mental Health Effects of COVID-19: The Need for At-Home Neurotherapeutics Is Now. Brain Stimul 2020. doi:10.1016/j.brs.2020.04.005.

| | |
|---|---|
| Age, years | 41.8 (12.2) |
| Women | 56.3% |
| Education, years | 17.1 (3.2) |
| Left-handed | 6.3% |
| Age of onset, years | 23.8 (10.8) |
| Length of current depressive episode, months | 51.7 (78.2) |
| Comorbid anxiety | 81.3% |
| Currently working | 64.6% |
| Baseline BDI-II | 35.0 (9.8) |
| Baseline HRSD-17 | 23.8 (5.3) |
| Receiving psychopharmacotherapy during treatment | 70.8% |
| Antidepressant | 58.3% |
| Antidepressant combination | 22.9% |
| Antipsychotic augmentation | 14.6% |
| Lithium augmentation | 4.2% |
| Psychostimulant augmentation | 14.6% |
| T3 augmentation | 2.1% |
| Benzodiazepine | 18.8% |
| Number of past antidepressant trials, lifetime | 5.2 (3.6) |
| ATHF total score | 4.2 (4.3) |
| ATHF number of trials, current episode | 1.4 (1.1) |
| ATHF highest score | 2.6 (1.6) |
| Data are mean (SD) or number of participants (% of total). BDI-II = Beck Depression Inventory-II, HRSD-17 = 17-item Hamilton Rating Scale for Depression, ATHF = Antidepressant Treatment History Form. | |

*Table 1*: Baseline demographic and clinical characteristics (n = 48)

| BDI-II | |
|---|---|
| Response *post-acute* | 12/48 (25.0%) |
| Response *post-taper* | 17/48 (35.4%) |
| Response *1-month* | 10/35 (28.6%) |
| Remission *post-acute* | 8/48 (16.7%) |
| Remission *post-taper* | 13/48 (27.1%) |

| BDI-II | |
|---|---|
| Remission *1-month* | 8/35 (22.9%) |
| Score baseline | 35.0 (9.8) |
| Score change *post-acute* | 25.8 (12.2) |
| Score change *post-taper* | 22.2 (13.1) |
| Score change *1-month* | 21.5 (12.1) |
| Percent improvement *post-acute* | 27.4% (27.8%) |
| Percent improvement *post-taper* | 37.9% (33.0%) |
| Percent improvement *1-month* | 36.1% (34.7%) |
| HRSD-17 | |
| Response *post-acute* | 11/48 (22.9%) |
| Response *post-taper* | 28/48 (58.3%) |
| Response *1-month* | 21/35 (60.0%) |
| Remission *post-acute* | 3/48 (6.3%) |
| Remission *post-taper* | 18/48 (37.5%) |
| Remission *1-month* | 13/35 (37.1%) |
| Score baseline | 23.8 (5.3) |
| Score change *post-acute* | 16.3 (6.6) |
| Score change *post-taper* | 12.2 (8.0) |
| Score change *1-month* | 11.3 (7.4) |
| Percent improvement *post-acute* | 31.4% (23.8%) |
| Percent improvement *post-taper* | 48.6% (31.9%) |
| Percent improvement *1-month* | 52.4% (28.5%) |

Data are mean (SD). For remission and response rates, data are n (% of participants assessed). BDI-II = Beck Depression Inventory-II, HRSD-17 = 17-item Hamilton Rating Scale for Depression.

*Table 2*: Response and remission rates, changes in depression severity scores and percent improvement throughout the study

| | |
|---|---|
| Remission *post-acute* | 12/31 (38.7%) |
| Remission *post-taper* | 15/31 (48.4%) |
| Remission *1-month* | 13/23 (56.5%) |
| HRSD-17 | |
| Remission *post-acute* | 16/38 (42.1%) |
| Remission *post-taper* | 20/38 (52.6%) |
| Remission *1-month* | 17/28 (60.7%) |
| Data are n (% of participants assessed). Suicidality item is #9 on the BDI-II and #3 on the HRSD-17. BDI-II = Beck Depression Inventory-II, HRSD-17 = 17-item Hamilton Rating Scale for Depression. | |

**Table 3**: Remission rates regarding suicidality

| | |
|---|---|
| Serious AE | 0/48 (0.0%) |
| AE total | 19/48 (39.6%) |
| Headache | 11/48 (22.9%) |

| | |
|---|---|
| Fatigue | 7/48 (14.6%) |
| Nausea | 2/48 (4.2%) |
| Scalp tenderness | 3/48 (4.2%) |
| Hand twitching | 1/48 (2.1%) |
| Insomnia | 3/48 (6.3%) |
| Dizziness | 1/48 (2.1%) |
| Anxiety | 3/48 (6.3%) |
| Jaw pain | 1/48 (2.1%) |
| Visual symptoms | 1/48 (2.1%) |
| First treatment pain VRS | 5.6 (2.0) |
| Last treatment pain VRS | 1.9 (1.9) |
| Number of days to reach target stimulation intensity | 3.8 (6.3) |
| Number of participants (n=48) reporting adverse events (AE - %). For pain, data mean (SD). VRS = Verbal Rating Scale. | |

**Table 4**: Adverse events

**Figure 1**: Clinical setup - 4 recliners equipped with AAAs holding rTMS coils, connected to the rTMS stimulators. Treatment was delivered in an open room setting, allowing up to 4 participants to receive treatment simultaneously, with the help of one or two technicians. AAA = Anchored Articulating Arm

**Figure 2**: Anchored Articulating Arm (AAA) schematic - AAAs are made out of inexpensive camera tripod equipment and are anchored at their base to the corner of the treatment chair. Coils are held securely by a clamp locking it to the AAA, maintaining position until treatment has been completed, at which time they can be quickly twisted out of the way.

**Figure 3**: Study timeline - Treatment consisted of an arTMS course of 6 sessions/day (50 min inter-sessions intervals) over 5 days (on weekdays), thus totalling 30 sessions. After a 1-week gap, arTMS was followed by a tapering course of once-daily stimulation (minimum of 20, maximum of 25 sessions), 3-5 days per week, to decrease the odds of relapse. Baseline visits were conducted the week prior to arTMS initiation and consisted of a clinical assessment by a trained research staff, navigation cap fitting and motor threshold calibration. Patients were clinically reassessed at 3 subsequent timepoints: *post-acute* visits were conducted during the one-week gap following arTMS; *post-taper* visits were conducted within 7 days of tapering termination; *1-month follow-up* visits were conducted 1-month (± 7 days) after the last tapering session. arTMS = accelerated repetitive transcranial magnetic stimulation, ATHF = Average Antidepressant Treatment History Form, MINI = Mini International Neuropsychiatric Interview, BDI-II = Beck Depression Inventory-II, HRSD-17 = 17-item Hamilton Rating Scale for Depression

**Figure 4**: Trial CONSORT flow diagram - arTMS = accelerated repetitive transcranial magnetic stimulation.

**Fig. 5.** Trajectories of improvement on the BDI-II. Responders showed rapid improvement during the accelerated course, having achieved response on average response post-acutely, and continued to show slow but steady additional improvement up to the end of the tapering course. Use of background shading delineates arTMS from tapering course. BDI-II = the Beck Depression Inventory – II, arTMS = accelerated repetitive transcranial magnetic stimulation, Tx = treatment (rTMS session)