Keen relationship between COVID_19 and food supply suggest some animal origin foods as excellent vehicle of SARS-Cov-2

Nazli Saeedi [1], Seyed Abbas Rafat[2*]

1 Research Center for Pharmaceutical Nanotechnology, Biomedicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran Email : saeedi@hotmail.fr, 2 Department of Animal Science, Faculty of Agriculture, University of Tabriz, Tabriz, Iran, Email;

abbasrafat@hotmail.com *Corresponding Author

Abstract

In recent months, nearly all countries tried to decrease human-to-human contact as the principal mode of transmission of SARS-CoV-2. However, other modes of transmission also need to be clarified in more depth, especially, the foodborne transmission. We assessed the effect of animal origin foods consumption on the pandemic of COVID-19. For this purpose, we studied the relationship among 20 food supply as independent variables, and the parameter of Total Cases as dependent variable. Here we show a relationship between a group of animal origin foods and total cases. Regression, Bayes, and Lasso results showed that eggs and fresh water fish have positive coefficient. So, among the transmission ways of COVID_19, the role of foodborne transmission should be more significant than previously thought. The possibility of animal origin foodborne transmission should be taken into more consideration. The perspective is to expand the surveillance of SARS-Cov-2 during the food production chain. In conclusion, the results of the present study indicate that one important vehicle for SARS-Cov-2 may be some of animal origin

22 foods. It is recommended that virologists examine the possibility of freshwater fish and chicken's eggs

23 being as excellent vehicles/preservatives for SARS-Cov-2.

Introduction

At the time of writing this manuscript, the number of total cases of COVID-19 past 28 mark.

During the pandemic, most countries around the world tried to minimize human-to-human

contact and travels(Merler et al., 2020). However, the prevalence of COVID-19 continued.

Despite efforts of many countries to reduce the transmission by person-to-person way the rapid

outbreak shows the necessity of research on other possible transmission routes. During early

pandemics, in Italy and the USA, for example, the chances of being surprised by pandemics were

low, but both countries faced epidemic dramatically. Among the various routes of transmission,

FBT is most likely. We suppose the role of COVID-19's transmission by other ways should be

more significant than previously thought. Transmission of MERS, a similar pathogenic virus to

SARS-Cov-2, through food has strengthened our hypothesis.

Furthermore, some researchers cited the possibility of FBT for SARS-Cov-2(Hi, 2020;

Kingsbury et al., 2020; Newell et al., 2010; Panel and Biohaz, 2011; Todd et al., 2020). Also,

FAO/WHO experts mentioned the data gap in the relationship between viruses and food borne

disease (FAO/WHO, 2008) . The motivation for the present study is to find the effects of human

food consumptions (independent variables) on COVID_19 (As dependent variable).

Material and Methods

Table 1 shows descriptive statistics of studied variables. We assessed the possible relationship

between the consumption of main foods and total cases od covid_19. Accordingly, we studied

total cases as a dependent variable and 20 food supply data as independent variables. The

primary purpose of the study was to find effect of anima origin food on total cases, however, we

considered six plant origin foods for comparison reasons. The statistical methods included

correlation, stepwise and LASSO regression, and Bayes analysis. Food supply (kg/capita/yr.)

was quantity in each country based on the latest available data of FAO. Adjusted R-square of

fitted models are presented in Table 2, respectively. Detail material and methods are included in

supplementary materials.

Table 1

Descriptive statistics of studied foods (n=20) besides total cases.

#	Variable (kg/capita/yr)	Abbreviation	N	Mean	Std Dev	Minimum	Maximum
1	**Pig meat**	pig	169	14.66	15.52	0.00	71.48
2	**Bovine Meat**	bov	174	11.20	9.00	0.78	55.48
3	**Cheese**	che	173	4.97	6.55	0.00	30.82
4	**Cream**	cre	159	1.21	2.76	0.00	18.45
5	**Eggs**	Egg	174	7.08	5.09	0.16	19.15
6	**Freshwater Fish**	freshf	174	3.38	4.54	0.00	34.00
7	**Honey**	hon	168	0.36	0.46	0.00	3.51
8	**Whole milk**	milk	174	59.17	48.93	0.96	258.44
9	**Poultry Meat**	poul	174	20.56	16.46	0.42	72.87
10	**Demersal Fish**	demer	173	3.40	4.93	0.00	33.45
11	**Cephalopods**	ceph	171	0.44	0.96	0.00	6.70
12	**Butter, Ghee**	butg	174	1.25	1.64	0.00	9.25
13	**Marine Fish, Other**	marf	172	2.74	4.87	0.00	28.19
14	**Aquatic Plants**	aqp	162	0.20	1.91	0.00	22.41
15	**Beverages, Alcoholic**	bev	171	3.20	3.87	0.00	26.90
16	**Rice(Milled Equivalent)**	rice	174	30.95	36.20	0.95	171.73
17	**Potatoes and products**	potato	174	34.89	33.65	0.05	183.16
18	**Maize and products**	maiz	170	21.71	28.52	0.00	158.40
19	**Wheat and products**	wheat	174	70.92	46.85	1.89	222.40
20	**wine**	wine	169	6.27	9.49	0.00	51.41
	Total Cases (Per Country)	TC	215	130744.74	619823.34	3	6558095

69 **Table 2**

70 **R-Square, Adjusted R-square, AIC and number of data in the fitted models**

71 **for food consumption.**

	Stepwise	LASSO	BAYES
R-Square	0.33	0.47	-
Adj-R Square	0.31	0.42	-
AIC	-	313	562
n	131	131	130

We estimated the relationship between 20 food supply and Total Cases of covid19. The methods include Stepwise Regression with Proc Reg, LASSO regression with GLMSELECT Proc, and Bayes analysis with Proc Genmod (SAS). Food supply was quantity (kg/capita/yr.). Fourteen foods of animal origin and six foods of plant origin were selected. The primary purpose of the study was to find AOF relationship with total cases, but some plant origin foods were chosen for comparison reason.

We started our analyses first time on March 2020. Then we repeated it several times with updated COVID_19 data on April, May, June, and September. Finally we presented the latest results.

Data

We downloaded the data the following websites:

COVID_19: https://www.worldometers.info/coronavirus/ 10 September 2020

Crops Primary Equivalent: http://www.fao.org/faostat/en/#data/CC

Food Supply - Livestock and Fish: http://www.fao.org/faostat/en/#data/CL

We used base-e logs, also known as natural logs for the transformation of total cases and independent variables. In all analyses, the accuracy of the model with non-transformed data was very low. So, all variables has been transformed to

normalize the distributions. With transformation of data, the accuracy of the model

increased to an acceptable level. In comparison of models, Lasso's results are more

reliable than stepwise regression and Bayes, because it's AIC was smaller and also

it had higher accuracy as R-square (Table 2). Visual examination of the trace plot

displays proper mixing in Bayesian analysis.

We provide the SAS code for an application of the used statistical methods (Rafat,

2020). We note that this work was posted as preprint on medrxiv (Rafat, seyed

abbas, 2020) .

Results and Discussion

Stepwise Regression, Bayes, and Lasso's results confirmed that eggs and fresh

water fish have high positive relevance with total cases. With our results in Table

3, implying the need for tracing of SARS-COV-2 in main AOF. Bayes results, e.g.,

show there is a 0.99 probability of a positive correlation between eggs and total

cases, adjusted for the other covariates. The number of AOF and their role in our

results varies depending on the type of statistical method used, but all results are

consistent. The results of all three statistical methods confirm each other in the

case of eggs. Negative high correlation between marine fish and total cases is

curious observation needs more study.

Table 3. The results of multivariate analyses. All variables are logarithmic transformed data.

Variable	Stepwise	Lasso	Bayes
Pig meat		-0.26	0.36
Bovine Meat		0.12	0.75
Cheese			0.35
Cream		-0.18	0.19
Eggs	**0.85**	**0.98**	**0.99**
Freshwater fish		**0.72**	**0.99**
Honey		-1.18	0.03
Whole milk			0.34
Poultry meat			0.38
Demersal fish			0.47
Cephalopods			0.36
Butter/ghee		0.19	0.84
Marine fish(other)	-1.49	-1.32	-1.00
Aquatic plants			0.38
Beverages(Alcoholic)			0.18
Rice(Milled Equivalent)		0.30	0.95
Potatoes and products		0.37	0.96
Maize and products		0.39	0.99
Wheat and products		0.34	0.91
Wine			0.64

119
120

The effect of a group of AOF on total cases was significant in our analysis. So, we suppose a noticeable effect of AOF as FBT/vehicle. At present, we are not able to interpret all significant effects and their coefficients. Interpretation of the factors could be the subject of future researches. Potential transmission ways of COVID-19 based on the results of our analysis and bibliography is summarized as Fig 1. Lake and Kingsbury (2020) provided a review about potential for FBT of COVID-19, but they could not guarantee the application of the data found on other coronaviruses to SARS-CoV-2(Kingsbury et al., 2020)**.** The persistence of viruses by refrigeration temperature(FAO/WHO, 2008) might be an explanation of the relationship between COVID-19 and AOF. The viral persistence of coronavirus in soft cheese (Panon et al., 1988) and raw milk(Hamdi, 2013) has been reported. We don't know what's going on from "Farm to Fork" about AOF. For example, SARS-CoV-2 may be more resistant to heat when cooking, as it is observed for poliovirus in beef(Pirtle, 1991). It is recommended not to consume raw or undercooked animal products(Moy, 2020). Kitchen utensils can transmit the virus among foods(Sarah C. P. Williams, n.d.), accordingly, AOF may be a suitable vehicle, and the virus inside them will survive better and longer.

Furthermore, the methods of diagnosing the virus in food are not complete, and there are many limitations(Bosch et al., 2018). In food science technology, the

primary attention paid to bivalve moullouses. Nowadays, it is necessary to expand

the detection of viruses to all ANIMAL ORIGIN FOOD.

The high-fat tissue in the body has effect on the pathogenesis of COVID-

19(Maffetone and Laursen, 2020). Although the probability of transmitting the

virus through wine is zero, we deliberately kept it in the initial model of our

analysis. Results showed wine is eliminated among significant variables of the

suited model (Table 3). So we concluded that just the energy content of food

should not keep a variable in the fitted model. So, in addition to energy content,

there must be something else in food-related COVID-19. We concentrate on the

transmission of COVID-19 by AOF, rather than the immunity response.

The paper of (Katy Askew, 2020) guided us to construct our hypothesis about the

essential relationship between COVID-19 and AOF include.

WHO's recommendations state that COVID_19 have not FBT, and suitable food

safety practices could prevent the probability of FBT. However, the realization of

proper food safety practices for all AOF is questionable. It is reported that

acquisition of SARS coronavirus by consumption of contaminated food is possible

(Yepiz 2013). Furthurmore, oral infection of SARS-Cov-2 could not be excluded

among transmission ways (FAO/WHO 2008). Promkuntod 2006 recommends

taking precautions to prevent the spread of the avian flu by consumption of quail

eggs(PROMKUNTOD et al., 2006). One explanation for our observation is the

trapping and accumulation of sars-cov-2 by AOF. Thereafter, along food

processing or preparation, AOF conserve SARS-Cov-2 infectively.

Conclusion

Despite the awareness and readiness of many countries for COVID-19 during early

2020, why did it spread so quickly? It seems FBT has received less attention

relative to other routes. The perspective is to expand the surveillance of SARS-

Cov-2 during the food production chain. It is recommended that virologists

examine the possibility of freshwater fish and chicken's eggs being as an excellent

vehicle/preservator for SARS-Cov-2.Some animal origin food may have been the

first cause of transmission of the SARS-Cov-2 from reservoirs populations to

humans.

ACKNOWLEDGEMENTS:

Author contributions: Conceptualization and Visualisation: N. S. K., Methodology: S. A. R.

Writing: Both authors. The authors have declared no competing interest. Data and materials

availability: All data are available in the manuscript or the supplementary materials.

References

Bosch, A., Gkogka, E., Le, F.S., Loisy-hamon, F., Lee, A., Lieshout, L. Van, Marthi, B.,

    Myrmel, M., Sansom, A., Charlotte, A., Winkler, A., Zuber, S., Phister, T., 2018.

    International Journal of Food Microbiology Foodborne viruses□: Detection , risk

    assessment , and control options in food processing. International Journal of Food

    Microbiology 285, 110–128. https://doi.org/10.1016/j.ijfoodmicro.2018.06.001

Bourgarel, M., Pfukenyi, D., Becquart, P., Morand, S., 2019. Next-Generation Sequencing on

    Insectivorous Bat Guano□: An Accurate Tool to Identify Arthropod Viruses of Potential

    Agricultural Concern. https://doi.org/10.3390/v11121102

Cavanagh, D., Cavanagh, D., 2007. Coronaviruses in poultry and other birds Coronaviruses in

    poultry and other birds 9457. https://doi.org/10.1080/03079450500367682

FAO/WHO, 2008. [Food and Agriculture Organization of the United Nations/World Health

    Organization]. Microbiological Risk Assessment Series 14, 151.

Geisbert, T.W., Mire, C.E., Geisbert, J.B., Chan, Y.-P., Agans, K.N., Feldmann, F., Fenton,

    K.A., Zhu, Z., Dimitrov, D.S., Scott, D.P., Bossart, K.N., Feldmann, H., Broder, C.C., 2014.

    Therapeutic treatment of Nipah virus infection in nonhuman primates with a neutralizing

    human monoclonal antibody. Science translational medicine 6, 242ra82-242ra82.

    https://doi.org/10.1126/scitranslmed.3008929

Gibbins, J.D., Hale, C., Silver, M., Fischer, M., Steinberg, J., Basler, C.A., Jacobs, J.R.,

    Kennedy, E.D., Tomasi, S., Trout, D., Hornsby-myers, J., 2020. COVID-19 Among

    Workers in Meat and Poultry Processing Facilities — 69, 557–561.

Guan, Y., Zheng, B.J., He, Y.Q., Liu, X.L., Zhuang, Z.X., Cheung, C.L., Luo, S.W., Li, P.H.,

Zhang, L.J., Guan, Y.J., Butt, K.M., Wong, K.L., Chan, K.W., Lim, W., Shortridge, K.F.,

Yuen, K.Y., Peiris, J.S.M., Poon, L.L.M., 2003. Isolation and characterization of viruses

related to the SARS coronavirus from animals in southern China. Science (New York, NY)

302, 276–278. https://doi.org/10.1126/science.1087139

Hamdi, A., 2013. DETECTION OF ROTA AND CORONA VIRUSES IN RAW MILK AND

MILK PRODUCTS. Benha veterinary medical journal 24, 79–85.

Hi, Z.S., 2020. REVIEW ARTICLE A brief review of foodborne zoonoses in China 1497–1504.

https://doi.org/10.1017/S0950268811000872

Kai Kupferschmidt, 2013. New Middle Eastern Virus Found in Camels.

Katy Askew, 2020. 2020 German slaughterhouses under fire over "insufficient precautionary

measures" to block COVID-19 spread.

Kingsbury, J., Lake, R., Hewitt, J., Smit, E., King, N., 2020. POTENTIAL FOR FOODBORNE

TRANSMISSION OF COVID-19☐: 1–51.

Maffetone, P.B., Laursen, P.B., 2020. The Perfect Storm☐: Coronavirus (Covid-19) Pandemic

Meets Overfat Pandemic 8, 1–6. https://doi.org/10.3389/fpubh.2020.00135

Maori, E., Trust, W., Reports, C., 2019. Discovery of RNA transfer through royal jelly could aid

development of honey bee vaccines 3–5. https://doi.org/10.1016/j.celrep.2019.04.073

Mcmahon, D.P., Wilfert, L., Paxton, R., Brown, M., Holloway, R., 2018. Emerging Viruses in

Bees☐: From. https://doi.org/10.1016/bs.aivir.2018.02.008

Merler, S., Pastore, A., Mu, K., Rossi, L., Sun, K., 2020. The effect of travel restrictions on the

spread of the 2019 novel coronavirus (COVID-19) outbreak 400, 395–400.

Miller, R.S., Sweeney, S.J., Slootmaker, C., Grear, D.A., Salvo, P.A. Di, Kiser, D., Shwiff, S.A., 2017. Cross-species transmission potential between wild pigs , livestock , poultry , wildlife , and humans□: implications for disease risk management in North America. Scientific Reports 1–14. https://doi.org/10.1038/s41598-017-07336-z

Moy, G.G., 2020. IUFoST / CIFST hold an Extraordinary Scienti fi c Roundtable on COVID-19 and Food Safety. npj Science of Food 1–2. https://doi.org/10.1038/s41538-020-0068-2

Newell, D.G., Koopmans, M., Verhoef, L., Duizer, E., Aidara-kane, A., Sprong, H., Opsteegh, M., Langelaar, M., Threfall, J., Scheutz, F., Giessen, J. Van Der, Kruse, H., 2010. International Journal of Food Microbiology Food-borne diseases — The challenges of 20 years ago still persist while new ones continue to emerge. International Journal of Food Microbiology 139, S3–S15. https://doi.org/10.1016/j.ijfoodmicro.2010.01.021

Panel, E., Biohaz, H., 2011. Scientific Opinion on an update on the present knowledge on the occurrence and control of foodborne viruses 1 9, 1–96. https://doi.org/10.2903/j.efsa.2011.2190.

Panon, G., Tache, S., Labie, C., 1988. Respective stability of rotavirus and coronavirus in bovine milk 68, 49–64.

Papineau, A., Berhane, Y., Wylie, T.N., Wylie, K.M., Sharpe, S., Lung, O., 2019. Genome Organization of Canada Goose Coronavirus , A Novel Species Identified in a Mass Die-off of Canada Geese. Scientific Reports 1–8. https://doi.org/10.1038/s41598-019-42355-y

Pirtle, E.C., 1991. Virus survival in the environment 10, 733–748.

PROMKUNTOD, N., ANTARASENA, C., PROMMUANG, PORNTIP, PROMMUANG,

PRAISON, 2006. Isolation of Avian Influenza Virus A Subtype H5N1 from Internal

Contents (Albumen and Allantoic Fluid) of Japanese Quail (Coturnix coturnix japonica)

Eggs and Oviduct during a Natural Outbreak. Annals of the New York Academy of

Sciences 1081, 171–173. https://doi.org/doi:10.1196/annals.1373.020

Rafat, S.A. , saeedi, N. 2020. Human food consumption patterns concerning COVID-19

pandemic. https://www.medrxiv.org/content/10.1101/2020.06.16.20132464v4.article-

metrics

Rafat, S.A., 2020. DATA of Preprint: Human food consumption patterns concerning COVID-19

pandemic. https://doi.org/10.13140/RG.2.2.30851.12321

Rizzo, F., Edenborough, K.M., Toffoli, R., Culasso, P., Zoppi, S., Dondo, A., Robetto, S.,

Rosati, S., Lander, A., Kurth, A., Orusa, R., Bertolotti, L., Mandola, M.L., 2017.

Coronavirus and paramyxovirus in bats from Northwest Italy 1–11.

https://doi.org/10.1186/s12917-017-1307-x

Sarah C. P. Williams, n.d. Kitchen Utensils Transfer Viruses.

Tao, Z., Tian, J., Pei, Y., Yuan, M., Zhang, Y., Dai, F., 2020. A new coronavirus associated with

human respiratory disease in China 579. https://doi.org/10.1038/s41586-020-2008-3

Todd, E.C.D., Greig, J.D., Todd, E., Llc, C., 2020. Viruses of foodborne origin □: a review 25–

46.

Figure captions

Fig. 1 Potential transmission ways of COVID-19 based on the results of our analysis and bibliography. It is necessary to study the exchange of human pathogen viruses in wildlife and livestock. Where and how suspicious foods become contaminated must be investigated. The potential spillover of pathogenic viruses from arthropods hosts to humans should be investigated(Tao et al., 2020). The cross-species transmission is reported(Miller et al., 2017). There may be a type of contact among flowers, bees, and bats. Infected humans could have transmitted MERS to the camels (Kai Kupferschmidt, 2013), similar situation is imaginable for SARS-Cov-2. SARS-CoV-2 and MERS-Co are among the most notable examples of spillover into humans(Guan et al., 2003). References for arrows: 1 (Bourgarel et al., 2019; Cavanagh and Cavanagh, 2007; Maori et al., 2019; Mcmahon et al., 2018; Moy, 2020; Rizzo et al., 2017)   2: (Miller et al., 2017; Moy, 2020)   3: (Miller et al., 2017; Papineau et al., 2019) 4:(Geisbert et al., 2014; Tao et al., 2020)   5: (Gibbins et al., 2020; Newell et al., 2010; Panon et al., 1988) 6 : (Gibbins et al., 2020; Hi, 2020; Katy Askew, 2020)