COVID_19's lost puzzle piece: the power of domestic animal reservoir

Nazli Saeedi [1], Seyed Abbas Rafat[2*]

1 Research Center for Pharmaceutical Nanotechnology, Biomedicine Institute, Tabriz

University of Medical Sciences, Tabriz, Iran Email : saeedi@hotmail.fr, 2 Department of

Animal Science, Faculty of Agriculture, University of Tabriz, Tabriz, Iran, Email;

abbasrafat@hotmail.com *Corresponding Author

Abstract

In recent months, despite being aware of the COVID-19 pandemic, many countries faced

increasing epidemic of this disease. Nearly all countries tried to decrease human-to-human

contact as the principal mode of transmission of SARS-CoV-2. Other modes of transmission also

need to be clarified in more depth, especially, the foodborne transmission. We assessed the effect

of animal origin foods (AOF) consumption on the pandemic of COVID-19. For this purpose, we

studied the relationship among 20 food supply as independent variables, and the parameter of

Total Cases dependent variable. Here we show a relationship between a group of AOF and total

cases. Stepwise Regression, Bayes, and Lasso results showed that eggs and fresh fish have

positive coefficient. So, among the transmission ways of COVID_19, the role of foodborne

transmission should be more significant than previously thought. The possibility of animal origin

foodborne transmission should be taken into more consideration. There is need to invent a

pasteurization-like-processes to implicate immediately before ingestion of food. Of course, there

will be challenges along the way of the revolution in the food preparation culture. The

perspective is to expand the surveillance of SARS-Cov-2 during the food production chain.

Author summary

At the time of writing this manuscript, the number of TC of COVID-19 past seven-million mark. During the pandemic, most countries around the world minimized human-to-human contact and travels. However, the prevalence of COVID-19 continued. Despite the awareness of many countries about the principal mode of person-to-person transmission, the rapid outbreak shows the necessity of research on other transmission routes. We suppose the role of FBT of COVID-19 should be more significant than previously thought. The effect of a group of AOF including eggs and fresh fish on total cases was significant in our analysis. At present, we are not able to interpret all significant effects and their coefficients. Interpretation of the factors could be the subject of future researches. In food science technology, it is necessary to expand the detection of viruses to all AOF. WHO's recommendations state that COVID_19 have not FBT, and suitable food safety practices could prevent the probability of FBT. However, the realization of proper food safety practices for all foods is questionable.

Introduction

At the time of writing this manuscript, the number of TC of COVID-19 past seven-million mark. During the pandemic, most countries around the world minimized human-to-human contact and travels[1]. However, the prevalence of COVID-19 continued. Despite the awareness of many countries about the principal mode of person-to-person transmission, the rapid outbreak shows the necessity of research on other transmission routes. In Italy and the USA, for example, the chances of being surprised were low, but both countries faced epidemic dramatically. Among the various routes of transmission, FBT is most likely. We suppose the role of FBT of COVID-19 should be more significant than previously thought. Transmission of MERS, a similar pathogenic virus to SARS-Cov-2, through food has strengthened our hypothesis.

Furthermore, some researchers cited the possibility of FBT for SARS-Cov-2[2–6]. Also,

FAO/WHO experts mentioned the data gap in the relationship between viruses and FB disease

[7] . The motivation for the present study is to find the relationship between COVID_19 and

human food consumption patterns.

Finding the relevance of TC with food

Material and methods are included in supplementary materials.

We assessed a relationship between the consumption of some foods and TC. Accordingly, we

studied TC as a dependent variable and 20 food supply data as independent variables. The

statistical methods included correlation, stepwise and LASSO regression, Principal Component

Analysis (PCA), and Bayes analysis. Food supply (kg/capita/yr.) was quantity in each country.

The primary purpose of the study was to find effect of AOF on TC, however, we considered six

plant origin foods for comparison. Histogram of the variables, and their logarithmic

transformation are presented in Figs S1 and S2. Diagnostics panels and selection criteria of

regression models are presented in Figs S3-S5. The scree plot of PCA is shown in Fig S6. Visual

examination of the trace plot in Bayesian analysis is shown in Fig S8. Adjusted R-square and

Geweke diagnostics of fitted models are presented in S1 and S2 Tables, respectively.

Furthermore, the relationship between TC and animal production in each country was estimated

using the same statistical methods.

Results and Discussion

Food consumption

Table 1 shows descriptive statistics of studied variables. Results of PCA show that AOF and TC

are mostly grouped (Fig 1 and S7 Fig). The highest positive Pearson correlation with TC belongs

to eggs. Stepwise Regression, Bayes, and Lasso results showed that eggs and fresh fish have

positive coefficient. With our results in Table 2, implying the need for tracing of SARS-COV-2

in some AOF. Bayes results, e.g., show there is a 0.99 probability of a positive correlation

between eggs and TC, adjusted for the other covariates. The number of AOF and their role in our

results varies depending on the type of statistical method used, but all results are consistent.

Table 1. Descriptive statistics of studied foods (n=20) besides total cases.

#	Variable (kg/capita/yr)	Abbreviation	N	Mean	Std Dev	Minimum	Maximum
1	**Pig meat**	pig	169	14.66	15.52	0.00	71.48
2	**Bovine Meat**	bov	174	11.20	9.00	0.78	55.48
3	**Cheese**	che	173	4.97	6.55	0.00	30.82
4	**Cream**	cre	159	1.21	2.76	0.00	18.45
5	**Eggs**	Egg	174	7.08	5.09	0.16	19.15
6	**Fresh water Fish**	freshf	174	3.38	4.54	0.00	34.00
7	**Honey**	hon	168	0.36	0.46	0.00	3.51
8	**Milk, Whole**	milk	174	59.17	48.93	0.96	258.44
9	**Poultry Meat**	poul	174	20.56	16.46	0.42	72.87
10	**Demersal Fish**	demer	173	3.40	4.93	0.00	33.45
11	**Cephalopods**	ceph	171	0.44	0.96	0.00	6.70
12	**Butter, Ghee**	butg	174	1.25	1.64	0.00	9.25
13	**Marine Fish, Other**	marf	172	2.74	4.87	0.00	28.19
14	**Aquatic Plants**	aqp	162	0.20	1.91	0.00	22.41
15	**Beverages, Alcoholic**	bev	171	3.20	3.87	0.00	26.90
16	**Rice(Milled Equivalent)**	rice	174	30.95	36.20	0.95	171.73
17	**Potatoes and products**	potato	174	34.89	33.65	0.05	183.16
18	**Maize and products**	maiz	170	21.71	28.52	0.00	158.40
19	**Wheat and products**	wheat	174	70.92	46.85	1.89	222.40
20	**wine**	wine	169	6.27	9.49	0.00	51.41
	Total Cases (Per Country)	TC	215	29247.33	138459.41	1	1837578

Table 2. The results of Pearson correlation and multivariate analyses. All variables are

logarithmic transformed data. For abbreviations, see Table 1.

	r, P Value, n	Stepwise	lasso	Bayes
pig	0.17, 0.03, 156	-0.52	-0.28	0.19
bov	0.26, 0.008, 161			0.63
Che	0.28, 0.0004, 161			0.29
cre	0.18, 0.03, 150			0.47
egg	0.43, <.0001, 161	1.84	**1.16**	**0.99**
freshf	0.31, <.0001, 161	1	**0.90**	**0.99**
hon	0.23, 0.004, 158			0.20
milk	0.21, 0.006, 161			0.48
poul	ns			0.31
demer	0.25, <.0001, 160		0.22	0.79
ceph	ns			0.51
Butg	0.34, <.0001, 161		0.08	0.80
marf	-0.45, <.0001, 159	-1.36	-1.29	-1.00
aqp	ns			0.50
bev	ns			0.16
rice	ns		0.30	0.98
potato	0.36, <.0001,161		0.22	0.88
maiz	ns		0.13	0.92
wheat	0.29, 0.0002, 161		0.32	0.90
wine	0.19, 0.02, 157			0.75

The effect of a group of AOF on TC was significant in our analysis. So, we suppose a noticeable effect of AOF in FBT. At present, we are not able to interpret all significant effects and their coefficients. Interpretation of the factors could be the subject of future researches. Potential transmission ways of COVID-19 based on the results of our analysis and bibliography is summarized as Fig 2. The transmission of coronaviruses through food has been reported[6]. Lake and Kingsbury (2020) provided a review about potential for FBT of COVID-19, but they could not guarantee the application of the data found on other coronaviruses to SARS-CoV-2[6]. The persistence of viruses by refrigeration temperature[7] might be an explanation of the relationship between COVID-19 and AOF. The viral persistence of coronavirus in soft cheese [8] and raw milk[9] has been reported. We don't know what's going on from "Farm to Fork" about AOF. For example, SARS-CoV-2 may be more resistant to heat when cooking, as it is observed for poliovirus in beef[10]. It is recommended not to consume raw or undercooked animal products[11]. Kitchen utensils can transmit the virus among foods[12], accordingly, AOF may be a suitable vehicle, and the virus inside them will survive better and longer. The human pathogenic virus, such as Enterovirus, can survive in honey, so it is imaginable similarly for SARS-CoV-2. Honey bees deposit viruses on flowers[13]. Maori showed horizontal RNA flow among honey bees by the ingestion of royal jelly[14]. Trade in raw honey bee products could pose pandemic risks for pathogen viruses of honey bees[15]. The pessimist hypothesis would be the transmission of SARS-CoV-2 by raw honey.

Furthermore, the methods of diagnosing the virus in food are not complete, and there are many limitations[16]. In food science technology, the primary attention paid to bivalve moullouses. Nowadays, it is necessary to expand the detection of viruses to all AOF.

117 WHO's recommendations state that COVID_19 have not FBT, and suitable food safety practices
118 could prevent the probability of FBT. However, the realization of proper food safety practices for
119 all AOF is questionable.

120 **Animal production**
121

The highest pair-wise Pearson correlations between the number of livestock and TC belong to
poultry (0.71), turkeys (0.68), pigs (0.62), horses (0.59), and beehives (0.57) with P-value of
(<0.0001) for all of them. The chosen variables in Bayesian analysis were the number of chicken,
turkeys, pigs, and goats, and in Lasso analysis they were the number of chickens, pigs, and
turkeys. The observed positive correlation between the number of beehives and TC should be
explained by circulating viruses between animal and human. As, some zoonotic viruses first
circulate in animals, and sometimes they infect humans. After that, they adapt to humans. A
recent zoonotic transmission event of SARS-CoV-2 is reported[17]. It is crucial to characterize
circulating viruses in the human working environment, especially where an interface with
wildlife[18]. Spread of influenza virus during commercial movement of eggs is reported by
Pillai[19]. Poultry production in the extensive system increases the chance of SARS-CoV-2
transmission among other species[20]. Feathers detached from domestic ducks reported as a
source of environmental contamination to pathogenic avian influenza virus[21]. Qiu et al.(2020)
identified pigs, cows, goats, and sheep among the most likely intermediate animal hosts for
SARS-CoV-2 [22]. The outbreak of COVID-19 highlights viral spillover from animals[23].
There is the potential for the emergence of new coronavirus diseases in domestic birds, from
both avian and mammalian sources. For example, a novel species of coronaviridae family has
been identified recently[24]. *Coronaviridae*'s family viruses are likely can cross the species
barrier of viruses and infect other species, fishes, cattle, pigs, and humans [25].

Conclusion

Despite the awareness and readiness of many countries for COVID-19 during early 2020, why did it spread so quickly? It seems FBT has received less attention relative to other routes. There is need to invent a pasteurization-like-processes to implicate immediately before ingestion of food. Of course, there will be challenges along the way of the revolution in the food preparation culture. The perspective is to expand the surveillance of SARS-Cov-2 during the food production chain.

Fig. 1 PCA results: the interrelationships between all variables. Some of AOF are closer to TC. For abbreviations, see Table 1.

Fig. 2 Potential transmission ways of COVID-19 based on the results of our analysis and bibliography. It is necessary to study the exchange of human pathogen viruses in wildlife and livestock. Where and how suspicious foods become contaminated must be investigated. The potential spillover of pathogenic viruses from arthropods hosts to humans should be investigated[23]. The cross-species transmission is reported[26].There may be a type of contact among flowers, bees, and bats. Infected humans could also have transmitted MERS to the camels [27]. SARS-CoV-2 and MERS-Co are among the most notable examples of spillover into humans[28]. References for arrows:

 1 [11,14,18,20,29,30]

2: [11,26]

3: [24,26]

4:[23,31]

5: [5,8,32]

6 : [4,32,33]

Material and Methods

Food consumption

We estimated the relationship between 20 food supply and Total Cases of covid19.

The methods include pair-wise correlation with Proc Corr, Stepwise Regression

with Proc Reg, LASSO regression with GLMSELECT Proc, Principal Component

Analysis with Proc Princomp, and Bayes analysis with Proc Genmod (SAS). Food

supply was quantity (kg/capita/yr.). Fourteen foods of animal origin and six foods

of plant origin were selected. The primary purpose of the study was to find animal

origin foods relationship with TC, but some plant origin foods were chosen to

compare.

We started our analyses first time on March 2020. Then we repeated it several

times with updated COVID_19 data on April, May, and June. Finally we presented

the results of June' data.

Data

COVID_19 https://www.worldometers.info/coronavirus/   1 June 2020

Live Animal production data: http://www.fao.org/faostat/en/#data/QA

Food Supply - Crops Primary Equivalent:

http://www.fao.org/faostat/en/#data/CC

Food Supply - Livestock and Fish Primary Equivalent:

http://www.fao.org/faostat/en/#data/CL

We used base-e logs, also known as natural logs (LOG in SAS) for the

transformation of TC and independent variables. In all analyses, the accuracy of

the model with non-transformed data was very low. So, all variables has been

transformed to normalize the distributions. With transformation of data, the

accuracy of the model increased to an acceptable level. In PCA, as a nonparametric

method, we utilized non-transformed data, in which the variables are treated

equally, i.e., that is, there are no dependent variables.

In comparison of models, Lasso's results are more reliable than stepwise regression

and Bayes, because it's AIC was smaller and had higher accuracy as R-square (S1

Table).

S1 Table. Adjusted R-square and AIC of fitted models for food consumption.

	Stepwise	LASSO	BAYES
R-Square	0.48	0.52	-
Adj-R_Square	0.47	0.48	-
AIC	-	301	551
n	131	131	131

Bayesian Analysis: Visual examination of the trace plot displays proper mixing, for one independent variable in presented in S8 Fig. The P-value in the Geweke Diagnostics (S2 Table) showed that the mean estimate of the Markov chain is stable over time. The positive probability that B1 greater than 0 is estimated.

213 **S2 Table Geweke diagnostics from Genmode analysis**. Pr > z shows

214 convergence for all variables. For abbreviations, see Table 1

Parameter[*]	z	Pr > |z|
Intercept	0.2734	0.7845
pig	0.6475	0.5173
bov	-0.2165	0.8286
che	-0.7437	0.4571
cre	-0.9484	0.3429
egg	-1.5173	0.1292
freshf	0.4354	0.6633
hon	0.7984	0.4246
milk	-0.0434	0.9654
poul	0.9524	0.3409
demer	-0.9509	0.3416
ceph	0.7935	0.4275
butg	1.898	0.0577
marf	-0.4608	0.6449
aqp	0.58	0.5619
bev	-1.1802	0.2379
rice	-0.9809	0.3266
potato	-1.0137	0.3107
maiz	-0.3024	0.7624
wheat	0.5066	0.6124
wine	-0.0076	0.994
Scale	0.0274	0.9781

* Logarithmic transformation for all parameters

Although the probability of transmitting the virus through wine is zero, we

deliberately kept it in the initial model of our analysis. Results showed wine is

eliminated among significant variables of the suited model (Table 2). So we

concluded that just the energy content of food should not keep a variable in the

fitted model. So, in addition to energy content, there must be something else in

food-related COVID-19. The high-fat tissue in the body affects on the

pathogenesis of COVID-19[34]. We concentrate FB transmission of COVID-19

rather than the immunity response.

Animal and crop production

We utilized the GLMSLECT procedure of SAS 9.2 with the method of least

absolute shrinkage and selector operator (LASSO). The dependent variable was

TC, and independent variables were beehives, cattle, chicken, pigs, goats, turkeys,

horse, rabbit, maize, barley, and apples. Maize, Barley, apples were deliberately

kept in a multi-variable model, to test whether plant material production remains in

the model compared to livestock production. We used the GENMOD procedure of

the SAS using the *BAYES* statement to obtain *Bayesian* analysis. Spearman and

Pearson correlation between stocks and TC calculated. Analyzes were performed

with log-transformed data. Due to the higher accuracy of the transformed data,

their results were used.

There are references[33] that guide us to construct our hypothesis about the

essential relationship between COVID-19 and AOF include.

We provide the SAS code for an application of the used statistical methods [35].

We note that this work was posted as preprint on medrxiv[36] .

ACKNOWLEDGEMENTS:

Author contributions: Conceptualization and Visualisation: N. S. K., Methodology: S. A. R.

Writing: Both authors.

The authors have declared no competing interest.

Data and materials availability: All data are available in the manuscript or the supplementary

materials.

Supporting information

S1 Fig. Histogram of variables of Total cases before (left) and after (right)

logarithmic transformation.

S2 Fig. Histogram of some independent variables before and after logarithmic

transformation. First row, from Left to right: cream, log cream, pigs, logpigs.

Second row: Bovine meat, log bovin meat, eggs, log of eggs.

S3 Fig. Diagnostics panel. Fit diagnostics and residual by regressors, in

multivariate regression with stepwise selection of variables.

S4 Fig. Other Criteria Suggest Stopping the Selection (GLMSELECT proc of

SAS). Progression of squared errors (left) and Fit criteria of LASSO analysis.

S5 Fig. Results of the GLMSELECT Proc of SAS. The standardized coefficients

of all the effects that are selected at some step of the elastic net method, plotted as

a function of the step number of LASSO analysis.

S6 Fig. Scree plot and percentage of variance explained by principal

componentsofPCA analysis.

S7 Fig. Results of PROC PRINCOMP, includes 1 by 3 (left)and 2 by 3(right)

component pattern plots. Principal component analysis results showed that AOF

are generally closer to TC in the comparison with other studied foods.

S8 Fig. Bayesian Analysis: Visual examination of the trace plot displays proper

mixing for Eggs.

References

1.   Merler S, Pastore A, Mu K, Rossi L, Sun K. The effect of travel restrictions on the spread

of the 2019 novel coronavirus (COVID-19) outbreak. 2020;400: 395–400.

2.   Panel E, Biohaz H. Scientific Opinion on an update on the present knowledge on the

occurrence and control of foodborne viruses 1. 2011;9: 1–96.

doi:10.2903/j.efsa.2011.2190.

3.   Todd ECD, Greig JD, Todd E, Llc C. Viruses of foodborne origin☐: a review. 2020; 25–

46.

4.   Hi ZS. REVIEW ARTICLE A brief review of foodborne zoonoses in China. 2020; 1497–

1504. doi:10.1017/S0950268811000872

5.   Newell DG, Koopmans M, Verhoef L, Duizer E, Aidara-kane A, Sprong H, et al.

International Journal of Food Microbiology Food-borne diseases — The challenges of 20

years ago still persist while new ones continue to emerge. International Journal of Food

Microbiology. 2010;139: S3–S15. doi:10.1016/j.ijfoodmicro.2010.01.021

6.   Kingsbury J, Lake R, Hewitt J, Smit E, King N. POTENTIAL FOR FOODBORNE

TRANSMISSION OF COVID-19☐: 2020; 1–51.

7.   FAO/WHO. [Food and Agriculture Organization of the United Nations/World Health

Organization]. Microbiological Risk Assessment Series. 2008;14: 151.

8.   Panon G, Tache S, Labie C. Respective stability of rotavirus and coronavirus in bovine

milk. 1988;68: 49–64.

9.   Hamdi A. Mohamed, Adham M. Abdou , Ekbal M. Adel, Sayed Ahmed H. Salem MIME-

H. DETECTION OF ROTA AND CORONA VIRUSES IN RAW MILK AND MILK

PRODUCTS. Benha veterinary medical journal. 2013;24: 79–85. Available:

http://www.bvmj.bu.edu.eg/abstract.php?ee=156

10.   Pirtle EC. Virus survival in the environment. 1991;10: 733–748.

11.   Moy GG. IUFoST / CIFST hold an Extraordinary Scienti fi c Roundtable on COVID-19

and Food Safety. npj Science of Food. 2020; 1–2. doi:10.1038/s41538-020-0068-2

12.   Sarah C. P. Williams. Kitchen Utensils Transfer Viruses. Available:

295   https://www.sciencemag.org/news/2012/12/kitchen-utensils-transfer-viruses

296   13.   Id SAA, Burnham PA, Boncristiani HF, Brody K. RNA virus spillover from managed

297   honeybees (Apis mellifera) to wild bumblebees (Bombus spp .). 2019; 1–13.

298   14.   Maori E, Trust W, Reports C. Discovery of RNA transfer through royal jelly could aid

299   development of honey bee vaccines. 2019; 3–5. doi:10.1016/j.celrep.2019.04.073

300   15.   Mutinelli F. The spread of pathogens through trade in honey bees and their products (

301   including queen bees and semen): overview and recent developments Honey bees.

302   2011;30: 257–271.

303   16.   Bosch A, Gkogka E, Le FS, Loisy-hamon F, Lee A, Lieshout L Van, et al. International

304   Journal of Food Microbiology Foodborne viruses□: Detection , risk assessment , and

305   control options in food processing. International Journal of Food Microbiology. 2018;285:

306   110–128. doi:10.1016/j.ijfoodmicro.2018.06.001

307   17.   Kujawski SA, Wong KK, Collins JP, Epstein L, Killerby ME, Midgley CM, et al. Clinical

308   and virologic characteristics of the first 12 patients with coronavirus disease 2019

309   (COVID-19) in the United States. Nature Medicine. 2020;26: 861–868.

310   doi:10.1038/s41591-020-0877-5

311   18.   Bourgarel M, Pfukenyi D, Becquart P, Morand S. Next-Generation Sequencing on

312   Insectivorous Bat Guano□: An Accurate Tool to Identify Arthropod Viruses of Potential

313   Agricultural Concern. 2019. doi:10.3390/v11121102

314   19.   Pillai SPS, Saif YM, Lee CW. Detection of influenza A viruses in eggs laid by infected

315   turkeys. Avian diseases. 2010;54: 830–833. doi:10.1637/9102-101209-Reg.1

316   20.   Cavanagh D, Cavanagh D. Coronaviruses in poultry and other birds Coronaviruses in

317   poultry and other birds. 2007;9457. doi:10.1080/03079450500367682

318   21.   Yamamoto Y, Nakamura K, Yamada M, Mase M. Persistence of Avian Influenza Virus (

319   H5N1) in Feathers Detached from Bodies of Infected Domestic Ducks □. 2010;76: 5496–

320   5499. doi:10.1128/AEM.00563-10

321   22.   Qiu Y, Zhao Y, Wang Q, Li J, Zhou Z, Liao C, et al. Predicting the angiotensin converting

322   enzyme 2 (ACE2) utilizing capability as the receptor of SARS-CoV-2. Microbes and

323   Infection. 2020;2. doi:10.1016/j.micinf.2020.03.003

324   23.   Tao Z, Tian J, Pei Y, Yuan M, Zhang Y, Dai F. A new coronavirus associated with human

325   respiratory disease in China. 2020;579. doi:10.1038/s41586-020-2008-3

326   24.   Papineau A, Berhane Y, Wylie TN, Wylie KM, Sharpe S, Lung O. Genome Organization

327   of Canada Goose Coronavirus , A Novel Species Identified in a Mass Die-off of Canada

328   Geese. Scientific Reports. 2019; 1–8. doi:10.1038/s41598-019-42355-y

329   25.   Leong JC. Fish Viruses. Encyclopedia of Virology. 2008; 227—234. doi:10.1016/b978-

330   012374410-4.00400-3

331   26.   Miller RS, Sweeney SJ, Slootmaker C, Grear DA, Salvo PA Di, Kiser D, et al. Cross-

332   species transmission potential between wild pigs , livestock , poultry , wildlife , and

333   humans□: implications for disease risk management in North America. Scientific Reports.

334   2017; 1–14. doi:10.1038/s41598-017-07336-z

335   27.   Kai Kupferschmidt. New Middle Eastern Virus Found in Camels. 2013. Available:

336 https://www.sciencemag.org/news/2013/11/new-middle-eastern-virus-found-camels

337 28. Guan Y, Zheng BJ, He YQ, Liu XL, Zhuang ZX, Cheung CL, et al. Isolation and

338   characterization of viruses related to the SARS coronavirus from animals in southern

339   China. Science (New York, NY). 2003;302: 276–278. doi:10.1126/science.1087139

340 29. Rizzo F, Edenborough KM, Toffoli R, Culasso P, Zoppi S, Dondo A, et al. Coronavirus

341   and paramyxovirus in bats from Northwest Italy. 2017; 1–11. doi:10.1186/s12917-017-

342   1307-x

343 30. Mcmahon DP, Wilfert L, Paxton R, Brown M, Holloway R. Emerging Viruses in Bees□:

344   From. 2018. doi:10.1016/bs.aivir.2018.02.008

345 31. Geisbert TW, Mire CE, Geisbert JB, Chan Y-P, Agans KN, Feldmann F, et al.

346   Therapeutic treatment of Nipah virus infection in nonhuman primates with a neutralizing

347   human monoclonal antibody. Science translational medicine. 2014;6: 242ra82-242ra82.

348   doi:10.1126/scitranslmed.3008929

349 32. Gibbins JD, Hale C, Silver M, Fischer M, Steinberg J, Basler CA, et al. COVID-19

350   Among Workers in Meat and Poultry Processing Facilities —. 2020;69: 557–561.

351 33. Katy Askew. 2020 German slaughterhouses under fire over "insufficient precautionary

352   measures" to block COVID-19 spread. 2020.

353 34. Maffetone PB, Laursen PB. The Perfect Storm□: Coronavirus (Covid-19) Pandemic

354   Meets Overfat Pandemic. 2020;8: 1–6. doi:10.3389/fpubh.2020.00135

355 35. Rafat SA. DATA of Preprint: Human food consumption patterns concerning COVID-19

356   pandemic. 2020. doi:10.13140/RG.2.2.30851.12321

357   36.   Rafat, S.A. saeedi N. Human food consumption patterns concerning COVID-19

358   pandemic. Available: https://www.medrxiv.org/content/10.1101/2020.06.16.20132464v1

376 Fig1

379   Fig 2

382   Fig s1

387   Fig s2

392   Fig s3

397   Fig s4

403   Fig s5

409   Fig s6

413   Fig s7

419   Fig s8