COVID_19's lost puzzle piece: the power of domestic animal reservoir

Nazli Saeedi [1], Seyed Abbas Rafat[2*]

1 Research Center for Pharmaceutical Nanotechnology, Biomedicine Institute, Tabriz

University of Medical Sciences, Tabriz, Iran Email : saeedi@hotmail.fr, 2 Department of

Animal Science, Faculty of Agriculture, University of Tabriz, Tabriz, Iran, Email;

abbasrafat@hotmail.com *Corresponding Author

Abstract

In recent months, despite being aware of the COVID-19 pandemic, many countries faced

increasing epidemic of this disease. Nearly all countries tried to decrease human-to-human

contact as the principal mode of transmission of SARS-CoV-2. Other modes of transmission also

need to be clarified in more depth, especially, the foodborne transmission. We assessed the effect

of animal origin foods (AOF) consumption on the pandemic of COVID-19. For this purpose, we

studied the relationship among 20 food supply as independent variables, and the parameter of

Total Cases dependent variable. Here we show a relationship between a group of AOF and total

cases. Stepwise Regression, Bayes, and Lasso results showed that eggs and fresh fish have

positive coefficient. So, among the transmission ways of COVID_19, the role of foodborne

transmission should be more significant than previously thought. The possibility of animal origin

foodborne transmission should be taken into more consideration. There is need to invent a

pasteurization-like-processes to implicate immediately before ingestion of food. Of course, there

will be challenges along the way of the revolution in the food preparation culture. The

perspective is to expand the surveillance of SARS-Cov-2 during the food production chain. In

conclusion, the results of the present study indicate that the reservoir of SARS-Cov-2 may be some of

the domestic animals. It should be noted that this study was conducted with data related to the time (2013)

before the effect of COVID_19 pandemic on food supply. It is recommended that virologists examine the

possibility of freshwater fish and chicken's eggs being reservoirs for SARS-Cov-2.

Author summary

At the time of writing this manuscript, the number of total cases of COVID-19 past seven-

million mark. During the pandemic, most countries around the world minimized human-to-

human contact and travels. However, the prevalence of COVID-19 continued. Despite the

awareness of many countries about the principal mode of person-to-person transmission, the

rapid outbreak shows the necessity of research on other transmission routes. We suppose the role

of FBT of COVID-19 should be more significant than previously thought. The effect of a group

of AOF including eggs and fresh fish on total cases was significant in our analysis. At present,

we are not able to interpret all significant effects and their coefficients. Interpretation of the

factors could be the subject of future researches. In food science technology, it is necessary to

expand the detection of viruses to all AOF. WHO's recommendations state that COVID_19 have

not FBT, and suitable food safety practices could prevent the probability of FBT. However, the

realization of proper food safety practices for all foods is questionable.

Introduction

At the time of writing this manuscript, the number of total cases of COVID-19 past seven-

million mark. During the pandemic, most countries around the world minimized human-to-

human contact and travels[1]. However, the prevalence of COVID-19 continued. Despite the

awareness of many countries about the principal mode of person-to-person transmission, the

rapid outbreak shows the necessity of research on other transmission routes. In Italy and the

USA, for example, the chances of being surprised were low, but both countries faced epidemic

dramatically. Among the various routes of transmission, FBT is most likely. We suppose the role

of FBT of COVID-19 should be more significant than previously thought. Transmission of

MERS, a similar pathogenic virus to SARS-Cov-2, through food has strengthened our

hypothesis.

Furthermore, some researchers cited the possibility of FBT for SARS-Cov-2[2–6]. Also,

FAO/WHO experts mentioned the data gap in the relationship between viruses and FB disease

[7] . The motivation for the present study is to find the relationship between COVID_19 and

human food consumption patterns.

Finding the relevance of total cases with food

Material and methods are included in supplementary materials.

We assessed a relationship between the consumption of some foods and total cases. Accordingly,

we studied total cases as a dependent variable and 20 food supply data as independent variables.

The statistical methods included correlation, stepwise and LASSO regression, Principal

Component Analysis (PCA), and Bayes analysis. Food supply (kg/capita/yr.) was quantity in

each country. The primary purpose of the study was to find effect of AOF on total cases,

however, we considered six plant origin foods for comparison. Histogram of the variables, and

their logarithmic transformation are presented in Figs S1 and S2. Diagnostics panels and

selection criteria of regression models are presented in Figs S3-S5. The scree plot of PCA is

shown in Fig S6. Visual examination of the trace plot in Bayesian analysis is shown in Fig S8.

Adjusted R-square and Geweke diagnostics of fitted models are presented in S1 and S2 Tables,

respectively.

Furthermore, the relationship between total cases and animal production in each country was

estimated using the same statistical methods.

Results and Discussion

Food consumption
72

Table 1 shows descriptive statistics of studied variables. Results of PCA show that AOF and total

cases are mostly grouped (Fig 1 and S7 Fig). The highest positive Pearson correlation with total

cases belongs to eggs. Stepwise Regression, Bayes, and Lasso results showed that eggs and fresh

fish have positive coefficient. With our results in Table 2, implying the need for tracing of

SARS-COV-2 in some AOF. Bayes results, e.g., show there is a 0.99 probability of a positive

correlation between eggs and total cases, adjusted for the other covariates. The number of AOF

and their role in our results varies depending on the type of statistical method used, but all results

are consistent.

Table 1. Descriptive statistics of studied foods (n=20) besides total cases.

#	Variable (kg/capita/yr)	Abbreviation	N	Mean	Std Dev	Minimum	Maximum
1	**Pig meat**	pig	169	14.66	15.52	0.00	71.48
2	**Bovine Meat**	bov	174	11.20	9.00	0.78	55.48
3	**Cheese**	che	173	4.97	6.55	0.00	30.82
4	**Cream**	cre	159	1.21	2.76	0.00	18.45
5	**Eggs**	Egg	174	7.08	5.09	0.16	19.15
6	**Freshwater Fish**	freshf	174	3.38	4.54	0.00	34.00
7	**Honey**	hon	168	0.36	0.46	0.00	3.51
8	**Whole milk**	milk	174	59.17	48.93	0.96	258.44
9	**Poultry Meat**	poul	174	20.56	16.46	0.42	72.87
10	**Demersal Fish**	demer	173	3.40	4.93	0.00	33.45
11	**Cephalopods**	ceph	171	0.44	0.96	0.00	6.70
12	**Butter, Ghee**	butg	174	1.25	1.64	0.00	9.25
13	**Marine Fish, Other**	marf	172	2.74	4.87	0.00	28.19
14	**Aquatic Plants**	aqp	162	0.20	1.91	0.00	22.41
15	**Beverages, Alcoholic**	bev	171	3.20	3.87	0.00	26.90
16	**Rice(Milled Equivalent)**	rice	174	30.95	36.20	0.95	171.73
17	**Potatoes and products**	potato	174	34.89	33.65	0.05	183.16
18	**Maize and products**	maiz	170	21.71	28.52	0.00	158.40
19	**Wheat and products**	wheat	174	70.92	46.85	1.89	222.40
20	**wine**	wine	169	6.27	9.49	0.00	51.41
	Total Cases (Per Country)	TC	215	29247.33	138459.41	1	1837578

96 **Table 2. The results of Pearson correlation (r) and multivariate analyses.** All variables are
97 logarithmic transformed data.
98

Variable	r, P Value, N	Stepwise	Lasso	Bayes
Pig meat	0.17, 0.03, 156	-0.52	-0.28	0.19
Bovine Meat	0.26, 0.008, 161			0.63
Cheese	0.28, 0.0004, 161			0.29
Cream	0.18, 0.03, 150			0.47
Eggs	0.43, <.0001, 161	1.84	**1.16**	**0.99**
Freshwater fish	0.31, <.0001, 161	1	**0.90**	**0.99**
Honey	0.23, 0.004, 158			0.20
Whole milk	0.21, 0.006, 161			0.48
Poultry meat	non-significant			0.31
Demersal fish	0.25, <.0001, 160		0.22	0.79
Cephalopods	non-significant			0.51
Butter/ghee	0.34, <.0001, 161		0.08	0.80
Marine fish(other)	-0.45, <.0001, 159	-1.36	-1.29	-1.00
Aquatic plants	non-significant			0.50
Beverages(Alcoholic)	non-significant			0.16
Rice(Milled Equivalent)	non-significant		0.30	0.98
Potatoes and products	0.36, <.0001, 161		0.22	0.88
Maize and products	non-significant		0.13	0.92

Wheat and products	0.29, 0.0002, 161	0.32	0.90
Wine	0.19, 0.02, 157		0.75

The effect of a group of AOF on total cases was significant in our analysis. So, we suppose a noticeable effect of AOF in FBT. At present, we are not able to interpret all significant effects and their coefficients. Interpretation of the factors could be the subject of future researches. Potential transmission ways of COVID-19 based on the results of our analysis and bibliography is summarized as Fig 2. The transmission of coronaviruses through food has been reported[6]. Lake and Kingsbury (2020) provided a review about potential for FBT of COVID-19, but they could not guarantee the application of the data found on other coronaviruses to SARS-CoV-2[6]. The persistence of viruses by refrigeration temperature[7] might be an explanation of the relationship between COVID-19 and AOF. The viral persistence of coronavirus in soft cheese [8] and raw milk[9] has been reported. We don't know what's going on from "Farm to Fork" about AOF. For example, SARS-CoV-2 may be more resistant to heat when cooking, as it is observed for poliovirus in beef[10]. It is recommended not to consume raw or undercooked animal products[11]. Kitchen utensils can transmit the virus among foods[12], accordingly, AOF may be a suitable vehicle, and the virus inside them will survive better and longer. The human pathogenic virus, such as Enterovirus, can survive in honey, so it is imaginable similarly for SARS-CoV-2. Honey bees deposit viruses on flowers[13]. Maori showed horizontal RNA flow among honey bees by the ingestion of royal jelly[14]. Trade in raw honey bee products could pose pandemic risks for pathogen viruses of honey bees[15]. The pessimist hypothesis would be the transmission of SARS-CoV-2 by raw honey.

Furthermore, the methods of diagnosing the virus in food are not complete, and there are many

limitations[16]. In food science technology, the primary attention paid to bivalve moullouses.

Nowadays, it is necessary to expand the detection of viruses to all AOF.

WHO's recommendations state that COVID_19 have not FBT, and suitable food safety practices

could prevent the probability of FBT. However, the realization of proper food safety practices for

all AOF is questionable.

Animal production
126

The highest pair-wise Pearson correlations between the number of livestock and total cases

belong to poultry (0.71), turkeys (0.68), pigs (0.62), horses (0.59), and beehives (0.57) with P-

value of (<0.0001) for all of them. The chosen variables in Bayesian analysis were the number of

chicken, turkeys, pigs, and goats, and in Lasso analysis they were the number of chickens, pigs,

and turkeys. The observed positive correlation between the number of beehives and total cases

should be explained by circulating viruses between animal and human. As, some zoonotic

viruses first circulate in animals, and sometimes they infect humans. After that, they adapt to

humans. A recent zoonotic transmission event of SARS-CoV-2 is reported[17]. It is crucial to

characterize circulating viruses in the human working environment, especially where an interface

with wildlife[18]. Spread of influenza virus during commercial movement of eggs is reported by

Pillai[19]. Poultry production in the extensive system increases the chance of SARS-CoV-2

transmission among other species[20]. Feathers detached from domestic ducks reported as a

source of environmental contamination to pathogenic avian influenza virus[21]. Qiu et al.(2020)

identified pigs, cows, goats, and sheep among the most likely intermediate animal hosts for

SARS-CoV-2 [22]. The outbreak of COVID-19 highlights viral spillover from animals[23].

There is the potential for the emergence of new coronavirus diseases in domestic birds, from

both avian and mammalian sources. For example, a novel species of *coronaviridae* family has

been identified recently[24]. *Coronaviridae*'s family viruses are likely can cross the species

barrier of viruses and infect other species, fishes, cattle, pigs, and humans [25].

Conclusion
147

Despite the awareness and readiness of many countries for COVID-19 during early 2020, why

did it spread so quickly? It seems FBT has received less attention relative to other routes. There

is need to invent a pasteurization-like-processes to implicate immediately before ingestion of

food. Of course, there will be challenges along the way of the revolution in the food preparation

culture. The perspective is to expand the surveillance of SARS-Cov-2 during the food production

chain. In conclusion, the results of the present study indicate that the reservoir of SARS-Cov-2 may be

some of the domestic animals. It should be noted that this study was conducted with data related to the

time (2013) before the effect of COVID_19 pandemic on food supply. It is recommended that virologists

examine the possibility of freshwater fish and chicken's eggs being reservoirs for SARS-Cov-2.

Material and Methods

Food consumption

We estimated the relationship between 20 food supply and Total Cases of covid19.

The methods include pair-wise correlation with Proc Corr, Stepwise Regression

with Proc Reg, LASSO regression with GLMSELECT Proc, Principal Component

Analysis with Proc Princomp, and Bayes analysis with Proc Genmod (SAS). Food

supply was quantity (kg/capita/yr.). Fourteen foods of animal origin and six foods

of plant origin were selected. The primary purpose of the study was to find animal

origin foods relationship with total cases, but some plant origin foods were chosen

to compare.

We started our analyses first time on March 2020. Then we repeated it several

times with updated COVID_19 data on April, May, and June. Finally we presented

the results of June' data.

Data

COVID_19 https://www.worldometers.info/coronavirus/   1 June 2020

Live Animal production data: http://www.fao.org/faostat/en/#data/QA

Food Supply - Crops Primary Equivalent:

http://www.fao.org/faostat/en/#data/CC

Food Supply - Livestock and Fish Primary Equivalent:

http://www.fao.org/faostat/en/#data/CL

We used base-*e* logs, also known as natural logs (LOG in SAS) for the

transformation of total cases and independent variables. In all analyses, the

accuracy of the model with non-transformed data was very low. So, all variables

has been transformed to normalize the distributions. With transformation of data,

the accuracy of the model increased to an acceptable level. In PCA, as a nonparametric method, we utilized non-transformed data, in which the variables are treated equally, i.e., that is, there are no dependent variables.

In comparison of models, Lasso's results are more reliable than stepwise regression and Bayes, because it's AIC was smaller and had higher accuracy as R-square (S1 Table).

S1 Table. R-Square, Adjusted R-square, AIC and number of data in the fitted models for food consumption.

	Stepwise	LASSO	BAYES
R-Square	0.48	0.52	-
Adj-R Square	0.47	0.48	-
AIC	-	301	551
N	131	131	131

Bayesian Analysis: Visual examination of the trace plot displays proper mixing, for one independent variable in presented in S8 Fig. The P-value in the Geweke Diagnostics (S2 Table) showed that the mean estimate of the Markov chain is stable over time. The positive probability that B1 greater than 0 is estimated.

S2 Table Geweke diagnostics from Genmode analysis. Pr > z shows

convergence for all variables. For abbreviations, see Table 1

Parameter*	z	Pr > |z|
Intercept	0.27	0.78
Pig meat	0.64	0.51
Bovine Meat	-0.21	0.82
Cheese	-0.74	0.45
Cream	-0.94	0.34
Eggs	-1.51	0.13
Freshwater fish	0.43	0.66
Honey	0.79	0.43
Whole milk	-0.04	0.97
Poultry meat	0.95	0.34
Demersal fish	-0.95	0.34
Cephalopods	0.79	0.43
Butter/ghee	1.89	0.06
Marine fish(other)	-0.46	0.65
Aquatic plants	0.58	0.56
Beverages(Alcoholic)	-1.18	0.24
Rice(Milled Equivalent)	-0.98	0.33
Potatoes and products	-1.01	0.31
Maize and products	-0.30	0.76
Wheat and products	0.50	0.61
Wine	-0.01	0.99
Scale	0.02	0.98

* Logarithmic transformation for all parameters

Although the probability of transmitting the virus through wine is zero, we

deliberately kept it in the initial model of our analysis. Results showed wine is

eliminated among significant variables of the suited model (Table 2). So we

concluded that just the energy content of food should not keep a variable in the

fitted model. So, in addition to energy content, there must be something else in

food-related COVID-19. The high-fat tissue in the body affects on the

pathogenesis of COVID-19[34]. We concentrate FB transmission of COVID-19

rather than the immunity response.

Animal and crop production

We utilized the GLMSLECT procedure of SAS 9.2 with the method of least

absolute shrinkage and selector operator (LASSO). The dependent variable was

total cases, and independent variables were beehives, cattle, chicken, pigs, goats,

turkeys, horse, rabbit, maize, barley, and apples. Maize, Barley, apples were

deliberately kept in a multi-variable model, to test whether plant material

production remains in the model compared to livestock production. We used the

GENMOD procedure of the SAS using the *BAYES* statement to obtain *Bayesian*

analysis. Spearman and Pearson correlation between stocks and TOTAL

CASEScalculated. Analyzes were performed with log-transformed data. Due to the

higher accuracy of the transformed data, their results were used.

There are references[33] that guide us to construct our hypothesis about the

essential relationship between COVID-19 and AOF include.

We provide the SAS code for an application of the used statistical methods [35].

We note that this work was posted as preprint on medrxiv[36] .

ACKNOWLEDGEMENTS:

Author contributions: Conceptualization and Visualisation: N. S. K., Methodology: S. A. R.

Writing: Both authors. The authors have declared no competing interest. Data and materials

availability: All data are available in the manuscript or the supplementary materials.

References

1.   Merler S, Pastore A, Mu K, Rossi L, Sun K. The effect of travel restrictions on the spread

of the 2019 novel coronavirus (COVID-19) outbreak. 2020;400: 395–400.

2.   Panel E, Biohaz H. Scientific Opinion on an update on the present knowledge on the

occurrence and control of foodborne viruses 1. 2011;9: 1–96.

doi:10.2903/j.efsa.2011.2190.

3.   Todd ECD, Greig JD, Todd E, Llc C. Viruses of foodborne origin□: a review. 2020; 25–

46.

4.   Hi ZS. REVIEW ARTICLE A brief review of foodborne zoonoses in China. 2020; 1497–

1504. doi:10.1017/S0950268811000872

5.   Newell DG, Koopmans M, Verhoef L, Duizer E, Aidara-kane A, Sprong H, et al.

International Journal of Food Microbiology Food-borne diseases — The challenges of 20

years ago still persist while new ones continue to emerge. International Journal of Food

Microbiology. 2010;139: S3–S15. doi:10.1016/j.ijfoodmicro.2010.01.021

6.   Kingsbury J, Lake R, Hewitt J, Smit E, King N. POTENTIAL FOR FOODBORNE

TRANSMISSION OF COVID-19☐: 2020; 1–51.

7.   FAO/WHO. [Food and Agriculture Organization of the United Nations/World Health

Organization]. Microbiological Risk Assessment Series. 2008;14: 151.

8.   Panon G, Tache S, Labie C. Respective stability of rotavirus and coronavirus in bovine

milk. 1988;68: 49–64.

9.   Hamdi A. Mohamed, Adham M. Abdou , Ekbal M. Adel, Sayed Ahmed H. Salem MIME-

H. DETECTION OF ROTA AND CORONA VIRUSES IN RAW MILK AND MILK

PRODUCTS. Benha veterinary medical journal. 2013;24: 79–85. Available:

http://www.bvmj.bu.edu.eg/abstract.php?ee=156

10.   Pirtle EC. Virus survival in the environment. 1991;10: 733–748.

11.   Moy GG. IUFoST / CIFST hold an Extraordinary Scienti fi c Roundtable on COVID-19

and Food Safety. npj Science of Food. 2020; 1–2. doi:10.1038/s41538-020-0068-2

12.   Sarah C. P. Williams. Kitchen Utensils Transfer Viruses. Available:

https://www.sciencemag.org/news/2012/12/kitchen-utensils-transfer-viruses

13.   Id SAA, Burnham PA, Boncristiani HF, Brody K. RNA virus spillover from managed

honeybees (Apis mellifera) to wild bumblebees (Bombus spp .). 2019; 1–13.

14.   Maori E, Trust W, Reports C. Discovery of RNA transfer through royal jelly could aid

development of honey bee vaccines. 2019; 3–5. doi:10.1016/j.celrep.2019.04.073

15.   Mutinelli F. The spread of pathogens through trade in honey bees and their products (

including queen bees and semen): overview and recent developments Honey bees.

2011;30: 257–271.

16.   Bosch A, Gkogka E, Le FS, Loisy-hamon F, Lee A, Lieshout L Van, et al. International

Journal of Food Microbiology Foodborne viruses□: Detection , risk assessment , and

control options in food processing. International Journal of Food Microbiology. 2018;285:

110–128. doi:10.1016/j.ijfoodmicro.2018.06.001

17.   Kujawski SA, Wong KK, Collins JP, Epstein L, Killerby ME, Midgley CM, et al. Clinical

and virologic characteristics of the first 12 patients with coronavirus disease 2019

(COVID-19) in the United States. Nature Medicine. 2020;26: 861–868.

doi:10.1038/s41591-020-0877-5

18.   Bourgarel M, Pfukenyi D, Becquart P, Morand S. Next-Generation Sequencing on

Insectivorous Bat Guano□: An Accurate Tool to Identify Arthropod Viruses of Potential

Agricultural Concern. 2019. doi:10.3390/v11121102

19.   Pillai SPS, Saif YM, Lee CW. Detection of influenza A viruses in eggs laid by infected

turkeys. Avian diseases. 2010;54: 830–833. doi:10.1637/9102-101209-Reg.1

20.   Cavanagh D, Cavanagh D. Coronaviruses in poultry and other birds Coronaviruses in

poultry and other birds. 2007;9457. doi:10.1080/03079450500367682

21.   Yamamoto Y, Nakamura K, Yamada M, Mase M. Persistence of Avian Influenza Virus (

H5N1) in Feathers Detached from Bodies of Infected Domestic Ducks □. 2010;76: 5496–

5499. doi:10.1128/AEM.00563-10

22.   Qiu Y, Zhao Y, Wang Q, Li J, Zhou Z, Liao C, et al. Predicting the angiotensin converting

284   enzyme 2 (ACE2) utilizing capability as the receptor of SARS-CoV-2. Microbes and

285   Infection. 2020;2. doi:10.1016/j.micinf.2020.03.003

286   23.   Tao Z, Tian J, Pei Y, Yuan M, Zhang Y, Dai F. A new coronavirus associated with human

287   respiratory disease in China. 2020;579. doi:10.1038/s41586-020-2008-3

288   24.   Papineau A, Berhane Y, Wylie TN, Wylie KM, Sharpe S, Lung O. Genome Organization

289   of Canada Goose Coronavirus , A Novel Species Identified in a Mass Die-off of Canada

290   Geese. Scientific Reports. 2019; 1–8. doi:10.1038/s41598-019-42355-y

291   25.   Leong JC. Fish Viruses. Encyclopedia of Virology. 2008; 227—234. doi:10.1016/b978-

292   012374410-4.00400-3

293   26.   Miller RS, Sweeney SJ, Slootmaker C, Grear DA, Salvo PA Di, Kiser D, et al. Cross-

294   species transmission potential between wild pigs , livestock , poultry , wildlife , and

295   humans□: implications for disease risk management in North America. Scientific Reports.

296   2017; 1–14. doi:10.1038/s41598-017-07336-z

297   27.   Kai Kupferschmidt. New Middle Eastern Virus Found in Camels. 2013. Available:

298   https://www.sciencemag.org/news/2013/11/new-middle-eastern-virus-found-camels

299   28.   Guan Y, Zheng BJ, He YQ, Liu XL, Zhuang ZX, Cheung CL, et al. Isolation and

300   characterization of viruses related to the SARS coronavirus from animals in southern

301   China. Science (New York, NY). 2003;302: 276–278. doi:10.1126/science.1087139

302   29.   Rizzo F, Edenborough KM, Toffoli R, Culasso P, Zoppi S, Dondo A, et al. Coronavirus

and paramyxovirus in bats from Northwest Italy. 2017; 1–11. doi:10.1186/s12917-017-1307-x

30. Mcmahon DP, Wilfert L, Paxton R, Brown M, Holloway R. Emerging Viruses in Bees⬚: From. 2018. doi:10.1016/bs.aivir.2018.02.008

31. Geisbert TW, Mire CE, Geisbert JB, Chan Y-P, Agans KN, Feldmann F, et al. Therapeutic treatment of Nipah virus infection in nonhuman primates with a neutralizing human monoclonal antibody. Science translational medicine. 2014;6: 242ra82-242ra82. doi:10.1126/scitranslmed.3008929

32. Gibbins JD, Hale C, Silver M, Fischer M, Steinberg J, Basler CA, et al. COVID-19 Among Workers in Meat and Poultry Processing Facilities —. 2020;69: 557–561.

33. Katy Askew. 2020 German slaughterhouses under fire over "insufficient precautionary measures" to block COVID-19 spread. 2020.

34. Maffetone PB, Laursen PB. The Perfect Storm⬚: Coronavirus (Covid-19) Pandemic Meets Overfat Pandemic. 2020;8: 1–6. doi:10.3389/fpubh.2020.00135

35. Rafat SA. DATA of Preprint: COVID_19's lost puzzle piece: the power of domestic animal reservoir https://www.medrxiv.org/content/10.1101/2020.06.16.20132464 2020 . DOI: 10.13140/RG.2.2.24000.56324

36. Rafat, S.A. saeedi N. Human food consumption patterns concerning COVID-19 pandemic. Available: https://www.medrxiv.org/content/10.1101/2020.06.16.20132464

Fig. 1 PCA results: the interrelationships between all variables. Some of AOF are closer to total cases. For abbreviations, see
Table 1.

Fig. 2 Potential transmission ways of COVID-19 based on the results of our analysis and bibliography. It is necessary to study the exchange of human
pathogen viruses in wildlife and livestock. Where and how suspicious foods become contaminated must be investigated. The potential spillover of pathogenic
viruses from arthropods hosts to humans should be investigated[23]. The cross-species transmission is reported[26]. There may be a type of contact among
flowers, bees, and bats. Infected humans could have transmitted MERS to the camels [27], similar situation is imaginable for SARS-Cov-2. SARS-CoV-2 and
MERS-Co are among the most notable examples of spillover into humans[28]. References for arrows: 1 [11,14,18,20,29,30]   2: [11,26]   3: [24,26]   4:[23,31]
5: [5,8,32] 6 : [4,32,33]

S1 Fig. Histogram of variables of Total Cases before (left) and after (right) logarithmic transformation.

S2 Fig. Histogram of some independent variables before and after logarithmic transformation.

First row, from Left to right: cream, log cream, pig meats, log of pig meats. Second row: Bovine meat, log of
bovine meat, eggs, log of eggs.

S3 Fig. Diagnostics panel. Fit diagnostics and residual by regressors, in multivariate regression with stepwise

selection of variables. logtot= logarithm of total cases

S4 Fig. Other Criteria Suggest Stopping the Selection (GLMSELECT proc of SAS). Progression of squared

errors (Left) and Fit criteria of LASSO analysis (Right). logtot= logarithm of total cases

357 **S6 Fig. Scree plot and percentage of variance explained by principal components of PCA analysis.**

361 **S7 Fig. Results of PROC PRINCOMP, includes 1 by 3 (left) and 2 by 3 (right) component pattern plots.**
362 Principal component analysis results showed that AOF are generally closer to total cases in the comparison with
363 other studied foods. For abbreviations, see Table 1.

366 **S8 Fig. Bayesian Analysis:** Visual examination of the trace plot displays proper mixing for Eggs.