Common genetic associations between age-related diseases

Handan Melike Dönertaş[1*], Daniel K. Fabian[1,2], Matías Fuentealba Valenzuela[1,2], Linda Partridge[2,3], Janet M. Thornton[1*]

[1]European Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge, CB10 1SD, UK.
[2]Institute of Healthy Aging, Department of Genetics, Evolution and Environment, University College London, London, UK
[3]Max Planck Institute for Biology of Aging, Cologne, Germany
*Correspondence: donertas.melike@gmail.com; thornton@ebi.ac.uk

Abstract

Age is a common risk factor in many diseases, but the molecular basis for this relationship is elusive. In this study we identified 4 disease clusters from 116 diseases in the UK Biobank data, defined by their age-of-onset profiles, and found that diseases with the same onset profile are genetically more similar, suggesting a common etiology. This similarity was not explained by disease categories, co-occurrences or disease cause-effect relationships. Two of the four disease clusters had an increased risk of occurrence from age 20 and 40 years respectively. They both showed an association with known aging-related genes, yet differed in functional enrichment and evolutionary profiles. Moreover, they both had age-related expression and methylation changes. We also tested mutation accumulation and antagonistic pleiotropy theories of aging and found support for both.

Keywords:

Aging, age-related disease, GWAS, UK Biobank, mutation accumulation, antagonistic pleiotropy

Introduction

Aging is associated with a time-dependent decrease in the functional integrity of organisms and an increase in susceptibility to pathologies[1]. The worldwide increase in lifespan has not been matched by an increase in healthspan, and there is a growing period of loss of function, and disease at the end of life[2]. Aging thus poses a significant global challenge, because it is the major risk factor for chronic conditions, including cardiovascular disease, cancer, and dementia[3]. Although these diseases involve different organs and pathologies, they all show a strong dependence on age[4] and could, therefore share common etiologies based upon the underlying mechanisms of aging. It is therefore important to understand if the aging process itself leads to different age-related conditions through common pathways, or if the age-dependency of different diseases has independent, time-dependent causes.

Despite the negative impact of aging on organismal fitness and functionality, it is widespread in the animal world as well as in humans[5] and has therefore been described as an evolutionary paradox[6]. Aging can nonetheless evolve as the force of natural selection weakens with age due to extrinsic hazard. Mutations that are deleterious only in later life can accumulate in the population through mutation pressure, because the force of natural selection eliminating them from the population declines with the age of onset of their effects (mutation accumulation theory of aging)[7]. Pleiotropic variants that are beneficial during early life but detrimental later in life can also become prevalent in populations through natural selection (antagonistic pleiotropy theory of aging)[8]. Thus, genome-wide germline genetic variants that increase the risk of diseases at old age may not be pruned by natural selection or may be associated with beneficial phenotypes earlier in life.

The risk of many **a**ge-**r**elated **d**iseases (ARDs) is influenced by genetic variation. **G**enome-**w**ide **a**ssociation **s**tudies (GWAS) have identified genetic variants that alter complex traits. Pleiotropy, where variants or genes influence multiple traits, is more prevalent than previously thought[9–12], indicating that different traits share common causal pathways[13]. Pleiotropy within the disease classification system[12] and in certain disease classes, such as immune-related diseases[14,15] and cancer[16], has been studied, but the understanding of pleiotropy in ARDs more broadly is limited. Some studies have investigated the common pathways between manually curated age-related traits[17–19]. Despite the challenges of combining results from different published datasets, these studies provided the first clues that at least some ARDs share common pathways, which are also related to a significant but limited number of longevity-regulating genes in model organisms. In this study, using disease age-of-onset profiles, we extend the previous efforts by providing the first data-driven classification of a large number of diseases according to their age-profile, followed by a genetic analysis in one of the largest and most comprehensive cohorts available. In this way, we provide a comparative genetic analysis between ARDs and non-ARDs and also between ARDs with different age profiles.

The **UK B**io**b**ank (UKBB)[20] includes genetic and health-related data for almost half a million participants. We extracted age-of-onset profiles for 116 diseases and identified unbiased clusters to define the relationship between disease incidence and age. We identified variants associated with each disease and compared the genetic associations between diseases based on these clusters. We first found that diseases with the same age profile share genetic associations, which cannot be explained by disease categories, co-occurrences, or mediated pleiotropy, and thus reflects a common etiology (Figure S1). We further characterized these shared associations compared to previously known longevity-associated genes, genes with age-dependent expression or methylation changes, and biological functions. Finally, we compared the variants associated with diseases that start to occur at different ages and identified different evolutionary characteristics, supporting the mutation accumulation and antagonistic pleiotropy theories of aging.

Results

Data

We used self-reported diseases and age-at-diagnosis covering 484,598 participants, and their genotypes in the UKBB[20]. Age-at-diagnosis for self-reported diseases ranges from the age of 0 to 70, and is less biased than age-at-diagnosis for ICD10 codes, which includes data for only after 1992[21]. Details of the UKBB data, quality control steps, and exploratory analyses are given in the Supplementary Information and Supplementary Figures S2-9. Self-reported diseases in the UKBB are hierarchically structured and the top nodes; such as cardiovascular or endocrine diseases, were considered as *disease categories* (Figure S7). We only analyzed common diseases (*i.e.* with at least 2,000 cases) that were not sex-limited (n=116 in 472 self-reported diseases). Importantly, we did not include cancer in our analyses as the interaction between genetic and environmental contributors is likely different from non-cancer illnesses, even though they may have a similar age-of-onset profile (for details, see Methods).

Age-of-onset clusters

*Figure 1: Age-of-onset profiles clustered by the PAM algorithm, using dissimilarities calculated with **t**emporal **cor**relation measure (CORT). The y-axis shows the number of individuals who were diagnosed with the disease at a certain age, divided by the total number of people having that disease. Values were calculated by taking the median value of 100 permutations of 10,000 people in the UKBB (see Methods). The x-axis shows the age-of-onset in years. Each line denotes one disease and is colored by disease categories. The heatmap in the right upper corner shows the percent overlap between categories and clusters. Numbers give the % of an age-of-onset cluster belonging to each category. Supplementary Figures S10-19 shows the distributions for each disease separately.*

Age is associated with increased risk of many diseases. In order to characterize the association between age and different diseases we first used age-at-diagnosis as a proxy to disease onset and derived disease age-of-onset profiles (Figure S10-19). On average, cardiovascular and endocrine diseases had a high median age-of-onset, while infections had the lowest age-of-onset (Figure S20). We then clustered diseases into 4 clusters (the optimum number determined by the gap statistic) using the PAM algorithm and disease dissimilarities calculated using CORT distance[22] (Figure 1, Table S1). Cluster 1 diseases (n=25) showed a

rapid increase with age after the age of 40; 11 were cardiovascular diseases, but the cluster
also included other diseases such as diabetes, osteoporosis, and cataract. Cluster 2 (n=51)
diseases started to increase in the population at an earlier age of 20, but had a slower rate of
increase with age; the diseases in this cluster were the most diverse, including 17
musculoskeletal, 13 gastrointestinal diseases, as well as others such as anemia, deep venous
thrombosis, thyroid problems, depression. Cluster 3 diseases (n=30) showed a low age
dependency with a mostly uniform distribution across ages, but with slight increases around
the ages of 10 and 60 years. This category included similar numbers of immunological,
neurological, musculoskeletal, gastrointestinal and respiratory diseases but all have an
'immune' component even if not classified in this way by the UKBB (e.g., inflammatory bowel
disease (gastrointestinal), asthma (respiratory), psoriasis (dermatology)). Cluster 4 (n=10) had
a peak at around 0-10 years of age and included respiratory diseases (n=5) and infections
(n=4). Notably, all infectious diseases were in this cluster.
129
Diseases in the same age-of-onset cluster show higher genetic similarity
Using linear mixed models implemented in BOLT-LMM[23], we performed GWAS for case
versus control on each disease separately and included approximately 10 million common
variants that pass quality control and have **m**inor **a**llele **f**requency (MAF) > 0.01 (see
Methods)[24,25]. Considering associations with the literature-standard p-value lower than 5×10^{-8}
as significant[26,27], we next quantified the associations for each disease, category, and age-
of-onset cluster (Figure S21). The **m**ajor **h**istocompatibility **c**omplex (MHC) region is excluded
from all analyses, as in the literature, because of its unusually high effect sizes and **l**inkage
disequilibrium (LD) patterns (chr6: 28,477,797 - 33,448,354)[28,29]. Out of 116 diseases, 36 had
no significant association and the total number of polymorphic sites with at least one significant
association was 93,817. The maximum number of significant associations per disease was
35,001 (hypertension) and the median and mean were 13.5 and 1389.3, respectively. We also
checked if diseases from different age-of-onset clusters vary in the number of associations.
Cluster 4 had hardly any significant associations (the disease with the maximum number of
associations had only 3 significant variants). Although cluster 1 had the highest number of
significant associations on average, the values across clusters 1, 2, and 3 were not
significantly different (Figure S21b). Moreover, endocrine, immunological, cardiovascular
diseases had the highest number of associations and infections had the lowest (Figure S21c).
Only 1% of the significant polymorphisms (n=932) were in coding regions, and of these 49%
(n=452) were missense and only 1% (n=10) were nonsense. We further found that 47% of
significant variants (n=43,810) were associated with multiple diseases, but only ~9% were
associated with multiple diseases from different categories (n=8,048) and again ~9% with
different age-of-onset clusters (n=8,801) (Figure S22).
153

154
Figure 2: (a) Network representation of the genetic similarities calculated by the overlaps between
significantly associated SNPs between diseases. Nodes (n=47) show diseases with a significant genetic
similarity to at least one disease and are colored by the age-of-onset cluster. Edges (n=167) are
weighted by the genetic similarity corrected by disease categories and co-occurrences. (b) The
difference between genetic similarity within and across the age-of-onset clusters. The y-axis shows
genetic similarity corrected by category and co-occurrence (raw values are available in Figure S23).
The x-axis groups similarities into different or same age-of-onset clusters. c) Network representation of
the causal relationships between diseases calculated using LCV. Each node (n=48) shows a disease,
colored by the age-of-onset cluster. Size of the nodes represent the number of significant causal
relationships between diseases, including both in and out degrees. Arrows show the causal relationship
between pairs with FDR corrected p≤0.01 and GCP>0.6. The inset bar plot shows the percent significant

We next sought to characterize the genetic similarities between diseases using a score that
shows the excess of overlapping associations between diseases, given the number of
significant associations for each disease (see Methods). Importantly, we calculated genetic
similarities between 80 diseases that have at least one significant association, excluding the
pairs that are vertically connected (*i.e.* ancestors to child) in the disease hierarchy (*e.g.*
essential hypertension and hypertension). We found 47 significant overlaps and diseases with
similar age-of-onset profiles showed a higher genetic similarity, even when controlled for
disease categories and co-occurrences (F-test $p=1.19 \times 10^{-8}$, Figure 2a-b). Moreover, this
trend was reproducible when each cluster was analyzed separately (Figure S24). While
correcting for the disease categories and co-occurrences, some true positive signals may be
removed from the analysis. However, this correction is necessary, as we used the same cohort
for multiple diseases and, thus, diseases that co-occur use the same set of samples.
Nevertheless, we retained a significant signal even after this correction, demonstrating that
diseases with a similar age-of-onset profile show increased genetic similarity compared to
those with different profiles, suggesting shared genetic associations (Figure S1).

We repeated the analysis using 1,703 previously defined LD blocks[30] instead of considering
all **s**ingle **n**ucleotide **p**olymorphisms (SNP) as independent. There was no significant genetic
similarity between diseases from different age-of-onset clusters (Figure S25) and the
similarities within the same age-of-onset cluster were not explained by the disease categories
(p=0.89) and co-occurrences (p=0.15). We further confirmed the results using **h**igh-**d**efinition
likelihood (HDL) inference of genetic correlations, which makes use of the genome-wide data
instead of the significant associations and also considers LD[31]. Similarity scores calculated by
SNP overlaps and HDL regression were significantly correlated ($\rho = 0.75$, p<2.2e-16), and the
similarities between diseases in the same age-of-onset cluster was significantly higher than
that of in different age-of-onset clusters (Figure S26), confirming the conclusion that diseases
with similar age-of-onset profiles show higher genetic similarity using another methodology.

Lastly, we checked whether economic status or diet is associated with our results and whether
that might drive the commonality between diseases. We found no association between the list
of shared SNPs involved in multiple diseases or multiple disease categories and SNPs
associated with the "Townsend deprivation index" or any type of special diet, except for gluten-
free diet[32]. There was a significant overlap between SNPs associated with gluten-free diet in
UKBB and SNPs associated with diseases spanning multiple categories in the age of onset
clusters 1 & 2, and 1 & 2 & 3. If two traits are associated, we would see an increased overlap
as the p-value cutoff is more stringent; however, this was not the case for any of the traits,
including the gluten-free diet (Figure S27).

Mediated pleiotropy does not explain higher genetic similarities within age-of-onset
clusters
We next asked if mediated pleiotropy, rather than a common etiology (Figure S1), may explain
higher similarity within age-of-onset clusters. Using a recent methodology developed by
O'Connor & Price, we tested for partial or fully causal relationship between diseases[33]. In
particular, the method identifies if a **l**atent **c**ausal **v**ariable (LCV) mediates the genetic
correlation between diseases. Using a genetic causality proportion, it assigns a causal
relationship if one of the diseases is more strongly correlated with the LCV. The authors report
that, unlike mendelian randomization, this method can distinguish between the correlation due
to common etiology and causation. We tested for potential causation between 60 diseases,
excluding the ones with less than 10 significant genetic variants and low heritability estimates
($Z_h<7$)[33]. Also, similar to genetic similarities, we did not calculate the causation between
diseases that are vertically connected in the disease hierarchy. Following the same
significance criteria proposed in the methods article (FDR corrected p≤0.01 and mean **G**enetic

Causality **P**roportion (GCP)>0.6), we found significant evidence for full or partial genetic
causality in 91 disease pairs between 48 out of 116 diseases in our analysis (Figure 2c, Table
S2). Using Fisher's exact test, we tested if mediated pleiotropy was more common between
diseases in certain age-of-onset clusters but did not find any significant difference (FDR
corrected p>0.1 for all comparisons, inset bar plot in Figure 2c). Thus, although we detected
mediated pleiotropy between some diseases, higher genetic similarities within the same age-
of-onset clusters (Figure 2a-b) were not explained in this way and were more likely to be driven
by common etiologies.
229
We also investigated the diseases with the highest involvement in mediated pleiotropy. **D**eep
vein **t**hrombosis (DVT) (n=14), venous thromboembolic disease (n=13), and pulmonary
embolism (n=9) had the highest number of *out* degrees, meaning they were found as causal
for multiple diseases, including all 3 age-of-onset clusters and 5 different disease categories.
Gastro-**o**esophageal **r**eflux (GORD)/gastric reflux (n=10) and esophageal disorder (n=8), on
the other hand, had the highest number of *in* degrees, meaning there are multiple diseases
detected as causal. These causal diseases spanned 5 disease categories and age-of-onset
clusters 1 and 2.
238
Genes associated with the age-dependent disease clusters overlap with known
longevity modulators and independent aging-related traits
We next mapped all variants to genes based on proximity or known **e**xpression **q**uantitative
trait **l**oci (eQTLs) using the GTEx (v8) eQTL associations[34] (see Methods). We first analyzed
the overlap between diseases and tissue-specific eQTLs (Figure S28). The results included
tissue-disease pairs that were plausible, such as liver and high cholesterol. However, overall,
the number of tissue-specific eQTLs were biased due to the varying number of samples in
each tissue and thus, we used eQTLs in all tissues to map SNPs to genes. We confirmed that
the SNPs associated with diseases spanning multiple disease categories have eQTLs in
multiple tissues (Figure S29). To assess the reproducibility of the genes identified, we
compared the significant hits with all those reported in the GWAS Catalog. We verified that
most of the diseases had significant overlaps with the same or associated traits in the GWAS-
Catalog (*e.g.* hypertension and blood pressure or osteoporosis and bone density), confirming
that our results were reproducible with independent data (Table S3).
253
We next compiled the genes associated with multiple diseases and multiple categories and
grouped them based on the age-of-onset cluster of the associated diseases (Table S4). In
particular we created two sets of genes, '*multidisease*' and '*multicategory*', for clusters 1, 2,
and 3. We excluded cluster 4 because the number of variants significantly associated with this
cluster was low (n=7 associated with 5 diseases), mapping to only 3 genes (*ZPBP2*, *NPC1L1,*
TARID). We also compiled genes associated with multiple diseases or categories in
combinations of different age-of-onset clusters. Importantly, genes associated with multiple
clusters are not in the gene sets for individual clusters as the latter only include the genes
specific to individual clusters, *i.e.* cluster 1, cluster 2 and cluster 1 & 2 genes included all
mutually exclusive sets.
264
Using these lists, we sought to understand if the common genes between diseases with the
same age-of-onset profile had previously been associated with aging. We compared the
multidisease and *multicluster* gene lists with the literature-based aging databases: GenAge
human (genes associated lifespan in humans or closely related species), human orthologs in
GenAge model organism (genes modulating lifespan in model organisms), CellAge (genes
regulating cellular senescence), DrugAge targets (drugs modulating lifespan in model
organisms), and all databases combined[35–37] (Figure 3a). In general, genes associated with
clusters 1 and 2, but not Cluster 3, showed significant enrichment with known aging-related
genes. The list of overlapping genes is given in Table S5. The CellAge database showed the
significant overlaps with genes associated with clusters 1, 2, and '1 & 3'. DrugAge targets had
a significant overlap with clusters 1, 2, and '1 & 2'. GenAge Human only had significant

association with genes associated with cluster '1 & 2'. GenAge model organism data significantly overlapped with genes associated with all clusters (1 & 2 & 3). And the combination of all aging-related gene sets showed significant overlaps with clusters 1, 2, 1 & 2, and 1 & 2 & 3. In conclusion, although the association is established through a small subset of genes as also reported in the literature[17,18], the clusters 1 and 2, constituting age-dependent profiles, shared a significant genetic component with known longevity- and senescence-modulators, while cluster 3 did not. We further confirmed the aging relevance of our cluster 1 and cluster 2 genes using independent GWAS results for 'aging' and 'longevity' traits as well as Alzheimer's, Parkinson's, and age-related macular degeneration diseases which are not represented in our data (Figure S30). The significant overlaps include longevity associated *APOC1*, *APOC1P1*, *TOMM40*, and aging associated *APOE*, *CHRNA3*, *DEF8*, *LINC01239*, *TRIM59* genes, some of which also overlap with Alzheimer's, Parkinson's and age-related macular degeneration.

Figure 3: a) Overlap between known aging-related genes in databases and genes associated with diseases in different clusters. The x-axis shows log2 enrichment score, and the y-axis shows the age-of-onset clusters. The numbers of genes in each cluster (for both Multidisease and Multicategory genes) are given. The size of the points shows the statistical significance (large points show marginal p-

Age-related changes in expression and methylation of the genes associated with
diseases
To further explore the association between the disease-associated SNPs and ageing, we
analyzed the age-related expression and methylation profiles of the genes associated with
SNPs identified in our study (Figures S31-33). Using tissue-specific eQTL-gene associations,
we first checked whether the disease-associated SNPs might induce gene expression
changes that mimic the changes that occur at the age of 40, which might explain the age-of-
onset distribution of the first cluster. The multicategory cluster 1 eQTLs that increase the
expression of genes indeed mimicked the age-related expression changes (Figure S31,
expression difference is positive at the age of 40 and is significantly different from 30 and 50).
They were enriched in telomere maintenance and organization, columnar epithelial cell
differentiation, glucose metabolic process, and multicellular organismal signaling ($p < 0.01$ but
n.s. after multiple testing correction). Interestingly, multicategory cluster 2 and 3 eQTLs that
decrease the expression of genes also mimicked the age-related changes (Figure S32,
expression difference is negative at the age of 40 and is significantly different from 30 and 50).
These genes were enriched in immune response-related categories (Table S6). Thus, the
overall picture is complex and requires further investigation, but it suggests age-related
increases in the expression of genes associated with multicategory cluster 1 SNPs might
influence the age-of-onset curve. Still, immune-related genes that decrease in expression may
also contribute to the occurrence of cluster 2 and 3 diseases in late ages.

We next analyzed the association between age-of-onset clusters and methylation by
comparing the multidisease and multicategory gene sets with i) DNA methylation
("GO:0006306", n = 29) and demethylation ("GO:0080111", n = 20) GO Terms, and ii)
differentially methylated genes reported in previously published studies[38,39]. Although DNA
demethylation category included none of the multidisease or multicategory genes, the DNA
methylation category had 4 overlapping genes. GNAS and HELLS are multidisease cluster 1
genes, and GATAD2A is a multicategory cluster 1 gene. Disruption in the ortholog of HELLS
gene in mice (Hells) was found associated with premature aging[35,40]. The fourth gene, CTCF,
is a multidisease cluster 2 gene with no known association with aging. Although the overlap is
only specific to clusters with age-dependent profiles, these GO terms were not significantly
enriched in the disease-associated genes (Figure 3). Next, using the published results of two
external methylation data analyses[38,39], we found a significant overlap between the genes that
show differential methylation with age and the clusters with age-dependent profiles, clusters
1, 2 and 1 & 2 (Figure S33). Overall, although there was no significant association with the
regulators of DNA methylation, we found age-related changes in gene expression and
methylation for the late-onset clusters.

Genes associated with different age-of-onset clusters have different functions
Gene **o**ntology (GO) enrichment analyses were applied to the gene lists, including **b**iological
process (BP), **m**olecular **f**unction (MF) and **c**ellular **c**omponent (CC) categories. Cluster 1 was
associated with many lipoprotein-related categories, cellular signaling, and blood pressure
(Figure 3b). Cluster 2 showed association to plasminogen activation and protein activation
cascade (Figure 3c). Cluster 3 and clusters `2 & 3` had associations to many immune-related
categories and cell adhesion (Figure 3d, e). Genes in clusters '1 & 3' did not have any
significant associations. Genes associated with cluster '1 & 2' were related to nucleosome
complex, glucose homeostasis, retinoic acid binding (Figure 3f). Genes associated with at

least one disease in all clusters ('1 & 2 & 3') showed association with interleukin-7 response, differentiation, telomere as well as nucleosome complex (Figure 3f). Since cluster 3 did not have an age-dependent profile, the association with nucleosome complex could represent pleiotropic genes in general. Here we listed the categories that are representative to all other significant functional groups. The full list is given in Table S7, and the procedure of selecting representatives is described in the Methods. Overall, these results suggest that, although cluster 1 and cluster 2 genes were both linked to previously identified aging-related genes, they have distinct functional profiles.

Evolution of aging and age-related diseases

Figure 4: Risk allele frequency distributions (y-axis) for different age-of-onset clusters (x-axis) in the UKBB for a) SNPs associated with one disease (excluding antagonistic associations), b) SNPs specific to one cluster (excluding antagonistic associations) (ns: p>0.05, *:p≤0.05, **: p≤0.01, ***: p≤0.001, ****: p≤0.00001), and c) SNPs that have antagonistic association with cluster 1 and 2 (excluding agonists between cluster 1 and 2). d) The same as panel c but for different 1000 Genomes super-populations (ALL: complete 1000 Genomes cohort, AFR: African, AMR: Ad Mixed American, EAS: East Asian, EUR: European, SAS: South Asian).

Lastly, we sought to understand the abundance of disease-associated variants in the population and their relationship with the evolutionary theories of aging. **M**utation **a**ccumulation (MA) theory explains the functional decline at later ages by relatively lower selection pressure on deleterious germline variants that are functional at later ages. Accordingly, we first hypothesized that SNPs associated with later-onset diseases (Cluster 1) would have a higher frequency than the SNPs associated with diseases that occur at earlier ages (Clusters 2 and 3), which are presumably under stronger selection pressure. Not finding any difference in the power to detect SNPs for different clusters by comparing median MAF and number of cases (Figure S34), we compared the allele frequencies associated with different age-of-onset clusters[41]. As SNPs which are close together in the genome are expected to have similar allele frequencies due to linkage, we calculated the median risk allele frequency for SNPs within previously defined LD blocks[30]. Supporting the MA, diseases of cluster 1 had significantly

higher risk allele frequencies than cluster 2, both for the SNPs associated with one disease
(Figure 4a, Wilcoxon test p=0.00033) or with one cluster (Figure 4b, Wilcoxon test p=0.0068,
also confirmed by bootstrapping n=100 loci for B=1,000; Figure S35). We further confirmed
that this trend is not specific to the UK population, as we obtained comparable results in all
super-populations of the 1000 Genomes Project[42] (Figure S36-37). Variants associated with
cluster 3, which includes immune-related diseases, were not significantly different from those
associated with cluster 1, although cluster 3 diseases can occur even at an earlier age.
Moreover, although the difference in the median allele frequencies was not significant, the
shapes of distributions were different, with a significant shift towards higher risk allele
frequencies only in cluster 1 (p_{cl1}=0.05, p_{cl3}=0.86 calculated using 10,000 permutations). High
minor allele frequencies, thus a higher variation we observed in cluster 3 is in line with the
previous suggestion that immune-related genes are under long-term balancing selection in
humans[43], although positive selection also influences immunity[44–46].
395
To test the **a**ntagonistic **p**leiotropy theory (AP), we first asked if the diseases with different
onsets have an excess of antagonistic SNPs. Similar to a previous study[41], we defined a
pleiotropic biallelic SNP as *agonistic* if the risk allele is the same for different diseases, and as
antagonistic if opposite alleles are associated with increased risk for different diseases. If one
of these diseases is under a stronger negative selection, then the risk allele of the other
disease could increase over time. Comparing the proportion of agonist and antagonist SNPs
within and between the age-of-onset clusters, we found that there is an excess of antagonistic
pleiotropy between diseases with different age-of-onset profiles (Fisher's exact test p<0.001,
Table S8). Next, we tested the differences in risk allele frequencies between the clusters, as
AP predicts a higher risk allele frequency for late-onset diseases. The risk allele frequencies
for antagonistic associations between cluster 1 and 3 or 2 and 3 did not show any significant
difference in the UKBB or 1000 Genome super-populations (Figures S38-39). Interestingly,
the difference between the risk allele frequencies for cluster 1 and cluster 2 was also not
significant for the UKBB population (Figure 4c). However, all 1000 Genome super-populations
except for Europeans had higher risk allele frequencies for cluster 1 diseases (Figure 4d). We
hypothesized that this is mainly due to increased power when testing the antagonistic
associations with frequency closer to 0.5. We thus investigated the allele frequency
differences for the significant variants with increased effect sizes. Indeed, associations with a
larger effect size showed the expected differences in allele frequencies, although the number
of independent loci was limited (Figure S40). We also examined the type of diseases and
genes associated with antagonistic pleiotropy. The main driver of the pattern was the loci with
ABCG8 and *ABCG5* genes, showing antagonistic relationship for high cholesterol (cluster 1)
and other lipid-related diseases in cluster 2, such as gallbladder disease and cholelithiasis.
Another locus included variants that show antagonistic relationship with cardiovascular
disease (cluster 1) and the cluster 2 diseases gout (*ADH1B*), osteoarthritis and joint disorder
(*SLC39A8*), and osteoarthritis (*BANK1*). Another potential candidate was a locus associated
with hypertension (cluster 1) and musculoskeletal diseases (cluster 2), but this locus included
multiple candidate genes (Table S9). Nevertheless, our comparison is between common
diseases that occur after the age of 20 and 40, which are both after the average age at first
reproduction and therefore the start of the decrease in the force of natural selection[47]. Thus,
a better comparison would include the mutations causing rare developmental diseases, which
are not available in the UKBB.
428
429 **Discussion**
430
The number and the incidence of diseases increase with age. In this study, we explored
whether this results from a common genetic component among ARDs, which might also be
linked to aging. We compared genetic associations and age-of-onset distributions of 116 self-
reported diseases in the UKBB and found shared variants, genes and pathways, which were
also associated with aging.

Using an unsupervised, data-driven approach to classify diseases based on their age-of-onset
profiles, we found 4 main clusters; i) diseases that rapidly increase after 40 years of age, ii)
diseases that increase after 20 years of age, iii) diseases with no age-related pattern, and iv)
diseases that peak at around 10 years of age. Notably, unlike previous studies[18,48], by using
this unsupervised approach, we detect a distinction between cluster 1 and cluster 2, which
both show age-dependency but distinct age-of-onset distributions. These two clusters were
associated with genes with different functional and evolutionary characteristics, although they
both overlap with known aging-related genes. Moreover, germline mutations may influence
late-onset diseases through affecting the function of genes that show differential expression
or methylation with age.

Based on genetic associations, diseases with similar age-of-onset profiles showed a higher
genetic similarity on average, compared to diseases in other clusters, even when controlled
for disease categories and co-occurrences (Figure 2a-b). Moreover, this similarity within age-
of-onset clusters was not explained by mediated pleiotropy, in which one of the diseases is
causal for the other one, suggesting instead a common etiology. We then studied the genes
involved and found that genes associated with clusters 1 and 2 (both constituting ARDs) are
enriched with known longevity- and senescence-modulators, while genes associated with
cluster 3, which does not show an age-dependent profile, did not show this enrichment. In
addition, we found that genes associated with different age-of-onset clusters have different
functions. Comparing the risk allele frequencies of variants associated with different age-of-
onset profiles, we found support for both mutation accumulation and antagonistic pleiotropy
theories of aging, although the number of independent loci supporting the second was limited.
Lastly, we provide a list of drugs that target the common genetic component between ARDs,
which may in the future be tested for their effects on multimorbidity and polypharmacy
associated with late life.

In this study, we had a limited age range, covering individuals up to 65 years old and thus,
could not analyze diseases of later ages. Neither did we consider the cancers or changes in
regulation of gene expression for the same cohort, which are affected not only by aging, but
also various environmental or intrinsic factors. Future cohorts with a broader age range and
spanning multi-omics data, somatic mutations, health outcomes, and lifestyle information, will
enable a better understanding of the genetic mechanisms of age-of-onset determination and
establishing the causal link with candidate genes. Despite these limitations, we present a novel
approach to study ARDs using an unbiased, data-driven approach and show that ARDs share
common genetic associations linked to aging. Our results suggest that targeting the common
pathways between multiple ARDs could offer compression of late life multimorbidity as well as
alleviating the effects of polypharmacy.

Software Availability
All the code used to perform analyses is available in GitHub:
https://github.com/mdonertas/ukbb_ageonset

Data Availability
The full set of GWAS results from this study can be accessed using BioStudies (S-BSST407)
and all results generated in the analysis are provided as Supplementary Datasets and Tables.

Ethics Statement
Conflict of interest
The authors declare that they have no conflict of interest.

Author Contributions

H.M.D conceived and designed the study with contributions from L.P. and J.M.T.. H.M.D analyzed the data with the help of D.K.F and M.F.V.. H.M.D. interpreted the results and wrote the manuscript with contributions from all authors. All authors read, revised and approved the final version of this manuscript.

Acknowledgment

This research has been conducted using the UK Biobank Resource (application no. 30688). The authors thank the GWAS-Catalog team for providing the list of studies using UK Biobank data; James Stephenson and Roman Laskowski for their help in running VarMap tool; and Mehmet Somel, Susan Ozanne, Pedro Beltrao, and Wolfgang Huber for fruitful discussions. H.M.D. is a member of Darwin College, University of Cambridge.

Funding Statement

This work is funded by EMBL (H.M.D., J.M.T.), the Wellcome Trust (098565/Z/12/Z; L.P., J.M.T), and Comisión Nacional de Investigación Científica y Tecnológica - Government of Chile (CONICYT scholarship; M.F.V.).

References

1. López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The hallmarks of aging. *Cell* **153**, 1194–1217 (2013).
2. Crimmins, E. M. Lifespan and Healthspan: Past, Present, and Promise. *Gerontologist* **55**, 901–911 (2015).
3. Partridge, L., Deelen, J. & Slagboom, P. E. Facing up to the global challenges of ageing. *Nature* **561**, 45–56 (2018).
4. Niccoli, T. & Partridge, L. Ageing as a risk factor for disease. *Curr. Biol.* **22**, R741-52 (2012).
5. Flatt, T. & Partridge, L. Horizons in the evolution of aging. *BMC Biol.* **16**, 93 (2018).
6. Medvedev, Z. A. An attempt at a rational classification of theories of ageing. *Biol. Rev. Camb. Philos. Soc.* **65**, 375–398 (1990).
7. Medawar, P. B. Unsolved problem of biology. *Med. J. Aust.* **1**, 854–855 (1953).
8. Williams, G. C. Pleiotropy, Natural Selection, and the Evolution of Senescence. *Evolution* **11**, 398–411 (1957).
9. Bulik-Sullivan, B. *et al.* An atlas of genetic correlations across human diseases and traits. *Nat. Genet.* **47**, 1236–1241 (2015).
10. Pickrell, J. K. *et al.* Detection and interpretation of shared genetic influences on 42 human traits. *Nat. Genet.* **48**, 709–717 (2016).
11. Cross-Disorder Group of the Psychiatric Genomics Consortium *et al.* Genetic relationship between five psychiatric disorders estimated from genome-wide SNPs. *Nat. Genet.* **45**, 984–994 (2013).
12. Cortes, A., Albers, P. K., Dendrou, C. A., Fugger, L. & McVean, G. Identifying cross-disease components of genetic risk across hospital data in the UK Biobank. *Nat. Genet.* **52**, 126–134 (2020).
13. Solovieff, N., Cotsapas, C., Lee, P. H., Purcell, S. M. & Smoller, J. W. Pleiotropy in complex traits: challenges and strategies. *Nat. Rev. Genet.* **14**, 483–495 (2013).
14. Ellinghaus, D. *et al.* Analysis of five chronic inflammatory diseases identifies 27 new associations and highlights disease-specific patterns at shared loci. *Nat. Genet.* **48**, 510–518 (2016).
15. Parkes, M., Cortes, A., van Heel, D. A. & Brown, M. A. Genetic insights into common pathways and complex relationships among immune-mediated diseases. *Nat. Rev. Genet.* **14**, 661–673 (2013).
16. Bien, S. A. & Peters, U. Moving from one to many: insights from the growing list of pleiotropic cancer risk genes. *Br. J. Cancer* **120**, 1087–1089 (2019).

17. Johnson, S. C., Dong, X., Vijg, J. & Suh, Y. Genetic evidence for common pathways in
human age-related diseases. *Aging Cell* **14**, 809–817 (2015).
18. Fernandes, M. *et al.* Systematic analysis of the gerontome reveals links between aging
and age-related diseases. *Hum. Mol. Genet.* **25**, 4804–4818 (2016).
19. Wang, J., Zhang, S., Wang, Y., Chen, L. & Zhang, X.-S. Disease-aging network reveals
significant roles of aging genes in connecting genetic diseases. *PLoS Comput. Biol.* **5**,
e1000521 (2009).
20. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data.
Nature **562**, 203–209 (2018).
21. UKB : Data-Field 41262. https://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=41262.
22. Chouakria, A. D. & Nagabhushan, P. N. Adaptive dissimilarity index for measuring time
series proximity. *Adv. Data Anal. Classif.* **1**, 5–21 (2007).
23. Loh, P.-R. *et al.* Efficient Bayesian mixed-model analysis increases association power in
large cohorts. *Nat. Genet.* **47**, 284–290 (2015).
24. Anderson, C. A. *et al.* Data quality control in genetic case-control association studies. *Nat.*
Protoc. **5**, 1564–1573 (2010).
25. Zhou, W. *et al.* Efficiently controlling for case-control imbalance and sample relatedness
in large-scale genetic association studies. *Nat. Genet.* **50**, 1335–1341 (2018).
26. Pe'er, I., Yelensky, R., Altshuler, D. & Daly, M. Estimation of the Multiple Testing Burden
for Genomewide Association Studies of Common Variants. *Nature Precedings* (2007)
doi:10.1038/npre.2007.359.1.
27. Panagiotou, O. A., Ioannidis, J. P. A. & Genome-Wide Significance Project. What should
the genome-wide significance threshold be? Empirical replication of borderline genetic
associations. *Int. J. Epidemiol.* **41**, 273–286 (2012).
28. MHC region of the human genome - Genome Reference Consortium.
https://www.ncbi.nlm.nih.gov/grc/human/regions/MHC?asm=GRCh37.
29. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from
polygenicity in genome-wide association studies. *Nat. Genet.* **47**, 291–295 (2015).
30. Berisa, T. & Pickrell, J. K. Approximately independent linkage disequilibrium blocks in
human populations. *Bioinformatics* **32**, 283–285 (2016).
31. Ning, Z., Pawitan, Y. & Shen, X. High-definition likelihood inference of genetic correlations
across human complex traits. *Nat. Genet.* (2020) doi:10.1038/s41588-020-0653-y.
32. UK Biobank — Neale lab. http://www.nealelab.is/uk-biobank/.
33. O'Connor, L. J. & Price, A. L. Distinguishing genetic correlation from causation across 52
diseases and complex traits. *Nat. Genet.* **50**, 1728–1734 (2018).
34. Gamazon, E. R. *et al.* Using an atlas of gene regulation across 44 human tissues to inform
complex disease- and trait-associated variation. *Nat. Genet.* **50**, 956–967 (2018).
35. Tacutu, R. *et al.* Human Ageing Genomic Resources: new and updated databases.
Nucleic Acids Res. **46**, D1083–D1090 (2018).
36. Barardo, D. *et al.* The DrugAge database of aging-related drugs. *Aging Cell* vol. 16 594–
597 (2017).
37. Avelar, R. A. *et al.* A multidimensional systems biology analysis of cellular senescence in
aging and disease. *Genome Biol.* **21**, 91 (2020).
38. Adelman, E. R. *et al.* Aging Human Hematopoietic Stem Cells Manifest Profound
Epigenetic Reprogramming of Enhancers That May Predispose to Leukemia. *Cancer*
Discov. **9**, 1080–1101 (2019).
39. Marttila, S. *et al.* Ageing-associated changes in the human DNA methylome: genomic
locations and effects on gene expression. *BMC Genomics* **16**, 179 (2015).
40. Sun, L.-Q. *et al.* Growth retardation and premature aging phenotypes in mice with
disruption of the SNF2-like gene, PASG. *Genes Dev.* **18**, 1035–1046 (2004).
41. Rodríguez, J. A. *et al.* Antagonistic pleiotropy and mutation accumulation influence human
senescence and disease. *Nat Ecol Evol* **1**, 55 (2017).
42. 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation.
Nature **526**, 68–74 (2015).
43. Bitarello, B. D. *et al.* Signatures of Long-Term Balancing Selection in Human Genomes.

597   *Genome Biol. Evol.* **10**, 939–955 (2018).

598 44. Kosiol, C. *et al.* Patterns of positive selection in six Mammalian genomes. *PLoS Genet.*
599    **4**, e1000144 (2008).

600 45. Nielsen, R. *et al.* A scan for positively selected genes in the genomes of humans and
601    chimpanzees. *PLoS Biol.* **3**, e170 (2005).

602 46. Shultz, A. J. & Sackton, T. B. Immune genes are hotspots of shared positive selection
603    across birds and mammals. *Elife* **8**, (2019).

604 47. Fisher, R. A. The genetical theory of natural selection. **272**, (1930).

605 48. Wolfson, M., Budovsky, A., Tacutu, R. & Fraifeld, V. The signaling hubs at the crossroad
606    of longevity and age-related disease networks. *Int. J. Biochem. Cell Biol.* **41**, 516–520
607    (2009).

Methods

UK Biobank Data

Data was downloaded using bash and following the guidelines provided by the UK Biobank.

Sample quality control

After excluding all samples from individuals who have withdrawn their data from UK Biobank, we first filtered out all samples without genotypes (N = 14,248). Then, we used the following criteria for the remaining 488,295 samples.

Discordant sex: Data includes two entries for sex: 1) self-reported and 2) genetic sex determined using the call intensities on sex chromosomes. There are multiple reasons why these two entries may not correspond, such as sample mishandling, errors in data input, transgender individuals, and sex chromosome aneuploidies[1,2]. Since we used sex as a covariate in our GWAS model, we preferred to be cautious about this issue and excluded all cases where the genetic sex and self-reported sex did not correspond and all cases where sex chromosome aneuploidy was detected. Specifically, we used the fields '31-0.0' (Sex) and '22001-0.0' (Genetic sex) to compile discordant information. There were 235 self-reported males being identified as female by the genetics, and 143 self-reported females being identified as males by the genetics. We excluded these 378 cases, 0.077% of the data. Moreover, field '22019-0.0' (Sex chromosome aneuploidy) is used to exclude cases with sex chromosome aneuploidy. There were 651 cases of aneuploidy, 0.133% of all data. 181 of these cases (27.80% of aneuploidy cases) were also detected as discordant information in the first step. This corresponds to 47.88% of discordant sex cases. Overall, we identified 848 samples to be excluded based on this criterion.

Genotype call rate & Heterozygosity: Genotype missingness and heterozygosity are widely used as a measure of DNA sample quality. For quality filtering based on missingness and heterozygosity we only used the suggested exclusions by UK Biobank. Specifically, we used the field '22010-0.0' (Recommended genomic analysis exclusions) and determined the cases with 'poor heterozygosity/missingness' (N = 469). We next used the field '22018-0.0' (Genetic relatedness exclusions) and noted down the cases with 'Participant self-declared as having a mixed ancestral background' (N = 692), and the cases with 'High heterozygosity rate (after correcting for ancestry) or high missing rate' (N = 840). Lastly, there were 968 cases that are suggested as outliers for heterozygosity or missing rate, field '22027-0.0' (Outliers for heterozygosity or missing rate). We then checked the scatter plots for logit(Missingness) vs. Heterozygosity for each Ethnic Background, in accordance with the identification of samples to exclude by the UK Biobank[2] (Figure S43). Logit transformation is used to linearize sigmoidal distribution of missingness. Investigation of heterozygosity can detect DNA sample contamination, inbreeding, or mixed ethnicity[1]. This quality check reveals when people with a mixed ethnicity tend to have a higher heterozygosity, even after correcting for PCs. We confirmed these are in accordance with the original article and excluded the samples suggested by the UK Biobank.

Overall, there were 3,697 samples excluded based on these two criteria. Please note that the numbers presented above may not add up to this number, because there were some samples excluded based on multiple criteria. The percent overlap across multiple criteria is given in Figure S44.

Preparing the trait data

Using the samples that passed the quality control (N=484,598), we subsetted the data so that it included only the baseline visit. Apart from the data that is already available in UK Biobank, we calculated some other values: *1) BMI:* Using the columns for 'Weight' and 'Standing height' we calculated BMI as: Weight / (Standing Height / 100)^2, *2) Parent Age at Death - Minimum:* The youngest age at which either parent died. *3) Parent Age at Death - Maximum:* The age of

death for the parent who lived longest. *4) Parent Age at Death - Average:* The average age of death for the two parents. If neither of the parents died, or if the data was unavailable, these values (2-4) were set to be NA. If only one parent died, we use the corresponding age as both the minimum, maximum, and average. *5) The number of self-reported non-cancer diseases:* The number of unique self-reported non-cancer illnesses each participant recorded in the baseline recruitment. *6) The number of self-reported cancers:* The number of unique self-reported cancers each participant recorded in the baseline recruitment. *7) Self-reported diseases after taking the disease hierarchy into consideration (Propagated disease data):* The self-reported diseases in UK Biobank are not independent, but rather are organized in a hierarchical manner. Using the relationship information between diseases, we propagated disease-participant associations, upwards, including terms higher up the tree. For example, if a person reports having "essential hypertension", we also annotate that person with "hypertension", and "cardiovascular disease". *8) Age at diagnosis for the self-reported diseases after taking the disease hierarchy into consideration (Propagated age at diagnosis data):* We re-defined age at diagnosis using the minimum age at diagnosis for all the diseases that were child term for a particular disease in the disease hierarchy. *9) The number of self-reported non-cancer diseases after taking the disease hierarchy into consideration (Propagated number of non-cancer diseases):* The number of unique self-reported diseases each participant records after taking into account the data propagation. *10) Age when the last deceased person died:* We calculated the age of each person when the last death entry in the UKBB happened. This value is used to calculate the proportion of people who died at a certain age interval in Figure S2c.

Selecting diseases to analyze
We calculated the disease occurrences for all self-reported diseases in UK Biobank. Specifically, among the cohort we used, we calculated how many participants and what proportion of males and females reported each disease. Since we analyzed the same set of SNPs that have MAF>=0.01 across multiple diseases, to decrease the false positive rate in GWAS, we limited the diseases to a subset with at least 2,000 cases (n = 129 out of 472)[3]. Moreover, we only focused on diseases that were common and not sex-limited, *i.e.* we only considered diseases that are seen in 1 in every 1,000 males and females (n = 189 out of 472). The intersection of these two conditions was 116 diseases and we excluded all others.

We only analyzed self-reported non-cancer diseases (field '20002') and did not combine self-reported cancers (field '20001'), mainly because i) the number of cases is low (45,224 compared to 384,906 for other diseases), ii) cancer is thought as a result of a complex interaction between germline and somatic mutations[4,5], whereas the evidence for the effect of somatic mutations in other diseases is limited to rare and neurological disorders[6,7], iii) the relationship between cancer and aging is complex, *e.g.* while telomere attrition and cellular senescence are thought to be evolved as a tumor suppressor mechanisms; aging-related changes in epigenomic landscape and genomic instability contribute to cancer occurrence[8]. Thus, although a similar analysis using cancers would be interesting, we only focused on non-cancer self-reported diseases in this study. Since we did not exclude the individuals with cancer, we also checked if there is a significant overlap in individuals with cancer with the other diseases we analyzed (Figure S45). However, there was no such association.

Disease co-occurrence calculations

Table 1: Contingency table for disease comorbidities.

	Disease B	No disease B	Total
Disease A	N_{ab}	N_{anb}	T_a
No disease A	N_{nab}	N_{nanb}	T_{na}

Relative risk (RR) score

Relative risk is an estimate of having the disease A, when already affected by disease B. Overall it measures if disease A co-occurs with disease B more frequently than expected if these diseases were independent in the population. It is calculated as a fraction between the number of patients diagnosed with both diseases and a random expectation based on disease prevalence[9]. Mathematically it can be expressed as follows, using the values from Table 1:

$$P_{exposed} = \frac{N_{ab}}{T_a}, \; P_{notexposed} = \frac{N_{nab}}{T_{na}}$$

$$RR = \frac{P_{exposed}}{P_{notexposed}}$$

$$CI = lnRR \pm 1.96 \sqrt{\frac{T_a - N_{ab}}{\frac{N_{ab}}{T_a}} + \frac{T_{na} - N_{nab}}{\frac{N_{nab}}{T_{na}}}}$$

ϕ value (Pearson correlation for binary variables)

The ϕ value measures the robustness of the association between diseases based on co-occurrences[10]. Mathematically, it can be expressed as:

$$\phi_{AB} = \frac{C_{AB}N - P_A P_B}{\sqrt{P_A P_B (N - P_A)(N - P_B)}}$$

N: the total number of individuals
P_A: Prevalence of disease A
C_{AB}: Number of patients with both diseases

ϕ ranges between -1 and 1, where the sign indicates the type of association.

Disease age-of-onset

Disease dissimilarity measure

Temporal correlation: In order to calculate dissimilarities among diseases, we use CORT[11] distance as included in R package TSclust[12]. Euclidean distance and dynamic time warping[13] are the two most widely used proximity measures for time series proximity. However, they are both calculated based on the closeness of the values and disregard the growth behavior. Correlation-based measures are also used to calculate the similarity between time series.

750 However, Pearson correlation overestimates the similarity because of the underlying temporal
751 dependency and Spearman correlation fails to consider the growth rate as it is based on ranks.
752 Chouakria et al., on the other hand, suggested a measure that also considers the proximity-
753 based on growth behavior, CORT[11]. Temporal correlation between two time series objects
754 $S_1=(u_1,u_2,...,u_p)$ and $S_2=(v_1,v_2,...,v_p)$ is calculated as follows:

755
$$CORT(S_1,S_2) = \frac{\sum_{i=1}^{p-1}(u_{(i+1)} - u_i)(v_{(i+1)} - v_i)}{\sqrt{\sum_{i=1}^{p-1}(u_{(i+1)} - u_i)^2}\sqrt{\sum_{i=1}^{p-1}(v_{(i+1)} - v_i)^2}}$$

756
757 CORT ranges between -1 and 1. A value of CORT = 1 implies that two time series increase
758 or decrease simultaneously with the same growth rate, whereas a value of -1 shows the same
759 growth rate but in opposite direction. If the value is 0, it means there is no temporal correlation
760 between the series.
761
762 _Dissimilarity Index:_ The dissimilarity index suggested by Chouakria et al.[11], is calculated based
763 on an automatic adaptive tuning function and considers similarity based on both values and
764 behavior, _i.e._ the strength of monotonicity and closeness of the growth rates as calculated by
765 CORT measure introduced in the previous section. They suggest a dissimilarity index D as
766 follows:
767
768
$$D(S_1,S_2) = f(CORT(S_1,S_2)).\delta_{conv}(S_1,S_2)$$
769
770 Where $f(x)$ is an exponential adaptive tuning function:
771
772
$$f(x) = \frac{2}{1 + exp(k\ x)}, k \geq 0$$
773
774 As k increases, the contribution of behavior increases. We use k = 2 and as a result behavior
775 (CORT) contributes 76.2% to D and values (δ_{conv}) contribute 23.8%. For δ_{conv} we used
776 conventional Euclidean distance.
777
Clustering diseases by age-of-onset
We clustered data using 'partition around medoids (PAM)' algorithm[14] based on the distance
measure calculated using the previous step. The aim of this algorithm is to minimize the
average distance (based on any dissimilarity measure) between the objects and their closest
selected medoid object. It works very similarly to k-means, except instead of defining arbitrary
points as the means, it defines medoids among the objects. Thus, it can incorporate any
distance measure instead of just using the mean distance between points (_i.e.,_ euclidean
distances). The algorithm first searches for k number of objects that represent the structure of
the data (Here the number k is assumed to be known a priori but see the next section for the
determination of k). After finding a set of k medoids, k clusters are constructed by assigning
each observation to the nearest medoid. Overall, the goal is to find k representative objects
such that the sum of dissimilarities of the observations to their closest representative is as
small as possible. After each assignment, medoid and non-medoid data points are swapped
and a cost (sum of distances of points to the new medoid) is calculated. If the total cost of
configuration is decreased, then the new configuration is maintained, otherwise, it is reversed.
We used 'pam' function in the 'cluster' package[15] in R to apply this algorithm.
794
Choosing the optimum number of clusters
The clustering algorithm we used, PAM, clusters data into k clusters, which is determined by
the user. So, even if there is no real structure in data, as we increase the number of clusters,
we can get more and more clusters. A potential way to decide on the number of clusters is

using the gap statistic[16]. This value is calculated by comparing the logarithm of the within-sum-
of-squares (WSS) to averages from simulated data without any structure.
801

$$WSS_k = \sum_{l=1}^{k} \sum_{x_i \in C_l} d^2(x_i, \overline{x_l})$$

803 k: number of clusters
804 C_l: objects in the l-th cluster
805 $\overline{x}$: the average point.
806
Calculating only WSS, however, is not enough as it would be minimized when each point has
its own cluster. Thus, we use the gap statistic which suggests calculating $log(WSS_k)$ for a
range of values of k and compare it to that obtained by WSS calculated based on simulated
data. So, after WSS is calculated for various values of k, the algorithm involves generating B
(we choose B=1,000) reference datasets, using Monte Carlo sampling from a homogeneous
distribution and re-calculate WSS for all k values. Using these values gap(k) statistic is
calculated as:
814

$$gap(k) = \overline{l_k} - log(WSS_k)$$

$$\overline{l_k} = \frac{1}{B} \sum_{b=1}^{B} log(W_{kb}^*)$$

818 If the clustering is good (i.e. WSS is small) we expect $\overline{l_k}$ to be higher than log(WSS). Thus,
819 gap statistic is mostly positive and we are interested in the highest value. Tibshirani et al.[16]
820 suggests using the smallest k such that,
821

$$gap(k) \geq gap(k+1) - s'_{k+1}$$

823
824 where

$$s'_{k+1} = sd_{k+1}\sqrt{1 + \frac{1}{B}} \text{ and } sd_k^2 = \frac{1}{B-1}\sum_{b=1}^{B}(log(W_{kb}^*) - \overline{l_k})^2$$

826
827 Using this approach, we determined k = 4.
828
829 **Genome wide association study**
830
831 *Preparing the files required for GWAS*
832
833 *Fixing FAM files:* In UK Biobank FAM files, the column for 'phenotype' includes batch that is
834 coded with characters. In order to use BOLT-LMM[17], we updated all the entries in this column
835 to numeric values[18].
836
837 *'Remove' files for BOLT-LMM:* BOLT-LMM accepts a list of individuals to be removed from the
838 analysis as an input. These files are called 'remove' files and are in the FAM format. We
839 prepared these files for i) withdrawn samples (n = 51), ii) samples that failed the quality control
840 (n = 3,779), iii) samples that have information in PLINK files but lack BGEN files (n = 968).
841

Calculating the SNP statistics: In order to apply a quality filter for SNPs, using PLINK[1], we calculated i) p-values for each SNP showing whether it deviates from Hardy-Weinberg equilibrium, and ii) Minor allele frequencies (MAF).

SNP Quality Control: We excluded SNPs that deviate from Hardy-Weinberg equilibrium (p ≤ 1e-6, n = 202,473) or with a minor allele frequency (MAF) smaller than 0.01 (n = 127,969). In total, we discarded 314,697 SNPs (Note that the numbers do not add up as these SNPs can overlap), resulting in 9,886,868 sites.

Phenotype File: We created a phenotype file that can be used as an input for BOLT-LMM, including the following fields: sex, age when attended assessment center, calculated BMI, assessment center, ethnicity, batch, first 20 PCs, and self-reported diseases (one column per disease).

GWAS run using BOLT-LMM
For each disease, we run GWAS separately using BOLT-LMM with the following inputs:
- We remove the samples that are in plink files but now in bgen; samples that did not pass our QC; samples from the individuals who have withdrawn their data from the UKBB
- We excluded the SNPs that deviate from Hardy-Weinberg equilibrium, and have minor allele frequency lower than 0.01.
- We used Sex, Age, BMI, assessment center, ethnicity, batch, and the first 20 PCs as covariates.
- To run the mixed-model, a reference LD score table is required. We used LD scores generated using 1000 Genomes European-ancestry samples, which is provided with the BOLT-LMM download.
- Genetic map for hg19 file provided in the BOLT-LMM website.
- We set 'bgenMinMAF' argument to 1e-2 and 'bgenMinINFO' parameter to 0.5 to only include SNPs that pass these criteria.

GWAS Results
We removed MHC region (chr6: 28,477,797 - 33,448,354) from the analysis and considered positions with a p-value lower than $5x10^{-8}$ as a significant association.

Coding Variants
We used VarMap[19] to map variants to proteins and domains. VarMap provides detailed information about coding variants, including annotations for the missense, synonymous, and nonsense variations. In our analysis, if a variant is not annotated as a coding variant in VarMap output, we assumed it is non-coding.

Genetic similarities between diseases
In order to calculate the overlap between diseases we used the number of SNPs that are significantly associated with both diseases, but corrected by the number that is expected by chance, if two diseases are independent:

$$Genetic\ Similarity\ =\ \frac{N_{common}}{N_{d1}\ x\ N_{d2}}\ x\ N_{total}$$

N_{common}: Number of SNPs in common.
N_{dx}: Number of SNPs associated with disease X.
N_{total}: Total number of SNPs analyzed in the study.

The statistical significance of these genetic similarities is calculated using the binomial test, and the similarity is only considered for downstream analysis if p<=0.01. Moreover, the value

895 is only calculated if two diseases do not have any hierarchical relationships in the disease
896 hierarchy.
897
898 In order to assess the genetic similarity within age-of-onset clusters, we further used linear
899 regression to correct log2 genetic similarity value by disease co-occurrences (risk ratios) and
900 disease categories (binary data showing whether two diseases are of the same category). The
901 'corrected genetic similarity' is the residuals from this linear model.
902
LD Blocks
In order to assess the similarity between different diseases we use overlaps across significant
associations and thus preferred not to do fine mapping. However, a significant challenge is
that genomic variations are not independent but instead linked in the genome. To understand
the effect of linkage disequilibrium or overcome it, we made use of linkage disequilibrium
blocks previously defined for human genome[20]. We repeated the analysis for genetic similarity
after collapsing all positions within an LD block and thus creating independent genomic loci (n
= 1,703). We use binary information for LD blocks, i.e. blocks with at least one significant
association are considered as a hit, and the rest are not.
912
High-definition likelihood (HDL) method to calculate genetic similarities
We repeated the genetic similarity analysis with HDL[21], which is a recent alternative to LDSC
to calculate genetic correlations. Using the code available in github
(https://github.com/zhenin/HDL/) we calculated the genetic similarities between diseases,
using the leading eigenvalues explaining 99 and their correspondent eigenvectors in LD score
matrix.
919
Comparison between SNPs associated with multiple diseases and economic status or
specific diet
We used the UK Biobank GWAS summary statistics provided by Neale Lab[22], and calculated
the overlap between SNPs associated with Townsend Deprivation Index (inverse-rank
normalized), or specific diet regimes (i.e. gluten-free, lactose-free, low calorie, vegan,
vegetarian, or other as specified in the UK Biobank) and the SNPs associated with multiple
diseases or multiple disease categories in each age-of-onset cluster.
927
Analysis of mediated pleiotropy between diseases
Using the LCV method developed by O'Connor & Price, we tested the causal relationships
between diseases[23]. We used the R functions developed by the authors and provided on
GitHub (github.com/lukejoconnor/LCV/). We calculated the genetic causality proportion (GCP)
between each disease pair, if the diseases have at least 10 significant variants and a
significant heritability estimate as suggested by the developers ($Z_h \geq 7$). We only calculated
GCP if the diseases are not vertically connected on the disease hierarchy. Following the
criteria applied by the developers, we considered pairs with FDR corrected $p \leq 0.01$ and mean
GCP>0.6 as significant.
937
938 **SNP to gene mapping**
939 We map all SNPs analyzed in GWAS to genes based on proximity and eQTL results.
940
Using proximity
Using VariantAnnotation[24], TxDb.Hsapiens.UCSC.hg19.knownGene[25], and
GenomicRanges[26] packages in R, we mapped the genomic coordinates for each SNP to
genes. Specifically, if a gene is within the coding region, intron, 5' or 3' UTR, or 1kb down- or
up-stream of the transcription start site, we annotated that SNP to the gene. As a result, we
had 4,443,872 SNP-gene associations for 4,236,176 SNPs and 22,228 Entrez gene IDs. We
used the Ensembl biomaRt[27] package in R to retrieve HGNC symbols (17,994), Ensembl Gene
IDs (20,507), and gene descriptions for the Entrez gene IDs obtained from
TxDb.Hsapiens.UCSC.hg19.knownGene database.

950
Using GTEx eQTL data
Using SNP-gene associations based on GTEx v8 eQTL data (accessed on 20.10.2020)[28], we
associated SNPs with the genes they could potentially regulate. We first mapped the hg38
variants in GTEx v8 to hg19 using 'rtracklayer', 'liftOver', and 'hg38ToHg19.over.chain'
packages in R. We generated a combined tissue list, which associates SNP to the gene if
there is at least one tissue in which there is a significant ($p <= 5e-8$) association. We chose to
create a combined tissue list due to following reasons: 1) disease-tissue matching is not
straightforward, especially for complex system-wide diseases[29] and 2) the power to detect
eQTLs depend heavily on the number of samples from each tissue. Thus relying on
incomparable set of eQTL sets from different tissues may bias our analysis.
961
962 As a result, there were 3,261,028 unique SNPs associated with 20,043 Ensembl Gene IDs.
963 We used the biomaRt[27] package in R to retrieve HGNC Symbols (16,068), Entrez IDs (14,574),
964 and gene descriptions.
965
Comparison of proximity and eQTL based mapping
Instead of only focusing on disease-associated SNPs, we first mapped all SNPs that we
analyzed to discover if there is a bias for certain genes (e.g. some genes could have many
more SNPs because they are longer, or because they are already associated with certain
traits and the chip is designed in that way). There were as much as 19,195 SNPs mapped to
one gene (CSMD1) by proximity, whereas there were 68 SNPs per gene on average (median).
The number of SNPs per gene was on average, higher for the mappings by eQTL (Figure
S46a). The maximum was 10720 SNPs for C4A gene and the median number of SNPs per
gene was 324. However, we did not consider MHC region in our downstream analysis and
thus this region is also excluded. The correlation between the number of SNPs per gene was
low (rho = 0.18, Figure S46b). Since the proximity-based mapping is by definition dependent
on the gene length, we also tested if there is a significant correlation between the number of
SNPs per gene and gene length. While the correlation is low for gene mappings by eQTL
(Spearman's correlation rho = 0.08, p = 2.2e-16), mappings by proximity show a high
correlation as expected (Spearman's correlation rho = 0.87, p < 2.2e-16). This also explains
the low correlation between eQTL and proximity-based mappings. We next checked the
correlation between the number of SNPs per gene mapped by proximity but only to promoter
region. The correlation between the number of SNPs and gene length decreased (rho = 0.21),
and the correlation with the number of SNPs by eQTL slightly increased but was still low (rho
= 0.14). Overall, we concluded that both eQTL data and proximity-based mapping could
capture different information and decided to use both for the downstream analyses.
987
GWAS Catalog analysis
We accessed the GWAS Catalog on 30-07-2019 and used v1.0.2 e96 dataset[30]. We excluded
all studies which used UK Biobank dataset (n = 190, data courtesy of GWAS Catalog team).
Using the associations with a p-value lower than $5x10^{-8}$, we compiled significant associations
between MAPPED_GENEs and MAPPED_TRAITs. We use GWAS catalog analysis to check
if our GWAS hits are supported by previous studies and applied a Fisher test between all traits
in GWAS catalog and the diseases in our study. P-values are corrected for multiple testing
using FDR correction.
996
Analysis of the association with aging
We downloaded GenAge human, GenAge model organism[31] and DrugAge[32] data on Aug 13,
2019 and CellAge[33] data on Oct 02, 2019 (CellAge data is kindly provided by Avelar et al.).
We used HGNC Symbols for GenAge and CellAge genes. In order to compile genes that are
targeted by the drugs in DrugAge database, using the drug names in DrugAge data, we first
compiled PubChem IDs using PubChem REST API[34]. Using UniChem[35], we mapped
PubChem IDs to ChEMBL IDs[36]. Next, using DGIdb[37], we compiled the genes targeted by
these ChEMBL IDs. As a result, we had 307 genes from GenAge human database, 902 genes

from GenAge model organism database, 279 genes from CellAge database, and 714 genes
targeted by DrugAge drugs. We next calculated the overlaps between these databases and
the genes associated with multiple diseases or multiple categories in different age-of-onset
clusters. To calculate the expected values and statistical significance, we used 10,000
permutations calculating the overlap for the same number of random genes among genes that
can be detected by GWAS. Then, an odds ratio is calculated by dividing the observed value
to the mean of expected values.
1012
Functional Enrichment Test
1013
Using the goseq package in R[38], which takes the gene length bias into account, we performed
a functional analysis of the genes associated with different age-of-onset clusters. Using GO
categories with more than 10 and less than 500 annotated genes, we applied an enrichment
test for the Gene Ontology (GO)[39,40] Biological Process (BP), Molecular Function (MF), and
Cellular Compartment (CC) categories. BY correction[41] is applied to the p-values for all tests
for all clusters and 3 GO Categories (BP, MF, and CC) combined. We considered associations
with a BY-corrected p-value lower than 0.05 as significant. For the ease of visualization and
comprehension we selected representative categories for significant associations as follows:
For each cluster and GO Ontology (*i.e.* BP, MF, CC) separately; i) Jaccard similarity index (*i.e.*
number of genes in common divided by the number of unique genes in each category
combined) is calculated between all significantly associated GO Categories; ii) Jaccard indices
are hierarchically clustered and cut to k number of groups, where k is the minimum number of
clusters which ensures median Jaccard similarity within a cluster is above 0.5; iii) The category
with the highest average similarity to other categories in the same cluster is assigned as the
representative.
1029
Gene Expression Analysis
1030
Using GTEx v8 data[42], we analyzed age-related changes in gene expression for the tissues
with at least 10 samples (n = 48). We filtered out the genes with median TPM lower than 1.
We next log2 transformed expression matrix and used linear model to correct for the effect of
sex and death. The normalized expression matrix is quantile normalized using
normalize.quantiles function from preprocessCore package[43] in R. We excluded any sample
that deviates 3 standard deviation from the first 4 principal components calculated for this
quantile normalized matrix using 'prcomp' function with scale = T argument and repeated the
procedure until there is no outlier in the data (32 tissues did not have any outliers, 14 had all
outliers removed in one round, 1 tissue required 2 rounds and 1 tissue - Cells-Cultured
fibroblasts – required 5 rounds of outlier removal). The whole analysis pipeline and quality
control steps and results are available in https://github.com/mdonertas/aging_in_GTEx_v8. In
order to match age-related expression changes with the variants, we used tissue-specific
eQTL data. More specifically, we identified the gene-tissue pairs that have significant eQTL
associations and matched their gene expression levels. We only considered genes with
eQTLs in the same direction. If disease associated variants are associated with different
direction of change for a gene, we excluded these cases from the analysis given in Figures
S31-32.
Methylation Analysis
1048
1049 Using the reported gene sets as differentially methylated during ageing from two studies[44,45],
1050 we checked the overlap with our gene sets. We repeated the same analysis applied to
1051 calculate the overlap with aging-related genes.
1052
Drug Repurposing
1053
We searched for the drugs that specifically target multicategory genes in cluster 1, cluster 2,
or cluster 1 and 2. Using the Fisher's exact test, we compiled the drugs in DGIdb[37] that
specifically target these genes (p≤0.01) and drugs that target only one gene in one of these
clusters. Importantly, we excluded all non-specific drugs (*i.e.* targeting more than 10 genes)
from the analyses. The interaction data is compiled from DGIdb, and the names, indications
and phases of the drugs are obtained from ChEMBL REST API[36].

Evolutionary Analysis

In order to test the mutation accumulation and antagonistic pleiotropy theories of aging we used the risk allele frequencies in UK Biobank and 1000 Genomes super-populations[46]. A risk allele is an allele that shows positive association with a disease. Since the SNPs are not independent and have similar allele frequencies in a given LD block, we analyzed LD blocks instead of individual SNPs and used the median risk allele frequency for a given LD block. We used only the biallelic SNPs for this analysis. Allele frequencies for UK Biobank are calculated using BOLT-LMM and the allele frequencies for 1000 Genome super-populations are obtained from the vcf file provided on the 1000 Genomes project website. To test the antagonistic pleiotropy excess, we calculated the proportion of antagonistic vs. agonist SNPs within the same vs. different age-of-onset clusters using Fisher's exact test. We considered pleiotropic SNPs as agonist if the risk allele for two or more diseases are the same, and antagonist if the risk alleles are opposite. We only tested the risk allele frequency differences between cluster 1 and cluster 2. Also, we excluded any SNPs that are antagonistic within an age-of-onset cluster and agonist between clusters.

References

1. Anderson, C. A. *et al.* Data quality control in genetic case-control association studies. *Nat. Protoc.* **5**, 1564–1573 (2010).

2. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203–209 (2018).

3. Zhou, W. *et al.* Efficiently controlling for case-control imbalance and sample relatedness in large-scale genetic association studies. *Nat. Genet.* **50**, 1335–1341 (2018).

4. Kanchi, K. L. *et al.* Integrated analysis of germline and somatic variants in ovarian cancer. *Nat. Commun.* **5**, 3156 (2014).

5. Khurana, E. *et al.* Role of non-coding sequence variants in cancer. *Nat. Rev. Genet.* **17**, 93–108 (2016).

6. Poduri, A., Evrony, G. D., Cai, X. & Walsh, C. A. Somatic mutation, genomic variation, and neurological disease. *Science* **341**, 1237758 (2013).

7. Zhang, L. & Vijg, J. Somatic Mutagenesis in Mammals and Its Implications for Human Disease and Aging. *Annu. Rev. Genet.* **52**, 397–419 (2018).

8. Finkel, T., Serrano, M. & Blasco, M. A. The common biology of cancer and ageing. *Nature* **448**, 767–774 (2007).

9. Sanchez-Valle, J. *et al.* Unveiling the molecular basis of disease co-occurrence: towards personalized comorbidity profiles. *bioRxiv* 431312 (2018) doi:10.1101/431312.

10. Gutiérrez-Sacristán, A. *et al.* comoRbidity: an R package for the systematic analysis of disease comorbidities. *Bioinformatics* **34**, 3228–3230 (2018).

11. Chouakria, A. D. & Nagabhushan, P. N. Adaptive dissimilarity index for measuring time series proximity. *Adv. Data Anal. Classif.* **1**, 5–21 (2007).

12. Montero, P. & Vilar, J. TSclust: An R Package for Time Series Clustering. *Journal of Statistical Software, Articles* **62**, 1–43 (2014).

13. Berndt, D. J. & Clifford, J. Using Dynamic Time Warping to Find Patterns in Time Series. in *Proceedings of the 3rd International Conference on Knowledge Discovery and Data Mining* 359–370 (AAAI Press, 1994).

14. Partitioning Around Medoids (Program PAM). in *Finding Groups in Data* (eds. Kaufman,
L. & Rousseeuw, P. J.) 68–125 (John Wiley & Sons, Inc., 1990).

15. Maechler, M., Rousseeuw, P., Struyf, A., Hubert, M. & Hornik, K. cluster: Cluster Analysis
Basics and Extensions. (2019).

16. Tibshirani, R., Walther, G. & Hastie, T. Estimating the number of clusters in a data set via
the gap statistic. *Journal of the Royal Statistical Society: Series B (Statistical*
Methodology) vol. 63 411–423 (2001).

17. Loh, P.-R. *et al.* Efficient Bayesian mixed-model analysis increases association power in
large cohorts. *Nat. Genet.* **47**, 284–290 (2015).

18. Loh,   P.-R.   *BOLT-LMM*   *v2.*   *3.1*   *User*   *Manual*.
https://data.broadinstitute.org/alkesgroup/BOLT-LMM/ (2017).

19. Stephenson, J. D., Laskowski, R. A., Nightingale, A., Hurles, M. E. & Thornton, J. M.
VarMap: a web tool for mapping genomic coordinates to protein sequence and structure
and retrieving protein structural annotations. *Bioinformatics* (2019)
doi:10.1093/bioinformatics/btz482.

20. Berisa, T. & Pickrell, J. K. Approximately independent linkage disequilibrium blocks in
human populations. *Bioinformatics* **32**, 283–285 (2016).

21. Ning, Z., Pawitan, Y. & Shen, X. High-definition likelihood inference of genetic correlations
across human complex traits. *Nat. Genet.* (2020) doi:10.1038/s41588-020-0653-y.

22. UK Biobank — Neale lab. http://www.nealelab.is/uk-biobank/.

23. O'Connor, L. J. & Price, A. L. Distinguishing genetic correlation from causation across 52
diseases and complex traits. *Nat. Genet.* **50**, 1728–1734 (2018).

24. Obenchain, V. *et al.* VariantAnnotation: a Bioconductor package for exploration and
annotation of genetic variants. *Bioinformatics* **30**, 2076–2078 (2014).

25. Carlson, M. & Maintainer, B. P. TxDb.Hsapiens.UCSC.hg19.knownGene: Annotation
package for TxDb object(s). (2015).

26. Lawrence, M. *et al.* Software for computing and annotating genomic ranges. *PLoS*
Comput. Biol. **9**, e1003118 (2013).

27. Durinck, S., Spellman, P. T., Birney, E. & Huber, W. Mapping identifiers for the integration

of genomic datasets with the R/Bioconductor package biomaRt. *Nat. Protoc.* **4**, 1184–

1191 (2009).

28. Gamazon, E. R. *et al.* Using an atlas of gene regulation across 44 human tissues to inform

complex disease- and trait-associated variation. *Nat. Genet.* **50**, 956–967 (2018).

29. Lage, K. *et al.* A large-scale analysis of tissue-specific pathology and gene expression of

human disease genes and complexes. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 20870–20875

(2008).

30. Buniello, A. *et al.* The NHGRI-EBI GWAS Catalog of published genome-wide association

studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res.* **47**, D1005–

D1012 (2019).

31. Tacutu, R. *et al.* Human Ageing Genomic Resources: new and updated databases.

Nucleic Acids Res. **46**, D1083–D1090 (2018).

32. Barardo, D. *et al.* The DrugAge database of aging-related drugs. *Aging Cell* **16**, 594–597

(2017).

33. Avelar, R. A. *et al.* A multidimensional systems biology analysis of cellular senescence in

aging and disease. *Genome Biol.* **21**, 91 (2020).

34. Kim, S. *et al.* PubChem 2019 update: improved access to chemical data. *Nucleic Acids*

Res. **47**, D1102–D1109 (2019).

35. Chambers, J. *et al.* UniChem: a unified chemical structure cross-referencing and identifier

tracking system. *J. Cheminform.* **5**, 3 (2013).

36. Gaulton, A. *et al.* The ChEMBL database in 2017. *Nucleic Acids Res.* **45**, D945–D954

(2017).

37. Cotto, K. C. *et al.* DGIdb 3.0: a redesign and expansion of the drug–gene interaction

database. *Nucleic Acids Res.* **46**, D1068–D1073 (2018).

38. Young, M. D., Wakefield, M. J., Smyth, G. K. & Oshlack, A. Gene ontology analysis for

RNA-seq: accounting for selection bias. *Genome Biol.* **11**, R14 (2010).

39. The Gene Ontology Consortium & The Gene Ontology Consortium. The Gene Ontology

Resource: 20 years and still GOing strong. *Nucleic Acids Research* vol. 47 D330–D338 (2019).

40. Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat. Genet.* **25**, 25–29 (2000).

41. Benjamini, Y. & Yekutieli, D. The control of the false discovery rate in multiple testing under dependency. *Ann. Stat.* **29**, 1165–1188 (2001).

42. GTEx Consortium. Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science* **348**, 648–660 (2015).

43. Bolstad, B. preprocessCore: A collection of pre-processing functions. (2020).

44. Adelman, E. R. *et al.* Aging Human Hematopoietic Stem Cells Manifest Profound Epigenetic Reprogramming of Enhancers That May Predispose to Leukemia. *Cancer Discov.* **9**, 1080–1101 (2019).

45. Marttila, S. *et al.* Ageing-associated changes in the human DNA methylome: genomic locations and effects on gene expression. *BMC Genomics* **16**, 179 (2015).

46. 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74 (2015).