Standardized Testing Demonstrates Altered Odor Detection Sensitivity and Hedonics in Asymptomatic College Students as SARS-CoV-2 Emerged Locally

Julie Walsh-Messinger, Ph.D.[1,2*] Sahar Kaouk, B.S.[1] Hannah Manis, B.S.[1]
Rachel Kaye, M.D.[3] Guillermo Cecchi, Ph.D.[5] Pablo Meyer, Ph.D.[5]
Dolores Malaspina, M.D., M.S.P.H[6]

[1]Department of Psychology, University of Dayton, Dayton, OH
[2]Department of Psychiatry, Wright State University Boonshoft School of Medicine, Dayton, OH
[3]Department of Otolaryngology-- Head & Neck Surgery, Rutgers New Jersey Medical Center, Newark, NJ
[4]IBM Thomas J. Watson Research Center, Yorktown Heights, NY
[5]Departments of Psychiatry, Neuroscience, and Genetics and Genomics, Icahn School of Medicine at Mount Sinai, New York, NY

*Corresponding Author:
Julie Walsh-Messinger, PhD
Department of Psychology
University of Dayton
300 College Park
Dayton, OH 45469-1430
Phone: (937) 229-1380
Fax: (937) 229-3900
E-mail: jmessinger1@udayton.edu

Keywords: Olfaction; SARS-CoV-2; COVID-19

ABSTRACT

Anosmia is a recognized symptom of COVID-19, but the relationship of SARS-CoV-2 exposure with olfactory dysfunction remains enigmatic. This report adds unique data from healthy students tested as the virus unknowingly emerged locally. Standardized and validated measures of odor detection, identification and hedonics were employed to test healthy university students at a mid-sized, private, Midwestern University. Data from asymptomatic students (N=22), tested as SARS-CoV-2 unknowingly emerged locally, were compared to students tested just prior to local virus transmission (N=25), and our normative sample (N=272) tested over the previous four years. The exposed cohort demonstrated significantly reduced odor detection sensitivity compared to the students in the prior group (P=.01; d=0.77), with a distribution skewed towards less detection sensitivity (P=.03). Categorically, the exposed group was significantly more likely to have hyposmia (OR=7.7), particularly the subgroup assessed in the final week before campus closure (OR=13.6). The exposed group also rated odors as less unpleasant (P<.001, CLES=0.77,) which also showed a skewed distribution (P=.005). The groups had similar odor identification performance. Psychometric measures of odor detection sensitivity and hedonics may detect early SARS-CoV-2 exposure in asymptomatic and pre-symptomatic persons with normal odor identification. Viral detection by nasal associated lymphoid tissue is known to trigger systemic immune effects, but its activation may also reduce smell sensitivity and shift perception of the environment towards unpleasant, increasing the social isolation that may mitigate viral infection or transmission. Regular testing of odor detection and hedonics may have value for identifying regional viral exposure.

INTRODUCTION

Coronaviruses (CoVs) are a group of enveloped single stranded RNA viruses associated with the common cold in adult humans and upper respiratory viruses in other mammals and birds (Paules et al., 2020). The viruses were considered to produce mild symptoms until the highly lethal SARS-Cov-1 presented in 2003, followed by Middle East respiratory syndrome (MERS-CoV) in 2012. The novel SARS-CoV-2 virus emerged in Wuhan in 2019 and rapidly became a worldwide pandemic (World Health Organization, 2020), challenging health services and economic stability worldwide.

Self-reports of diminished smell (hyposmia/anosmia) and taste (ageusia/dysguesia) led to the inclusion of chemosensory impairment as recognized COVID-19 symptoms by the World Health Organization (WHO) and Center for Disease Control (CDC), fueling efforts to identify the pathobiology, particularly for the olfactory symptoms (Parma et al., 2020). While the proportion of cases with olfactory complaints varies by ascertainment and assessment approach and with sample age and sex (Speth et al., 2020), there is a consensus that olfactory symptoms can present early in the course of the infection, sometimes as the only symptom (Jang et al., 2020), and they appear to be a better predictor of COVID-19 positivity than other symptoms such as fever and cough (Gerkin et al., 2020). Early olfactory impairment may be associated with better outcomes (Lechien et al., 2020), although a recent report suggests that there may be a subgropwith COVID-19 smell loss that experiences longer duration of symptoms (Gerkin et al., 2020).

SARS-Cov-2 may be spread through respired droplets, even by asymptomatic persons (Luo et al., 2020); thus, indicators of latent infection are of crucial interest. Entry of SARS-CoV-2 into cells involves spike (S) protein binding to angiotensin-converting enzyme 2 (ACE2) along with host proteases, principally transmembrane serine protease 2 (TMPRSS2). Olfactory receptor neurons do not express these receptors (Hoffman et al., 2016), so the earliest olfactory changes likely reflect non-neuronal olfactory epithelial cell entry, including cells that initiate ultra-rapid immune responses (Butowt & Bilinska, 2020). The nasal epithelium contains innate immune-associated tissue, the nasal-associated lymphoid tissue. This phylogenetically ancient system is the first line of immune defense against inspired or ingested viruses and other pathogens, organizing systemic inflammatory processes (Sepahi & Salinas, 2016) that may be protective against more severe pulmonary involvement. As diminished odor sensitivity in rhinosinusitis is correlated with cytokine responses (Chidambaram & Goh, 2017; Schlosser et al., 2016), diminished sense of smell may not entail overt pathology to olfactory receptor neurons.

Discerning the mechanism for the earliest loss of olfactory sensitivity may be a key to early detection of SARS-CoV-2 exposure and infection, although prospective studies on the earliest olfactory impairment is limited. Most studies have relied on self-reported symptoms, which are significantly less sensitive to dysfunction than standard olfactory measures. To date, no studies have comprehensively assessed odor detection, identification, and hedonic judgments in asymptomatic and pre-symptomatic persons exposed to COVID-19. Odor detection and identification involve different olfactory pathways, but to our knowledge, they have not yet been distinguished in the COVID-19 literature. Additionally, no studies have

compared olfactory function in exposed persons to a large normative sample of age-appropriate and comorbidity-matched participants, nor had the availability of comparison measures from a cohort tested prior to the emergence of the pandemic.

We have applied our unique data, drawn from four years of systematic olfactory function assessments in healthy undergraduate students, to address these significant gaps in the literature. Data from an asymptomatic cohort of students exposed to SARS-CoV-2 virus as it became endemic were compared to data from a cohort tested in the weeks prior to the arrival of the virus and to ecologically valid comparison groups.

METHOD

Participants

Olfaction Tested Cohorts

Undergraduate students (N=367) participated in one of four research studies of odor detection sensitivity and identification conducted at a private Midwestern university from 2016–2020. Participants were asked to not wear perfume, cologne, or strongly scented lotions, smoke or vape at least four hours prior to their appointment or consume food or beverages 15 minutes prior to testing. The studies were all approved by the University of Dayton Research Review and Ethics Committee (RREC), which operates under the Institutional Review Board (IRB). All participants provided voluntary written informed consent, which was obtained in accordance to the Declaration of Helsinki, and they earned research experience class credit for their participation.

We retrospectively separated participants into cohorts (see Figure 1a) based on campus exposures to SARS-CoV-2, exclusive of 64 students who reported or exhibited nasal congestion, common cold, or other illness symptoms (N=50) or smoked cigarettes or vaped (N=14) within four hours of testing. Students (Cohort IV), tested February 14–26, 2020 (N=9), when COVID-19 exposure is uncertain, were excluded from all analyses except where noted. Cohort I (N=25) was comprised of those tested before local transmission of the virus, from January to early February 2020. Cohort II (N=22) included students tested from late February to early March 2020, as the virus unfolded in the area. A subset of this group (Cohort IIa) was tested immediately before social distancing was enforced on campus. Publicly available health records confirm that SARS-CoV-2 was endemic to the area as early as January 13, 2020 and became more widespread by the end of February (Ohio Department of Health, 2020a).

To assess the potential impact of seasonal factors, comparisons were made to the students tested late February and early March 2016–2019 (Cohort III; N=52). Normative performance was determined from an ecologically valid comparison group of all students (N=272) tested in this laboratory prior to February 7, 2020.

Health Survey

Separate RREC approval was obtained to conduct an online health survey following reports that anosmia/hyposmia was potentially linked to COVID-19. Of the 130 students who signed-up for the survey to earn research experience credit for class, 113 (87%) provided written informed consent and subsequently completed the survey administered via Qualtrics (Figure 1b). A subset of students from the aforementioned cohorts completed this survey

(N=9), including four students from Cohort II, all of whom provided consent to link their survey

and olfaction data.

Measures

Olfaction

All measures were administered by systematically trained undergraduate and graduate

research assistants. The *Sniffin' Sticks n-Butanol Threshold Test* (Hummel et al., 2007) assessed

odor detection sensitivity, using a single-staircase forced-choice design. The test utilizes pen-

like odor dispensing devices, 32 of which are blanks (no odor) and 16 of which contain varying

concentrations of n-butanol (alcohol). The test employs a single-staircase forced-choice design

in which the test administrator presents odor triplets directly below the blindfolded

participant's nostril (two blanks, one with n-butanol) one at a time in random order for five

seconds each. After each triplet presentation, the participant was asked to identify the pen that

contained the odor. Scores range from 0 (low) to 16 (high). Participants were classified as

anosmic, hyposmic, or normosmic based on published data (Hummel et al., 2007).

The *Sniffin' Sticks Identification Test* (Hummel et al., 2007) is comprised of 16 pen-like

devices that deliver common odorants (e.g. orange, garlic, fish, etc.). The test administrator

presented each odor pen directly under the participant's nostril, one at a time for 3-4 seconds.

After each presentation, the test administrator read four response choices and the participant

choose the one that best described the odor. Scores range from 0 (low) to 16 (high).

Participants then rated each the pleasantness and unpleasantness of each odor on two five-

point Likert scales ranging from "0" (not pleasant/unpleasant) to "4" (extremely

pleasant/unpleasant). Responses across all 16 odors were summed to yield total pleasantness

and unpleasantness scores, with scores for each scale ranging from "0" (not

pleasant/unpleasant) to 64 (extremely pleasant/unpleasant).

Health Survey

Participants were surveyed between April 14[th] and May 10[th], 2020 as to whether they

had experienced fever, sore throat, lymphadenopathy, cough, dyspnea, nasal congestion,

myalgias, fatigue, diarrhea, and/or emesis (excluding alcohol related incidents) since January 1,

2020. For each symptom endorsed, follow-up queries assessed severity, timing, and frequency.

Additional questions asked chronic health conditions and whether they suspected or formally

diagnosed with infectious diseases (influenza, COVID-19, strep pharyngitis, mononucleosis,

and/or common cold) during this period, followed-up with questions regarding test results (if

any), timing, concurrent symptoms, and

A modified version of the *Self-Report Inventory to Measure Olfactory Dysfunction and*

Quality of Life (Pusswald et al., 2012) assessed changes in the sense of smell during a one-

month period and the impact of any change on daily functions, modified to also assess changes

in taste, including sweet, salty, bitter, and sour. Participants were asked whether they

experienced any changes in their sense of smell or taste during January, February, March, or

April 2020; if yes, they were asked to complete the measure for that month.

Data analysis

Data analyses were conducted with SPSS (version 23.0). Participants with missing data

were excluded from analyses, as detailed in the legends of Tables 1 and 2. Independent sample

t-tests and chi-squares were used to test for cohort differences in demographic characteristics. As the data was normally distributed, independent sample t-tests were employed to compare the olfaction performance of Cohorts I and II. Due to unequal sample sizes Mann-Whitney U tests were used to compare the olfaction performance of Cohort IIa to Cohorts I and III and to Compare Cohort II to Cohort III.

To ensure that potential group differences were not the result of anomalous Cohort I data, a two-sample Kolmogorov-Smirnov (K-S) test was used to compare the frequency distributions of Cohorts II and IIa to the normative sample distribution. Distribution fits were applied as follows: Normative distribution gaussian fit mean=8.26113 [8.01615, 8.50612], sigma=1.95481 [1.79628, 2.14426]; Cohort II Gamma distribution fit mean=4.91205, a=5.99561 [3.37288, 10.6578], b = 1.22059 [0.669847, 2.22415]. One-sample t-tests were used to compare the exposed cohorts to published norms. Cohen's d, Common Language Effect Size (CLES; (McGraw & Wong, 1992) and 95% confidence intervals [CI] were computed using the Psycometrica online calculator (Lenhard & Lenhard, 2016). Odds ratios were employed to examine risk of hyposmia over time in the weeks leading up to the closure of campus.

RESULTS

Odor Detection

Means and standard deviations for the olfaction measures are presented in Table 2. Significantly reduced odor detection sensitivity was demonstrated by Cohort II, the exposed cohort, compared to Cohort I, which was assessed prior to SARS-CoV-2's emergence in the area (t=2.60; P=.01; d=0.77; CI, 0.17, 1.36). Cohort IIa, the subsample tested in the final week before

social isolation, was also significantly impaired (U=58.50; P=.007; CLES=0.77; CI, 0.31, 1.81).

Comparisons of odor sensitivity thresholds to those of students during the same time period in the previous four years (Cohort III) confirmed the absence of a seasonal explanation, with Cohort IIa showing less odor detection sensitivity (U=162.50; P=.03; CLES=0.75; CI, 0.30, 1.64), and a similar trend for the entire Cohort II (U=411.50; P=.06; CLES=0.63; CI=-.06, 0.95).

To compare Cohorts II and IIa to our normative sample distribution fits were applied as follows: Normative distribution gaussian fit mean=8.26113 [8.01615, 8.50612], sigma=1.95481 [1.79628, 2.14426]; Cohort II Gamma distribution fit mean=4.91205, a=5.99561 [3.37288, 10.6578], b = 1.22059 [0.669847, 2.22415]. The results showed significant rightward (positive) skewing, indicating reduced odor detection sensitivity for Cohorts II (D=0.38; P=.005; Figure 2) and IIa (D=0.52, P=.007). Similarly, Cohorts II (t=-2.92., P=.008) and IIa (t=-3.39 P=.007) had diminished odor detection in comparison to published normative data (Hummel et al., 2007).

Smell Diagnosis

Categorical diagnoses were examined to define the proportion of students with hyposmia in comparison to our normative sample. From mid-February to March, which included Cohort II and Cohort-IV, the latter assessed between February 14–26, hyposmia was 4-fold more likely than in our normative sample (OR=4.13; CI, 1.8, 9.50). This increased by 7.7-fold for Cohort II (OR=7.74; CI, 3.1, 19.40) and by 13.6-fold for Cohort IIa (OR=13.61; CI, 3.40, 35.66; Figure 3).

Odor Identification and Hedonics

There were no significant differences in odor identification (all P's >.13) or odor pleasantness ratings (all P's >.92). Cohort II rated odors as significantly less unpleasant than Cohort III (U=257.0, P<.001, CLES=0.77, CI, 0.51, 1.56). The distribution of unpleasantness ratings for Cohort II was also positively skewed compared to the normative sample (D=0.429, P=.001). Odor detection threshold and unpleasantness ratings were not correlated across the entire sample or within any cohort (all P's >.16).

Health Survey

Approximately 15% (N=16) of participants who completed the health survey reported being tested for influenza by clinicians between December 31, 2019 and March 9, 2020. Four had a positive test for influenza (25%), none of whom reported chemosensory changes. Six participants (38%) with negative influenza tests did report chemosensory changes at the time of their flu-like symptoms, consistent with SARS-CoV-2. The last positive influenza test was resulted on February 22, 2020. The only participant formally diagnosed with COVID-19 by a physician reported having anosmia and ageusia two weeks prior other recognized COVID-19 symptoms. Six additional participants suspected they were infected with SARS-CoV-2 but did not obtain testing and four students reported sudden onset anosmia in the absence of other COVID-19 symptoms.

Four Cohort II participants completed the survey, two of whom showed the greatest impairments in odor detection sensitivity (scores of 4 and 4.25). Neither of them reported any subjective change in olfaction, but one experienced nasal congestion and a sore throat a week

after olfactory testing, followed by a cough a week later. Another participant (odor detection sensitivity=7) reported the onset of persistent muscle aches almost a month after olfaction testing.

DISCUSSION

This is the first report of significantly reduced odor detection sensitivity and altered hedonic ratings of odors in asymptomatic students assessed prior to knowledge that the SARS-CoV-2 virus was spreading in the local area. The study employed a comprehensive, psychometrically validated olfaction testing battery administered prior to and during the unique window in which the virus emerged. These findings suggest that olfactory impairments are the earliest manifestation of SARS-CoV-2 exposure and infection and may be useful for early identification of affected persons to limit contagion.

These findings were detected in asymptomatic healthy students. The follow-up health questionnaire confirmed that some participants who subsequently developed COVID-19 symptoms had shown the most severe sensitivity decrements yet had no subjective awareness of olfactory impairment. This finding highlights the superiority of psychometric olfaction tests over self-reports, particularly in asymptomatic persons. Also noteworthy is the intact odor identification of the exposed cohort, showing spared peripheral olfactory processing despite diminished detection sensitivity.

Most studies of COVID-19 patients that have employed objective tests to assess smell identification (but not odor detection) have found significant decrements (Hornuss et al., 2020; Lechien et al., 2020; Moein et al., 2020). The only other study to date that has assessed odor

detection sensitivity in the context of COVID-19 showed reduced odor detection sensitivity with intact odor discrimination (Vaira et al., 2020), consistent with our findings. Odor identification requires accurate transmission of olfactory stimuli to higher processing regions, indicating intact olfactory receptor neuron function and the fidelity of transmission through the olfactory bulb. Later viral infection and inflammatory pathology may damage the peripheral olfactory pathways, impacting these higher order processes, but our findings suggest that the earliest blunting of smell may have different underpinnings. Further supporting this hypothesis, the results of a recent large-scale multinational study show two distinct subgroups of COVID-19 patients with olfactory impairment, a group that recovers in under 40 days, and a second group that displays more variable and prolonged recovery (Gerkin et al., 2020).

While olfactory receptor neurons do not express the ACE2 receptor and TMPRSS2 that are requisite for viral entry into cells, these are expressed in the epithelial support cells (Brann et al., 2020). In particular, ACE2 protein expression is found in the basal layer of the squamous epithelium of the nasal mucosa and nasopharynx (Hamming et al., 2004). Sensory cells in the nasal mucosa detect viral pathogens and very rapidly organize countering immune responses (Sepahi et al., 2019) through the activation of nasopharynx-associated lymphoid tissue (NALT). This ancient, innate immune system is the first line of immune defense against airborne pathogens and can activate B and T cell receptor expression.

Humans can also detect sickness in other individuals through an olfactory route, even when they are not ill themselves, activating a specific multisensory threat circuitry that involves the primary and secondary olfactory areas, including entorhinal/piriform cortex, orbitofrontal cortex, inferior frontopolar gyrus, middle frontal gyrus, and thalamus

(Regenbogen et al., 2017). Activation of this pathway would involve the amygdala, which is consistent with altered perceptions of unpleasantness in the hedonic odor assessments. A resultant reduction of smell and taste sensitivity may be a protective response to pathogen exposure rather than pathology. Along with a shift in the perception of the environment as more unpleasant, behavioral alterations could act to limit social interactions and feeding, preventing exposure to a more pathogenic load of viral particles or transmitting the virus to conspecifics (Dantzer, 2001). Social withdrawal is a known component of sickness behavior directed through the central nervous system, but protection against environmental threats through decreasing odor sensitivity is unexplored.

Our findings are consistent with the involvement of some pathways previously associated with immune activation and sickness behavior. It is notable that females and younger persons, who have enhanced chemodetection and immune responses compared to their male and older counterparts, also have better COVID-19 outcomes (Scully et al., 2020; Sorokowski et al., 2019). Better detection of airborne pathogens may preempt more severe illness though personal immune activation and social withdrawal to prevent viral spread through the social group. Although correlation does not imply causation, a subset of COVID-19 positive individuals with olfactory or gustatory dysfunction appear to have a milder course of illness (Gerkin et al., 2020; Wang et al., 2020). Potentially, viral detection in the environment may preclude infection through behavioral and inflammatory mechanisms.

However, without restoration of homeostasis this system could drive inflammation, social dysfunction, and contribute to psychiatric conditions. Our laboratory has shown reduced odor detection sensitivity in association with social withdrawal in males and females with

schizophrenia, bipolar disorder, and in a non-clinical sample of persons with olfactory complaints (Cieslak et al., 2015; Hardy et al., 2012; Keller & Malaspina, 2013; Malaspina et al., 2012). A coordinated immune and behavioral response is not unexpected as infectious diseases account for the greatest amount of human mortality over evolution (Inhorn & Brown, 1990), highlighted again by the COVID-19 pandemic. Viral threats are estimated to underlie a third of mammalian genetic adaptations (Enard et al., 2016).

The uniqueness of this report rests in the natural experiment wherein SARS-CoV-2 arrived on a campus already having ongoing studies of olfactory function. This research will never be replicated; yet it is not without limitations. The sample sizes of the 2020 cohorts are small, although we still found moderate to large effects and statistical significance. Although our subjects were healthy and without nasal symptoms, we did not explicitly assess COVID-19 symptoms at the time of olfaction testing as the studies were not designed to test hypotheses related to infectious diseases. The absence of SARS-CoV-2 positivity confirmation also limits our ability to definitively draw the conclusion about exposure to SARS-CoV-2 at the time Cohort II participated. The retrospective nature of our health survey and lack of local and national access to COVID-19 testing for young adults also limits our health survey results. The State of Ohio, where at almost half of the students reside, only allowed access to COVID-19 testing for non-high-risk persons <60 and non-healthcare workers on June 11, 2020 (Ohio Department of Health, 2020b).

This report adds a scientifically rigorous perspective to the early clinician and patient reports associating anosmia/hyposmia and ageusia/dysgeusia with COVID-19 (Bienkov, 2020; Gale, 2020; Hopkins & Kumar, 2020; Rabin, 2020; Stone, 2020) which propelled the American

Academy of Otolaryngology to advocate for the CDC and WHO to add these symptoms to COVID-19 screening and urge precautionary isolation for individuals experiencing chemosensory impairment. Crucially, we found that odor detection can be significantly impaired without nasal obstruction or subjective awareness, as participants did not report impaired olfaction at the time of their assessment. While it may not be realistic to administer Sniffin' Sticks on a population level, daily tracking of odor detection through self-administered protocols could aid in identifying COVID-19 infection in asymptomatic persons and also contribute to our understanding of how olfactory changes may be associated with infection severity.

Role of the Funding Source

This work was supported by the National Institute of Mental Health [grant number R01MH110418 to DM] and the University of Dayton Research Council [grant numbers IGRU16 and IGRV16 to JWM];

Conflict of Interest

The authors report no financial conflicts of interest.

Acknowledgements: Thank you to the following students for their assistance with data collection, entry, and verification: Brooke Lipnos, Madison Degnan, Rhiannon Gibbs, Wyatt Kaiser, Jenna Kilian, Michael A. Lee, Russell Mach, Maia McLin, Julie Mitchell, Lauren Olson, Piper Sereno, Madeline Scherer, Lisa Stone, Lucia Zook. We also thank Michelle Phares for her

assistance in executing online data collection, and Drs. Lee Dixon and Jackson Goodnight for their support in developing the health survey.

REFERENCES

Bienkov, A. (2020, March 22, 2020). Sudden loss of smell could indicate "hidden carriers" of coronavirus, say UK experts. *Science Alert*. https://www.sciencealert.com/mild-covid-19-might-cause-a-lost-of-smell-or-taste

Brann, D., Tsukahara, T., Weinreb, C., Logan, D. W., & Datta, S. R. (2020). Non-neural expression of SARS-CoV-2 entry genes in the olfactory epithelium suggests mechanisms underlying anosmia in COVID-19 patients. *bioRxiv*, 2020. https://doi.org/10.1101/2020.03.25.009084

Butowt, R., & Bilinska, K. (2020, 2020/05/06). SARS-CoV-2: Olfaction, Brain Infection, and the Urgent Need for Clinical Samples Allowing Earlier Virus Detection. *ACS Chemical Neuroscience, 11*(9), 1200-1203. https://doi.org/10.1021/acschemneuro.0c00172

Chidambaram, S., & Goh, E. L. (2017). The Link Between Cytokine Levels and Loss of Olfaction in Chronic Rhinosinusitis. *JAMA Otolaryngology–Head & Neck Surgery, 143*(2), 195-195. https://doi.org/10.1001/jamaoto.2016.2424

Cieslak, K., Walsh-Messinger, J., Stanford, A., Vaez-Azizi, L., Antonius, D., Harkavy-Friedman, J., Goetz, D., Goetz, R. R., & Malaspina, D. (2015, Nov). Olfactory performance segregates effects of anhedonia and anxiety on social function in patients with schizophrenia. *J Psychiatry Neurosci, 40*(6), 387-393. https://doi.org/10.1503/jpn.140268

Dantzer, R. (2001, 2001/03/01). Cytokine-Induced Sickness Behavior: Mechanisms and

    Implications. *Annals of the New York Academy of Sciences, 933*(1), 222-234.

    https://doi.org/10.1111/j.1749-6632.2001.tb05827.x

Enard, D., Cai, L., Gwennap, C., & Petrov, D. A. (2016, 2016/05/17). Viruses are a dominant

    driver of protein adaptation in mammals. *eLife, 5*, e12469.

    https://doi.org/10.7554/eLife.12469

Gale, J. (2020, March 23, 2020). NBA player's loss of smell highlights unusual marker of Covid-

    19. 2020. *Bloomberg*. https://www.bloomberg.com/news/articles/2020-03-23/nba-

    player-s-loss-of-smell-highlights-unusual-marker-of-covid-19

Gerkin, R. C., Ohla, K., Veldhuizen, M. G., Joseph, P. V., Kelly, C. E., Bakke, A. J., Steele, K. E.,

    Farruggia, M. C., Pellegrino, R., Pepino, M. Y., Bouysset, C., Soler, G. M., Pereda-Loth, V.,

    Dibattista, M., Cooper, K. W., Croijmans, I., Di Pizio, A., Ozdener, M. H., Fjaeldstad, A.

    W., Lin, C., et al. (2020). The best COVID-19 predictor is recent smell loss: a cross-

    sectional study. *medRxiv*, https://doi.org/10.1101/2020.07.22.20157263

Hamming, I., Timens, W., Bulthuis, M. L. C., Lely, A. T., Navis, G. J., & van Goor, H. (2004,

    2004/06/01). Tissue distribution of ACE2 protein, the functional receptor for SARS

    coronavirus. A first step in understanding SARS pathogenesis. *The Journal of Pathology,*

    203(2), 631-637. https://doi.org/10.1002/path.1570

Hardy, C., Rosedale, M., Messinger, J. W., Kleinhaus, K., Aujero, N., Silva, H., Goetz, R. R., Goetz,

    D., Harkavy-Friedman, J., & Malaspina, D. (2012, Feb). Olfactory acuity is associated with

    mood and function in a pilot study of stable bipolar disorder patients. *Bipolar Disord,*

    14(1), 109-117. https://doi.org/10.1111/j.1399-5618.2012.00986.x

Health, O. D. o. (2020a). *COVID-19 Dashboard*. Retrieved June 14, 2020 from

https://coronavirus.ohio.gov/wps/portal/gov/covid-19/dashboards/overview

Health, O. D. o. (2020b). *COVID-19 UPDATE: Testing Expansion, R0 in Ohio, Place of Worship*

Best Practices, New Role for Dr. Acton, ODJFS Employment/Training Gran

https://coronavirus.ohio.gov/wps/portal/gov/covid-19/resources/news-releases-news-

you-can-use/testing-expansion

Hoffman, H. J., Rawal, S., Li, C.-M., & Duffy, V. B. (2016, 2016/06/01). New chemosensory

component in the U.S. National Health and Nutrition Examination Survey (NHANES):

first-year results for measured olfactory dysfunction. *Reviews in Endocrine and*

Metabolic Disorders, 17(2), 221-240. https://doi.org/10.1007/s11154-016-9364-1

Hopkins, C., & Kumar, N. (2020). *Loss of sense of smell as marker of COVID-19 infection*.

https://www.entuk.org/sites/default/files/files/Loss%20of%20sense%20of%20smell%20

as%20marker%20of%20COVID.pdf

Hornuss, D., Lange, B., Schröter, N., Rieg, S., Kern, W. V., & Wagner, D. (2020). Anosmia in

COVID-19 patients. *Clinical microbiology and infection : the official publication of the*

European Society of Clinical Microbiology and Infectious Diseases, S1198-

1743X(1120)30294-30299. https://doi.org/10.1016/j.cmi.2020.05.017

Hummel, T., Kobal, G., Gudziol, H., & Mackay-Sim, A. (2007, 2007/03/01). Normative data for

the "Sniffin' Sticks" including tests of odor identification, odor discrimination, and

olfactory thresholds: an upgrade based on a group of more than 3,000 subjects.

European Archives of Oto-Rhino-Laryngology, 264(3), 237-243.

https://doi.org/10.1007/s00405-006-0173-0

Inhorn, M. C., & Brown, P. J. (1990). The Anthropology of Infectious Disease. *Annual Review of Anthropology, 19*, 89-117. www.jstor.org/stable/2155960

Jang, Y., Son, H.-J., Lee, S., Lee, E. J., Kim, T. H., & Park, S. Y. (2020). Olfactory and taste disorder: The first and only sign in a patient with SARS-CoV-2 pneumonia. *Infection Control & Hospital Epidemiology*, 1-1. https://doi.org/10.1017/ice.2020.151

Keller, A., & Malaspina, D. (2013, 2013/07/23). Hidden consequences of olfactory dysfunction: a patient report series. *BMC Ear, Nose and Throat Disorders, 13*(1), 8. https://doi.org/10.1186/1472-6815-13-8

Lechien, J. R., Chiesa-Estomba, C. M., Hans, S., Barillari, M. R., Jouffe, L., & Saussez, S. (2020). Loss of Smell and Taste in 2013 European Patients With Mild to Moderate COVID-19. *Annals of Internal Medicine*. https://doi.org/10.7326/M20-2428

Lenhard, W. & Lenhard, A. (2016). *Calculation of Effect Sizes.* Retrieved from: https://www.psychometrica.de/effect_size.html. Dettelbach (Germany): Psychometrica. https://doi.org/DOI: 10.13140/RG.2.2.17823.92329

Luo, Y., Trevathan, E., Qian, Z., Li, Y., Li, J., Xiao, W., Tu, N., Zeng, Z., Mo, P., Xiong, Y., & Ye, G. (2020). Asymptomatic SARS-CoV-2 Infection in Household Contacts of a Healthcare Provider, Wuhan, China. *Emerging infectious diseases, 26*(8), 1930-1933. https://doi.org/10.3201/eid2608.201016

Malaspina, D., Goetz, R., Keller, A., Messinger, J. W., Bruder, G., Goetz, D., Opler, M., Harlap, S., Harkavy-Friedman, J., & Antonius, D. (2012, Mar). Olfactory processing, sex effects and heterogeneity in schizophrenia. *Schizophr Res, 135*(1-3), 144-151. https://doi.org/10.1016/j.schres.2011.11.025

McGraw, K. O., & Wong, S. P. (1992). A common language effect size statistic. *Psychological Bulletin, 111*(2), 361–365. https://doi.org/https://doi.org/10.1037/0033-2909.111.2.361

Moein, S. T., Hashemian, S. M. R., Mansourafshar, B., Khorram-Tousi, A., Tabarsi, P., & Doty, R. L. (2020, 2020/04/17). Smell dysfunction: a biomarker for COVID-19. *International Forum of Allergy & Rhinology, n/a*(n/a). https://doi.org/10.1002/alr.22587

Parma, V., Ohla, K., Veldhuizen, M. G., Niv, M. Y., Kelly, C. E., Bakke, A. J., Cooper, K. W., Bouysset, C., Pirastu, N., Dibattista, M., Kaur, R., Liuzza, M. T., Pepino, M. Y., Schöpf, V., Pereda-Loth, V., Olsson, S. B., Gerkin, R. C., Rohlfs Domínguez, P., Albayay, J., Farruggia, M. C. et al. (2020). More than smell - COVID-19 is associated with severe impairment of smell, taste, and chemesthesis. *Chem Senses*. https://doi.org/10.1093/chemse/bjaa041

Paules, C. I., Marston, H. D., & Fauci, A. S. (2020). Coronavirus Infections—More Than Just the Common Cold. *JAMA, 323*(8), 707-708. https://doi.org/10.1001/jama.2020.0757

Pusswald, G., Auff, E., & Lehrner, J. (2012, 12/01). Development of a Brief Self-Report Inventory to Measure Olfactory Dysfunction and Quality of Life in Patients with Problems with the Sense of Smell. *Chemosensory Perception, 5*. https://doi.org/10.1007/s12078-012-9127-7

Rabin, R. C. (2020, March 22, 2020). Lost sense of smell may be peculiar clue to coronavirus infection. *The New York Times*.

Regenbogen, C., Axelsson, J., Lasselin, J., Porada, D. K., Sundelin, T., Peter, M. G., Lekander, M., Lundström, J. N., & Olsson, M. J. (2017). Behavioral and neural correlates to multisensory detection of sick humans. *Proceedings of the National Academy of Sciences, 114*(24), 6400-6405. https://doi.org/10.1073/pnas.1617357114

Schlosser, R. J., Mulligan, J. K., Hyer, J. M., Karnezis, T. T., Gudis, D. A., & Soler, Z. M. (2016).

Mucous Cytokine Levels in Chronic Rhinosinusitis–Associated Olfactory Loss. *JAMA*

Otolaryngology–Head & Neck Surgery, 142(8), 731-737.

https://doi.org/10.1001/jamaoto.2016.0927

Scully, E. P., Haverfield, J., Ursin, R. L., Tannenbaum, C., & Klein, S. L. (2020, 2020/07/01).

Considering how biological sex impacts immune responses and COVID-19 outcomes.

Nature Reviews Immunology, 20(7), 442-447. https://doi.org/10.1038/s41577-020-

0348-8

Sepahi, A., Kraus, A., Casadei, E., Johnston, C. A., Galindo-Villegas, J., Kelly, C., García-Moreno,

D., Muñoz, P., Mulero, V., Huertas, M., & Salinas, I. (2019). Olfactory sensory neurons

mediate ultrarapid antiviral immune responses in a TrkA-dependent manner.

Proceedings of the National Academy of Sciences, 116(25), 12428-12436.

https://doi.org/10.1073/pnas.1900083116

Sepahi, A., & Salinas, I. (2016, 2016/01/01/). The evolution of nasal immune systems in

vertebrates. *Molecular Immunology, 69*, 131-138.

https://doi.org/https://doi.org/10.1016/j.molimm.2015.09.008

Sorokowski, P., Karwowski, M., Misiak, M., Marczak, M. K., Dziekan, M., Hummel, T., &

Sorokowska, A. (2019, 2019-February-13). Sex Differences in Human Olfaction: A Meta-

Analysis [Systematic Review]. *Frontiers in Psychology, 10*(242).

https://doi.org/10.3389/fpsyg.2019.00242

Speth, M. M., Singer-Cornelius, T., Oberle, M., Gengler, I., Brockmeier, S. J., & Sedaghat, A. R.

(2020). Olfactory Dysfunction and Sinonasal Symptomatology in COVID-19: Prevalence,

Severity, Timing, and Associated Characteristics. *Otolaryngology--head and neck surgery: Official journal of American Academy of Otolaryngology-Head and Neck Surgery, 163*(1), 114-120. https://doi.org/10.1177/0194599820929185

Stone, J. (2020, March 20, 2020). There's an unexpected loss of smell and taste in coronavirus patients. *Forbes*. https://www.forbes.com/sites/judystone/2020/03/20/theres-an-unexpected-loss-of-smell-and-taste-in-coronavirus-patients/#36842f765101

Vaira, L. A., Deiana, G., Fois, A. G., Pirina, P., Madeddu, G., De Vito, A., Babudieri, S., Petrocelli, M., Serra, A., Bussu, F., Ligas, E., Salzano, G., & De Riu, G. (2020, 2020/06/01). Objective evaluation of anosmia and ageusia in COVID-19 patients: Single-center experience on 72 cases. *Head & Neck, 42*(6), 1252-1258. https://doi.org/10.1002/hed.26204

Wang, Y., Liu, Y., Liu, L., Wang, X., Luo, N., & Li, L. (2020). Clinical Outcomes in 55 Patients With Severe Acute Respiratory Syndrome Coronavirus 2 Who Were Asymptomatic at Hospital Admission in Shenzhen, China. *The Journal of Infectious Diseases, 221*(11), 1770-1774. https://doi.org/10.1093/infdis/jiaa119

World Health Organization (2020). *Coronavirus disease (COVID-19) pandemic*. Retrieved 06/29/2020 from https://www.who.int/emergencies/diseases/novel-coronavirus-2019

Table 1. Means, Standard Deviations for the Cohorts and Results of Cohort Comparisons

	Cohort II (exposed)	Normative Sample			Cohort II vs Cohort I		Cohort II vs Cohort III		Cohort II vs Normative Sample	
	N=22	All‡ N=243	Cohort I N=25	Cohort III† N=52	t	p	U	p	D	p
Age, M ± SD	19.45 ± 1.23	19.28 ± 1.11	19.08 ± 1.08	19.31 ± 1.09	0.93	.36	550.50	.79	1.92	.83
Menstrual cycle day, M ± SD §	15.41 ± 8.56	17.31± 8.20	14.59 ± 8.51		-0.97	.35				
					X^2	p	X^2	p	X^2	p
Gender — number (%)					0.00	.99	0.482	.49	0.49	.83
Male	7 (32)	83 (34)	8 (32)	21 (40)						
Female	15 (68)	160 (66)	17 (68)	31 (60)						
Racial/Ethnicity — number (%)					0.41	0.52				
Asian or Asian American	1 (4.5)	(12)	1 (4)							
Black or African American	2 (9)	(3)	3 (12)							
Hispanic White	2 (9)	(12)	4 (16)							
Non-Hispanic White	16 (73)	(71)	16 (64)							
Other	1 (4.5)	(3)	1 (4)							
Cigarette Use N (%)					0.39	.53	0.64	.43	0.94	.33
None	19 (86)	250 (92)	22 (88)	48 (92)						
Yes, but not daily	1 (5)	14 (5)	1 (4)	3 (6)						
Yes, daily	2 (5)	4 (2)	2 (8)	1 (2)						
Use of Vaping Devices N (%)					2.02	0.37				
None	17 (77)	(82)	18 (72)							
Yes, but not daily	3 (14)	(14)	2 (8)							
Yes, daily	2 (9)	(3)	5 (20)							

Tests statistics refer to: t=Independent Samples t-test; U=Mann-Whitney U-test; D=Two Samples Kolmogorov–Smirnov test

§Females only; converted to a 28-day cycle; Cohort I and II N=15 females; data missing from Cohort I females (N=2)

‡Data missing: race/ethnicity and use of vaping devices (N=209); menstrual cycle in females (N=137); Counts were not reported and comparative analyses were not conducted due to the large percentage of missing data.

†Data missing: race/ethnicity and use of vaping devices (N=47); menstrual cycle in females (N=26); Counts and percentages were not reported and comparative analyses were not conducted due to the large percentage of missing data.

Table 2. Means, standard deviations, and diagnostic classification for olfaction measures, by cohort.

	Cohort II		Normative Sample		
	All	**Cohort IIa**	**All**	**Cohort I**	**Cohort III**
	N=22	N=11	N = 272	N=25	N=52
Detection Threshold	7.32 ± 3.21	6.50 ± 2.76	8.40 ± 2.15	9.79 ± 3.24	8.26 ± 19.31
% Correctly Identified	72.44 ± 12.74	67.05 ± 11.21	71.22 ± 10.26	71.25 ± 11.98	72.36 ± 9.37
Pleasantness Ratings	25.10 ± 7.86	23.20 ± 9.03	24.55 ± 7.23	25.20 ± 7.62	24.79 ± 7.77
Unpleasantness Ratings	12.05 ± 6.18	13.55 ± 5.99	18.54 ± 7.42	14.52 ± 4.07	19.37 ± 7.39
Smell Diagnosis — no. (%)					
Hyposmia	11 (50)	7 (64)	31 (11)	2 (8)	2 (4)
Normosmia	11 (50)	4 (36)	241 (89)	23 (92)	44 (96)

Unless otherwise indicated, data displayed are M ± SD
†Odor intensity ratings missing from Cohort I (N=3), Cohort II (N=8), and the normative sample (N=249)

Figure 1. Sample sizes and assessment time periods for **a)** olfaction tested cohorts, and **b)** health survey participants.

Figure 2: Cumulative distributions of odor detection sensitivity for the normative sample (blue) and Cohort II (orange), fit to a gamma distribution (dotted line), and normal distribution (solid line). Distribution fits were applied as follows: Normative distribution gaussian fit mean=8.26113 [8.01615, 8.50612], sigma=1.95481 [1.79628, 2.14426]; Cohort II Gamma distribution fit mean=4.91205, a=5.99561 [3.37288, 10.6578] ,b=1.22059 [0.669847, 2.22415] Inset: histograms for odor sensitivity threshold for the normative sample (blue; normal fit) and for Cohort II (orange; gamma fit).

Figure 3. Temporal plot of the percentage of hyposmic participants (orange line) compared to normative population (blue dot).