Title Page

Title

Hospital based autonomous pre-clinical screening of COVID-19: An emergency triage

using a vital signs recording system, Paris-Ile de France region.

Authors

Albert Brizio MD (1), Valérie Faure MD (2), Franck Baudino MD (3), Arnaud Wilmet MD
(4), ORCID: http://orcid.org/0000-0003-3063-1770   Jean-Paul Gonzalez MD (5)
8

Affiliations

(1) SMUR, Hôpital Delafontaine, Centre hospitalier de Saint Denis, , Saint Denis, France.

brizioalbert@hotmail.com; (2) Emergency Department Hôpital privé du Vert Galant Groupe

Ramsey, Tremblay en France, France. Email : Valerie.Faure@ramsaygds.fr; (3) Health for

Development, Paris, France, Email : franck.baudino@h4d.com; (4) Health for Development,

Paris, France, Email : arnaud.wilmet@h4d.com ; (5) School of Medicine, Georgetown

University, Department of Microbiology & Immunology, Washington, DC 20057 USA.

Commonwealth Trade Partners Inc. Centaurus Biotech LLC. Email:

Jean.Paul.Gonzalez@georgetown.edu

Corresponding author

Albert Brizio, 14 Place Etienne Pernet, 75015, Paris, France. brizioalbert@hotmail.com

Acknowledgments

This work and the study behind has been possible with the exceptional support of RAMSAY

SANTE, Health for Development and the participation of the staff of the Vert Galant

Hospital, Madame Sandrine Legras, Dr. François-Xavier Peccaud, and Dr. Florian Gontier.

We want to thank M. Marc Souris for map design and Ms. Soukaina Gonzalez for writing

assistance, technical editing and language editing.

ABSTRACT

The COVID-19 pandemic, has had a dramatic toll worldwide on the populations but also has

been essentially supported by the existing public health system, particularly hospital-based

emergency wards and intensive care units.

In France, the first cases were identified on the 24th of January 2020. The first epidemic

sprout emerged in the Eastern part of the country and spread in two weeks towards the center

to the Paris-region where it peaked on the 14th of April 2020. In Paris and the region around

it, the intensity of the epidemic has increased significantly to have a strong impact on all

public and private hospital systems in a few weeks.

During France's 2020 COVID-19 epidemic, a private hospital went into a major

organizational change of its Emergency Department which mainly included the use of a

Telemedicine Booth for vitals automatic recording during triage procedures.

The present study describes the organizational scheme adopted by the hospital and discusses

the data of 1,844 patients that attended the facility over a month. While among them, 766

patients were engaged in a automated triage process supported by a Telemedicine Booth.

Patients' clinical characteristics are comparable to those found in international literature

during the COVID-19 pandemic.

The use of the Telemedicine Booth as a screening process facilitated patients' flow. It

usefully participated in the patient rapid orientation, relieving the hospital emergency

department, actively contributes in a safe and secure environment highly trusted by the

hospital staff and health workers. To our nowledge, the Telemedicine Booth use as a

screening process during an epidemic constitutes a first contribution to such an innovative

approach.

Competing Interest Statement

The authors have declared no competing interest.

Funding Statement

No external funding was received for this work

Authors Declarations

The authors confirm that all relevant ethical guidelines have been followed, and any necessary

IRB and/or ethics committee approvals have been obtained.

The details of the IRB/oversight body that provided approval or exemption for the research

described are given below: NA

All necessary patient/participant consent has been obtained and the appropriate institutional

forms have been archived.

The authors understand that all clinical trials and any other prospective interventional studies

must be registered with an ICMJE-approved registry, such as ClinicalTrials.gov. I confirm

that any such study reported in the manuscript has been registered and the trial registration ID

is provided (note: if posting a prospective study registered retrospectively, please provide a

statement in the trial ID field explaining why the study was not registered in advance).

The authors have followed all appropriate research reporting guidelines and uploaded the

relevant EQUATOR Network research reporting checklist(s) and other pertinent material as

supplementary files, if applicable.

INTRODUCTION

The World Health Organization (WHO) declared, on March 11[th] 2020, the Coronavirus

disease 2019 (COVID-19) as a pandemic, pointing to the increasing number of infections

among more than 110 countries and territories worldwide, and the risk of further global

spread. Considered by WHO, Europe had become by that time the epicenter of the pandemic,

the level three of the COVID epidemic emergency was officially declared by the French

authorities on March 14[th] 2020[1].

According to the local governmental instructions, the Vert Galant Hospital (Fig. 1), as a

private health facility within Paris metropolitan area, shifted from normal activity to a crisis

management on March 19[th]. The newly developed engagement involved the whole hospital

aiming principally to facilitate the influx of new patients and avoid overwhelming the

potential of the Emergency Department (ED) (i.e. inclusive of the Intensive Care Unit). This

article describes, analyzes this innovative organizational scheme and discusses the outcomes

of its development and the impact over a month's period of time (March 19[th] to April 19[th],

2020) during the intense inception of COVID-19 epidemic within Paris metropolitan area.

PERSONNEL, METHOD, AND MATERIAL

Study site. The Vert Galant Hospital is a 100 bed private hospital with an intensive care unit

(ICU) of 14 beds of an emergency department (ED) with a potential to manage a mean flux of

hundred patients a day, as well as other medical wards including: gynecology and obstetrics;

orthopedics; oncology; cardiology; gastroenterology; pulmonology; urology; and

endocrinology. It essentially covers a low-medium income population of the Seine Saint-

Denis, 93 Department (French District Unit), wich is one of lowest income areas in France

with a poverty rate of 26,9% as compared to the 14,3% of continental France[2]. The hospital

accepts all patient with national health insurance as well as irregular migrants without

insurance.

The Vert Galant emergency department was designated by its hospital board to manage the

increasing flow of patients due to the intensive rise of the COVID-19 epidemic. Therefore, by

setting up a new organizational scheme, measures were taken to lessen risks for the health

workers without reducing the efficiency of patient care.

Telemedicine Booth. One of the main measures of the new hospital strategy during the

epidemic crisis was to implement the use of a Telemedicine Booth (H4D ConsultStation®)

which allow, among other health data collection, the measurement and the automatic

recording of the patients' essential vital signs. The Telemedicine Booth is also a

teleconsultation facility allowing, besides vital sign collection, an interactive video

conferencing with a medical practitionner when needed, the latter was not used in the present

context of the management. The Telemedicine Booth is a Class 2 certified medical device and

can be described as a fiberglass cylinder with a seat, a video and audio system plus a touch

video screen for patient-machine interaction, printer, and a series vital signs' measurement

devices (Table 1).

Health workers and administrative staff. The ED, on a daily basis, is staffed by two (2)

senior medical practitioners (MP), three (3) nurses, three (3) aids/stretcher-bearers, two (2)

secretaries, and during the night shift by one MP (1), two (2) nurses, one aid/stretcher-

bearers, and one (1) secretary.

Management and organizational scheme. The ED management was and is under the

responsibility of the MP and the director of the hospital. According to emergency plan, a

COVID-19 crisis unit was set up while routine hospital activity was reduced or suspended for

some non-essential sectors. Two-thirds of the surgery wards were transformed in COVID-19

wards and the number of ICU beds was increased by two fold (i.e. 28 beds). One of the ED

MP was mandated as coordinator in charge of the organizational change and supported by the

chief nurse. Technical choices were team verdicts involving the ED MP, the Chief Nurse, and

a crisis unit representative.

The COVID-19 ED system was set up for the time being, it was designed to focus on the

implementation, to follow existing recommendations as far as possible as well as previous

experiences. Records were kept of all activities mainly using existing procedures. Ultimately,

the principal objectives where: 1/ To create a specific Covid-19 patient flow in the existing

facilities; 2/ To maintain a separate non-COVID-19 flow; 3/ To manage the patient and record

their vital signs in view of reducing infectious risk (i.e. infection control) and protecting the

health workers and the patients. In order to achieve these objectives, the capacity of the ICU

increased by two-fold, and patients' flow was redesigned. The outpatient activity and facilities

were canceled, such facilites, situated in the immediate proximity of the ED, were designated

for non-COVID-19 emergency space. Also, it was decided to keep as the single public

entrance to the hospital, the ED entrance, and to create an external temporary initial triage

station in order to separate potential COVID-19 from non-COVID-19 patients before entering

the facility. The parking area, in front of the ED entrance, was utilized to set up a tent (fig.1).

Strategic engagement. Triage the act of prioritizing patients' access and determining

severity, usually performed by a trained nurse, was considered strategical. The pre-existing

ED triage room was small, with open entry with the secretaries' room. To avoid any risk of

transmission for nurses during the triage, it was decided to use a Telemedicine Booth

allowing an autonomous recording of the vital signs in association with a questionnaire.

Human supervision was considered essential to speed up the triage process and to provide the

necessary help if needed. It was decided to install the Telemedicine Booth in the ED waiting

room and train a group of stretcher-bearers for this novel role. Health workers received two

(2) hour training focusing on the COVID-19 infection risk including: The use of personal

protective equipment; the technical management of the Telemedicine Booth; infection control

(hands cleaning, Telemedicine Booth disinfection and safe disposal); the recognition of

respiratory distress as well as other potentially severe conditions.

The triage itself was modified during daily activity by replacing the FRENCH 2018 triage

grid[3] for non-COVID-19 patients maintained in the blue sector, by a three steps procedure for

COVID-19 patients inclduing: A first step at the tent, a questionnaire was intended to identify

COVID-19 patients; a second step, in the red sector, the vital signs taken at the Telemedicine

Booth intended to first evaluate the severity of clinical presentation; and a third step for the

nurse to re-evaluate the syndrome severity while contextualizing it with the help of medical

history before the medical examination by the MP.

During night hours the organization shifted while the night attendance rate did not justify the

Telemedicine Booth facilitator's use and did not include it for screening.

Patient pathway. Outside of the hospital premises, the parking access was controlled, while

immediately front of the entrance the "first triage" tent was set up. A medical student or a

nurse fully dressed with personal protective equipment (PPE) sat at the entrance for: Guiding

the patient to disinfect his hands with hydroalcoholic gel and to accurately fit a provided

surgical mask; applying a questionary on symptoms and reasons to attend. The questionary

was based on available clinical guidelines and a formal list of symptoms to identify suspected

COVID -19 cases[4].

Patients waited in a single row at 1,5m distance one to each other following marks on the

ground. Once oriented to his destination, the patient was invited to cross the tent and enter the

building through a glass sliding door. The pathway used by each patient was directed through

a color-coded system (i.e. red Covid-19 suspected; blue non-Covid-19) from the time he

entered the hospital to the time he left the triage and was redirected based on the data acquired

(Fig. 2). Non-COVID-19 patients followed blue signs leading to blue doors giving access to

the "blue" sector. The "blue" patients would simply proceed straightforward, enter the blue

zone, and be attended. When needed, they would attend the blue sector of radiology, if

dismissed, would exit through a dedicated door. COVID-19 suspected patients would follow

the red signs leading to the "red" ED sector. The "red" patients, had to turn left, go through a

transparent curtain, then an automatic sliding door leading to the waiting room, before being

welcomed by the stretcher-bearer fully dressed in the appropriate PPE and in charge of the

Telemedicine Booth. In the autonomous "check-up" modality, the patient was guided through

the Telemedicine Booth by a video and audio tutorial. At the end of the process, the

Telemedicine Booth print out a report. This process took usualy seven (7) minutes to the

patient with no external help.

Patients transported by ambulance were bypassing the tent and entering directly to the

dedicated wards. During the night, the first triage was maintained in the tent although in the

red sector, the Telemedicine Booth was bypassed, and, once registered the patient was

directly admitted to the examination room where the vital signs were measured by the nurse.

In order to maintain security distances one out of two seats of the waiting room was marked

with tape crosses for not to be used. Once the administrative formalities finished at the

registration desk, the patient was invited to enter the Telemedicine Booth, having first

disinfected his hands. After the vital signs recording, the patient stayed in the waiting room

under the supervision of the specially trained stretcher-bearer unless the saturation level was

below 95%. This would lead to immediate admittance of the medical examination. Once

admitted to the examination room, a nurse took the patient's personal history, measured

oxygen saturation and respiratory rate. Followed the medical examination. If required, the

patient was accompanied or redirected to the COVID-19 dedicated CT-SCAN (Computed

tomography Scanner) for lung examination following a specific "red" path.

The health worker entered from a separate entrance in the "clean" part of the ward. A buffer

zone, called dressing/undressing zone, separates it from the red zone. In the clean zone, the

health worker dressed with PPE before entering the red one. During their activity in the red

zone, overcoats, aprons, gloves, and masks, replacements were available. While returning in

the clean zone, undressing was mandatory as was the disinfection of the hands following

infection control guidelines. The red and blue sectors initially had two seperate teams, but

organizational needs led to frequent shifts of health workers from one team to the other.

Except for MP, two separate health worker groups covered the 12 hours day and night shifts.

Activities evaluation. Evaluation procedures were not specifically included in the project

design. The register at the tent, the patient management software, and the Telemedicine Booth

activity software ensured data collection with accuracy. All software complied with existing

French regulations and laws. An *a posteriori* survey was conducted among the personnel of

the ED for qualitative feedback including a questionary concerning their perception of

workload, contagious risk, and difficulties encountered with the Telemedicine Booth were

submitted.

RESULTS

Quantitative indicators of activity. Over the study period, which corresponds to the

epidemiological peak of COVID-19 in France[5], 1,844 patients attended the Hopital du Vert

Galant ED. 766 of them (41.5%) were triaged at the tent as suspected COVID-19 cases and

were directed to the "red" sector (RS). The total number of admissons in the RS is higher

(838) due to a 9.3 % readmissions. The population's mean age was 47.6 years (46.0 yo

female, 49.0 yo male). 61% (51.9 % males) of the patients were diagnosed as COVID-19

positive, either clinically or with the support of RT-CPR and Chest CT-SCAN (Computed

Tomography Scanner). Out of the 469 diagnosed COVID-19 positive, 50.9% had a diagnostic

confirmation by RT-CPR and/or Chest CT-SCAN. 13.4% of the total screened patients were

hospitalized (among them 67% males) including 2.5% in ICU (70% males).

88% attended the ED between 8 AM and 8 PM and, among them, 68% passed through the

Telemedicine Booth for autonomous vital signs recording. The data collected from 47 patients

allowed to calculate the mean time between the first triage at the tent and the registration by

the secretary (15,6 minutes). The registration to Telemedicine Booth time was 12,1 minutes.

The mean time spent by the patients in the Telemedicine Booth, calculated for all patients,

lasted for only 4,2 minutes (Table 2).

The new organization set up took three days and needed the workload of ten (10) health

workers. 21 full-time nurses and nurse-assistants and, 31 individual contractors working for

the ED during this period.

Out of the 21 full-time health workers, 15 answered to the survey evaluating their subjective

perception of the new organizational scheme: 7 found their workload increased, 4 unchanged,

and 4 reduced, while 12 did not find difficulties related to the use of the Telemedicine Booth

and 9 felt that the use of the booth increased their safety. Among the full-time health workers,

2 only tested positive by RT-CPR without proven origin (domestic or profesional) of the

infection.

DISCUSSION

In comparison to the same time period in 2019, at the Vert Galant Hopital, there has been a

43.0% reduction of the ED's attendance. This reflects a global phenomenon during the

COVID-19 crises[6] while unexpectedly, the COVID-19 suspected patients were less than the

half of the usual frequentation, nevertheless, it was about 6 times higher than the COVID-19

attendance rate of the 93 Department's public hospitals (41,5% % versus 6,8%)[7]. This could

be explained due to close-by population considering the hospital as a "COVID hospital"

refraining them to attend the hospital for other medical reasons. Altogether, these figures

show the importance of maintaining adequate services for non-COVID-19 cases or any

concurrent critical epidemic.

The risk of focusing clinical inquiry on COVID-19 symptoms concerned mainly the patients

presenting atypical symptoms of the disease and false positives. Therefore, the initial triage

not only has to avoid under-triage for the contagious disease but also under-triage of the other

concurrent conditions not related to it. Possibly, the multi step triage that has been used during

the study, appears to be the most appropriate approach to avoid such risk. By assuming the

accuracy of clinical diagnosis, an over-triage of 39% was estimated. Also, an over-triage of

25 to 50%[8] is accepted for traumas, the present over-triage may be acceptable for effectively

reducing the potential risk of COVID-19 positive patients admittance to non-COVID-19

areas. However, over triage in epidemic conditions exposes false positive patients to

infectious risk. With this perspective, the 39% figure may be considered excessive but limited

data are available on this matter from the medical litterature.

The 13.4% COVID-19 hospitalization rate is half of the one registered by the department's

public hospitals (27,2%)[9] and this could be partially explained by the relatively young age of

the population who have attended the Vert Galant hospital facilities.

Although males represented half of the COVID-19 positive patients, they denote 67% of the

one hospitalized and 70 % of the one ICU admitted. This is in agreeemnt with other produced

elsewere of an increased occurrence of severe COVID-19 cases incidence in males[10].

Moderate underuse of the Telemedicine Booth is probably due to the speeding-up of the flow.

However, to avoid any delay, when the Telemedicine Booth was in use and an examination

room was available, the patient was directly admitted to the room thus bypassing the

Telemedicine Booth. Nonetheless, about 35 nurse-time hours were spared over a one month

period.

The time between registration and the vital signs taking by the Telemedicine Booth is about

nine (9) minutes less than the time between registration and vitals measurement by the nurse

in normal times (22 minutes). The average total time from the gate triage to the Telemedicine

Booth was 27.7 minutes. The mean vital signs' recording time at the Telemedicine Booth was

61%, less than the 7 minutes needed for a complete autonomous video-guided vital signs

recording (used for check-up in a non-epidemic context). This difference is due to the

dedicated personnel that actively assisted patients and often skipped the video tutorial used

for patient self-guidance. Generally speaking, patient's security was enhanced by the new

system by reducing the vital signs recording time and ultimately improved patient's

supervision.

Despite the consistent decrease of patients' attendance, the majority of the personnel did not

perceive their workload as reduced because the organizational constraints linked to the new

triage, dressing, disinfection procedures, and patients' anxiety. The majority did not consider

the Telemedicine Booth as a burden when more than half of them considered that its use

increased their own safety. If a few expressed some concern, it was always associated with

human error.

The incidence of COVID-19 among non-medical health workers can be evaluated with

difficulty because of the small sample size and the numerous confounding factors: shifts

between red and blue zones, the absence of routine testing, and the absence of a baseline. To

our knowledge, statistical data concerning health workers being not yet available for France,

the 14% of symptomatic COVID 19 health workers found in this study (beside a non-proven

infection origin) seems in agreement with the 10% infection rate among nurses found in

Lombardy, Italy[11].

The importance to adjust the facility to the epidemic context and to protect the health workers

from infection was used as an opportunity for the patient's health education. Therefore,

necessary time was taken for patients' guidance, in particular to accurately practice hands

disinfection and masks fitting upon admission for triage at the tent and Telemedicine Booth

use.

CONCLUSION

This case study illustrates how a major change in organizational habits centering on the

introduction of an automatic vital signs recorder in an emergency department can very well

possible with a limited disadvantage. The presence of the Telemedicine Booth did not slow

down the patients' triage process, and potentially improved it by an efficient triage and a

better trust of the health worker adherence to protective measures.

Although, more research would be needed to fully identify the pros and cons induced by the

use of automated vital signs recording Telemedicine Booth during an epidemic in hospital

settings. This experience constitutes a first contribution to the exploration of this innovative

approach.

REFERENCES

1. France. Ministère des Affaires Sociales et de la Santé. Circulaire interministérielle
DGS/BOP/DGSCGC no 2012-420 du 17 décembre 2012 relative au plan national
de prévention et de lutte « pandémie grippale ». Available from: https://solidarites-
sante.gouv.fr/fichiers/bo/2013/13-04/ste_20130004_0100_0063.pdf
313
314   2. Quelques statistiques récentes sur la pauvreté à Saint-Denis – Saint Denis Ma
315   Ville [Internet]. [cited 2020 Jun 17]. Available from:
316   https://www.saintdenismaville.com/quelques-statistiques-recentes-sur-la-
317   pauvrete-a-saint-denis/

318
319   3. Société Française de Médecine d'Urgence. Le triage en structure des urgences.
320   Recommandations formalisées d'experts . Paris SFMU 2013. Available from :
321   https://www.sfmu.org/upload/referentielsSFMU/2019%2012%2006%20FREN
322   CH%20pour%20PDF%20A4%20V1.2%20site.pdf
323
324   4. Guan W-J, Ni Z-Y, Hu Y, Liang W-H, Ou C-Q, He J-X, et al. Clinical Characteristics of
325   Coronavirus Disease 2019 in China. N Engl J Med. 2020 30;382(18):1708–20.

326
5. INFOGRAPHIES. Coronavirus : morts, hospitalisations, âge des malades... Suivez
l'évolution de l'épidémie en France et dans le monde [Internet]. Franceinfo. [cited
2020 Jun 17]. Available from:
https://www.francetvinfo.fr/sante/maladie/coronavirus/infographies-covid-19-
morts-hospitalisations-age-malades-l-evolution-de-l-epidemie-en-france-et-
dans-le-monde-en-cartes-et-graphiques.html

333
334   6. ESRI, Brick A, Walsh B, ESRI, Keegan C, ESRI, et al. COVID-19 and emergency
335   department attendances in Irish public hospitals [Internet]. ESRI; 2020 May
336   [cited 2020 Jun 17]. Available from: https://www.esri.ie/publications/covid-19-
337   and-emergency-department-attendances-in-irish-public-hospitals
338
339   7. Géodes - Santé publique France - Indicateurs : cartes, données et graphiques
340   [Internet]. [cited 2020 Jun 17]. Available from:
341   https://geodes.santepubliquefrance.fr/index.php#c=indicator&f=0&i=sursaud_c
342   orona_hebdo.prop_corona_pass_sau&s=2020-S11&t=a01&view=map2
343
344   8. DO DEH, MD JLB, editors. Disaster Medicine. Second edition. Philadelphia: LWW;
345   2007. 512 p.
346
347   9. Taux d'hospitalisation en France- COVID-19 - Médecine d'urgence - Urgences
348   médicales [Internet]. [cited 2020 Jun 17]. Available from: https://urgences-
349   serveur.fr/taux-d-hospitalisation-en-france-covid-19.html
350
10. Plaçais L, Richier Q. COVID-19 : caractéristiques cliniques, biologiques et
radiologiques chez l'adulte, la femme enceinte et l'enfant. Une mise au point au
cœur de la pandémie. La Revue de Médecine Interne. 2020 May 1;41(5):308–18.

11. Italia. Ministero della Sanità. Istituto Superiore di Sanità. Epidemia COVID-19
Aggiornamento nazionale 28 aprile 2020. Available from :
https://www.epicentro.iss.it/coronavirus/bollettino/Bollettino-sorveglianza-
integrata-COVID-19_28-aprile-2020.pdf
358

360 Tab 1: The vitals recorded by the hospital based Telemedicine Booth during the COVID-19

361 epidemic in the Paris metropolitan area (may-April 2020)

Vital Signs	Recording device
SpO2*	Pulse oxymeter
Cardiac frequency	Pulse oxymeter
Temperature	Infrared thermometer
Blood pressure	Electronic sphygmomanometer on the left side
Body mass index	Inbuilt scale and height gauge

363 * Sp02 = Peripheral oxygen saturation

365   Table 2 : Main quantitative data acquired during the during the COVID-19 epidemic in the

366   Paris metropolitan area (may-April 2020)

Data Recording during period	Patients screened in the Red Sector « Covid-19» hospital *		
	Total	Female	Male
Admissions	838	411	427
Patients	766	382	384
Mean ages	47.6	46.0	49.2
Positive COVID-19 **	469	227	242
Positive COVID-19 PCR &/or CT-SCAN	239	107	132
Positive COVID-19 clinically diagnosed	230	120	110
Deceased in ED	1	1	0
Hospitalized in intensive care (number)	20	6	14
Hospitalized in medicine wards	83	30	53
Dismissed (Exits)	662	345	317
Registered between 8 AM and 8 PM	673	338	335
Telemedicine Booth use between 8 AM and 8 PM total patients	458	241	217
Telemedicine Booth use between 8 AM and 8 PM total admissions	524	272	252
Triage tent to Registration (secretary) [$]	15.6	/	/
Registration to Telemedicine Booth [$]	12.1	/	/
Time spent in Telemedicine Booth [$]	4.2	/	/

367   *= Patient hospitalized; ** = clinically & rtPCR/CT-SCAN; [$] = Mean time in minutes

369 Figure 1 : Incidence of COVID-19 cases and localization of Paris and the Vert Galant

370 Hospital in the Seine Saint Denis Departement (modified from Santé publique France)

371
372
373

374   Fig 2: COVID-19 patient pathway upon admission to the Emergency Department of the

375   hospital during the epidemic of Paris Metropolitan area (March-May 2020).

377   Legend:

Health worker in Personal Protective Equipment (PPE)

Patient online distancing marks

Temporary wall

Curtain

Barrier

No seating

PPE stock point