

# 1 Asynchronicity of endemic and emerging mosquito-borne disease 2 outbreaks in the Dominican Republic

3  
4 Mary E. Petrone<sup>1\*</sup>, Rebecca Earnest<sup>1\*</sup>, José Lourenço<sup>2</sup>, Moritz U.G. Kraemer<sup>2</sup>, Robert Paulino-Ramirez<sup>3</sup>,  
5 Nathan D. Grubaugh<sup>1\*</sup>, Leandro Tapia<sup>3\*</sup>

6  
7 <sup>1</sup>Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven, CT 06510, USA

8 <sup>2</sup>Department of Zoology, University of Oxford, Oxford, United Kingdom

9 <sup>3</sup>Instituto de Medicina Tropical & Salud Global, Universidad Iberoamericana, Santo Domingo, Dominican Republic

10 \*These authors contributed equally

11 Correspondence: [mary.petrone@yale.edu](mailto:mary.petrone@yale.edu) (M.E.P.); [L.tapia@prof.unibe.edu.do](mailto:L.tapia@prof.unibe.edu.do) (L.T.)

## 12 **ABSTRACT**

13  
14 Mosquito-borne viruses pose a perpetual public health threat to countries and territories in the Caribbean  
15 due to the region's tropical climate and seasonal reception of international tourists. Outbreaks of the  
16 emerging viruses chikungunya and Zika in 2014 and 2016, respectively, demonstrated the rapidity with  
17 which these viruses can spread between islands. At the same time, the number of reported dengue fever  
18 cases, caused by the endemic dengue virus, has steadily climbed over the past decade, and a large dengue  
19 outbreak that began sweeping through this region in 2019 continues in 2020. Sustainable disease and  
20 mosquito control measures are urgently needed to quell virus transmission in the long term and prevent  
21 future outbreaks from occurring. To improve upon current surveillance methods, we analyzed temporal and  
22 spatial patterns of chikungunya, Zika, and dengue outbreaks reported in the Dominican Republic between  
23 2012 and 2018. The viruses that caused these outbreaks are transmitted by *Aedes* mosquitoes, which are  
24 sensitive to seasonal climatological variability. In this study, we evaluated whether climate and the spatio-  
25 temporal dynamics of past dengue outbreaks could inform when and where future emerging disease  
26 outbreaks might occur. We found that the temporal and spatial distribution of emerging disease outbreaks  
27 did not conform to those of seasonal dengue outbreaks. Rather, the former occurred when climatological  
28 conditions were suboptimal for *Aedes* activity. Provincial dengue attack rates did not correspond to those of  
29 emerging diseases. Our study also provides evidence for under-reporting of dengue cases, especially  
30 following the 2016 Zika outbreak. We advocate for the implementation of a sustainable and long-term  
31 surveillance system to monitor the spread of known mosquito-borne viruses and to identify emerging threats  
32 before they cause outbreaks. Specifically, we recommend the use of febrile illness incidence, case fatality  
33 rates, and serosurveys during inter-outbreak periods to better understand rates of transmission and  
34 asymptomatic infection.

## 35 **INTRODUCTION**

36 Emerging and endemic mosquito-borne viruses are a constant public health concern in the Caribbean (Cao-  
37 Lormeau, 2016; Mavian et al., 2018; Musso et al., 2018). This region is especially vulnerable to the spread  
38 of the former due to its tropical climate and large tourism industry, which attracts visitors from all across the  
39 globe. The recent outbreaks of chikungunya (Cassadou et al., 2014; Dorléans et al., 2018; Halstead, 2015)  
40 and Zika (Faria et al., 2016b; Grubaugh et al., 2018; Metsky et al., 2017) in 2014 and 2016, respectively,  
41 demonstrated that viruses once associated with mild illness can re-emerge and cause devastating health  
42 outcomes. Reports that Mayaro virus, which has already been detected in the Caribbean, can be transmitted  
43 by the urbanized mosquito species *Aedes* indicate that future outbreaks of emerging mosquito-borne  
44 diseases may be on the horizon (Ganjian & Riviere-Cinamond, 2020; Hotez & Murray, 2017; Long et al.,  
45 2011; Lorenz et al., 2019; Weaver & Reisen, 2010). In addition to this threat, dengue virus is endemic to

46 many Caribbean countries and territories and has caused outbreaks with increased frequency over the past  
47 decade. Large outbreaks, which began in 2019 but have continued through 2020 (Pan-American Health  
48 Organization (PAHO), 2020a), extend an alarming trend of a rising number of dengue cases reported  
49 annually in the Americas in recent years (PAHO, 2014). However, the danger of these viruses and the  
50 diseases they cause lie not only in their debilitating and sometimes life-threatening symptoms. The  
51 unpredictability of when and where a new outbreak will occur precludes preparedness. Outbreak response  
52 strategies are inherently reactionary and, in their transience, disrupt public health systems when new  
53 initiatives are introduced and subsequently phased out. This is especially problematic in resource-limited  
54 settings where the strategic allocation of resources should be prioritized to maximize the impact of disease  
55 control efforts. New approaches centered around sustainable, long-term surveillance are needed to curtail  
56 the potential for future public health crises in the Caribbean.

57 One such approach is the use of climate data to evaluate the risk of viral spread. The transmission of  
58 mosquito-borne viruses, both emerging and endemic, should adhere to similar temporal and spatial patterns  
59 such that the dynamics of past outbreaks can inform those of future outbreaks. Many flaviviruses, like  
60 dengue and Zika viruses, and alphaviruses, like chikungunya virus, are transmitted by *Aedes* mosquitoes,  
61 which are sensitive to climatological variability (Kraemer et al., 2019; Kraemer, et al., 2015; Obolski et al.,  
62 2019). Temporal concordance between seasonal weather patterns and transmission of yellow fever virus  
63 (flavivirus) has been documented on the African continent (Hamlet et al., 2018). Similar associations have  
64 been reported for dengue fever incidence in Hanoi, Vietnam and in Myanmar (Do et al., 2014; Perez-  
65 Guzman et al., 2018) and for chikungunya and Zika incidence in the Americas (Obolski et al., 2019).  
66 However, there is evidence to suggest that this relationship does not extend to the transmission dynamics  
67 of viruses during their first year of introduction into a new population (Lourenço et al., 2017; Faria et al.,  
68 2016). Therefore, we considered whether, while climate may be a useful indicator for future endemic virus  
69 outbreaks, other factors including population size, demographics, and the timing of introduction should be  
70 considered when developing strategies to prevent future emerging disease outbreaks.

71 To answer these questions in the context of the Caribbean, we analyzed dengue, chikungunya, and Zika  
72 cases reported daily in the Dominican Republic between 2012 and 2018. We found that emerging disease  
73 outbreaks (chikungunya and Zika) occurred earlier in the year than dengue outbreaks, and the timing and  
74 location of introductions of emerging viruses impacted when and where corresponding outbreaks occurred.  
75 Moreover, the spread of chikungunya and Zika viruses was tolerant to sub-optimal climates for transmission  
76 by *Aedes* mosquitoes, likely due to the large size of the susceptible human population. Predicted mosquito  
77 abundance was similarly uninformative for the spatial distribution of emerging disease attack rates and force  
78 of infection. Moreover, provincial-level dengue attack rates were consistent between dengue outbreaks, but  
79 they did not correspond to local attack rates of chikungunya and Zika. Taken together, we demonstrate that  
80 dengue virus may not always be an appropriate model to prepare for future emerging mosquito-borne  
81 disease outbreaks. Instead, a sustainable and long-term mosquito-borne disease surveillance system is  
82 needed to facilitate proactive responses to emerging threats and to track the continued spread of known  
83 diseases including dengue, chikungunya, and Zika. We specifically propose the use of two indicators,  
84 incidence of febrile illness cases and dengue case fatality rates, to monitor surveillance performance and  
85 identify potential emerging threats.

## 86 RESULTS

### 87 Multiple endemic and emerging mosquito-borne virus outbreaks in the Dominican Republic

88 Between 2012 and 2018, the Dominican Republic, a country that shares the Caribbean island La Hispaniola  
89 with Haiti (**Fig. 1a**), experienced five disease outbreaks caused by mosquito-borne viruses (**Fig. 1b,c**). We

90 delineated this period into six seasons (Seasons 1-6), each beginning in April of every year between 2012  
 91 - 2018, coincident with the start of the rainy season. Three of the five outbreaks were caused by dengue  
 92 virus (Seasons 1, 2, and 4; **Fig. 1b,c**). Sufficient serosurvey data for the Dominican Republic are not  
 93 available to determine the predominant dengue virus serotype(s) during each outbreak, but the Pan-  
 94 American Health Organization reports that serotypes 1, 2, and 4 were prevalent in the Caribbean between  
 95 2012 and 2014, and serotypes 2, 3, and 4 were circulating in the region in 2015 (PAHO, 2020b). The other  
 96 two outbreaks were caused by emerging viruses, chikungunya and Zika (Seasons 3 and 5, respectively;  
 97 **Fig. 1b,c**) (Bowman et al., 2018). Those outbreaks were the first and only to be reported of either disease  
 98 in the country, and the number of reported cases in the Caribbean as a whole plummeted in subsequent  
 99 years (Lindsey et al., 2018; Zingman et al., 2017; PAHO, 2020c). After the season dominated by Zika virus  
 100 (Season 5), only a small number of dengue cases were diagnosed (Season 6).

101 Clinical and demographic characteristics differed between reported dengue, chikungunya, and Zika cases.  
 102 The majority of dengue cases experienced fever (98.3%), were hospitalized for their condition (93.7%), and  
 103 were between the ages of 0 and 15 (62.3%), suggesting immunity in the adult population (**Table 1**). Patients  
 104 diagnosed with chikungunya and Zika were older and predominantly female. Rash and/or arthralgia in  
 105 conjunction with fever are typical clinical manifestations of chikungunya and Zika infections (Duffy et al.,  
 106 2009; He et al., 2017; Langsjoen et al., 2016; Martinez et al., 2019; Pineda et al., 2016; Rosario et al., 2015).  
 107 Although lower than rates reported elsewhere, arthralgia rates among chikungunya cases in our dataset  
 108 were consistent with previously documented rates among individuals diagnosed with chikungunya in the  
 109 Dominican Republic (Langsjoen et al., 2016).

110 **Table 1: Epidemiological and clinical characteristics of cases**

Season and Disease	Year	Number of Cases	Median Age of Cases (IQR)	% Hospitalized	% Female	% Reported Fever	% Reported Rash	% Reported Arthralgia
Season 1 (dengue)*	2012-2013	13,666	11 (11)	94.3	44.0	98.7	15.0	31.0
Season 2 (dengue)*	2013-2014	15,079	11 (11)	92.9	46.3	97.4	5.6	16.4
Season 3 (dengue)	2014-2015	5,236	13 (18)	88.7	45.0	96.1	8.0	23.2
Season 4 (dengue)*	2015-2016	19,619	12 (13)	93.9	46.0	98.9	4.0	17.6
Season 5 (dengue)	2016-2017	3,093	15 (25)	85.5	42.4	98.1	7.1	24.9
Season 6 (dengue)	2017-2018	1,437	18 (24)	87.3**	38.5	98.3	4.0	27.4
Chikungunya	2014	6,461	28 (32)	32.0	60.8	89.5	26.0	<b>63.0</b>
Zika	2016	5,161	30 (21)	16.1	73.9	74.2	<b>70.7</b>	8.5

111 IQR: interquartile range

112 \* Outbreak reported

113 \*\* The type of medical attention received by 22.9% of the reported cases during this period was not reported and was assumed to be hospital  
 114 admissions

115

## 116 **Chikungunya and Zika outbreak dynamics did not conform to seasonal dengue patterns**

117 Implementing effective disease prevention and control measures requires knowledge of when their  
 118 etiological viruses are most likely to emerge, re-emerge, and spread (Grubaugh, Ladner, et al., 2019).  
 119 Dengue, chikungunya, and Zika are caused by RNA viruses transmitted by the mosquito vectors *Aedes*  
 120 *aegypti* and *Aedes albopictus*. The abundance and capacity of these vectors are sensitive to climatological  
 121 factors including temperature, rainfall, and humidity (Kraemer et al., 2019; Kraemer, et al., 2015), such that  
 122 transmission of all three viruses has been shown to fluctuate in kind (Do et al., 2014; Hamlet et al., 2018;


123 Obolski et al., 2019; Perez-Guzman et al., 2018; Faria et al., 2016a). We therefore hypothesized that the  
124 timing of the five outbreaks should concord with seasonal weather patterns.

125 We found that the emerging disease outbreaks (chikungunya [Season 3] and Zika [Season 5]) occurred  
126 earlier in the year compared to endemic outbreaks (dengue). The chikungunya and Zika outbreaks peaked  
127 15 and 26 weeks earlier, respectively, than the averaged dengue peak (epidemiological week 41; **Fig. 1b**).  
128 While both chikungunya and dengue cases began to rise around epidemiological week 15, a rise in Zika  
129 cases was observed during the first weeks of the year.

130 Because the implementation of national-level reporting of chikungunya and Zika cases influenced when  
131 these outbreaks were detected (**Fig 1c**, arrows), we calculated the number of weeks elapsed between the  
132 first reported case of each outbreak and the peak number of cases within provinces (**Fig. 1d**). We assumed  
133 that climatological factors did not vary widely between provinces during a given season. Our reasoning  
134 followed that the rate of viral transmission is limited by the extrinsic incubation period (EIP) of the virus in  
135 the mosquito vector (Liu-Helmersson et al., 2014), and the EIP for *Aedes* mosquitoes is mainly influenced  
136 by temperature (Chan & Johansson, 2012; Mordecai et al., 2017; Winokur et al., 2020). Therefore, because  
137 we did not expect transmission rates to vary widely between provinces, provincial outbreaks that peaked  
138 soon after the reporting system was implemented would indicate that transmission in those provinces  
139 preceded the establishment of this system.

140 Our analysis identified one such instance. The majority of provincial chikungunya and Zika outbreaks peaked  
141 within 12 and 25 weeks, and the mean time to peak across provinces did not significantly differ (**Fig. 1d,e**).  
142 This was in contrast to the substantial heterogeneity of the timing of the three dengue outbreaks. The lack  
143 of uniformity across these outbreaks may have been due to a slower rate of spread between regions  
144 because of a pre-existing and spatially heterogeneous herd immunity landscape. Five provinces, four of  
145 which are located in the south-western part of the country, reported peak numbers of Zika cases in January  
146 and February of 2016 (**Fig. 1d**). For the reasons stated above and because Zika virus had been circulating  
147 in the Americas for at least two years before it was reported in the Dominican Republic (Faria et al., 2016,  
148 2017), this observation suggested that Zika virus was already circulating in those provinces before the  
149 national reporting system was implemented at the beginning of January. We hypothesized that early Zika  
150 virus transmission could have occurred in the south-western region because of the region's proximity to  
151 Haiti. In particular, this region shares a border with Haiti, is connected to the Haitian capital Port-au-Prince  
152 via a major roadway, and is home to a large binational market (**Fig. 1e, map, star**). A temporal comparison  
153 of the Zika outbreaks reported in this region, the rest of the country, and Haiti revealed that the regional  
154 outbreak was more consistent with the 2016 Zika outbreak reported in Haiti (**Fig. 1e, inset**). These findings  
155 indicate that the Dominican Republic experienced two geographically and temporally distinct Zika outbreaks.  
156 From reported case counts alone we cannot conclude whether Zika virus was introduced into the Dominican  
157 Republic from Haiti or vice versa; however, it is clear that binational coordination is an essential component  
158 of local mosquito-borne disease control because the viruses these vectors transmit do not recognize  
159 international borders.

160



161


162 **Figure 1: Two outbreaks of emerging disease, chikungunya and Zika, and three endemic dengue outbreaks were reported**  
 163 **between 2012 and 2018 in the Dominican Republic.** (a) The Dominican Republic is a country in the Caribbean that shares the  
 164 island of La Hispaniola with Haiti. (b) Cases in the Dominican Republic reported by the Ministry of Health during each outbreak per  
 165 epidemiological week for five seasons. (c) Weekly reported cases from 2012-2018. Arrows indicate when the first and last cases  
 166 were reported for emerging pathogens. (d) Number of weeks elapsed between the first case reported nationally and peak cases  
 167 reported in 32 provinces (each data point represents a different province). The chikungunya outbreak began during epidemiological  
 168 week (EW) 6, the Zika outbreak during EW 1, and the three dengue outbreaks during EW 14. The mean time to peak for each  
 169 outbreak were compared using an unpaired t test. The qualitative statistical result of the analysis did not change when the five early  
 170 Zika provinces were excluded. (e) Geographic and temporal comparisons of Zika cases reported by four provinces in the south-  
 171 western region of the Dominican Republic. A major roadway connects the Haitian capital Port-au-Prince to the southwestern region  
 172 of the Dominican Republic (grey line, map). A large binational market is located in the province Pedernales (star, map). Similar to  
 173 Haiti, provinces in the southwest experienced an earlier outbreak of Zika compared to the remaining 28 provinces (inset).  
 174

175 **Initial outbreaks of emerging mosquito-borne diseases can occur during periods of sub-optimal**  
 176 **climatological conditions**

177 We considered two possible explanations for the asynchronicity we observed between outbreaks of the  
 178 emerging and endemic viruses (**Fig. 1**). (1) Either climate patterns differed between seasons and the  
 179 emerging outbreaks preceded those of dengue due to seasonal stochasticity, or (2) the spread of emerging

180 viruses was less sensitive to climatological factors compared to dengue virus. By analyzing daily  
181 climatological data collected over the duration of our study period, we found support for the latter (**Fig. 2**).

182 To investigate the relationship between climate and case incidence, we used temperature and humidity time  
183 series data to estimate the mosquito-borne transmission potential throughout our study period (**Fig. 2**). For  
184 this analysis, we used *Index P*, a metric that is calculated by incorporating climate and entomological data  
185 into a Bayesian framework to estimate the transmission potential of individual female mosquitoes (Obolski  
186 et al., 2019). We calculated transmission potential for each week of our study period using data reported in  
187 six of thirty-two provinces. We selected these six provinces based on their representativeness of the  
188 country's sub-climates and data availability. On average, transmission potential fluctuated seasonally, rising  
189 between April and November of each year, coincident with the rainy season, and fell shortly thereafter (**Fig.**  
190 **2**). When we compared the temporal dynamics of transmission potential to reported disease incidence, we  
191 found that the number of emerging disease cases reported weekly peaked before transmission potential  
192 had reached a seasonal maximum for both outbreaks (chikungunya and Zika), while the number of dengue  
193 cases reported weekly peaked after this point for all three dengue outbreaks (**Fig. 2a**). Similar climatological  
194 patterns persisted on the provincial level (**Fig. 2b**). Some provinces experienced clear seasonal fluctuations  
195 in transmission potential, whereas seasonal peaks and troughs were less defined in Distrito Nacional and  
196 María Trinidad Sánchez, which have low to moderately humid sub-climates. Despite this variation, the timing  
197 of provincial outbreaks conformed to the trend observed on the national level: dengue outbreaks peaked  
198 after a period characterized by high transmission potential, just as transmission potential was beginning to  
199 wane; in contrast emerging disease outbreaks occurred concurrently with increasing transmission potential.  
200 Moreover, we cannot attribute climatic anomalies to the decline in reported dengue cases in Seasons 3, 5,  
201 and 6.



202

203

204

205

206

207

208

**Figure 2. Emerging disease outbreaks occurred during periods of sub-optimal climate.** (a) Average weekly transmission potential (*Index P*) and nationally reported cases. *Index P* describes the reproductive potential of an individual female *Aedes* mosquito, meaning its absolute value should be interpreted biologically rather than epidemiologically. Relative fluctuations in *Index P* reflect seasonal changes in the expected rate of mosquito-borne virus transmission. (b) *Index P* and reported cases for six provinces. Temperature and humidity data were retrieved from the National Meteorology Office (ONAMET) database (saip.gob.do) for Distrito Nacional, Barahona, Puerto Plata, María Trinidad Sánchez, and Santiago. Hourly climatological data for Monseñor Nouel


209 were obtained from Open Weather Map. Reported ([openweathermap.org](http://openweathermap.org)) cases of dengue, chikungunya, and Zika were organized  
210 by province based on the individual's place of residence.

211

## 212 **The sizes of the susceptible human population influenced the speed at which emerging viruses** 213 **spread**

214 Variability in seasonal weather patterns and vectorial capacity did not account for differences in the timing  
215 of emerging disease outbreaks (**Fig. 2**). Rather, we hypothesized that the patterns observed for dengue  
216 were likely to have been influenced by a pre-existing and spatially heterogeneous herd immunity landscape.  
217 This would mandate that transmission potential must remain high for an extended period before epidemic  
218 growth is achieved each season.

219 To test the hypothesis that local susceptibility influenced the timing of the outbreaks, we examined the  
220 relationship between the size of the susceptible human population and each outbreak's doubling time (**Fig.**  
221 **3**). We estimated that the median estimated effective reproduction number ( $R_{\text{eff}}$ ) for the three dengue  
222 outbreaks was between 1.3 and 1.4, while those of the chikungunya and Zika outbreaks ranged from 1.6  
223 to 2.45 (**Fig. 3a**). These findings are consistent with the epidemiology of the three diseases and previous  
224 estimates of  $R_{\text{eff}}$  for chikungunya and Zika (Bowman et al., 2018; Cauchemez et al., 2014; Nishiura et al.,  
225 2016). Published estimates of the basic reproduction number ( $R_0$ ) for dengue, equivalent to  $R_{\text{eff}}$  in a wholly  
226 susceptible population, vary widely (0.97-65) due to the parameter's sensitivity to the size of the local  
227 susceptible population (Liu et al., 2020). We observed a slightly longer doubling time (smaller  $R_{\text{eff}}$ ) during  
228 the third dengue outbreak, consistent with increasing levels of herd immunity from the two previous  
229 outbreaks. In our dataset, reported dengue cases were primarily in the 0-15 age group, indicating older age  
230 groups had high levels of pre-existing immunity and were not susceptible to disease (**Table 1; Fig. 3b**). In  
231 contrast, we speculated that the entire population was susceptible to chikungunya and Zika, allowing these  
232 viruses to spread despite sub-optimal weather conditions, facilitating an earlier outbreak peak, and affecting  
233 a much wider range of ages (**Table 1; Fig. 3b**).



234

235

236

237

238

239

**Figure 3. The size of the susceptible population was larger for emerging diseases than dengue.** (a) The effective reproduction number ( $R_{eff}$ ) of chikungunya and Zika far exceeded that of dengue.  $R_{eff}$  was calculated using the growth rate of each outbreak (Lipsitch et al., 2003) and estimates of the incubation period and serial interval of each disease (Table S2). (b) The median age of reported cases during emerging disease outbreaks significantly exceeded that of the three dengue outbreaks. The median age group of the national population is 20-25 as calculated from population data reported in the 2010 census.

240

241

**Spatial distribution of dengue attack rates is a poor indicator for initial outbreaks of emerging disease**

242

243

244

245

246

Given that the timing of emerging disease outbreaks (chikungunya and Zika) did not conform to that of dengue outbreaks (Fig. 1,2), we suspected that the spatial distribution of dengue cases would be equally uninformative for chikungunya and Zika outbreaks. Specifically, we hypothesized that the relative burden of dengue within provinces during individual outbreaks would correlate with the burden of dengue during subsequent outbreaks but not with the relative burden of emerging disease cases.

247

248

249

250

To address this question, we measured the age- and sex-adjusted attack rates by province for each of the outbreaks. We found that the attack rates for individual provinces across outbreaks were well correlated between dengue outbreaks and between chikungunya and Zika outbreaks, but not across the endemic and emerging viruses (Fig. 4a, b).

251

252

253

254

255

256


257

258

Next, we investigated the role of climate and land-use in perpetuating this trend, reasoning that larger mosquito populations would facilitate higher attack rates (Kraemer, et al., 2015; Rodriguez-Barraquer, Salje, et al., 2019). To this end, we compared an *Aedes aegypti* suitability score (AaS; Fig. 4c), a metric that incorporates ecological variables not included in our transmission potential estimates such as vegetation levels and precipitation, to population-adjusted attack rates of disease (Fig. S1, Fig. 4a). Also unlike our transmission potential estimates, AaS is an aggregated estimate of suitability per month in any given year. Interestingly, AaS was not well correlated with the adjusted provincial attack rates or with mean age of infection (Fig. S1). Moreover, provinces in which the highest levels of mosquito abundance were predicted

259 did not consistently report the highest burden of disease (**Fig. 4a,c**). One reason for these inconsistencies  
 260 is that AaS relies on historical climate data from 1970 to 2000 to estimate mosquito abundance. Therefore,  
 261 our estimates do not capture anomalous climatological events such as the hurricane that made landfall in  
 262 the Dominican Republic in October 2012. AaS also assumes a high level of stability in land use and minimal  
 263 urbanization since the thirty year period from which the estimates were made. Regardless of the true  
 264 underlying reason for these discrepancies, the spatial distributions between endemic (dengue) and  
 265 emerging (chikungunya and Zika) mosquito-borne virus diseases are not very well correlated.

266



267

268 **Figure 4: The spatial distribution of dengue cases is not well correlated with that of emerging disease cases or with**  
 269 **estimated mosquito abundance. (a)** Population-, age-, and sex-adjusted attack rates were significantly correlated within provinces  
 270 across dengue outbreaks and between Zika and chikungunya outbreaks. One province, Elias Piña, was excluded because of  
 271 assumed substantial under-reporting: the province reported 1 dengue case during Season 1 and 1 chikungunya case in a population  
 272 of 63,250. **(b)** R<sup>2</sup> values for dengue vs dengue (D vs D), dengue vs emerging (D vs E), and emerging vs emerging (E vs E)  
 273 by linear regression. **(c)** Mosquito abundance (AaS) was normalized across provinces.


274 **Clinical outcomes of dengue cases and trends in febrile illness cases are indicators of disease**  
275 **prevalence**

276 Up to this point, our findings demonstrate that emerging mosquito-borne disease outbreaks are not confined  
277 to the temporal and spatial patterns of endemic disease outbreaks in the Dominican Republic (**Figs. 1-4**).  
278 While similar inconsistencies in outbreak dynamics have been observed elsewhere in the Americas (**Fig.**  
279 **S2**) (Lourenço et al., 2017; Faria et al., 2016a), it is difficult to discern if our findings are due to differences  
280 in epidemiology or in underlying reporting biases. Therefore, we used general patterns of febrile illness and  
281 clinical characteristics to show that reporting biases did not influence our previous conclusions (**Fig. 5**).

282 Because the vast majority of cases in our dataset reported fever independent of disease (**Table 1**), we  
283 hypothesized that febrile illness incidence should reflect disease incidence reported during our study period.  
284 When we examined the number of febrile illness cases reported in the Dominican Republic per  
285 epidemiological week per season, we found that temporal trends in febrile illness cases in Season 3 (2014-  
286 2015) and Season 5 (2016-2017; **Fig. 5a**) were consistent with those we observed in our chikungunya and  
287 Zika case data, respectively (**Fig. 1a**). In Season 4 (2015-2016, dengue), febrile illness cases peaked in  
288 week 44, coincident with the peak of the dengue outbreak (**Fig. 1a**). To ensure that the spatial relationships  
289 we identified in **Figure 4** were not purely the product of differential reporting practices, we compared  
290 population-adjusted attack rates of mosquito-borne disease and febrile illness reported by individual  
291 provinces during each season and found no significant correlation (**Fig. S3**). Equally pronounced seasonal  
292 peaks of reported febrile illness were not observed in Season 1 and Season 2 despite the known dengue  
293 outbreaks that occurred within that period. We speculate that changes made in reporting protocols for febrile  
294 illness cases during the chikungunya outbreak in Season 3 prompted a sustained increase in national  
295 reporting thereafter (Ministerio de Salud Pública, 2014a).

296 We then investigated why reported febrile illness cases peaked seasonally in Seasons 5 and 6 despite an  
297 apparent decline in coincident dengue cases by comparing case-fatality rates (CFRs) of dengue cases  
298 across seasons. Specifically, we considered the possibility that this decline was due to under-reporting of  
299 dengue cases during the interim periods between dengue outbreaks (Seasons 3, 5, and 6). The average  
300 CFR during these seasons was 1.87% (range: 1.60%-2.40%), while the average CFR of reported cases  
301 during the three large dengue outbreaks in Seasons 1, 2, and 4 was 0.8% (range: 0.48%-1.20%) (**Fig. 5b**).  
302 A wide range of dengue CFRs is reported in the literature, with a mean of 1.62% (95% CI: 0.64–4.02%)  
303 (Low et al., 2018), and without serosurveillance data, we are unable to rule out the possibility that the  
304 circulation of different dengue serotypes resulted in differences in seasonal CFRs. However, if we assume  
305 that fatality among cases during the third dengue outbreak (Season 4) were well reported, dengue cases  
306 were under-reported by 33% in Season 3 and Season 5, and by 2-fold in Season 6. These are likely  
307 conservative estimates as the current surveillance system is passive and therefore does not capture  
308 asymptomatic cases. As a result, Seasons 3, 5, and 6 may represent periods characterized by a large  
309 number of mild cases that were not captured by the surveillance system. It is also likely that the number of  
310 deaths due to chikungunya was substantially under-reported; however, our calculation of 0.19%  
311 corresponds to a corrected, post-hoc estimate of this rate (0.15%) for the Dominican Republic (Freitas et  
312 al., 2018).

313



314  
315 **Figure 5. Reported febrile illness cases and case fatality rates of dengue cases during inter-dengue-outbreak periods**  
316 **exceeded those reported during dengue outbreaks.** (a) Temporal trends in weekly febrile illness cases reflect temporal trends  
317 in reported dengue, chikungunya, and Zika cases. Acute febrile illness cases reported by epidemiological week per municipality  
318 were obtained from the National Statistics Directorate of the Ministry of Health (saip.gob.do). Weekly cases were aggregated to  
319 the national level. Chikungunya and Zika outbreak peaks are denoted by the gray box. (b) Percent case fatality rates (CFR) for  
320 dengue cases rose during seasons in which no dengue outbreak was reported. CFRs were calculated for six seasons of reported  
321 dengue cases (S1-S6) and two seasons of emerging disease cases (chikungunya (C) and Zika (Z)).  
322

## 323 DISCUSSION

324 Our study demonstrates that, even when transmitted by the same mosquito vector, viruses are not beholden  
325 to the same temporal and spatial outbreak dynamics. Instead, when and where the new virus is introduced,  
326 the size of the susceptible human population, and the capacity of local surveillance systems determine these  
327 dynamics are detected. In short, dengue epidemiology cannot be used to anticipate the location and timing  
328 of future emerging mosquito-borne disease outbreaks in the Dominican Republic, and likely in other  
329 Caribbean countries and territories. Instead, consistent and sustainable surveillance methods should be  
330 implemented to limit disease and prevent future outbreaks. These methods could include serosurveillance  
331 of the population during periods between outbreaks (Imai et al., 2015), testing local mosquito vectors for  
332 viral infections (Fauver et al., 2017, 2018; Grubaugh et al., 2015), and monitoring health outcomes of  
333 travelers who visit the country (Grubaugh, et al., 2019).

334 Maintaining a sustainable surveillance system is critical for preventing the silent transmission of viruses that  
335 can fuel large outbreaks. Other countries in the Americas reported subsequent outbreaks of chikungunya  
336 and Zika after their initial outbreaks. We cannot conclusively determine whether the Dominican Republic  
337 experienced a similar pattern because surveillance data for chikungunya and Zika are not available for  
338 seasons following the initial outbreaks of these diseases. Elucidating whether dengue, chikungunya, and  
339 Zika are co-circulating in the country will be critical for triaging and providing appropriate clinical care to  
340 patients who present with febrile illness (Vogels et al., 2019), especially if chikungunya and Zika virus  
341 transmission is now in sync with dengue transmission (Bisanzio et al., 2018; L. P. Freitas et al., 2019).

342 Equally important is the identification of viruses that could spread across international borders. In our study,  
343 we observed an epidemiological link between Zika outbreaks in Haiti and the south-west region of the  
344 Dominican Republic. This connection likely extends to other mosquito-borne viruses such as Mayaro virus  
345 and West Nile virus, which have been reported in Haiti (Blohm et al., 2019; Lednicky et al., 2016;  
346 Weppelmann et al., 2017). The former can be transmitted by *Aedes* mosquitoes (Kantor et al., 2019; Long  
347 et al., 2011; Pereira et al., 2020; Wiggins et al., 2018), and the latter has been isolated from zoonotic  
348 reservoirs in the Dominican Republic (Komar et al., 2005). Similarly, reports of international travelers  
349 carrying mosquito-borne viruses from the Dominican Republic back to their home countries demonstrate  
350 that gaps in surveillance have global implications (Barzon et al., 2016; Díaz-Menéndez et al., 2018; Duijster  
351 et al., 2016; Goncé et al., 2018; Millman et al., 2016; Requena-Méndez et al., 2014). Therefore,

352 understanding the transmission patterns of viruses and developing a unified, international plan to combat  
353 them before they cause an outbreak will help mitigate the potential for such an event.

354 Our findings suggest that dengue cases were under-reported following the Zika outbreak in 2016; however,  
355 there are a number of possible explanations for the ostensible decline in dengue cases that should be  
356 explored. Wide-spread mosquito control measures motivated by the Zika outbreak could have limited the  
357 spread of dengue later that year. Although plausible, this line of reasoning does not explain why a similar  
358 post-Zika decline was observed in other countries in the Americas (Perez et al., 2019), nor why there was a  
359 resurgence of dengue cases in 2019. A second explanation is that Zika infections confer some level of  
360 temporary immunity to subsequent dengue infections (Gordon et al., 2019; Ribeiro et al., 2018, 2020;  
361 Rodriguez-Barraquer, Costa, et al., 2019). This theory cannot account for the small number of dengue cases  
362 reported in 2014 following the chikungunya outbreak, as the etiological agent of that disease is an  
363 alphavirus, and it assumes very high attack rates and extensive under-reporting of Zika in 2016 to have  
364 achieved sufficient levels of herd immunity. Our findings instead suggest that prior Zika infection protects  
365 against symptomatic dengue infections because we observed significant positive correlation between  
366 dengue attack rates within provinces across outbreaks. Given that cross-reacting immunity between dengue  
367 serotypes is well documented (Katzelnick et al., 2016; Tsang et al., 2019), our data suggest that a similar  
368 relationship between dengue and Zika would not result in a wide-spread decline of cases. Rather, dengue  
369 transmission could have reasonably persisted undetected if most of those infected were not hospitalized. If  
370 true, this hypothesis would explain why seasonal peaks in reported febrile illness cases persisted in 2017  
371 and 2018, and why CFRs among reported dengue cases appeared to be elevated in Seasons 5 and 6 if the  
372 true number of cases was under-reported. To better understand these complex interactions, the collection  
373 of serotype information should be incorporated into current dengue surveillance efforts.

374 There are a few important limitations to our study. First, our dataset included chikungunya and Zika case  
375 data from the initial wave of each disease, and we cannot therefore compare temporal and spatial dynamics  
376 of these diseases across seasons. After these initial outbreaks, diagnostic testing for these diseases has  
377 largely ceased. While the number of cases of these diseases reported in the Dominican Republic has  
378 declined to zero the true burden of disease is unknown. Future studies should investigate whether these  
379 viruses have continued to circulate undetected in the country and whether their spatiotemporal dynamics  
380 have since synchronized with that of dengue virus. Second, the reporting system for suspected chikungunya  
381 cases differed from that used for suspected Zika cases. During the chikungunya outbreak, most febrile  
382 illnesses cases without apparent cause were initially classified as suspected chikungunya cases. For this  
383 reason, the number of cases reported by the Pan-American Health Organization (PAHO) and the Ministry  
384 of Health was significantly larger than those which we have reported here (Ministerio de Salud Pública,  
385 2014b). Our chikungunya case data contains a disproportionate number of children in the <1 year age group,  
386 indicating that the dissemination of diagnostic testing may have been skewed towards high-risk groups.  
387 Third, our findings demonstrate that an epidemiological relationship existed between the Dominican  
388 Republic and Haiti during the Zika epidemic in 2016, but we cannot determine the directionality of cross-  
389 border virus movement without virus genomic data. However, given that this relationship exists and that  
390 mosquitoes do not recognize political boundaries, it can be assumed that bi-directional spillover of mosquito-  
391 borne diseases will occur in the future unless appropriate bi-national surveillance and control measures are  
392 implemented. Finally, our analysis primarily focused on virus transmission by *Aedes aegypti* mosquitoes,  
393 but it is possible that other mosquito vectors contributed to the propagation of the outbreaks we investigated.  
394 Specifically, *Aedes albopictus* may have played a key role in chikungunya transmission (Lamballerie et al.,  
395 2008; Vazeille et al., 2007). Limited data are available on the distribution of relevant mosquito vectors in the  
396 country (González et al., 2019, 2020), and broader entomological surveillance is needed to better address  
397 this question.

398 Taken together, our study demonstrates that surveillance for mosquito-borne diseases should be sustained  
399 during periods when transmission appears to be low because patterns in reported dengue cases are poor  
400 indicators of future emerging mosquito-borne virus outbreak dynamics. Reported symptoms and case  
401 demographics may be useful for identifying shifts in disease prevalence, but many clinical features,  
402 especially fever, that we have noted are likely a function of the reporting and diagnostic algorithms used  
403 during an outbreak. Active reporting of new dengue, chikungunya, and Zika cases and the broader  
404 deployment of diagnostics for newly emerged diseases are needed to ascertain more accurate case profiles.  
405 Outbreaks of emerging tropical diseases are a threat to the public health of the Caribbean, and endemic  
406 diseases such as dengue precipitate health crises with increasing frequency. Given the pervasiveness of  
407 mosquito-borne diseases in tropical climates, sustainable surveillance systems rather than reactionary  
408 disease control measures should be implemented to prevent future crises.

## 409 410 **MATERIALS AND METHODS**

### 411 412 *Description of data*

413 Data for suspected and confirmed cases of dengue, chikungunya, and Zika reported between 2012-2018  
414 were extracted from the Dominican Republic Ministry of Health Weekly Reports ([digepi.gob.do](http://digepi.gob.do)).  
415 Depersonalized demographic and clinical characteristics of cases were solicited from the National Statistics  
416 Directorate of the Ministry of Health (MoH) ([saip.gob.do](http://saip.gob.do)). Data were organized by date of onset of symptom  
417 report.

418 The National Epidemiology Directorate of the MOH collects dengue case reports by passive surveillance in  
419 a weekly manner in its Epi 1 form within their Sistema Nacional de Vigilancia Epidemiológica (SINAVE)  
420 Digital Platform for every healthcare setting across the country. These data were collected in weekly reports  
421 submitted to their website ([digepi.gob.do](http://digepi.gob.do)) and organized by province and week of reported cases. The data  
422 collected by the MoH includes age, sex, province and municipality of residence, date of symptom onset,  
423 clinical outcome, and symptomatology. The MoH included mandatory reports from suspected and confirmed  
424 chikungunya infection through 2014 and Zika virus throughout 2016, without continuing to do so thereafter.

425 Province- and municipality-level weekly acute febrile illness data from 2012-2018 were solicited from the  
426 National Epidemiology Directorate of the Ministry of Health ([saip.gob.do](http://saip.gob.do)), which collects the data as part of  
427 a passive surveillance system. In 2014, the Ministry of Health increased its efforts to identify febrile illness  
428 cases (Ministerio de Salud Pública, 2014a).

429 Daily climatological variables for mean temperature and relative humidity for 5 cities (Santo Domingo,  
430 Distrito Nacional; Barahona, Barahona; La Union, Puerto Plata; Cabrera, María Trinidad Sánchez; Santiago,  
431 Santiago) was collected from the National Meteorology Office (ONAMET) database ([saip.gob.do](http://saip.gob.do)) from  
432 January 2012 through December 2018. Hourly climatological data for Bonao, Monseñor Nouel for the same  
433 timeframe were obtained from [openweathermap.org](http://openweathermap.org).

434 Population data were extracted from the 9th National Population and Household Census. This census was  
435 conducted in 2010 by the National Statistics Office (National Statistics Office, 2012).

436  
437 Reported cases of dengue, chikungunya, and Zika for Haiti, Bolivia, Jamaica, and Venezuela were extracted  
438 from the Pan-American Health Organization website and PLISA database (PAHO, 2020d). Dengue and Zika  
439 cases, which were reported by epidemiological week, were aggregated by month to allow for a direct  
440 comparison to chikungunya cases, which were reported by month.

### 441 442 *Time to outbreak peak*

443 We calculated the time to peak for provincial outbreaks by first identifying the epidemiological week in which  
444 the first case of the national outbreak was reported (chikungunya: EW 6, Zika: EW 1, dengue (3): EW 14)  
445 and then counting the number of weeks elapsed until each of the 32 provinces reported the maximum  
446 number of cases for the corresponding outbreak. The mean time to peak for each outbreak was compared  
447 using an unpaired T test implemented in Prism v8.4.2.

#### 448 *Transmission potential*

449 We calculated weekly transmission potential (*Index P*) with the Bayesian approach developed by Obolski et  
450 al. (Obolski et al., 2019). Briefly, we extracted daily average temperature and relative humidity for 6 cities  
451 as described above. Rarely, temperature or humidity were not available for a given day. In these cases, we  
452 averaged the respective variable from the same date across the remaining 6 years. We did this for 14 days  
453 for Barahona and Puerto Plata, 3 days for Santiago, and 1 day for María Trinidad Sánchez.

454 We then used the R package MVSE and entomological and epidemiological priors documented in the  
455 literature (**Table S1**) to calculate daily transmission potential, which we then aggregated by week (Lourenço,  
456 2019). We found that the model was reasonably robust to a range of priors for each of the parameters and  
457 therefore elected to use short human incubation and infectious period estimates to inform the model. We  
458 adopted our prior estimate of human life expectancy to the Dominican Republic based on estimates by the  
459 United Nations Development Program (UNDP, 2019), but otherwise used the same priors as reported by  
460 Oboloski et al. (**Table S1**) (Obolski et al., 2019).

461

#### 462 *Effective reproduction number ( $R_{eff}$ ) estimates*

463 We estimated  $R_{eff}$  for the five outbreaks using the method developed by Lipsitch et al. (Lipsitch et al., 2003)  
464 to fit a linear regression model to our case data (**Eq. 1**).

465

$$466 R_{eff} = r^2 * (1 - f) * f * v^2 + r * v + 1 \quad \text{Eq. 1}$$

467

468 We used a range of values for latent and infectious periods that were well reported in the literature to  
469 calculate  $f$ , the proportion of the serial interval in the latent period, and  $v$ , the serial interval (**Table S2**). We  
470 obtained  $r$ , the epidemic growth rate, by extracting the slope of our model after fitting it to our case data. All  
471 calculations were done using R v4.0.0.

472

#### 473 *Adjusted attack rates and linear regression*

474 Province-level attack rates by outbreak were age- and sex-adjusted using the direct standardization method,  
475 with the national population as the reference population.

476

477 We calculated Pearson's R correlation coefficient for province-level attack rates between various pairs of  
478 outbreaks (**Fig. 4a,b**). An outlier analysis showed that the removal of outliers did not substantially affect the  
479 size or significance of the correlation coefficients, so we included all data points.

480

#### 481 *Aedes aegypti* suitability score (*AaS*)

482 *AaS* was calculated for each month using both climate and land-use data on a 5km x 5 km scale collected  
483 from 1970 and 2000 and compiled by WorldClim as described by Kraemer et al. (Kraemer, et al., 2015).  
484 Monthly suitability scores were extracted and averaged by province in Rv4.0.0.

485

## 486 **ACKNOWLEDGEMENTS**

487 We thank C. Vogels, A. Brito, J. Fauver, C. Kalinich, I. Ott, S. Lapidus, K. Gangavarapu, J. Pack, and S.  
488 Taylor for feedback and/or assistance with the methodology. The study was funded in part by the Hecht  
489 Global Health Faculty Network Award provided to N.D.G. from the Yale Institute of Global Health.

490

## 491 **COMPETING INTERESTS**

492 The authors declare no competing interests.

493

## 494 **REFERENCES**

- 495 Aldstadt, J., Yoon, I.-K., Tannitisupawong, D., Jarman, R. G., Thomas, S. J., Gibbons, R. V., Uppapong, A., Iamsirithaworn, S.,  
496 Rothman, A. L., Scott, T. W., & Others. (2012). Space-time analysis of hospitalised dengue patients in rural Thailand reveals  
497 important temporal intervals in the pattern of dengue virus transmission. *Tropical Medicine & International Health: TM & IH*,  
498 17(9), 1076–1085. <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-3156.2012.03040.x>
- 499 Barzon, L., Pacenti, M., Berto, A., Sinigaglia, A., Franchin, E., Lavezzo, E., Brugnaro, P., & Palù, G. (2016). Isolation of infectious  
500 Zika virus from saliva and prolonged viral RNA shedding in a traveller returning from the Dominican Republic to Italy, January  
501 2016. *Euro Surveillace: Bulletin Europeen Sur Les Maladies Transmissibles = European Communicable Disease Bulletin*,  
502 21(10), 30159. <https://doi.org/10.2807/1560-7917.ES.2016.21.10.30159>
- 503 Bisanzio, D., Dzul-Manzanilla, F., Gomez-Dantés, H., Pavia-Ruz, N., Hladish, T. J., Lenhart, A., Palacio-Vargas, J., González  
504 Roldan, J. F., Correa-Morales, F., Sánchez-Tejeda, G., Kuri Morales, P., Manrique-Saide, P., Longini, I. M., Halloran, M. E., &  
505 Vazquez-Prokopec, G. M. (2018). Spatio-temporal coherence of dengue, chikungunya and Zika outbreaks in Merida, Mexico.  
506 *PLoS Neglected Tropical Diseases*, 12(3), e0006298. <https://doi.org/10.1371/journal.pntd.0006298>
- 507 Blohm, G., Elbadry, M. A., Mavian, C., Stephenson, C., Loeb, J., White, S., Telisma, T., Chavannes, S., Beau De Rochar, V. M.,  
508 Salemi, M., Lednicky, J. A., & Morris, J. G., Jr. (2019). Mayaro as a Caribbean traveler: Evidence for multiple introductions  
509 and transmission of the virus into Haiti. *International Journal of Infectious Diseases: IJID: Official Publication of the*  
510 *International Society for Infectious Diseases*, 87, 151–153. <https://doi.org/10.1016/j.ijid.2019.07.031>
- 511 Bowman, L. R., Rocklöv, J., Kroeger, A., Olliaro, P., & Skewes, R. (2018). A comparison of Zika and dengue outbreaks using  
512 national surveillance data in the Dominican Republic. *PLoS Neglected Tropical Diseases*, 12(11), e0006876.  
513 <https://doi.org/10.1371/journal.pntd.0006876>
- 514 Cao-Lormeau, V.-M. (2016). Tropical Islands as New Hubs for Emerging Arboviruses. *Emerging Infectious Diseases*, 22(5), 913–  
515 915. <https://doi.org/10.3201/eid2205.150547>
- 516 Cassadou, S., Boucau, S., Petit-Sinturel, M., Huc, P., Leparç-Goffart, I., & Ledrans, M. (2014). Emergence of chikungunya fever  
517 on the French side of Saint Martin island, October to December 2013. *Euro Surveillace: Bulletin Europeen Sur Les Maladies*  
518 *Transmissibles = European Communicable Disease Bulletin*, 19(13). <https://doi.org/10.2807/1560-7917.es2014.19.13.20752>
- 519 Cauchemez, S., Ledrans, M., Poletto, C., Quenel, P., de Valk, H., Colizza, V., & Boëlle, P. Y. (2014). Local and regional spread of  
520 chikungunya fever in the Americas. *Euro Surveillace: Bulletin Europeen Sur Les Maladies Transmissibles = European*  
521 *Communicable Disease Bulletin*, 19(28), 20854. <https://doi.org/10.2807/1560-7917.es2014.19.28.20854>
- 522 Chan, M., & Johansson, M. A. (2012). The incubation periods of Dengue viruses. *PloS One*, 7(11), e50972.  
523 <https://doi.org/10.1371/journal.pone.0050972>
- 524 Díaz-Menéndez, M., de la Calle-Prieto, F., Montero, D., Antolín, E., Vazquez, A., Arsuaga, M., Trigo, E., García-Bujalance, S., de  
525 la Calle, M., Seco, P. S., de Ory, F., & Arribas, J. R. (2018). Initial experience with imported Zika virus infection in Spain. In  
526 *Enfermedades Infecciosas y Microbiología Clínica* (Vol. 36, Issue 1, pp. 4–8). <https://doi.org/10.1016/j.eimc.2016.08.003>
- 527 Dorléans, F., Hoen, B., Najioullah, F., Herrmann-Storck, C., Schepers, K. M., Abel, S., Lamaury, I., Fagour, L., Césaire, R.,  
528 Guyomard, S., Troudard, R., Adélaïde, Y., Romagne, M.-J., Davidas, M., Rochais, S., Boa, S., de Saint-Alary, F., Preira, A.,  
529 Saint-Martin, P., ... Cabié, A. (2018). Outbreak of Chikungunya in the French Caribbean Islands of Martinique and  
530 Guadeloupe: Findings from a Hospital-Based Surveillance System (2013–2015). *The American Journal of Tropical Medicine*  
531 *and Hygiene*, 98(6), 1819–1825. <https://doi.org/10.4269/ajtmh.16-0719>
- 532 Do, T. T. T., Martens, P., Luu, N. H., Wright, P., & Choisy, M. (2014). Climatic-driven seasonality of emerging dengue fever in  
533 Hanoi, Vietnam. *BMC Public Health*, 14, 1078. <https://doi.org/10.1186/1471-2458-14-1078>
- 534 Duffy, M. R., Chen, T.-H., Hancock, W. T., Powers, A. M., Kool, J. L., Lanciotti, R. S., Pretrick, M., Marfel, M., Holzbauer, S.,  
535 Dubray, C., Guillaumot, L., Griggs, A., Bel, M., Lambert, A. J., Laven, J., Kosoy, O., Panella, A., Biggerstaff, B. J., Fischer, M.,  
536 & Hayes, E. B. (2009). Zika virus outbreak on Yap Island, Federated States of Micronesia. *The New England Journal of*  
537 *Medicine*, 360(24), 2536–2543. <https://doi.org/10.1056/NEJMoa0805715>
- 538 Duijster, J. W., Goorhuis, A., van Genderen, P. J. J., Visser, L. G., Koopmans, M. P., Reimerink, J. H., Grobusch, M. P., van der  
539 Eijk, A. A., van den Kerkhof, J. H. C. T., Reusken, C. B., Hahné, S. J. M., & The Dutch ZIKV study team. (2016). Zika virus  
540 infection in 18 travellers returning from Surinam and the Dominican Republic, The Netherlands, November 2015–March 2016.  
541 *Infection*, 44(6), 797–802. <https://doi.org/10.1007/s15010-016-0906-y>

- 542 Faria, N.R., Lourenço, J., Marques de Cerqueira, E., Maia de Lima, M., Pybus, O., & Carlos Junior Alcantara, L. (2016).  
543 Epidemiology of Chikungunya Virus in Bahia, Brazil, 2014-2015. *PLoS Currents*, 8.  
544 <https://doi.org/10.1371/currents.outbreaks.c97507e3e48efb946401755d468c28b2>
- 545 Faria, N. R., Azevedo, R. do S. da S., Kraemer, M. U. G., Souza, R., Cunha, M. S., Hill, S. C., Thézé, J., Bonsall, M. B., Bowden,  
546 T. A., Rissanen, I., Rocco, I. M., Nogueira, J. S., Maeda, A. Y., Vasami, F. G. da S., Macedo, F. L. de L., Suzuki, A.,  
547 Rodrigues, S. G., Cruz, A. C. R., Nunes, B. T., ... Vasconcelos, P. F. C. (2016). Zika virus in the Americas: Early  
548 epidemiological and genetic findings. *Science*, 352(6283), 345–349. <https://doi.org/10.1126/science.aaf5036>
- 549 Faria, N. R., Quick, J., Claro, I. M., Thézé, J., de Jesus, J. G., Giovanetti, M., Kraemer, M. U. G., Hill, S. C., Black, A., da Costa, A.  
550 C., Franco, L. C., Silva, S. P., Wu, C.-H., Raghwan, J., Cauchemez, S., du Plessis, L., Verotti, M. P., de Oliveira, W. K.,  
551 Carmo, E. H., ... Pybus, O. G. (2017). Establishment and cryptic transmission of Zika virus in Brazil and the Americas.  
552 *Nature*, 546(7658), 406–410. <https://doi.org/10.1038/nature22401>
- 553 Fauver, J. R., Gendernalik, A., Weger-Lucarelli, J., Grubaugh, N. D., Brackney, D. E., Foy, B. D., & Ebel, G. D. (2017). The Use of  
554 Xenosurveillance to Detect Human Bacteria, Parasites, and Viruses in Mosquito Bloodmeals. *The American Journal of*  
555 *Tropical Medicine and Hygiene*, 97(2), 324–329. <https://doi.org/10.4269/ajtmh.17-0063>
- 556 Fauver, J. R., Weger-Lucarelli, J., Fakoli, L. S., 3rd, Bolay, K., Bolay, F. K., Diclario, J. W., 2nd, Brackney, D. E., Foy, B. D.,  
557 Stenglein, M. D., & Ebel, G. D. (2018). Xenosurveillance reflects traditional sampling techniques for the identification of  
558 human pathogens: A comparative study in West Africa. *PLoS Neglected Tropical Diseases*, 12(3), e0006348.  
559 <https://doi.org/10.1371/journal.pntd.0006348>
- 560 Ferguson, N. M., Cucunubá, Z. M., Dorigatti, I., Nedjati-Gilani, G. L., Donnelly, C. A., Basáñez, M.-G., Nouvellet, P., & Lessler, J.  
561 (2016). EPIDEMIOLOGY. Countering the Zika epidemic in Latin America. *Science*, 353(6297), 353–354.  
562 <https://doi.org/10.1126/science.aag0219>
- 563 Freitas, A. R. R., Alarcón-Elbal, P. M., Paulino-Ramírez, R., & Donalisio, M. R. (2018). Excess mortality profile during the Asian  
564 genotype chikungunya epidemic in the Dominican Republic, 2014. *Transactions of the Royal Society of Tropical Medicine and*  
565 *Hygiene*, 112(10), 443–449. <https://doi.org/10.1093/trstmh/try072>
- 566 Freitas, L. P., Cruz, O. G., Lowe, R., & Sá Carvalho, M. (2019). Space–time dynamics of a triple epidemic: dengue, chikungunya  
567 and Zika clusters in the city of Rio de Janeiro. *Proceedings of the Royal Society B: Biological Sciences*, 286(1912),  
568 20191867. <https://doi.org/10.1098/rspb.2019.1867>
- 569 Ganjian, N., & Riviere-Cinamond, A. (2020). Mayaro virus in Latin America and the Caribbean. *Revista Panamericana de Salud*  
570 *Publica = Pan American Journal of Public Health*, 44, e14. <https://doi.org/10.26633/RPSP.2020.14>
- 571 Goncé, A., Martínez, M. J., Marbán-Castro, E., Saco, A., Soler, A., Alvarez-Mora, M. I., Peiro, A., Gonzalo, V., Hale, G.,  
572 Bhatnagar, J., López, M., Zaki, S., Ordi, J., & Bardají, A. (2018). Spontaneous Abortion Associated with Zika Virus Infection  
573 and Persistent Viremia. *Emerging Infectious Diseases*, 24(5), 933–935. <https://doi.org/10.3201/eid2405.171479>
- 574 González, M. A., Rodríguez, M. A., Vásquez, Y. E., del Carmen Rosario, E., Durán, J. C., & Alarcón-Elbal, P. M. (2020). A survey  
575 of tire-breeding mosquitoes (Diptera: Culicidae) in the Dominican Republic: considerations about a pressing issue.  
576 *Biomédica*, 40(3). <https://revistabiomedica.org/index.php/biomedica/article/view/5200>
- 577 González, M. A., Sosa, M. A. R., Bautista, Y. E. V., Fernández, L. D., de Prada, M. B., Guerrero, K. A., & Alarcón-Elbal, P. M.  
578 (2019). Micro-environmental features associated to container-dwelling mosquitoes (Diptera: Culicidae) in an urban cemetery  
579 of the Dominican Republic. In *Revista de Biología Tropical* (Vol. 67, Issue 1). <https://doi.org/10.15517/rbt.v67i1.33158>
- 580 Gordon, A., Gresh, L., Ojeda, S., Katzelnick, L. C., Sanchez, N., Mercado, J. C., Chowell, G., Lopez, B., Elizondo, D., Coloma, J.,  
581 Burger-Calderon, R., Kuan, G., Balmaseda, A., & Harris, E. (2019). Prior dengue virus infection and risk of Zika: A pediatric  
582 cohort in Nicaragua. *PLoS Medicine*, 16(1), e1002726. <https://doi.org/10.1371/journal.pmed.1002726>
- 583 Grubaugh, N. D., Faria, N. R., Andersen, K. G., & Pybus, O. G. (2018). Genomic Insights into Zika Virus Emergence and Spread.  
584 *Cell*, 172(6), 1160–1162. <https://doi.org/10.1016/j.cell.2018.02.027>
- 585 Grubaugh, N. D., Ladner, J. T., Lemey, P., Pybus, O. G., Rambaut, A., Holmes, E. C., & Andersen, K. G. (2019). Tracking virus  
586 outbreaks in the twenty-first century. *Nature Microbiology*, 4(1), 10–19. <https://doi.org/10.1038/s41564-018-0296-2>
- 587 Grubaugh, N. D., Saraf, S., Gangavarapu, K., Watts, A., Tan, A. L., Oidtman, R. J., Ladner, J. T., Oliveira, G., Matteson, N. L.,  
588 Kraemer, M. U. G., Vogels, C. B. F., Hentoff, A., Bhatia, D., Stanek, D., Scott, B., Landis, V., Stryker, I., Cone, M. R., Kopp, E.  
589 W., 4th, ... Andersen, K. G. (2019). Travel Surveillance and Genomics Uncover a Hidden Zika Outbreak during the Waning  
590 Epidemic. *Cell*, 178(5), 1057–1071.e11. <https://doi.org/10.1016/j.cell.2019.07.018>
- 591 Grubaugh, N. D., Sharma, S., Krajacich, B. J., Fakoli, L. S., III, Bolay, F. K., Diclario, J. W., II, Johnson, W. E., Ebel, G. D., Foy, B.  
592 D., & Brackney, D. E. (2015). Xenosurveillance: a novel mosquito-based approach for examining the human-pathogen  
593 landscape. *PLoS Neglected Tropical Diseases*, 9(3), e0003628. <https://doi.org/10.1371/journal.pntd.0003628>
- 594 Halstead, S. B. (2015). Reappearance of chikungunya, formerly called dengue, in the Americas. *Emerging Infectious Diseases*,  
595 21(4), 557–561. <https://doi.org/10.3201/eid2104.141723>
- 596 Hamlet, A., Jean, K., Perea, W., Yactayo, S., Biey, J., Van Kerkhove, M., Ferguson, N., & Garske, T. (2018). The seasonal  
597 influence of climate and environment on yellow fever transmission across Africa. *PLoS Neglected Tropical Diseases*, 12(3),  
598 e0006284. <https://doi.org/10.1371/journal.pntd.0006284>
- 599 He, A., Brasil, P., Siqueira, A. M., Calvet, G. A., & Kwatra, S. G. (2017). The Emerging Zika Virus Threat: A Guide for

- 600 Dermatologists. *American Journal of Clinical Dermatology*, 18(2), 231–236. <https://doi.org/10.1007/s40257-016-0243-z>
- 601 Hotez, P. J., & Murray, K. O. (2017). Dengue, West Nile virus, chikungunya, Zika-and now Mayaro? *PLoS Neglected Tropical*
- 602 *Diseases*, 11(8), e0005462. <https://doi.org/10.1371/journal.pntd.0005462>
- 603 Hugo, L. E., Jeffery, J. A. L., Trewin, B. J., Wockner, L. F., Nguyen, T. Y., Nguyen, H. L., Nghia, L. T., Hine, E., Ryan, P. A., &
- 604 Kay, B. H. (2014). Adult survivorship of the dengue mosquito *Aedes aegypti* varies seasonally in central Vietnam. *PLoS*
- 605 *Neglected Tropical Diseases*, 8(2), e2669. <https://doi.org/10.1371/journal.pntd.0002669>
- 606 Imai, N., Dorigatti, I., Cauchemez, S., & Ferguson, N. M. (2015). Estimating dengue transmission intensity from sero-prevalence
- 607 surveys in multiple countries. *PLoS Neglected Tropical Diseases*, 9(4), e0003719.
- 608 <https://doi.org/10.1371/journal.pntd.0003719>
- 609 Kantor, A. M., Lin, J., Wang, A., Thompson, D. C., & Franz, A. W. E. (2019). Infection Pattern of Mayaro Virus in *Aedes aegypti*
- 610 (Diptera: Culicidae) and Transmission Potential of the Virus in Mixed Infections With Chikungunya Virus. *Journal of Medical*
- 611 *Entomology*, 56(3), 832–843. <https://doi.org/10.1093/jme/tjy241>
- 612 Katzelnick, L. C., Montoya, M., Gresh, L., Balmaseda, A., & Harris, E. (2016). Neutralizing antibody titers against dengue virus
- 613 correlate with protection from symptomatic infection in a longitudinal cohort. *Proceedings of the National Academy of*
- 614 *Sciences of the United States of America*, 113(3), 728–733. <https://doi.org/10.1073/pnas.1522136113>
- 615 Komar, O., Robbins, M. B., Contreras, G. G., Benz, B. W., Klenk, K., Blitvich, B. J., Marlenee, N. L., Burkhalter, K. L., Beckett, S.,
- 616 González, G., Peña, C. J., Townsend Peterson, A., & Komar, N. (2005). West Nile Virus Survey of Birds and Mosquitoes in
- 617 the Dominican Republic. In *Vector-Borne and Zoonotic Diseases* (Vol. 5, Issue 2, pp. 120–126).
- 618 <https://doi.org/10.1089/vbz.2005.5.120>
- 619 Kraemer, M. U. G., Reiner, R. C., Jr, Brady, O. J., Messina, J. P., Gilbert, M., Pigott, D. M., Yi, D., Johnson, K., Earl, L., Marczak,
- 620 L. B., Shirude, S., Davis Weaver, N., Bisanzio, D., Perkins, T. A., Lai, S., Lu, X., Jones, P., Coelho, G. E., Carvalho, R. G., ...
- 621 Golding, N. (2019). Past and future spread of the arbovirus vectors *Aedes aegypti* and *Aedes albopictus*. *Nature*
- 622 *Microbiology*, 4(5), 854–863. <https://doi.org/10.1038/s41564-019-0376-y>
- 623 Kraemer, M. U. G., Sinka, M. E., Duda, K. A., Mylne, A. Q. N., Shearer, F. M., Barker, C. M., Moore, C. G., Carvalho, R. G.,
- 624 Coelho, G. E., Van Bortel, W., Hendrickx, G., Schaffner, F., Elyazar, I. R. F., Teng, H.-J., Brady, O. J., Messina, J. P., Pigott,
- 625 D. M., Scott, T. W., Smith, D. L., ... Hay, S. I. (2015). The global distribution of the arbovirus vectors *Aedes aegypti* and *Ae.*
- 626 *albopictus*. *eLife*, 4, e08347. <https://doi.org/10.7554/eLife.08347>
- 627 Kraemer, M. U. G., Sinka, M. E., Duda, K. A., Mylne, A., Shearer, F. M., Brady, O. J., Messina, J. P., Barker, C. M., Moore, C. G.,
- 628 Carvalho, R. G., Coelho, G. E., Van Bortel, W., Hendrickx, G., Schaffner, F., Wint, G. R. W., Elyazar, I. R. F., Teng, H.-J., &
- 629 Hay, S. I. (2015). The global compendium of *Aedes aegypti* and *Ae. albopictus* occurrence. *Scientific Data*, 2, 150035.
- 630 <https://doi.org/10.1038/sdata.2015.35>
- 631 Lamballerie, X. de, de Lamballerie, X., Leroy, E., Charrel, R. N., Ttsetsarkin, K., Higgs, S., & Gould, E. A. (2008). Chikungunya
- 632 virus adapts to tiger mosquito via evolutionary convergence: a sign of things to come? In *Virology Journal* (Vol. 5, Issue 1, p.
- 633 33). <https://doi.org/10.1186/1743-422x-5-33>
- 634 Langsjoen, R. M., Rubinstein, R. J., Kautz, T. F., Auguste, A. J., Erasmus, J. H., Kiatty-Figueroa, L., Gerhardt, R., Lin, D., Hari, K.
- 635 L., Jain, R., Ruiz, N., Muruato, A. E., Silfa, J., Bido, F., Dacso, M., & Weaver, S. C. (2016). Molecular Virologic and Clinical
- 636 Characteristics of a Chikungunya Fever Outbreak in La Romana, Dominican Republic, 2014. *PLoS Neglected Tropical*
- 637 *Diseases*, 10(12), e0005189. <https://doi.org/10.1371/journal.pntd.0005189>
- 638 Lednicky, J., De Rochars, V. M. B., Elbadry, M., Loeb, J., Telisma, T., Chavannes, S., Anilis, G., Cella, E., Ciccozzi, M., Okech,
- 639 B., Salemi, M., & Morris, J. G., Jr. (2016). Mayaro Virus in Child with Acute Febrile Illness, Haiti, 2015. *Emerging Infectious*
- 640 *Diseases*, 22(11), 2000–2002. <https://doi.org/10.3201/eid2211.161015>
- 641 Lessler, J., Ott, C. T., Carcelen, A. C., Konikoff, J. M., Williamson, J., Bi, Q., Kucirka, L. M., Cummings, D. A., Reich, N. G., &
- 642 Chaisson, L. H. (2016). Times to key events in Zika virus infection and implications for blood donation: a systematic review.
- 643 *Bulletin of the World Health Organization*, 94(11), 841–849. <https://doi.org/10.2471/BLT.16.174540>
- 644 Li, M. I., Wong, P. S. J., Ng, L. C., & Tan, C. H. (2012). Oral susceptibility of Singapore *Aedes* (*Stegomyia*) *aegypti* (Linnaeus) to
- 645 Zika virus. *PLoS Neglected Tropical Diseases*, 6(8), e1792. <https://doi.org/10.1371/journal.pntd.0001792>
- 646 Lindsey, N. P., Staples, J. E., & Fischer, M. (2018). Chikungunya Virus Disease among Travelers-United States, 2014–2016. *The*
- 647 *American Journal of Tropical Medicine and Hygiene*, 98(1), 192–197. <https://doi.org/10.4269/ajtmh.17-0668>
- 648 Lipsitch, M., Cohen, T., Cooper, B., Robins, J. M., Ma, S., James, L., Gopalakrishna, G., Chew, S. K., Tan, C. C., Samore, M. H.,
- 649 Fisman, D., & Murray, M. (2003). Transmission dynamics and control of severe acute respiratory syndrome. *Science*,
- 650 300(5627), 1966–1970. <https://doi.org/10.1126/science.1086616>
- 651 Liu-Helmersson, J., Stenlund, H., Wilder-Smith, A., & Rocklöv, J. (2014). Vectorial capacity of *Aedes aegypti*: effects of
- 652 temperature and implications for global dengue epidemic potential. *PloS One*, 9(3), e89783.
- 653 <https://doi.org/10.1371/journal.pone.0089783>
- 654 Liu, Y., Lillepold, K., Semenza, J. C., Tozan, Y., Quam, M. B. M., & Rocklöv, J. (2020). Reviewing estimates of the basic
- 655 reproduction number for dengue, Zika and chikungunya across global climate zones. *Environmental Research*, 182, 109114.
- 656 <https://doi.org/10.1016/j.envres.2020.109114>
- 657 Long, K. C., Ziegler, S. A., Thangamani, S., Hausser, N. L., Kochel, T. J., Higgs, S., & Tesh, R. B. (2011). Experimental

- 658 transmission of Mayaro virus by *Aedes aegypti*. *The American Journal of Tropical Medicine and Hygiene*, 85(4), 750–757.  
659 <https://doi.org/10.4269/ajtmh.2011.11-0359>
- 660 Lorenz, C., Freitas Ribeiro, A., & Chiaravalloti-Neto, F. (2019). Mayaro virus distribution in South America. *Acta Tropica*, 198,  
661 105093. <https://doi.org/10.1016/j.actatropica.2019.105093>
- 662 Lourenço, J., de Lima, M. M., Faria, N. R., Walker, A., Kraemer, M. U. G., Villabona-Arenas, C. J., Lambert, B., de Cerqueira, E.  
663 M., Pybus, O. G., Alcantara, L. C. J., & Recker, M. (2017). Epidemiological and ecological determinants of Zika virus  
664 transmission in an urban setting. In *eLife* (Vol. 6). <https://doi.org/10.7554/elife.29820>
- 665 Lourenço, J. MVSE: Estimator of climate-driven, mosquito-borne viral suitability index. January 14, 2019. Accessed: June 15,  
666 2020.  
667 <https://sourceforge.net/projects/mvse/>
- 668 Low, G. K.-K., Ogston, S. A., Yong, M.-H., Gan, S.-C., & Chee, H.-Y. (2018). Global dengue death before and after the new World  
669 Health Organization 2009 case classification: A systematic review and meta-regression analysis. *Acta Tropica*, 182, 237–245.  
670 <https://doi.org/10.1016/j.actatropica.2018.03.014>
- 671 Majumder, M. S., Cohn, E., Fish, D., & Brownstein, J. S. (2016). Estimating a feasible serial interval range for Zika fever. *Bulletin*  
672 *of the World Health Organization*, 10, 1–6. [https://www.who.int/bulletin/online\\_first/16-171009.pdf](https://www.who.int/bulletin/online_first/16-171009.pdf)
- 673 Martinez, J. D., Garza, J. A. C. la, & Cuellar-Barboza, A. (2019). Going Viral 2019: Zika, Chikungunya, and Dengue. *Dermatologic*  
674 *Clinics*, 37(1), 95–105. <https://doi.org/10.1016/j.det.2018.07.008>
- 675 Mavian, C., Dulcey, M., Munoz, O., Salemi, M., Vittor, A. Y., & Capua, I. (2018). Islands as Hotspots for Emerging Mosquito-Borne  
676 Viruses: A One-Health Perspective. *Viruses*, 11(1). <https://doi.org/10.3390/v11010011>
- 677 Metsky, H. C., Matranga, C. B., Wohl, S., Schaffner, S. F., Freije, C. A., Winnicki, S. M., West, K., Qu, J., Baniecki, M. L.,  
678 Gladden-Young, A., Lin, A. E., Tomkins-Tinch, C. H., Ye, S. H., Park, D. J., Luo, C. Y., Barnes, K. G., Shah, R. R., Chak, B.,  
679 Barbosa-Lima, G., ... Sabeti, P. C. (2017). Zika virus evolution and spread in the Americas. *Nature*, 546(7658), 411–415.  
680 <https://doi.org/10.1038/nature22402>
- 681 Millman, A. J., Esposito, D. H., Biggs, H. M., Decentecio, M., Klevos, A., Hunsperger, E., Munoz-Jordan, J., Kosoy, O. I.,  
682 McPherson, H., Sullivan, C., Voorhees, D., Baron, D., Watkins, J., Gaul, L., Sotir, M. J., Brunette, G., Fischer, M., Sharp, T.  
683 M., & Jentes, E. S. (2016). Chikungunya and Dengue Virus Infections Among United States Community Service Volunteers  
684 Returning from the Dominican Republic, 2014. *The American Journal of Tropical Medicine and Hygiene*, 94(6), 1336–1341.  
685 <https://doi.org/10.4269/ajtmh.15-0815>
- 686 Ministerio de Salud Pública (2014a) Plan de preparación y respuesta frente a brotes de Fiebre Chikungunya. 1st ed. Santo  
687 Domingo, Distrito Nacional: Ministerio de Salud Pública y Asistencia Social.  
688 Ministerio de Salud Pública (2014b). República Dominicana, Boletín Especial No. 13., August 20, 2014.  
689 [https://www.paho.org/dor/  
690 images/stories/archivos/chikungunya/boletin\\_chikv\\_no-13\\_2014\\_8\\_20.pdf?ua=1](https://www.paho.org/dor/images/stories/archivos/chikungunya/boletin_chikv_no-13_2014_8_20.pdf?ua=1)
- 691 Mordecai, E. A., Cohen, J. M., Evans, M. V., Gudapati, P., Johnson, L. R., Lippi, C. A., Miazgowiec, K., Murdock, C. C., Rohr, J.  
692 R., Ryan, S. J., Savage, V., Shocket, M. S., Stewart Ibarra, A., Thomas, M. B., & Weikel, D. P. (2017). Detecting the impact of  
693 temperature on transmission of Zika, dengue, and chikungunya using mechanistic models. *PLoS Neglected Tropical*  
694 *Diseases*, 11(4), e0005568. <https://doi.org/10.1371/journal.pntd.0005568>
- 695 Musso, D., Rodriguez-Morales, A. J., Levi, J. E., Cao-Lormeau, V.-M., & Gubler, D. J. (2018). Unexpected outbreaks of arbovirus  
696 infections: lessons learned from the Pacific and tropical America. *The Lancet Infectious Diseases*, 18(11), e355–e361.  
697 [https://doi.org/10.1016/S1473-3099\(18\)30269-X](https://doi.org/10.1016/S1473-3099(18)30269-X)
- 698 National Statistics Office (2012). IX CENSO NACIONAL DE POBLACIÓN Y VIVIENDA. vol. 1. Santo Domingo, Distrito Nacional.
- 699 Nishiura, H., Kinoshita, R., Mizumoto, K., Yasuda, Y., & Nah, K. (2016). Transmission potential of Zika virus infection in the South  
700 Pacific. *International Journal of Infectious Diseases: IJID: Official Publication of the International Society for Infectious*  
701 *Diseases*, 45, 95–97. <https://doi.org/10.1016/j.ijid.2016.02.017>
- 702 Obolski, U., Perez, P. N., Villabona-Arenas, C. J., Thézé, J., Faria, N. R., & Lourenço, J. (2019). MVSE : An R-package that  
703 estimates a climate-driven mosquito-borne viral suitability index. *Methods in Ecology and Evolution / British Ecological*  
704 *Society*, 10(8), 1357–1370. <https://doi.org/10.1111/2041-210X.13205>
- 705 Pan-American Health Organization (2014). Five-fold increase in dengue cases in the Americas over the past decade. Accessed:  
706 June 15, 2020. [https://www.paho.org/hq/index.php?option=com\\_content&view=article&id=9657:2014-los-casos-dengue-americas-  
707 quintuplicaron-diez-anos-segun-nuevos-datos-opsoms&Itemid=1926&lang=en](https://www.paho.org/hq/index.php?option=com_content&view=article&id=9657:2014-los-casos-dengue-americas-quintuplicaron-diez-anos-segun-nuevos-datos-opsoms&Itemid=1926&lang=en)
- 708 Pan-American Health Organization (PAHO) (2020a). Suspected dengue cases by epidemiological week for countries and  
709 territories of the Americas. Accessed: June 15, 2020. [https://www.paho.org/data/index.php/en/mnu-topics/indicadores-  
710 dengue-en/dengue-  
711 nacional-en/252-dengue-pais-ano-en.html?start=2](https://www.paho.org/data/index.php/en/mnu-topics/indicadores-dengue-en/dengue-nacional-en/252-dengue-pais-ano-en.html?start=2)
- 712 Pan-American Health Organization (PAHO) (2020b). Dengue serotypes by year for countries and territories of the Americas.  
713 Accessed: June 15, 2020. [https://www.paho.org/data/index.php/es/temas/indicadores-dengue/dengue-nacional/549-dengue-  
714 serotypes-es.html](https://www.paho.org/data/index.php/es/temas/indicadores-dengue/dengue-nacional/549-dengue-serotypes-es.html)
- 715 Pan-American Health Organization (PAHO) (2020c). Cases of Zika virus disease. Accessed: June 15, 2020.

- 716 [https://www.paho.org/data/](https://www.paho.org/data/index.php/en/mnu-topics/zika/524-zika-weekly-en.html)  
717 [index.php/en/mnu-topics/zika/524-zika-weekly-en.html](https://www.paho.org/data/index.php/en/mnu-topics/zika/524-zika-weekly-en.html)
- 718 Pan-American Health Organization (PAHO) (2020d). PLISA Health Information Platform for the Americas. Accessed: June 15,  
719 2020. <https://www.paho.org/data/index.php/en/>
- 720 Pereira, T. N., Carvalho, F. D., De Mendonça, S. F., Rocha, M. N., & Moreira, L. A. (2020). Vector competence of *Aedes aegypti*,  
721 *Aedes albopictus*, and *Culex quinquefasciatus* mosquitoes for Mayaro virus. *PLoS Neglected Tropical Diseases*, *14*(4),  
722 e0007518. <https://doi.org/10.1371/journal.pntd.0007518>
- 723 Perez, F., Llau, A., Gutierrez, G., Bezerra, H., Coelho, G., Ault, S., Barbiratto, S. B., Resende, M. C., Cerezo, L., Kleber, G. L.,  
724 Pacheco, O., Perez, O. L., Picos, V., Rojas, D. P., Siqueira, J. B., Suarez, M. F., Harris, E., Castellanos, L. G., Espinal, C., &  
725 Martin, J. L. S. (2019). The decline of dengue in the Americas in 2017: discussion of multiple hypotheses. In *Tropical*  
726 *Medicine & International Health* (Vol. 24, Issue 4, pp. 442–453). <https://doi.org/10.1111/tmi.13200>
- 727 Perez-Guzman, P. N., Carlos Junior Alcantara, L., Obolski, U., de Lima, M. M., Ashley, E. A., Smithuis, F., Horby, P., Maude, R.  
728 J., Lin, Z., Kyaw, A. M. M., & Lourenço, J. (2018). Measuring Mosquito-borne Viral Suitability in Myanmar and Implications for  
729 Local Zika Virus Transmission. *PLoS Currents*, *10*.  
730 <https://doi.org/10.1371/currents.outbreaks.7a6c64436a3085ebba37e5329ba169e6>
- 731 Pineda, C., Muñoz-Louis, R., Caballero-Urbe, C. V., & Viasus, D. (2016). Chikungunya in the region of the Americas. A challenge  
732 for rheumatologists and health care systems. *Clinical Rheumatology*, *35*(10), 2381–2385. [https://doi.org/10.1007/s10067-016-](https://doi.org/10.1007/s10067-016-3390-y)  
733 [3390-y](https://doi.org/10.1007/s10067-016-3390-y)
- 734 Requena-Méndez, A., Garcia, C., Aldasoro, E., Vicente, J. A., Martínez, M. J., Pérez-Molina, J. A., Calvo-Cano, A., Franco, L.,  
735 Parrón, I., Molina, A., Ruiz, M., Álvarez, J., Sánchez-Seco, M. P., & Gascón, J. (2014). Cases of chikungunya virus infection  
736 in travellers returning to Spain from Haiti or Dominican Republic, April-June 2014. *Euro Surveillance: Bulletin European Sur*  
737 *Les Maladies Transmissibles = European Communicable Disease Bulletin*, *19*(28), 20853. [https://doi.org/10.2807/1560-](https://doi.org/10.2807/1560-7917.es2014.19.28.20853)  
738 [7917.es2014.19.28.20853](https://doi.org/10.2807/1560-7917.es2014.19.28.20853)
- 739 Ribeiro, G. S., Hamer, G. L., Diallo, M., Kitron, U., Ko, A. I., & Weaver, S. C. (2020). Influence of herd immunity in the cyclical  
740 nature of arboviruses. *Current Opinion in Virology*, *40*, 1–10. <https://doi.org/10.1016/j.coviro.2020.02.004>
- 741 Ribeiro, G. S., Kikuti, M., Tauro, L. B., Nascimento, L. C. J., Cardoso, C. W., Campos, G. S., Ko, A. I., Weaver, S. C., Reis, M. G.,  
742 Kitron, U., & Salvador Arboviral Research Group. (2018). Does immunity after Zika virus infection cross-protect against  
743 dengue? *The Lancet. Global Health*, *6*(2), e140–e141. [https://doi.org/10.1016/S2214-109X\(17\)30496-5](https://doi.org/10.1016/S2214-109X(17)30496-5)
- 744 Riou, J., Poletto, C., & Boëlle, P.-Y. (2017). A comparative analysis of Chikungunya and Zika transmission. *Epidemics*, *19*, 43–52.  
745 <https://doi.org/10.1016/j.epidem.2017.01.001>
- 746 Rodriguez-Barraquer, I., Costa, F., Nascimento, E. J. M., Nery, N., Júnior, Castanha, P. M. S., Sacramento, G. A., Cruz, J.,  
747 Carvalho, M., De Olivera, D., Hagan, J. E., Adhikarla, H., Wunder, E. A., Jr, Coêlho, D. F., Azar, S. R., Rossi, S. L., Vasilakis,  
748 N., Weaver, S. C., Ribeiro, G. S., Balmaseda, A., ... Ko, A. I. (2019). Impact of preexisting dengue immunity on Zika virus  
749 emergence in a dengue endemic region. *Science*, *363*(6427), 607–610. <https://doi.org/10.1126/science.aav6618>
- 750 Rodriguez-Barraquer, I., Salje, H., & Cummings, D. A. (2019). Opportunities for improved surveillance and control of dengue from  
751 age-specific case data. *eLife*, *8*. <https://doi.org/10.7554/eLife.45474>
- 752 Rosario, V., Munoz-Louis, R., Valdez, T., Adames, S., Medrano, J., Paulino, I., Paula, J., & Alba-Férez, R. (2015). Chikungunya  
753 infection in the general population and in patients with rheumatoid arthritis on biological therapy. *Clinical Rheumatology*,  
754 *34*(7), 1285–1287. <https://doi.org/10.1007/s10067-015-2979-x>
- 755 Staples, J. E., Breiman, R. F., & Powers, A. M. (2009). Chikungunya fever: an epidemiological review of a re-emerging infectious  
756 disease. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*, *49*(6), 942–948.  
757 <https://doi.org/10.1086/605496>
- 758 Trpis, M., & Hausermann, W. (1986). Dispersal and other population parameters of *Aedes aegypti* in an African village and their  
759 possible significance in epidemiology of vector-borne diseases. *The American Journal of Tropical Medicine and Hygiene*,  
760 *35*(6), 1263–1279. <https://doi.org/10.4269/ajtmh.1986.35.1263>
- 761 Trpis, M., Häusermann, W., & Craig, G. B. (1995). Estimates of Population Size, Dispersal, and Longevity of Domestic *Aedes*  
762 *aegypti aegypti* (Diptera: Culicidae) by Mark–Release–Recapture in the Village of Shauri Moyo in Eastern Kenya. In *Journal*  
763 *of Medical Entomology* (Vol. 32, Issue 1, pp. 27–33). <https://doi.org/10.1093/jmedent/32.1.27>
- 764 Tsang, T. K., Ghebremariam, S. L., Gresh, L., Gordon, A., Halloran, M. E., Katzelnick, L. C., Rojas, D. P., Kuan, G., Balmaseda,  
765 A., Sugimoto, J., Harris, E., Longini, I. M., Jr, & Yang, Y. (2019). Effects of infection history on dengue virus infection and  
766 pathogenicity. *Nature Communications*, *10*(1), 1246. <https://doi.org/10.1038/s41467-019-09193-y>
- 767 United Nations Human Development (UNDP) (2019). Dominican Republic: Human Development Indicators. Accessed: June 15,  
768 2020. <http://hdr.undp.org/en/countries/profiles/DOM>
- 769 Vazeille, M., Moutailler, S., Coudrier, D., Rousseaux, C., Khun, H., Huerre, M., Thiria, J., Dehecq, J.-S., Fontenille, D.,  
770 Schuffenecker, I., Despres, P., & Failloux, A.-B. (2007). Two Chikungunya isolates from the outbreak of La Reunion (Indian  
771 Ocean) exhibit different patterns of infection in the mosquito, *Aedes albopictus*. *PloS One*, *2*(11), e1168.  
772 <https://doi.org/10.1371/journal.pone.0001168>
- 773 Vogels, C. B. F., Rückert, C., Cavany, S. M., Perkins, T. A., Ebel, G. D., & Grubaugh, N. D. (2019). Arbovirus coinfection and co-

- 774 transmission: A neglected public health concern? *PLoS Biology*, 17(1), e3000130.  
775 <https://doi.org/10.1371/journal.pbio.3000130>
- 776 Weaver, S. C., & Reisen, W. K. (2010). Present and future arboviral threats. *Antiviral Research*, 85(2), 328–345.  
777 <https://doi.org/10.1016/j.antiviral.2009.10.008>
- 778 Weppelmann, T. A., Burne, A., von Fricken, M. E., Elbadry, M. A., Beau De Rochars, M., Boncy, J., & Okech, B. A. (2017). A Tale  
779 of Two Flaviviruses: A Seroepidemiological Study of Dengue Virus and West Nile Virus Transmission in the Ouest and Sud-  
780 Est Departments of Haiti. *The American Journal of Tropical Medicine and Hygiene*, 96(1), 135–140.  
781 <https://doi.org/10.4269/ajtmh.16-0422>
- 782 Wiggins, K., Eastmond, B., & Alto, B. W. (2018). Transmission potential of Mayaro virus in Florida *Aedes aegypti* and *Aedes*  
783 *albopictus* mosquitoes. *Medical and Veterinary Entomology*, 32(4), 436–442. <https://doi.org/10.1111/mve.12322>
- 784 Winokur, O. C., Main, B. J., Nicholson, J., & Barker, C. M. (2020). Impact of temperature on the extrinsic incubation period of Zika  
785 virus in *Aedes aegypti*. *PLoS Neglected Tropical Diseases*, 14(3), e0008047. <https://doi.org/10.1371/journal.pntd.0008047>
- 786 Wong, P.-S. J., Li, M.-Z. I., Chong, C.-S., Ng, L.-C., & Tan, C.-H. (2013). *Aedes* (*Stegomyia*) *albopictus* (Skuse): a potential vector  
787 of Zika virus in Singapore. *PLoS Neglected Tropical Diseases*, 7(8), e2348. <https://doi.org/10.1371/journal.pntd.0002348>
- 788 World Health Organization (WHO) Zika virus: Fact Sheet. July 20, 2018. Accessed: June 15, 2020. [https://www.who.int/news-  
789 room/fact-sheets/detail/zika-virus](https://www.who.int/news-room/fact-sheets/detail/zika-virus)
- 790 Yasuno, M., & Tonn, R. J. (1970). A study of biting habits of *Aedes aegypti* in Bangkok, Thailand. *Bulletin of the World Health*  
791 *Organization*, 43(2), 319–325. <https://www.ncbi.nlm.nih.gov/pubmed/5312528>
- 792 Zingman, M. A., Paulino, A. T., & Payano, M. P. (2017). Clinical manifestations of chikungunya among university professors and  
793 staff in Santo Domingo, the Dominican Republic. *Revista Panamericana de Salud Publica = Pan American Journal of Public*  
794 *Health*, 41, e64. <https://www.ncbi.nlm.nih.gov/pubmed/28902277>