

1 Behavioral changes before lockdown, and decreased
2 retail and recreation mobility during lockdown,
3 contributed most to the successful control of the
4 COVID-19 epidemic in 35 Western countries

5 Koen Deforche,^{1*} Jurgen Vercauteren,² Viktor Müller³, Anne-Mieke Vandamme^{2,4*}

¹Emweb bv, Herent, Belgium

²KU Leuven, Department of Microbiology, Immunology and Transplantation,
Rega Institute for Medical Research, Clinical and Epidemiological Virology,
Institute for the Future, Leuven, Belgium

³Institute of Biology, Eötvös Loránd University, Budapest, Hungary,

⁴Center for Global Health and Tropical Medicine, Unidade de Microbiologia,
Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa,
Lisbon, Portugal

*To whom correspondence should be addressed; E-mail: koen.deforche@gmail.com,
annemie.vandamme@kuleuven.be.

6 **The COVID-19 pandemic has prompted a lockdown in many countries to con-**
7 **trol the exponential spread of the SARS-CoV-2 virus (1, 2). This resulted in**
8 **curbing the epidemic by reducing the time-varying basic reproduction num-**
9 **ber (R_t) to below one (3, 4). Governments are looking for evidence to balance**
10 **the demand of their citizens to ease some of the restriction, against the fear**
11 **of a second peak in infections. More details on the specific circumstances that**
12 **promote exponential spread (i.e. $R_t > 1$) and the measures that contributed**
13 **most to a reduction in R_t are needed. Here we show that in 33 of 35 West-**

1 **ern countries (32 European, plus Israel, USA and Canada), R_t fell to around**
2 **or below one during lockdown (March – May 2020). One third of the effect**
3 **happened already on average 6 days before the lockdown, with lockdown itself**
4 **causing another major drop in transmission. Country-wide compulsory usage**
5 **of masks was implemented only in Slovakia 10 days into lockdown, and on its**
6 **own reduced transmission by half. During lockdown, decreased mobility in re-**
7 **tail and recreation was an independent predictor of lower R_t during lockdown,**
8 **while changes in other types of mobility were not. These results are consistent**
9 **with anecdotal evidence that large recreational gatherings are super-spreading**
10 **events (5, 6), and may even suggest that infections during day-to-day contact**
11 **at work are not sufficient to spark exponential growth. Our data suggest mea-**
12 **sures that will contribute to avoiding a second peak include a tight control on**
13 **circumstances that facilitate massive spread such as large gatherings especially**
14 **indoors, physical distancing, and mask use.**

15 In February-March 2020, it became clear that the COVID-19 pandemic had been introduced
16 in Europe and Northern America (7) and governments and societies have responded in various
17 ways to slow down the spread of the virus. In the containment phase, testing, contact-tracing
18 and individual quarantine of (suspected) infected individuals are crucial to prevent a widespread
19 epidemic. If this fails, mitigation measures consist mainly of mobility restrictions and encour-
20 aged or mandated physical distancing to everyone regardless of symptoms or testing results
21 (jointly called lockdown).

22 Failing containment, most European countries managed to curb the epidemic by May through
23 lockdown measures (timeline and details of measures in the different countries (1)), however
24 without fully understanding which of the different measures had the most impact. Because of
25 economic and social implications of a lockdown, governments are cautiously relaxing measures

1 while trying to avoid a second peak by closely following the effect on the epidemics through
2 testing and monitoring hospitalizations and casualties (8). Coming out of lockdown entails a
3 return to containment measures, with testing, contact tracing and quarantine again being crucial.

4 Epidemiological models have been widely used to make predictions about the evolution of
5 the COVID-19 epidemics, to predict the outcome of testing strategies and intervention scenar-
6 ios, and to estimate epidemiological parameters (9–13). Epidemic spread is determined by both
7 biological properties of the virus as well as the behavior of the host population, and this is
8 reflected in the epidemiological parameters. One important parameter, the time-varying basic
9 reproductive number R_t , determines whether an epidemic is exponentially growing ($R_t > 1$)
10 or declining ($R_t < 1$). The goal of both containment and mitigation measures is to bring R_t
11 below 1 by changing the behavior of the infected, respectively, susceptible population through
12 reducing their contact networks.

13 In Western countries the population was developing an increased awareness as soon as me-
14 dia reported on the epidemic unfolding in Lombardy (Italy), and this led to a gradual response
15 ranging from voluntary physical distancing to government imposed mandatory self-isolation
16 upon showing symptoms, banning of public events, and encouraging physical distancing, with
17 ultimately a drastic lockdown when many businesses and schools were closed and mobility
18 was severely restricted (14). The diversity of measures and their timing gives us the opportu-
19 nity to investigate which differences were correlated with a better outcome. However, reliable
20 data quantifying the impact on human behavior of various country-dependent specific lockdown
21 measures is currently lacking.

22 Attempts to create such data sets face challenges such as variation in compliance to imposed
23 measures, and the unknown level of measures taken by individuals or organizations in absence
24 or anticipation of government regulations. Rather than an account of the ordered measures in
25 each country, we used mobility changes extracted from smartphone location data as an objective

1 indication and quantification of the lockdown (Suppl 1).

2 From incidence data of deaths and diagnosed cases in 35 countries (ECDC data down-
3 loaded on 3 June 2020, see Suppl 2), we estimated the change in transmission over time, while
4 differentiating between the effect of measures that preceded the lockdown, and the effect of the
5 lockdown itself. Since incidence data of deaths is considered more reliable than of diagnosed
6 cases (which is also highly dependent on testing policy and test availability), we estimated suit-
7 able values for the dispersion parameters by estimating variance from the fits and then use a
8 slightly more cautious value (Suppl 3). For case data we then simply decreased the value to
9 give a higher weight to death data, favoring better fits of death data over case data. Only data
10 up to 60 days into the lockdown (up to about mid-May in European countries) were used to
11 avoid the confounding effect of the gradual lifting of restrictions, which started in many Eu-
12 ropean countries in May. Given that time between infection and reported death was estimated
13 as 24 (95% IQR 18 - 32) days, potential lifting of measures in the beginning of May for some
14 countries could not have confounded the estimates.

15 In order to obtain comparable estimates of R_t , a simple SEIR compartment model was used,
16 with most epidemiological parameters kept constant. Parameters that model transmission rates
17 were allowed to change from an initial estimated value $R_{t,0}$ during a transition period also
18 estimated from the data, to $R_{t,1}$ until the day that mobility changes started, and then to $R_{t,2}$
19 during the lockdown identified on the basis of mobility data, using a piecewise linear model
20 (Figure 1a and methods). The estimated models fitted well the reported incidence data for most
21 countries (see Suppl 4). Across all countries, the median of posterior estimates for $R_{t,0}$ was
22 3.6 (95% IQR 2.5 – 4.9) (see Figure 2a). Before changes in mobility were observed (d_1), the
23 reproduction number was reduced to 2.3 (95% IQR 1.7 – 3.0). During lockdown, transmission
24 was further reduced to 0.77 (95% IQR 0.58 – 1.04). Only for Belarus and Moldova, the estimate
25 of $R_{t,2}$ was higher than 1 ($> 95\%$ credible), while all other countries have the $R_{t,2}$ value below




Figure 1: Model for changes of the time-varying reproduction number R_t as a piece-wise linear function. Dates d_1 and d_2 were estimated from mobility data. Date d_0 and values for $R_{t,0} - R_{t,2}$ were estimated from incidence data on diagnosed cases and deaths. a. Model used for all countries; b. Model additionally used for Slovakia, which allowed an extra change during lockdown with dates d_0 , d_3 and d_4 , and values for $R_{t,0} - R_{t,3}$ estimated from incidence data on diagnosed cases and deaths.

1 or around 1. Estimates for $R_{t,0}$ are expected to vary depending on setting, methodology and
2 assumptions on parameters (especially duration of infectious period and generation time) and
3 assumptions on how the number may vary over time (15). We find that our estimated values
4 for $R_{t,0}$ tend to be higher, and estimated values for $R_{t,1}$ lower, compared to estimates in other
5 studies for R_0 of around 2.7. Our estimates for $R_{t,0}$ are similar to R_0 estimates obtained using
6 models that also consider interventions that preceded a full lockdown (14).

7 We estimated that the date d_0 of the first decline of transmission preceded observed mobility
8 changes (d_1) with on average 6 days (95% IQR 0 – 22) (Figure 2b). During this period less inva-
9 sive measures such as the mandatory self-isolation upon showing symptoms, banning of public
10 events, and encouraging physical distancing may have contributed to a decline in transmission
11 (there was no substantial decline in mobility during this period). The earliest of these dates
12 (for Italy, Ireland, Germany, France, and Belgium) were all estimated around 20 February (95%
13 IQR 12 – 25 February), which might have been a result of awareness raised by the discovery of
14 the first European cluster around that time in Lombardy, Italy. Although the semantic meaning
15 given to d_0 (first date of decline of R_0 , before mobility changes), assumed that it would be esti-
16 mated before d_1 , this order was not enforced. For Slovakia, the estimated model placed this date
17 around 1 April (95% IQR March 22 – April 7), later than the dates d_1 and d_2 that mark the mo-
18 bility changes period (10 – 17 March). At the same time, a suspiciously low value of 1.7 (95%
19 IQR 0.8 – 2.6) for $R_{t,0}$ was estimated. Both observations indicated that the assumption that
20 mobility changes leading to the lockdown were the final measure to control transmission, was
21 not applicable to data of Slovakia. We therefore re-estimated a model which allowed a further
22 reduction in transmission to a value $R_{t,3}$ (assuming a prior distribution $R_{t,3} - R_{t,2} = \mathcal{N}(0, 1)$
23 expressing no change with respect to $R_{t,2}$), with a linear transition between co-estimated dates
24 d_3 and d_4 (see Figure 1b). As prior distributions for d_0 , $R_{t,0} - R_{t,2}$, and infection-to-death dura-
25 tion μ_d , overall estimates obtained across all countries were used. This resulted in an additional

Variable	Univariate estimate	Multivariate estimate
Retail and recreation	0.04 +/- 0.01 ($p < 0.004$)	0.07 +/- 0.02 ($p < 0.004$)
Grocery and pharmacy	0.02 +/- 0.02	-0.04 +/- 0.03
Parks	0.004 +/- 0.006	-0.018 +/- 0.009 ($p < 0.05$)
Transit stations	0.03 +/- 0.02	
Workplaces	0.06 +/- 0.02 ($p < 0.008$)	0.04 +/- 0.03
Residential	-0.11 +/- 0.04 ($p < 0.005$)	
$R_{t,1}$	0.05 +/- 0.06	0.02 +/- 0.05

Table 1: Univariate and multivariate associations of mobility changes during lockdown (per 10% mobility change) with the basic reproduction number $R_{t,2}$ during lockdown. Mobility data related to transit stations and residential places were left out from the multivariate analysis since these variables were highly correlated with mobility data related to workplaces.

1 estimated transition period for reduction in R_t between d_3 of 30 March (95% IQR 18 March
2 – 14 April), and d_4 of 4 April (95% IQR 21 March – 18 April, Figure 3). This followed im-
3 mediately after the introduction of mandatory face masks use in Slovakia on 25 March (16).
4 This would imply that using face masks brought R_t from 0.87 (95% IQR 0.56 – 1.26) to 0.50
5 (95% IQR 0.30 – 0.70), or thus a 43% (95% IQR 0 – 64%) reduction in transmission. In further
6 statistical analyses, the parameters estimated for this latter model were used for Slovakia.

7 To investigate how the reproduction number during lockdown $R_{t,2}$ related to mobility changes,
8 Google Mobility report data was used. For all six mobility categories, mobility during lock-
9 down was significantly different compared to baseline (Wilcoxon paired test $p < 0.01$, data
10 not shown). Lower $R_{t,2}$ values during lockdown were significantly associated with a larger
11 mobility reduction related to retail, recreation, and workplaces, and a lower mobility reduction
12 related to residential (Figure 4), Table 1). The association of mobility reduction related to retail
13 and recreation remained significant in a multivariate model which explained 40% of variance
14 (adjusted R^2) of the $R_{t,2}$ value during lockdown ($p < 0.002$). This suggests that, in 35 West-
15 ern countries, reductions of mobility related to retail and recreation during lockdown caused a
16 mean reduction of R_t of 0.50 (95% CI 0.18 – 0.81), or thus an average reduction of 22% (95%




Figure 2: a. Posterior estimates of the initial basic reproduction numbers ($R_{t,0}$), the reproduction number at start of lockdown ($R_{t,1}$) and during lockdown ($R_{t,2}$). b. Estimated median values (95% IQR) for d_0 (date of first reduction in transmission, presumably due to physical distancing, estimated from incidence data of deaths and diagnosed cases); and lockdown transition start and end dates d_1 and d_2 , estimated from mobility data (see also Suppl 1).




Figure 3: Estimated time-varying basic reproduction number R_t for Slovakia using a model that allowed an additional reduction of transmission rate at a co-estimated date during lockdown. The orange and red lines mark d_1 and d_2 as estimated from google mobility data. For reference, the date at which mandatory mask wearing was introduced (25 March) is indicated in blue.

1 CI 8 – 35) in transmission. Increased mobility related to parks may be an indicator of social
2 restrictions in visiting friends or family, and explained up to 4% (95% CI 0 – 8) of additional
3 reduction in transmission. The remaining reduction of R_t during lockdown may still be related
4 to mobility changes of other types of mobility currently not reported by Google (for example
5 mobility related to schools) or would require further refinement of categories to become observ-
6 able. Alternatively, other types of changes that coincided in time with these mobility changes
7 could be responsible.

8 By explaining only variation of $R_{t,2}$, instead of the reduction from $R_{t,1}$ to $R_{t,2}$, the above
9 model side-stepped to a large extent the colinearity that exists between changes for different
10 mobility categories, which showed the same trend in all countries. To verify that the reduction
11 in R_t as a result of the lockdown (comparing $R_{t,1}$ before lockdown to $R_{t,2}$ during lockdown)
12 was also associated with mobility changes, a multivariate model that directly predicted the
13 percentage change of R_t due to lockdown using average mobility changes (comparing mobility
14 data from before d_1 to after d_2), was also estimated (data not shown). This model had a high




Figure 4: Average percentage change in mobility, compared to baseline, for six location Google mobility categories. Colour reflects basic reproduction number R_t during lockdown. Retail and Recreation: restaurants, cafes, shopping centers, theme parks, museums, libraries, and movie theaters; Grocery and Pharmacy: grocery markets, food warehouses, farmers markets, specialty food shops, drug stores, and pharmacies; Parks: local parks, national parks, public beaches, marinas, dog parks, plazas, and public gardens; Transit Stations: public transport hubs such as subway, bus, and train stations; Workplaces: places of work; Residential: places of residence.




Figure 5: Summary of estimated contributions to the reduction of transmission in 35 Western countries. Initial basic reproduction numbers ($R_{t,0}$), the reproduction number at start of lockdown ($R_{t,1}$) and during lockdown ($R_{t,2}$), and percentage reductions are shown as median values and 95% IQR. Estimates for individual countries (Italy, Spain and Slovakia) are shown as median posterior value and 95% cri.

1 explanatory power ($R^2 = 0.95$) but as expected suffered from high colinearity. Nevertheless,
 2 in agreement with the above finding, this model also attributed an estimated 31% (95% CI 11 –
 3 50) of the drop in R_t during lockdown to a reduction in mobility related to retail and recreation
 4 ($p < 0.003$).

5 Our findings are summarized in Figure 5 and trends were found to be robust to different
 6 assumptions of latent period duration (in the range of 2 to 4 days), and different assumptions of
 7 generation time (in the range of 4.5 to 5.9 days), see Suppl 3. Although the estimated value for
 8 $R_{t,2}$ for the epidemic in Sweden was found to be relatively high (1.02, 95% cri 0.99 – 1.06), it
 9 was equally well explained based on mobility changes, suggesting that although not enforced,
 10 in practice the country behaved the same as other Western countries in a lockdown.

11 The fact that mobility changes linked to retail and recreation were significantly associated
 12 with R_t both in univariate and multivariate analyses, suggests that activities related to visiting
 13 malls, bars, restaurants, or museums are linked to increased transmission and releasing those
 14 mobility restrictions should be done with care since they may carry a high risk for reigniting
 15 the epidemic. This is in line with mounting evidence of transmission being promoted by (loud)
 16 speaking or singing (bars, choir (17)), and longer time spent in densely populated indoor loca-




Figure 6: Structure of a standard SEIR compartment model with four compartments: susceptible (S), exposed (E), infectious (I), and removed (recovered or deceased, R). Susceptible individuals become latently infected by infectious individuals, with transmission rate β . Latently infected become infectious at rate σ . Infectious people are removed at rate γ .

1 tions with low air circulation (bars, restaurants, malls, events with mass gatherings ($I8$)). In the
2 presence of general 'physical distancing' guidelines, which were active in all countries during
3 lockdown, mobility changes related to workplaces (which correlated with transit stations and
4 residential places), did not have a clear impact on R_t , suggesting that easing of restrictions re-
5 lated to work may be more easily manageable. The lack of association of these other mobility
6 parameters may suggest that rather than being associated with an individual, superspreading
7 is mainly associated with the circumstances of the interaction. This is adding weight to the
8 arguments that superspreading events, such as large indoor gatherings, need to be avoided.

9 This study has several limitations. The use of a compartment model in conjunction with a
10 piecewise linear model for R_t is necessarily an approximation, and more complicated temporal
11 changes in transmission may have occurred in reality, but still it served the purpose of examining
12 the impact of different measures over time. Epidemiological parameter estimates may have been
13 impacted by shifts in age distribution of cases over time, given that COVID-19 has a distinct age
14 risk profile, and also by variation in data accuracy between countries. By limiting the study to
15 Western countries which coincided more or less in time, differences related to culture, climate,
16 and google mobility data interpretation, may have been less of a confounding factor.

17 **1 Methods**

1 The following equations describe the dynamics of individuals in each of the four compartments
2 of a standard SEIR model (see Figure 6):

$$\begin{aligned}\frac{dS}{dt} &= -\frac{\beta IS}{N} \\ \frac{dE}{dt} &= \frac{\beta IS}{N} - \sigma E \\ \frac{dI}{dt} &= \sigma E - \gamma I \\ \frac{dR}{dt} &= \gamma I\end{aligned}$$

3 The differential equations of the SEIR compartment models were numerically integrated
4 using `deSolve` (19). The duration of the latent period $T_{\text{lat}} = 1/\sigma$ was fixed to 3 days, and the
5 infectious period $T_{\text{inf}} = 1/\gamma$ to 5.2 days based on literature (20, 21) (robustness analyses for
6 other parameter values were conducted, see Suppl 3).

7 The dates d_1 and d_2 which mark the transition period for mobility changes were estimated
8 from mobility data reports (22), by fitting a step function with a linear transition period through
9 the sum of mobility changes related to transit stations and workplaces (see Suppl 1). This was
10 motivated by the adoption of teleworking as a measure in most countries, even in those (like
11 Sweden) which had the lightest lockdown regimen. The value for β was modeled as a piece-
12 wise linear function of time (Figure 1).

13 We wanted a model that does not require assumptions on timing or number of introductions
14 per country. Instead, the models were seeded with an initial single exposed individual and it
15 was assumed that the estimated date d_0 , marking the first change in value of β , was linked to a
16 co-estimated threshold for cumulative deaths. This method had the benefit of avoiding any bias
17 on estimated R_0 values because of assumptions on introduction time and numbers, while being
18 more efficient to sample than time of first infection.

19 Suppl 3 Table 1 lists all parameters used in the model and during the estimation from data,
1 with their values (either a constant, or a prior distribution for parameters that were estimated).

2 To estimate the model parameters, incidence data of diagnosed cases and deaths was used
3 (ECDC, <https://opendata.ecdc.europa.eu/covid19/casedistribution/csv>,
4 accessed on 6 June), within a Bayesian framework. The daily incidence of diagnosed cases and
5 deaths was evaluated against the prediction based on the simulation of the SEIR model, using a
6 negative binomial distribution. The posterior distributions of model parameters were estimated
7 in R using MCMC with Metropolis coupling (20 chains at different temperatures) (23–26).

8 Univariate and multivariate linear models were used for quantifying the effect of the average
9 values of Google Mobility data (22) over the time period prior to lockdown to $R_{t,1}$ and during
10 lockdown to $R_{t,2}$. Colinearity was assessed by calculating covariance-inflation factors (27).
11 Statistical analysis was done in RStudio (28).

12 All data files, R scripts and analysis steps are described in Suppl 2.

13 **2 Acknowledgments**

14 The resources and services used in this work were provided by the VSC (Flemish Supercom-
15 puter Center), funded by the Research Foundation - Flanders (FWO) and the Flemish Govern-
16 ment. VM was supported by the ELTE Institutional Excellence Program (NKFIH-1157-8/2019-
17 DT) and a COVID-19 grant (2020-2.1.1-ED-00001) of the National Research, Development and
18 Innovation Office.

19 **3 Author contributions**

20 K.D. contributed to the Bayesian analyses and conception of the study; J.V. contributed to the
21 statistical analyses; V.M. contributed to the Bayesian analyses and epidemiological interpretation;

22 A.M.V. contributed to the virological and epidemiological interpretation and revisions of the
1 manuscript.

2 **4 Competing interests**

3 The authors declare no competing interests.

4 **References**

- 5 1. ACAPS COVID-19 government measures update VII. [https://www.acaps.org/
6 country/world/special-reports#container-1476](https://www.acaps.org/country/world/special-reports#container-1476). Accessed: 2020-06-
7 04.
- 8 2. de Bruin], Y. B. *et al.* Initial impacts of global risk mitigation measures taken
9 during the combatting of the COVID-19 pandemic. *Safety Science* **128**, 104773
10 (2020). URL [http://www.sciencedirect.com/science/article/pii/
11 S0925753520301703](http://www.sciencedirect.com/science/article/pii/S0925753520301703).
- 12 3. Lonergan, M. & Chalmers, J. D. Estimates of the ongoing need for social distancing
13 and control measures post-“lockdown” from trajectories of COVID-19 cases and mor-
14 tality. *European Respiratory Journal* (2020). URL [https://erj.ersjournals.
15 com/content/early/2020/05/26/13993003.01483-2020](https://erj.ersjournals.com/content/early/2020/05/26/13993003.01483-2020). [https:
16 //erj.ersjournals.com/content/early/2020/05/26/13993003.
17 01483-2020.full.pdf](https://erj.ersjournals.com/content/early/2020/05/26/13993003.01483-2020.full.pdf).
- 18 4. Salje, H. *et al.* Estimating the burden of SARS-CoV-2 in france. *Sci-*
19 *ence* (2020). URL [https://science.sciencemag.org/content/early/
20 2020/05/12/science.abc3517](https://science.sciencemag.org/content/early/2020/05/12/science.abc3517). [https://science.sciencemag.org/
21 content/early/2020/05/12/science.abc3517.full.pdf](https://science.sciencemag.org/content/early/2020/05/12/science.abc3517.full.pdf).

- 22 5. Al-Tawfiq, J. A. & Rodriguez-Morales, A. J. Super-spreading events and contribution to
1 transmission of MERS, SARS, and SARS-CoV-2 (COVID-19). *Journal of Hospital In-*
2 *fection* **105**, 111–112 (2020). URL [https://doi.org/10.1016/j.jhin.2020.](https://doi.org/10.1016/j.jhin.2020.04.002)
3 04.002.
- 4 6. James, A. *et al.* High COVID-19 attack rate among attendees at events at a church —
5 arkansas, march 2020. *MMWR Morb Mortal Wkly Rep* **69**, 632–635 (2020).
- 6 7. Worobey, M. *et al.* The emergence of sars-cov-2 in europe and the us.
7 *bioRxiv* (2020). URL [https://www.biorxiv.org/content/early/2020/05/](https://www.biorxiv.org/content/early/2020/05/23/2020.05.21.109322)
8 23/2020.05.21.109322. [https://www.biorxiv.org/content/early/](https://www.biorxiv.org/content/early/2020/05/23/2020.05.21.109322.full.pdf)
9 2020/05/23/2020.05.21.109322.full.pdf.
- 10 8. Cohen, J. & Kupferschmidt, K. Countries test tactics in ‘war’ against covid-19. *Science*
11 **367**, 1287–1288 (2020). URL [https://science.sciencemag.org/content/](https://science.sciencemag.org/content/367/6484/1287)
12 367/6484/1287. [https://science.sciencemag.org/content/367/](https://science.sciencemag.org/content/367/6484/1287.full.pdf)
13 6484/1287.full.pdf.
- 14 9. Chowell, G., Sattenspiel, L., Bansal, S. & Viboud, C. Mathematical models to char-
15 acterize early epidemic growth: A review. *Physics of Life Reviews* **18**, 66 – 97
16 (2016). URL [http://www.sciencedirect.com/science/article/pii/](http://www.sciencedirect.com/science/article/pii/S1571064516300641)
17 S1571064516300641.
- 18 10. Park, M., Cook, A. R., Lim, J. T., Sun, Y. & Dickens, B. L. A systematic review of covid-
19 19 epidemiology based on current evidence. *Journal of Clinical Medicine* **9** (2020). URL
20 <https://www.mdpi.com/2077-0383/9/4/967>.

- 21 11. Wangping, J. *et al.* Extended sir prediction of the epidemics trend of covid-19 in italy
1 and compared with hunan, china. *Frontiers in Medicine* **7**, 169 (2020). URL <https://www.frontiersin.org/article/10.3389/fmed.2020.00169>.
2
- 3 12. Liu, Y., Gayle, A. A., Wilder-Smith, A. & Rocklöv, J. The reproductive number of COVID-
4 19 is higher compared to SARS coronavirus. *Journal of Travel Medicine* **27** (2020). URL
5 <https://doi.org/10.1093/jtm/taaa021>. Taaa021, [https://academic.](https://academic.oup.com/jtm/article-pdf/27/2/taaa021/32902430/taaa021.pdf)
6 [oup.com/jtm/article-pdf/27/2/taaa021/32902430/taaa021.pdf](https://academic.oup.com/jtm/article-pdf/27/2/taaa021/32902430/taaa021.pdf).
- 7 13. Anastassopoulou, C., Russo, L., Tsakris, A. & Siettos, C. Data-based analysis, modelling
8 and forecasting of the covid-19 outbreak. *PLOS ONE* **15**, 1–21 (2020). URL <https://doi.org/10.1371/journal.pone.0230405>.
9
- 10 14. Estimating the number of infections and the impact of nonpharmaceutical inter-
11 ventions on COVID-19 in 11 european countries. [https://www.imperial.](https://www.imperial.ac.uk/mrc-global-infectious-disease-analysis/covid-19/report-13-europe-npi-impact/)
12 [ac.uk/mrc-global-infectious-disease-analysis/covid-19/](https://www.imperial.ac.uk/mrc-global-infectious-disease-analysis/covid-19/report-13-europe-npi-impact/)
13 [report-13-europe-npi-impact/](https://www.imperial.ac.uk/mrc-global-infectious-disease-analysis/covid-19/report-13-europe-npi-impact/). Accessed: 2020-06-04.
- 14 15. Liu, Y., Gayle, A. A., Wilder-Smith, A. & Rocklöv, J. The reproductive number of COVID-
15 19 is higher compared to SARS coronavirus. *Journal of Travel Medicine* **27** (2020). URL
16 <https://doi.org/10.1093/jtm/taaa021>. Taaa021, [https://academic.](https://academic.oup.com/jtm/article-pdf/27/2/taaa021/32902430/taaa021.pdf)
17 [oup.com/jtm/article-pdf/27/2/taaa021/32902430/taaa021.pdf](https://academic.oup.com/jtm/article-pdf/27/2/taaa021/32902430/taaa021.pdf).
- 18 16. Technical report: Using face masks in the community.
19 [https://www.ecdc.europa.eu/en/publications-data/](https://www.ecdc.europa.eu/en/publications-data/using-face-masks-community-reducing-covid-19-transmission)
20 [using-face-masks-community-reducing-covid-19-transmission](https://www.ecdc.europa.eu/en/publications-data/using-face-masks-community-reducing-covid-19-transmission)
21 (2020). Accessed: 2020-06-04.

- 22 17. ... High SARS-CoV-2 attack rate following exposure at a choir practice — skagit county,
1 washington, march 2020. *MMWR Morb Mortal Wkly Rep* **69**, 606–610 (2020).
- 2 18. Lu, J. *et al.* Covid-19 outbreak associated with air conditioning in restaurant, guangzhou,
3 china, 2020. *Emerging Infectious Disease journal* **26**, 1628 (2020). URL [https://](https://wwwnc.cdc.gov/eid/article/26/7/20-0764_article)
4 wwwnc.cdc.gov/eid/article/26/7/20-0764_article.
- 5 19. Soetaert, K., Petzoldt, T. & Setzer, R. W. Solving differential equations in R: Package deS-
6 olve. *Journal of Statistical Software* **33**, 1–25 (2010). URL [http://www.jstatsoft.](http://www.jstatsoft.org/v33/i09)
7 [org/v33/i09](http://www.jstatsoft.org/v33/i09).
- 8 20. Ganyani, T. *et al.* Estimating the generation interval for coronavirus disease
9 (covid-19) based on symptom onset data, march 2020. *Eurosurveillance* **25**
10 (2020). URL [https://www.eurosurveillance.org/content/10.2807/](https://www.eurosurveillance.org/content/10.2807/1560-7917.ES.2020.25.17.2000257)
11 [1560-7917.ES.2020.25.17.2000257](https://www.eurosurveillance.org/content/10.2807/1560-7917.ES.2020.25.17.2000257).
- 12 21. He, X. *et al.* Temporal dynamics in viral shedding and transmissibility of covid-
13 19. *Nature Medicine* **26**, 672–675 (2020). URL [https://doi.org/10.1038/](https://doi.org/10.1038/s41591-020-0869-5)
14 [s41591-020-0869-5](https://doi.org/10.1038/s41591-020-0869-5).
- 15 22. Google COVID-19 community mobility reports”. [https://www.google.com/](https://www.google.com/covid19/mobility/)
16 [covid19/mobility/](https://www.google.com/covid19/mobility/). Accessed: 2020-05-04.
- 17 23. Verity, B. & Winskill, P. *drjacob: Flexible Markov Chain Monte Carlo via Reparameter-*
18 *ization* (2020). R package version 1.0.1.
- 19 24. Li, Q. *et al.* Early transmission dynamics in wuhan, china, of novel coron-
20 avirus–infected pneumonia. *New England Journal of Medicine* **382**, 1199–1207 (2020).
21 URL <https://doi.org/10.1056/NEJMoa2001316>. [https://doi.org/](https://doi.org/10.1056/NEJMoa2001316)
22 [10.1056/NEJMoa2001316](https://doi.org/10.1056/NEJMoa2001316).

- 23 25. Flegal, J. M., Hughes, J., Vats, D. & Dai, N. *mcmcse: Monte Carlo Standard Errors for*
1 *MCMC*. Riverside, CA, Denver, CO, Coventry, UK, and Minneapolis, MN (2020). R
2 package version 1.4-1.
- 3 26. Makowski, D., Ben-Shachar, M. S. & Lüdecke, D. bayestestr: Describing effects and their
4 uncertainty, existence and significance within the bayesian framework. *Journal of Open*
5 *Source Software* **4**, 1541 (2019). URL [https://joss.theoj.org/papers/10.](https://joss.theoj.org/papers/10.21105/joss.01541)
6 [21105/joss.01541](https://joss.theoj.org/papers/10.21105/joss.01541).
- 7 27. Fox, J. & Weisberg, S. *An R Companion to Applied Regression* (Sage, Thousand Oaks
8 CA, 2019), third edn. URL [https://socialsciences.mcmaster.ca/jfox/](https://socialsciences.mcmaster.ca/jfox/Books/Companion/)
9 [Books/Companion/](https://socialsciences.mcmaster.ca/jfox/Books/Companion/).
- 10 28. RStudio Team. *RStudio: Integrated Development Environment for R*. RStudio, Inc.,
11 Boston, MA (2019). URL <http://www.rstudio.com/>.